

THE MINERAL SPRINGS
OF JAPAN

R. ISHIZU

J2592

\$17.50

JAPAN

ERRATA.

- Pt. II, pp. 1-55, Electric conductivity, for $\chi_{18} \times 10^{-4}$, read $\chi_{18} \times 10^4$.
" p. 18, Table 16, No. 2, Classification, for sulphur, read acid vitriol.
" " " No. 3, Temp. of spring in C., for 100,0°, read 103,0°.
" p. 29, Table 27, Emanation per litre of water in 10^{-10} curies, for 21,05, read 12,05.
" p. 49, The hot spring "Goza-no-yu," Temperature, for 53,9° C., read 63,9° C.
" p. 61, The cold spring "Anamori-Kōsen," Calcium ion, for 0,7183 grm., read 0,1873 grm.
Pt. III, p. 9, line 6 from top, for 1,135, read 4,135.
" p. 30, line 9 from top, for 6,500, read 65,000.

THE MINERAL SPRINGS OF JAPAN

with

Tables of Analyses, Radio-activity,
Notes on Prominent Spas and List of Seaside Resorts
and Summer Retreats

specially edited

for

THE PANAMA-PACIFIC INTERNATIONAL EXPOSITION

By

Dr. R. ISHIZU,

Expert of the Imperial Hygienic Laboratory



DH

TOKYO IMPERIAL HYGIENIC LABORATORY

1915

All rights reserved

Published by
SANKYO KABUSHIKI KAISHA,
Maromachi, Nihonbashi-ku, Tokyo.

Printed by The Tokyo Tsukiji Type Foundry, Tsukiji, Tokyo.

PREFACE

The scenic beauty of Japan is known abroad only by sacred Mt. Fuji, the grand Nikkō temples, beautiful Hakone, and other such world-famous sights. These renowned beauty spots are truly typical of the characteristic charms of the Empire and it is little wonder foreign tourists never fail to visit any of these spots, but the fact that these places are so widely known is mostly due to their easy means of access and the better accommodation procurable. If you are willing to leave the beaten track and are ready for out-of-the-ordinary tours, you will find not a few fair sights and scenes, some even excelling the aforesaid places in their charms and beauty. This is especially the case with the simple rustic settings and surroundings of mineral springs. We find in remote districts not easy of access and out of reach of the bustle and noise of city life, by far the most excellent hot springs for therapeutic purposes, both on account of their good quality and healthy geographical situation. We cannot say that the medicinal value of mineral springs is in proportion to their popular fame, for in many instances the reverse seems to hold good. Accordingly it is not too much to claim, that almost all really efficacious mineral springs are hidden in distant places and unknown to ordinary tourists. But if physicians at large kept abreast of the progress of physical therapy and made full use of mineral springs with a thorough knowledge of their individual merits, so that they might be able to point to particularly suitable ones and induce patients and health-seekers to try them, the springs hitherto neglected would certainly be improved and provided with adequate facility of approach, accommodation, etc., and we believe the time will soon arrive for winning fame for Japan as an ideal health resort.

Japan is among the few countries endowed with an abundance of mineral springs, and no statistics are available to show exactly the total number of them now in existence. The investigation of Japanese mineral springs was made for the first time by the Government and reported in the International Exposition held in Frankfort, Germany, in 1881, some diagrams based on the data collected from the prefectural governments being exhibited. These materials were compiled in book form and published in Japanese under the title "The Mineral Springs of Japan." Since then our Imperial Hygienic Laboratory has continued, at the request of the prefectural authorities or private individuals, to experiment on the origins of springs or analyze sample mineral waters submitted to their investigation. The mineral springs, thus experimented upon, number more than one thousand. With the recent introduction of radio-therapy, public attention has been gradually drawn to the radio-activity of mineral springs, and since 1913 the Imperial Hygienic Laboratories, both of Tokyo and Osaka, have been making researches about the radio-activity of famous mineral springs, which number up to now more than one hundred and fifty, with more than one thousand origins of springs.

The present volume was compiled as an exhibit in the Panama-Pacific International Exposition in 1915, in order to make known to the world the general conditions of Japanese mineral springs, and consists chiefly of the analyses of famous mineral springs in Japan and reports of experiments on their radio-activity, both based on the materials above referred to, and some other items, with an appendix of descriptions of principal hot spring resorts, their locations, communications, geography, places of interest in the neighbourhood, and some adjacent sea bathing and climatic health resorts. This book, therefore, may lay claim to be the most authentic and up to date on the subject as yet attempted or compiled in English. But there are so many mineral springs in Japan that we cannot investigate all, some have been experimented upon imperfectly and others left out altogether. Under such circumstances the scope of the present work, it is needless to add, still leaves much to be desired.

The analyses shown in this volume were mostly made by our Imperial Hygienic Laboratory, and some by the prefectural hygienic laboratories, military garrison hospitals, etc., the names of assayers and dates of experiments being shown for each in the latter case.

For the articles on "the Scenery of Japan" and "Volcanoes and Mineral Springs in Japan," in the "Introductory Remarks," I am indebted to Dr. N. Yamazaki, Professor of the Tokyo Imperial University, and T. Katō, Professor of the Tokyo Higher Normal School; that on "Geology" were based on the publications of the Imperial Geological Survey of the Department of Agriculture and Commerce.

The article on "the Climate of Japan" was specially compiled by the Central Meteorological Observatory of Japan, and that on "the Radio-active Minerals in Japan," by Prof., K. Jimbō of the Tokyo Imperial University, from whom I have had most kind and valuable help also in dealing with subjects bearing on mineralogy and geology.

Not a few medical references I owe to R. Fortescue Fox's "Medical Hydrology," Dawson Turner's "Radium, its Physics and Therapeutics" (2nd edition), and "Deutsches Bäderbuch." The article on "the Geyser in Atami" in Chapter XII was taken from the Physical Review, Vol. XXII., No. 5, 1906, with consent of the authors. For the information on "the Geyser in Onikōbe" given in the Appendix, I am indebted to Prof. K. Honda, who has supplied me with suggestions and data on the subject.

For the descriptions of hot spring resorts at the end of this work, and the "Map of Imperial Government Railways" inserted herein, I am indebted to "An Official Guide to Eastern Asia," published by the Imperial Government Railways of Japan in 1914.

The compiler has pleasure in expressing his thanks to the above-named Government Institutions and individuals for their valuable help and assistance. I wish also to acknowledge my indebtedness to the reports of Mr. D. Ishitani, Professor of Physics in the Imperial Peers' College, who started the investigation of the radio-activity of mineral springs in Japan for the first time. In particular, I desire to express my sincere thanks to my assistants, Mr. Y. Kinugasa and Mr. T. Saitō, for the great care and trouble they have taken in the correction of proof sheets, and for undertaking the laborious task of checking a number of quotations from various authorities which are included in the volume.

Compiled by order of N. Nakagawa, the Director of the Sanitary Bureau, the Department of Home Affairs, to constitute part of the exhibits of the Imperial Hygienic Laboratory at the Panama Exposition.

RISAKU ISHIZU.

TOKYO IMPERIAL HYGIENIC LABORATORY,
April, 1915.

EXPLANATORY REMARKS.

1. The locations of mineral springs are here grouped by the prefectures and arranged in geographical order from the north-east to the south-west, thus beginning with Hokkaidō and running through to Kagoshima, after which come those of Taiwan (Formosa) and ending with Chōsen (Korea). As no data are available for Karafuto (Saghalien), no mention is made of it.

2. It is not always easy to exactly determine the geological structure of the spot where a spring gushes out; we mentioned only the general geological feature in most cases.

3. As it is sometimes very difficult owing to the conditions of spring origins to measure exactly the quantities of their flow of water, such figures herein shown had better to be considered as rough estimates. Nevertheless, we have done our best to point out possible faults in these rough figures, on account of their importance from a hygienic estimation.

4. The composition of each mineral spring is shown herein, according to the results of analyses made by our Imperial Hygienic Laboratory or prefectural governments concerned. The temperatures of original springs are of the latest date available.

5. The classification of mineral springs is generally made according to their chemical composition, based on the ion table. They are divided roughly into six groups, *i. e.*, simple, thermal, acid, carbonated, salt, sulphur, and iron springs. Their minor sub-divisions are shown in the methods of classification of mineral springs at the commencement of this volume, to which the reader is referred. Temperatures, as a rule, are Centigrade, but Fahrenheit is also used to indicate atmospheric temperatures in the appendix at the end of this volume. The distinction between cold and hot springs is drawn at the average bodily temperature 37°C .; those above 37°C . at their origin being classed into hot springs and those below into cold springs. Those springs of some definite radium emanation content are considered mineral springs, even if containing less than one gramme of solid constituents to the kilogramme of water.

6. As analyses are generally made with mineral waters taken from original springs and sent to the laboratory, such ingredients as carbon dioxide, hydrogen sulphide, ferrous salts, bicarbonates, etc., are found of less quantity than in the actual springs, while in some cases no trace even is left of these. Accordingly we cannot presume the classification of mineral springs treated under such analytic conditions to be absolutely or scientifically accurate.

7. "Total residue" given in the tables of analyses denotes the quantity of solid ingredients in one kilogramme of water, which was determined by evaporating to dryness in a weighed platinum dish, first over a lamp, then on the water-bath and finally exposing the residue, in the air-bath, to a temperature of about 180° , until no further diminution of weight took place.

8. The radio-activity of mineral waters is shown herein according to the results recently obtained by chemists of our laboratory, and mostly taken on the actual spot. Samples of water to be examined for radio-activity were taken, not from bath-tubs, but at the source of springs. In some instances specially mentioned, experiments were made, at the request of outsiders, on samples of mineral water sent up to our laboratory. As for the methods adopted, we rely on the accuracy of the experiments in which the Schmidt's electrometer was used; but only nominal value is expected for the results of simpler experiments made by the fontactoscope as used in order to save time by assayers who were despatched to the springs. In these instances comparatively big experimental errors are inevitable; and the reader is requested to grant due allowance for the above fact when comparing one experiment with another. As for the figures quoted, those in the Mache's unit are calculated down to two places of decimals but the second figures after the decimal point are not reliable.

9. The measurement of electric conductivity was made invariably on the spot at the original springs.

10. In the descriptions of hot spring resorts, the numbers of visitors in the year 1909 imply people who stopped overnight, excluding passing visitors. The populations and numbers of houses quoted are also for the same year.

11. Although every effort was made to investigate and point out the traditions regarding the medicinal powers of the mineral springs, no mention is made of any which are said to be simply

efficacious for all diseases and not specially adapted for any particular ailment. The dates of discovery, though often doubtful, are quoted as traditional.

12. Not a few springs are known by more than one name, but to avoid confusion the names most popular or generally understood are adopted.

13. The following abbreviations are used in the present volume to indicate the results of experiments on radio-activity:—

Schmidt's electrom. Schm.	}	Schmidt's electrometer
Kohl.-Löw. fontact. K. & L.	}	Kohlraush-Löwenthal's fontactoscope
Eng.-Siev. fontact. E. & S.	}	Engler-Sieveking's fontactoscope

14. In this work, all Japanese names are translated according to the rules made by the Society for Romanizing the writing of Japanese. Thus, all vowels are pronounced as in Italian, and all consonants as in usual English spelling. Special care must be taken to distinguish long vowels marked with —, as in Mr. Ōta and Mr. Ota, which are two entirely different family names.

15. Japanese words often appearing in geographical and other denominations as prefixes and suffixes, might be mentioned and found of assistance to the reader:—

- <i>ken</i>	Prefecture	- <i>toke</i>	mountain pass
- <i>gun</i>	Prefectural district	- <i>michi</i>	road or path
- <i>machu</i> }		- <i>kaido</i>	highway
- <i>chō</i> }	town	- <i>saka</i> }	
- <i>mura</i>	village or hamlet	- <i>saka</i> }	ascent or slope
- <i>yama</i> }		- <i>umi</i> }	
- <i>san</i> }		- <i>maru</i> }	sea
- <i>rin</i> }	mountain or hill	- <i>wan</i>	bay
- <i>take</i> }		- <i>fuchi</i>	pool; deep pool
- <i>dake</i> }		- <i>shima</i> }	
- <i>oka</i>	low hill	- <i>jima</i> }	islet or island
- <i>kawa</i> }		- <i>ko</i>	lake
- <i>sawa</i> }	river or stream	- <i>sata</i>	lagoon
- <i>tani</i> }		- <i>ike</i>	pond or small lake
- <i>dani</i> }	valley or gorge	- <i>numa</i>	marshy pond
- <i>taki</i> }		- <i>sawa</i> }	
- <i>daki</i> }	waterfall or cascade	- <i>swa</i> }	swamp or marsh
- <i>onsen</i> }		0-	large or larger
- <i>yu</i> }	hot spring	Kō-	small or smaller
- <i>sen</i> }		Moto-	original or former
- <i>ke-yu</i>	hot spring of...	Furu-	old
- <i>jinja</i> }		Shin-	new
- <i>jinja</i> }	Shintō shrine	Kami-	upper
- <i>miya</i> }		Naka-	middle
- <i>tera</i> }		Shimo-	lower
- <i>dera</i> }	Buddhist temple	Mae-	front
- <i>ji</i> }		Koen	park
- <i>do</i>	small temple	- <i>re</i> }	
- <i>hara</i> }		- <i>keian</i> }	
- <i>hara</i> }	plain or prairie	- <i>ryokuan</i> }	common suffixes to names
- <i>iza</i>	rock	- <i>ken</i> }	of hotels, inns, etc.
- <i>ishi</i>	stone	- <i>en</i> }	
- <i>mine</i>	peak		

CONTENTS

PART I

Introduction.	1
Chapter I. Scenery of Japan.	3
Chapter II. Volcanoes and Mineral Springs in Japan.	6
Chapter III. Geology of the Japanese Empire.	9
Chapter IV. Climate of Japan with Special Reference to Health Resorts in Japan.	12
Chapter V. The Origin of Climate Therapeutics, Balneo-therapy, and Sea Bathing in Japan.	23
Chapter VI. General Remarks about Hot Springs in Japan.	25
* Chapter VII. Japanese Mineral Springs and their Therapeutical Factors.	30
Chapter VIII. Short Remarks about Emanation Therapy.	31
Chapter IX. Determination of Radium Emanation and Electric Conductivity.	37
Chapter X. The Sinter Deposits of Mineral Springs.	39
Chapter XI. A Few Words about the Radio-active Minerals in Japan.	42
Chapter XII. On the Geyser in Atami.	47
Chapter XIII. A. Classification of Mineral Springs.	52
* B. Kinds of Medicinal Springs and their Indications.	53
C. Kinds of Baths and their Use.	59
D. Notable Mineral Springs arranged according to Chemical Composition.	64
Appendix A. On the Geyser at Onikōbe.	73
Appendix B. List of Notable Mineral Springs in Other Countries.	78

PART II

(I) The Tables of Analyses, Radio-activity, Electric Conductivity, Flow of Water, etc., of Mineral Springs, arranged geographically.	1
(II) The Tables of Mineral Springs arranged according to:	163
I. Radio-activity.	163
<i>Radio-activity of Notable Mineral Springs in Europe and in the United States of America.</i>	190
II. Temperature.	195
III. Flow of Water.	196
IV. The Quantity of Solid Ingredients.	197
(III) A. The Tables of Hot Spring Spas arranged according to:	198
V. Altitude.	198
VI. The Number of Visitors.	199
B. The Tables of Summer Resorts and Watering Places arranged geographically.	202

PART III

Notes on Prominent Spas, Seaside Resorts, and Summer Retreats.	1
General Index.	

List of Maps

1.	Imperial Government Railways	 facing	Pt. I, p. 3
2.	Distribution of Mineral Springs and Volcanoes in Japan	 „ „	6
3.	Ena Mineral Springs, Gifu-ken	 „ „	30
4.	Masutomi Mineral Springs, Yamanashi-ken	 „ „	30
5.	Distribution of Mineral Springs in San-in District	 „ „	30
6.	Distribution of Mineral Springs in Central Japan	 „	Pt. III, p. 12
7.	Toba and Environs	 „ „	50

List of Illustrations

Large	I 28		36	Tenryūkyō
markings	I 28		37	Nakabusa Mountain Spa
Hokutōlīte, radiograph of,	I 44		38	Ena Mineral Springs
„ photograph of,	I 44		39	„
Plate			40	Futami Hot Springs
1	Sinter deposited in wooden conduit of Hot Springs (<i>frontispiece</i>)		41	Oiwayama Summer Retreat
2	Sinter-coated Leaves and Roots, Sinter-Balls (<i>frontispiece</i>)		42	Ōmaki Hot Springs
3	Hokutōlīte (<i>frontispiece</i>)		43	Yamanaka Hot Springs
4	Sinter Cones („)		44	Komono (Yunoyama) Hot Springs
5*	Sources of Hot Springs		45	„
6*	Shigaku Spa, Yumura Spa (Shimane-ken), way to Kusatsu Spa, and source of Tsubame Hot Springs		46	Toba Seaside Resort
7*	Hot Waterfalls for Massage		47	Shiraya Mineral Springs
8*	Utilisations of hot springs		48	Sedo-no-kanayama (Yuzaki) Hot Springs
9*	Time Bath at Kusatsu		49	Yunomine (Shimura) Spa
10*	Measurement of Radium emanation of Spring waters		50	Sotonoyu Hot Springs
11*	Misasa Hot Springs		51	Hamadera Seas de Resort
11 _a *	The most noted Summer Resorts in Japan		52	The Principal Mineral Water bottled for Sale and Export (<i>facing p. 52</i>)
12	Noboribetsu Spa		53	The Clifford-Wilkinson Tansan Mineral Water Works
13	„		54	„
14	„		55	The Imperial Mineral Water Co.
15	Ōyu Hot Springs (Akita-ken)		56	„
16	Innai-Yunosawa Hot Springs		57	The Nunobiki Tansan Factory
17	Atsumi Hot Springs		58	Arima Spa
18	Onogawa Hot Springs		59	Kinosaki Hot Springs
19	Nomegawa Hot Springs		60	Misasa Spa
20	Goshiki Hot Springs		61	Ushio Hot Springs
21	Senami Spa		62	Port Tomo and the Island of Sensui-tō
22	Murasugi Spa		63	Dōgo Spa
23	Tochiomata Hot Springs		64	„
24	Matsunoyama Hot Springs		65	Funagoya Mineral Springs
25	Nasu-Yumoto Hot Springs		66	„
26	Yumoto Hot Springs, Nikkō		67	Shibaseki Hot Springs
27	Isobe Spa		68	Umi-Jigoku at Kannawa Spa
28	Nashiki-Kōsen		69	Hot Sand Baths at Beppu Spa
29	Hiraiso Seaside Resort		70	Karatsu Seaside Resort
30	Ōarai Seaside Resort		71	„
31	Hayama Seaside Resort		72	Inns at Karatsu
32	„		73	Takeo Hot Springs
33	Oiso Seaside Resort		74	„
34	Atami Spa		75	Jujō Hot Springs (Chōsen)
35	Masutomi Mineral Springs		76	On-yō Hot Springs („)
			77	Kwanshirei Hot Springs (Taiwan)

Fig. 1



Am. 3



1911



Fig. 4. Saturated vapor pressure of the Alkyl-Pol-Siloxane-Fluoropolymer (B) (1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 83

© 1997 by The Society for the Study of Inequalities and Applications. Printed in the United States of America.

1. 2. 3. 4. 5. 6. 7. 8. 9. 10. 11. 12. 13. 14. 15. 16. 17. 18. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100. 101. 102. 103. 104. 105. 106. 107. 108. 109. 110. 111. 112. 113. 114. 115. 116. 117. 118. 119. 120. 121. 122. 123. 124. 125. 126. 127. 128. 129. 130. 131. 132. 133. 134. 135. 136. 137. 138. 139. 140. 141. 142. 143. 144. 145. 146. 147. 148. 149. 150. 151. 152. 153. 154. 155. 156. 157. 158. 159. 160. 161. 162. 163. 164. 165. 166. 167. 168. 169. 170. 171. 172. 173. 174. 175. 176. 177. 178. 179. 180. 181. 182. 183. 184. 185. 186. 187. 188. 189. 190. 191. 192. 193. 194. 195. 196. 197. 198. 199. 200. 201. 202. 203. 204. 205. 206. 207. 208. 209. 210. 211. 212. 213. 214. 215. 216. 217. 218. 219. 220. 221. 222. 223. 224. 225. 226. 227. 228. 229. 230. 231. 232. 233. 234. 235. 236. 237. 238. 239. 240. 241. 242. 243. 244. 245. 246. 247. 248. 249. 250. 251. 252. 253. 254. 255. 256. 257. 258. 259. 260. 261. 262. 263. 264. 265. 266. 267. 268. 269. 270. 271. 272. 273. 274. 275. 276. 277. 278. 279. 280. 281. 282. 283. 284. 285. 286. 287. 288. 289. 290. 291. 292. 293. 294. 295. 296. 297. 298. 299. 300. 301. 302. 303. 304. 305. 306. 307. 308. 309. 310. 311. 312. 313. 314. 315. 316. 317. 318. 319. 320. 321. 322. 323. 324. 325. 326. 327. 328. 329. 330. 331. 332. 333. 334. 335. 336. 337. 338. 339. 340. 341. 342. 343. 344. 345. 346. 347. 348. 349. 350. 351. 352. 353. 354. 355. 356. 357. 358. 359. 360. 361. 362. 363. 364. 365. 366. 367. 368. 369. 370. 371. 372. 373. 374. 375. 376. 377. 378. 379. 380. 381. 382. 383. 384. 385. 386. 387. 388. 389. 390. 391. 392. 393. 394. 395. 396. 397. 398. 399. 400. 401. 402. 403. 404. 405. 406. 407. 408. 409. 410. 411. 412. 413. 414. 415. 416. 417. 418. 419. 420. 421. 422. 423. 424. 425. 426. 427. 428. 429. 430. 431. 432. 433. 434. 435. 436. 437. 438. 439. 440. 441. 442. 443. 444. 445. 446. 447. 448. 449. 450. 451. 452. 453. 454. 455. 456. 457. 458. 459. 460. 461. 462. 463. 464. 465. 466. 467. 468. 469. 470. 471. 472. 473. 474. 475. 476. 477. 478. 479. 480. 481. 482. 483. 484. 485. 486. 487. 488. 489. 490. 491. 492. 493. 494. 495. 496. 497. 498. 499. 500. 501. 502. 503. 504. 505. 506. 507. 508. 509. 510. 511. 512. 513. 514. 515. 516. 517. 518. 519. 520. 521. 522. 523. 524. 525. 526. 527. 528. 529. 530. 531. 532. 533. 534. 535. 536. 537. 538. 539. 540. 541. 542. 543. 544. 545. 546. 547. 548. 549. 550. 551. 552. 553. 554. 555. 556. 557. 558. 559. 560. 561. 562. 563. 564. 565. 566. 567. 568. 569. 570. 571. 572. 573. 574. 575. 576. 577. 578. 579. 580. 581. 582. 583. 584. 585. 586. 587. 588. 589. 590. 591. 592. 593. 594. 595. 596. 597. 598. 599. 600. 601. 602. 603. 604. 605. 606. 607. 608. 609. 610. 611. 612. 613. 614. 615. 616. 617. 618. 619. 620. 621. 622. 623. 624. 625. 626. 627. 628. 629. 630. 631. 632. 633. 634. 635. 636. 637. 638. 639. 640. 641. 642. 643. 644. 645. 646. 647. 648. 649. 650. 651. 652. 653. 654. 655. 656. 657. 658. 659. 660. 661. 662. 663. 664. 665. 666. 667. 668. 669. 670. 671. 672. 673. 674. 675. 676. 677. 678. 679. 680. 681. 682. 683. 684. 685. 686. 687. 688. 689. 690. 691. 692. 693. 694. 695. 696. 697. 698. 699. 700. 701. 702. 703. 704. 705. 706. 707. 708. 709. 710. 711. 712. 713. 714. 715. 716. 717. 718. 719. 720. 721. 722. 723. 724. 725. 726. 727. 728. 729. 730. 731. 732. 733. 734. 735. 736. 737. 738. 739. 740. 741. 742. 743. 744. 745. 746. 747. 748. 749. 750. 751. 752. 753. 754. 755. 756. 757. 758. 759. 760. 761. 762. 763. 764. 765. 766. 767. 768. 769. 770. 771. 772. 773. 774. 775. 776. 777. 778. 779. 780. 781. 782. 783. 784. 785. 786. 787. 788. 789. 790. 791. 792. 793. 794. 795. 796. 797. 798. 799. 800. 801. 802. 803. 804. 805. 806. 807. 808. 809. 810. 811. 812. 813. 814. 815. 816. 817. 818. 819. 820. 821. 822. 823. 824. 825. 826. 827. 828. 829. 830. 831. 832. 833. 834. 835. 836. 837. 838. 839. 840. 84

[illegible]

Fig. 3. $\sigma_{\text{eff}} = \sigma_{\text{eff}}(\sigma_{\text{eff}})$ for various values of the parameter σ_{eff} of the system (3.1). For $\sigma_{\text{eff}} = 0$, $\sigma_{\text{eff}} = 0.1$, $\sigma_{\text{eff}} = 0.2$, $\sigma_{\text{eff}} = 0.3$, $\sigma_{\text{eff}} = 0.4$, $\sigma_{\text{eff}} = 0.5$, $\sigma_{\text{eff}} = 0.6$, $\sigma_{\text{eff}} = 0.7$, $\sigma_{\text{eff}} = 0.8$, $\sigma_{\text{eff}} = 0.9$, $\sigma_{\text{eff}} = 1.0$.




Fig. 1



Fig. 2




Fig. 3

Fig. 1. Sinter-coated leaves found in a valley near Khimkent, the Mountain, Kirgiz Province. (Pl. I, p. 42; *Pl. Ill.*, p. 42).

Fig. 2. Plant roots coated with sinter, found at the Hot Waters of the Same at the Hot Springs, Kirgiz Province. (Pl. I, p. 42; *Pl. Ill.*, p. 42).

Fig. 3. Sinter balls, formed in a pool near the Hot Waters of Osh. (Pl. I, p. 42; *Pl. Ill.*, p. 42).

Fig. 1




Fig. 2




Fig. 3



- Fig. 1. Upper Drusy Surface of a Holmshite Crust. (Pl. I, p. 43; *Pl. III*, p. 70.)
 Fig. 2. The Same Specimen in Natural Orientation (Horizontal View), showing Deposit of Holmshite Above the Water Line. (Pl. I, p. 43; *Pl. III*, p. 70.)
 Fig. 3. Holmshite Deposit on the Entire Breadth of the Creek Bed. (Pl. I, p. 43; *Pl. III*, p. 70.)

Fig. 1.



Fig. 2.

Fig. 3.




Fig. 4.



Fig. 1. Sinter Cone formed by the Geyser of Olana, near Union. (Pl. I, p. 40; 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538, 539, 540, 541, 542, 543, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569, 570, 571, 572, 573, 574, 575, 576, 577, 578, 579, 580, 581, 582, 583, 584, 585, 586, 587, 588, 589, 590, 591, 592, 593, 594, 595, 596, 597, 598, 599, 600, 601, 602, 603, 604, 605, 606, 607, 608, 609, 610, 611, 612, 613, 614, 615, 616, 617, 618, 619, 620, 621, 622, 623, 624, 625, 626, 627, 628, 629, 630, 631, 632, 633, 634, 635, 636, 637, 638, 639, 640, 641, 642, 643, 644, 645, 646, 647, 648, 649, 650, 651, 652, 653, 654, 655, 656, 657, 658, 659, 660, 661, 662, 663, 664, 665, 666, 667, 668, 669, 670, 671, 672, 673, 674, 675, 676, 677, 678, 679, 680, 681, 682, 683, 684, 685, 686, 687, 688, 689, 690, 691, 692, 693, 694, 695, 696, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 707, 708, 709, 710, 711, 712, 713, 714, 715, 716, 717, 718, 719, 720, 721, 722, 723, 724, 725, 726, 727, 728, 729, 730, 731, 732, 733, 734, 735, 736, 737, 738, 739, 740, 741, 742, 743, 744, 745, 746, 747, 748, 749, 750, 751, 752, 753, 754, 755, 756, 757, 758, 759, 760, 761, 762, 763, 764, 765, 766, 767, 768, 769, 770, 771, 772, 773, 774, 775, 776, 777, 778, 779, 780, 781, 782, 783, 784, 785, 786, 787, 788, 789, 790, 791, 792, 793, 794, 795, 796, 797, 798, 799, 800, 801, 802, 803, 804, 805, 806, 807, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 824, 825, 826, 827, 828, 829, 830, 831, 832, 833, 834, 835, 836, 837, 838, 839, 840, 841, 842, 843, 844, 845, 846, 847, 848, 849, 850, 851, 852, 853, 854, 855, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 869, 870, 871, 872, 873, 874, 875, 876, 877, 878, 879, 880, 881, 882, 883, 884, 885, 886, 887, 888, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 911, 912, 913, 914, 915, 916, 917, 918, 919, 920, 921, 922, 923, 924, 925, 926, 927, 928, 929, 930, 931, 932, 933, 934, 935, 936, 937, 938, 939, 940, 941, 942, 943, 944, 945, 946, 947, 948, 949, 950, 951, 952, 953, 954, 955, 956, 957, 958, 959, 960, 961, 962, 963, 964, 965, 966, 967, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 978, 979, 980, 981, 982, 983, 984, 985, 986, 987, 988, 989, 990, 991, 992, 993, 994, 995, 996, 997, 998, 999, 1000)

PART I

Introductory Remarks.

INTRODUCTION.

The Empire of Japan is situated at the eastern extremity of Asia and consists of a group of islands, known as the Japanese archipelago, which stretch in a long curve in the north-western corner of the Pacific Ocean, and the Korean Peninsula (now called Chōsen) projecting from the Asiatic Continent. The said group of islands and the continent are separated by such marginal seas as the Sea of Okhotsk, Sea of Japan, and East China Sea. The islands themselves are again divided into the following smaller groups;—the Chishima (Kuriles), Japan Proper, Ryūkyū (Loochoo), etc, each forming a crescent-shaped curve, and thus present, as it were, the appearance of festoons suspended along the eastern shores of the Asiatic Continent. The largest of these islands is the Main Island (Honshū) located about in the middle, which is adjoined by the islands of Shikoku and Kyūshū in the south and the main island of Hokkai-dō in the north. Farther north than Hokkai-dō, there is Karafuto (Saghalien), of which the northern half is Russian territory. Down in the south, the island of Taiwan (Formosa) faces across the Bashee Channel the Phillippines belonging to the United States of America.

Generally speaking, all these islands are mountainous and carry two prominent mountain systems, meeting each other in the central part of the Main Island. In addition, there runs across these systems some volcanic chains, such as the Chishima-Zone, the Kirishima-Zone, and the Fuji-Zone; the last one intersects the middle portion of Honshū. Such being the case, the central part of Honshū is not only its broadest part but also the most complex and variegated in geographical conditions; and here are by far the highest and steepest mountains of the Empire.

As might be expected owing to the topographical nature of the country and the prevalence of volcanoes, mineral springs gush out all over Japan and number more than one thousand. Most of these hot springs are located amid beautiful scenery and the temperature and air leaves little to be desired, so that, combining climatic and therapeutic advantages, nature has adapted them to form ideal health resorts.

In observing the geographical features of Japan carefully, the following conditions may be noticed:—

- (1) The abundance of water; every valley among the mountains has a stream of clear and lucid water, constantly flowing and never dried up.
- (2) The diversity and density of vegetation; especially needle-leaved and foliage trees thrive everywhere and sometimes form very thick forests, thus contributing greatly to harmonize atmospheric humidity and adjust the flow of water in rivers.
- (3) The variety of geographical structures and consequently of sceneries.

These three elements as a national asset do much to enhance the distinctive charms peculiar to Japan. To cite some examples of special merit and admiration we might mention as typical the beach of Suma and Maiko (both in the neighbourhood of Kōbe) with its beautiful white sand dotted with green pines of quaint picturesque shapes of various age and size; the shady Mt. Arashi-yama in the suburbs of Kyōto with a rippling stream at its foot; the famous hot spring resorts, e.g., Hakone, Atami, Shuzenji, Ikao, Kusatsu, Arima, Beppu, etc.;—all these charming spots may be counted among the choicest of Japanese pleasure resorts. The land is fertile in cultivation and rich in a continual seasonable supply of domestic fresh vegetables, and moreover the country being seagirt, many kinds of fish are obtainable everywhere. This wealth of vegetables and fish forms a light diet every visitor finds satisfying and reasonably cheap. Furthermore the mild climate and fine weather generally prevailing make Japanese visits always enjoyable. The last, but not the least, significant among objects of interest, are the historic remains and monuments scattered all over

the country, whereby visitors with an inclination for antique study may find ample interest in tracing Japanese history. Indeed the combination of healthy climate, historic interest, and scenic beauty is nowhere so well-blended and complete as here in Japan. Thus each year brings more tourists to these shores and it goes without saying that this great increase in the number of the visitors is due to the inducements offered by the natural attractions of this country. Some tourists of international experience speak comparatively very highly of Japan and go so far as to remark that although Switzerland and Italy may well be proud of the majesty and grandeur of their scenery, neither country can excel Japan in the subtly soothing display of nature, its lichen and moss clothed rocks, its incomparable blossom, its terraced fields and pastures, its musical streams diapasoned with rumbling water-wheels, its thatched homesteads and peaceful landscapes characterized by a delicate gentle picturesqueness. Such a statement cannot be considered simply as a compliment, for indeed Japan in justice deserves the name of the Park of the World. From a climatic point of view, the whole of Japan is a great health resort. The only drawback is that our hot springs, as well as our mountain and seaside resorts leave much to be desired in point of various equipments either for enjoyment or accommodation, and owing to a past seclusive reserve their natural excellence and attractive merits have not been turned to the best advantage nor their rich resources fully utilized. But the Government Authorities are considering the organization of a society for the investigation of Japanese health resorts, in order to carry out their study in a scientific manner, will further provide some of those promising places with adequate arrangements and perfect their efficiency for medical purposes. Therefore it is confidently anticipated that all our hot spring, mountain, and seaside health resorts will in the near future be so developed in all that pertains to creature comfort and enjoyment, and creation of a greater health, and that there will be sufficient commodious hotels and hydros suitably equipped on modern lines to accommodate any demands the foreign visitor may make.



ENT RAILWAYS



EXPLANATIONS

- Railway Lines.
- Prefectural or Provincial Boundaries.
- Rly. Lines)
- Noted Rivers.
- Always' Steamer Lines.
- Noted Mountains.
- The names of Prefectures are shown in red.



IMPERIAL GOVERNMENT MAP OF RAILWAYS

Chapter I

SCENERY OF JAPAN.

A cursory glance at any map of the Northern Pacific shows us a long chain of islands hanging like a jewelled festoon on the aqueous frontage of the Asiatic continent and extending like guardian sentinels, the enormous length of about 2,500 miles from tropical seas to the subarctic waters. Adding to these the projection of the large peninsula of Chōsen (Korea) from the continent, these form our Japanese Empire.

The drift of a warm equatorial current washes the east coast, acting similarly as the Gulf Stream of the Atlantic, and summer and winter monsoons bring with them a plentiful precipitation of rain and snow throughout the year. When the traveller becomes tired of the monotonous fortnights' voyage across the Pacific Ocean or the tedious travel through the sameness of Siberian plains and sombre forests he will hail with a sense of relief, the first glance of the contrasting leaves of the book of nature, the picturesque Japan, its isles and islets and mountains covered by beautiful leafy forests. As one's steamer nears the coast one sees white silvery shining sandy beaches lined with beautiful pine-trees and dotted with fishermens huts and seaside villas, while in summer these beaches resound with the joy and laughter of the thousands who come to breathe the fresh exhilarating air and to swim in the warm genial rolling surfs. When one steps farther inland one will find everywhere clear and lucid hot springs and neatly equipped bathing houses, and that all these seas, mountains, and forests, play an important rôle in the beautiful scenery of Japan and form that base for the undieing native love of the homeland the visitor at once can understand and feel in sympathy with.

Situated on the oceanic border of the Asiatic Continent, the Japanese Islands and Korean Peninsula have been influenced enormously by several geological agencies. The present features of these lands are the results of the repeated modifications and transformations from the remotest archæan era to the recent. Especially during the palæozoic and tertiary age were these changes exceedingly violent, though the general outlines of the islands were constructed mainly during the tertiary period. The archæan gneiss and granite in Japan which lie under the paleozoic rocks are the prolongation of that system found in Central China, particularly do we find these well developed in the Korean Peninsula with its smooth downs and gentle escarpments. The deeply engrained valleys of crystalline schists, slate and sandstone of palæozoic formation in Kyūshū and Shikoku, and the rounded hills of granite in Chūgoku along the Inland Sea, form a distinct scenery characteristic of Southern Japan. In Northern Japan these palæozoic mountains have been cut into several blocks, which were afterwards surrounded by the tertiary sea. Its tufaceous deposits then were elevated and form at present extensive subdued mountains and hills, which are a characteristic of that district. There are small patches of the mesozoic deposits, but they have not any topographic peculiarities. During the post-tertiary age violent eruptions of volcanoes took place, these modifying strongly the configurations of land features throughout the country, especially in Kyūshū and Northern Japan. Numerous volcanoes were formed in several trails along the long axis of islands. Besides these trails there is another large one known as the Fuji-Zone across the Main Island, Honshū. On the northern boundary of Korea, there is an extensive area of lava plateau, extending further into Kirin Province in Manchuria. Though it has not come into popular notority, the scenery of this plateau land is found to be of an excellent type. The terraced deposits of the old quaternary epoch are developed very well in Northern Japan where volcanic activity has been powerful. Both these terraces and younger fluvial flats afford scope for the development of agriculture and industry, and attract a prosperous inhabitation.

Now let us briefly describe our land, namely the characteristics of its mountain ranges, volcanic cones, valleys, lakes and seas. The lofty and wide extending elevation of the principal mountain chains are naturally caused by the orogenetic actions, while their rugged peaks, gentle hills, deep gorges and wide valleys, all such features are detailed by the erosion, which is especially effective in such a country as Japan on account of much rainfall. Tectonically our islands consist of two arcs of mountain systems: the northern system extends from the Island of Saghalien southwards,

while the other stretches itself from China north-eastwards. They run along the northern and southern halves of the islands, and meet together in the central part of the main island of Honshū, where the most mountainous and high tracts are formed. Hida and Akaishi mountains so called the "*Japanese Alps*," which are now popular among our mountaineers, are the most eminent ridges in this district. Only one day's tour from Tōkyō by the Central Railway is required to find ourselves at the foot of the snow-clad peaks or in the alpine forests of these lofty mountains. Within their lonely valleys are some rural and simple bathing places, such as *Shirahone*, *Kamikōchi*, *Nakabusa*, etc. Not only by the mountaineers but by the people of the neighbouring provinces, these places are celebrated as summer resorts. The eminent needle of the Japanese Matterhorn, *Yari-ga-take*, and the rugged peak of our Jungfrau, *Hotaka-yama* with their perpetual snow fields, tranquil pools, or roaring torrents in their foregrounds, never fail to make impressions on the memory and Kodak films of the nature-loving tourist.

The numerous volcanoes we may divide into two types with regard to their forms,—one being represented by the simple cone of *Mt. Fuji*, and the other by the truncated cones with large caldera and central cones similar to *Mt. Hakone*. Without exaggeration the lofty cone of *Mt. Fuji* is, "*Mons excelsus et singularis*" rising in majestic grandeur 12,467 ft. above the sea-level, on the picturesque Bay of Suruga. Its perfect and splendid conical peak is clothed in snow nearly all the year round, snow remaining for a few weeks in summer only in shady nooks, hollows and crevices, and during that season the mountain is visited by multitudes of pilgrims climbing its sacred slopes as well as the feat-loving sportsman and worshippers of the sublime in nature. Not only to see the blush of sunrise illuminate the noble brow of this our "National Mountain," to the student and savant she is well known as offering many scientific attractions. On her sweeping skirts extend the pleats and ruffles of many lava flows, decorated further in folds and hollows with the translucent liquid sparkling gems of several silvery lakes. One of these lava streams forced its way down about fifteen miles along the valley of *Katsura-gawa*, until it stopped at the canyon of *Enkyō*, which is here spanned by the well-known bridge of the same name. Many parasitic cones, lava tunnels, and especially a great explosion crater on the southern flank, offer geological problems and studies much valued by the erudite scholar, whilst it is also very interesting not only for the botanical specialist but for all classes of mountain climbers to observe in one day's tour the varied flora changes found at different altitudes. The thicker forests and denser undergrowth of shrub and bush of the temperate zone found at the foot of the mountain change into conifer woods as we ascend a few thousand feet, and arriving half-way up the mountain we meet only such grasses as are usual in far northern districts, and even these soon disappear, and on the bleaker heights we see only naked blocks or lava, sparsely specked with some rarer varieties of lichen. A mountain observatory found on the summit is easily recognized by the indications of a vane and anemometer, restlessly turning on the top of a rocky cliff of the crater wall. Other mountains such as "*Kaimon*" in Kyūshū, "*Nantai-san*" (Nikkō), "*Iwaki-san*" and "*Ganju-san*" in Honshū, "*Makkarinupri*" or the so-called *Yezo-Fuji* answer much to the *Mt. Fuji* example, and we find these are also frequented by many people during the summer period when routes are open and climbing practicable.

In contrast to the lofty peak of the *Fuji* type comes a second attraction with its mighty *somma* (old crater) a yawning subsidence of mammoth proportions and another kind of world wonder. A vast caldera or amphitheatre where nature's forces have fought and wrestled in the lists of time is now peacefully turned into fine meadows, brooks and lakes, and forms a natural park, and moreover here are often found charming and comfortable bathing places. Of such, a typical example is *Mt. Hakone*, to reach which is only a three hours' journey in train and tram-car from Tōkyō. Here the beautiful *Lake Ashi* reflecting on its placid surface the inverted image of *Fuji* seen in the background, and the grand panorama of the mountains and seas as viewed from any point on the *somma* are splendid and beyond the power of human pen or word to describe. Many beautiful places, well known from mediæval times by the name of "Seven Hot Springs" are found excellent ones in this district, and now boast and justly so the best equipped foreign and native hotels. The grandeur of the caldera of *Mt. Aso* in Kyūshū is far more remarkable, its diameters are ten miles N.-S. and fifteen miles E.-W., its walls around stand at least 3,000 ft. high and the highest peak "*Saka-dake*" reaching to a height of 5,310 ft. Twenty thousand people living in about eighty villages are here sheltered and sustained in the lap of mother earth, a basin that is computed by scientists to undoubtedly be the largest crater ever found or known to man on earth. This elaborated

example and incredible display of immensity of volcanic action still retains a smoldering life, and of the five central cones, one is still found active while a number of hot springs are scattered at their feet. Among the many other volcanoes, especially noteworthy are "*Asama*" with its ascending column of smoke, "*Bandai-san*" with its tremendous explosion crater, "*Nikkō*" with its magnificent water-falls and lakes, "*Kirishima*" with some dozen craters and hot springs, and "*Unzen*" also with many springs and charming coast scenery, "*Sakura-jima*" in the Bay of Kagoshima came also recently to be well known by its sensational great eruption in January, 1914.

Some extinct volcanoes have suffered through the ages such powerful or strange erosion from weather action that they present peculiar mountain formation. In the vicinity of Karuizawa, the famous summer resort, there is a volcanic region, which is so much dissected, that even geologically it is very difficult to trace its original structure. *Mt. Myōgi* in this region is a most remarkable relic of volcanic grotesqueness, its natural arch, rock spines, pinnacles, numerous freak structures, and other wonderful views characterize this type of volcano, another example of which is to be seen in "*Yabakei*," Kyūshū. In the region of the above mentioned volcanic areas we find many hot springs being utilized by the people for bathing and other useful purposes and on which we shall devote a few lines in the next chapter.

As stated previously the current of the mountain streams are often very rapid, and to punt down such torrents over churning rapids and through rocky channels and narrow defiles affords a thrill of adventure and gives a very interesting enterprise and enjoyment to mountain tours. Usually the water which is accumulated in some intermontane basin at a high altitude, flows down to the piedmont flat making deep gorges on the border of upland. The well-known rapid, *Hozu*, near the old capital of Kyōto, is a deep cutting on the western fault-scap of the Tamba Plateau. The *River Tenryū*, *Fuji*, and *Kiso*, in Central Honshū, and the *River Kuma* in Kyūshū are also examples of the same kind but of larger and grander proportions. These rapids are commonly so swift that even the skilled crew often come dangerously near having their boats dashed against the rocks on either side or swamping them in the water. Reverting to the volcanic region, we can always find wonderful water-falls as "*Kegon*" in Nikkō with its rainbow hued mists, *Chinda-no-taki* on Mt. Aso, etc. The majority of well-known water-falls in Japan are nothing but the water leaping over the edge of some lava flow, but the setting and pictorial mounting of each is an undoubted call on one's highest admiration.

Of course in such mountainous islands as Japan, no one can expect to see the development of large plains, most of our plains are the deltas in the mouths of large rivers or if otherwise they are but narrow flat lands extending some miles along the river. At the foot of volcanoes there usually extend grassy plains with gentle slope, and when these are situated in higher districts, they command a view over a wide stretch of country and on account of the fine bracing air they attract many people in summer for invigorating recuperation such places as "*Karuizawa*," "*Gotemba*," or "*Fujimi*," being resorts of this class, popular, and worthy of mention.

Lake sceneries are not by any means rare in this country, and on volcanoes we find often their old craters filled with snowy water forming lucid emerald lakes. *Lake Ashi* in Hakone, already referred to, or *Lake Haruna* on the volcano of the same name, are the typical examples of these. But after all, *Lake Biwa* is the most popular and well known for its scenic beauty, being also the biggest depression lake in this country. Situated only a few miles from the old metropolis, Miyako, or Kyōto, and the main highway and railroad passing along its shore, it has been since long ages past a favourite resort theme, subject and inspiration for poets and artists. The eight choice views of this pear-shaped sheet of water have become interwoven into soul of the nation through their praises becoming familiarized in song, verse, and illustration.

Along the coast of the sea there are many lagoons embraced by the sand-dunes and sand-spits. *Hachirō-gata* in northern Honshū and the twin lakes of *Shinji-ko* and *Naka-no-umi* in Chūgoku belong to this type. The latter is often called the "*Lac Lemane of Japan*" and the finely situated town of *Matsue* is compared to Geneva for its pretty view. We must not neglect to mention these sand-spits as being important elements composing the scenic beauty of the coast, often covered with pine groves, some of those sand-spits stretch long arms for miles into the sea and make such typical beauty spots as *Ama-no-Hashidate* or "*Heavenly Bridge*," seen in the lovely Bay of Miyazu on the coast of the Sea of Japan, as also *Miho-no-Matsubara* or the "*Pine Groves of Miho*" which adorn the beach of the charming Bay of Suruga at the foot of Mt. Fuji. Along these sandy

stretches are found many summer resorts and that never failing enjoyment of swimming in the surfs gently rolling in from the ocean. Along the shore of the Sagami Bay near Tōkyō there are many such places, among them “*Kamakura*,” “*Zushi*,” “*Ōiso*,” etc., are very famous, while “*Suma*” and “*Akashi*” near Kōbe are also well known.

Besides these flat sandy coasts, another type of the shore seen all over Japan, especially in Kyūshū and Shikoku, is the rocky coast with many indentations, accompanied by numerous islands and rocks, giving diversified beauty. But after all the scenic beauty of the sea may be said to culminate in the world-wide known *Seto-uchi* or the Inland Sea. Enclosed from the tempestuous influence of the ocean, the spirit of peace seems to waft her angelic wand over the calm smooth glassy surface, which seldom shows a ripple, and mirrors the quaint forms of sailing craft, the green pine crested sunlit rocks and islands, and azure cloud flecked dome above in such vivid truthfulness, while the temple-strewn headlands and straw-thatched fishing hamlets nestling in nooks give the deck-bound passenger an ever changing panoramic feast, no wonder the impression given is indelible, the revel in colour seemingly a dream and the traveller who comes to the Far East and misses taking in this lovely passage certainly is robbed of an inestimable pleasure. The world tourist wearies of the rounds of abbeys and cathedrals, and becomes nauseated with the medley of museums and art-galleries, but the harmony of the Inland Sea hums on the finest silken heart strings and once seen vibrates on the mental vision to eternity in colours that shame earthly pigments. Did you see the Inland Sea? is the one question asked. Don't forget to see the Inland Sea is the advice given.

Chapter II

VOLCANOES AND MINERAL SPRINGS IN JAPAN.

From information to hand we have already been able to state that there are in Japan more than one thousand and five hundred mineral springs, already chemically analyzed. Of course there are many practically unknown, or at least not yet described, and valuable ones, especially in the lonely mountain region of Korea and volcanic districts of Hokkaidō are still to be reported on. It is most noticeable that the volcanoes and mineral springs have a consanguineous relationship here as everywhere else, and that their distributions are very intimately associated also here in Japan.


As the accompanying map shows, we find two main volcanic belts in Hokkai-dō which are the prolongations of the volcanic chains of the Kurile and Honshū ones. Accompanying the numerous active volcanoes “*Koma-ga-take*,” “*Usu*,” “*Tarumae*,” “*Tokachi*,” “*Meakan*,” etc., there are also many springs, but on account of the inconvenience of communication, only some of them are known at present, as those springs of Noboribetsu and Jōzankei. *Noboribetsu*, not far from the harbour of Muroran on the southern coast, is famous for its solfatara in full activity with numerous jets and formerly with a geyser. *Jōzankei* in a beautiful valley near the city of Sapporo is a spring well frequented.

In northern Honshū there are two main volcanic belts running along the long axis of the islands, one of these, the *Nasu-Zone* named after the famous active volcano, forms the backbone of the island of northern Honshū, and is characterized by its compact arrangement of volcanic cones accompanied by innumerable hot springs. The other belt lies on the coast of the Sea of Japan and similarly has well-developed conical mountains, a representative being *Mt. Chōkai*. Among the many hot springs in the northern end of Honshū, *Asamushi* with its railway communications and superior coast scenic attraction is a spring highly popular and widely known, while further inland we can find not only many valuable medicinal springs, but the famous geyser of *Fuki-age* situated in this district; but these latter springs being rather out of the way or a half-day's journey from the nearest railway station, visitors show a preference for the more conveniently connected springs as at Asamushi. Isolated springs only await the recognition of their merits but referring to the question of convenient situation in the vicinity of large cities and towns, there are many celebrated bathing places, e. g., “*Aone*” near Sendai, “*Kaminoyama*” near Yamagata, “*Iizaka*” near Fukushima, “*Higashiyama*” near Wakamatsu, etc., and further southwards more well-known places as “*Nasu*,” “*Shiobara*,” “*Nikkō*,” and “*Ikao*.” *Nasu* is popularly known by the tradition named *Sesshō-seki* or “Death

Distribution of Mineral Springs and V



Distribution of Mineral Springs and Volcanoes in Japan.



Stone," a feature connected with its solfatara. Shiobara has a series of springs and is most celebrated for its remarkable autumnal foliage, the brilliant maples, etc., seen in its glens and mountain ravines. No one can fail to notice the Imperial villa on the banks of the blue gorge, fringed as it is in prominent relief by dark forests. Nikkō known world-wide for its splendid mausolean temples and scenery, has its bathing place, "*Yumoto*," situated some few miles higher in the mountains and during the summer we find it well frequented. Ikao on the eastern flank of the beautiful volcano Haruna is another famous summer resort. These four springs may be reached from Tōkyō after a few hours' run by trains, tram-cars and *jinrikisha*. The above mentioned volcanic zones of northern Honshū meet with the Fuji-Zone in the central part of Honshū, forming there a congested volcanic district having many cones. Passengers on the Central and Shin-etsu Railway Lines successively, find a courteous and welcome treatment and every help by officials in reaching any desired point. The active volcanoes "*Asama*" and "*Shirane*," and the gigantic cone of "*Myōkō*" and "*Yatsu-gatake*" are most famous, while we further note other lofty volcanoes like "*On-take*" and "*Norikura*" on the ridge of the Japanese Alps. Accompanying these enumerated volcanoes there are found springs of various kinds, of which such well-known ones as "*Isobe*" near Mt. Myōgi, the most famous sulphur springs of "*Kusatsu*" at the foot of the Mt. Shirane, "*Shibu*," "*Bessho*," "*Asama*" (not associated with the volcano of the same name), and "*Suwa*" may be called to your notice. The latter one is on the shore of the beautiful *Lake Suwa*, which affords the best skating area for contests and sports during the winter. Strange to say Fuji has no share of hot springs, but in Hakone and some dissected volcanoes in the peninsula of Izu, in her zone, meet with numerous bathing places. The so-called "Seven Hot Springs" previously spoken of are found on the banks of the river Haya-kawa, which flows from the crater lake "*Ashi*" deeply cutting its way through the ring of mountains. "*Yumoto*" and "*Tōnosawa*" along the lower valley are simple thermal springs, while "*Ashi-no-yu*" and "*Owakidani*" on the higher reaches are remarkably strong in sulphur. "*Miyanoshita*" with its social life is the bathing place especially popular all the year round with foreigners, and "*Sokokura*" also a bathing resort being situated near, both these springs being of intermediate chemical nature. Near Hakone, on the eastern coast of Izu Peninsula, there is the celebrated geyser of "*Atami*." During the last few years, on account of the careless boring of wells in the vicinity of the geyser, the periodical discharge of hot water has become more or less irregular. Within a day's trip from Atami we can find numerous well-visited bathing places, as "*Izusan*," "*Yugawara*," "*Itō*" and "*Shuzenji*."

Turning to the western half of Honshū we have various hot springs, though they are less in number than in the Central and Northern Japan. Along the coast of the Sea of Japan there is one volcanic belt with the volcanoes "*Hakusan*," "*Daisen*," "*Sambe*," etc., which passes over to the island of Kyūshū. Another belt is found on the Inland Sea, or "*Seto-uchi*," but its volcanic life is almost extinct. The coast of the Sea of Japan is also not without its hot springs, of which the most noted are "*Wakura*" in Noto Peninsula, "*Yamanaka*" and "*Yamashiro*" in Kaga, and "*Kinosaki*" in Tajima, this latter with a most famous basalt cave "*Gembu-dō*" in its environs.

In the island of Shikoku, where we have no historic record of volcanic eruptions, there are but a few bathing places worthy of note apart from the celebrated "*Dōgo*," whereas the neighbouring great island of Kyūshū, is very richly besprinkled with volcanoes and hot springs, the distribution of these being on the two main zones of "*Aso*" and "*Kirishima*." The former connects the famous volcanoes "*Aso*," "*Unzen*," etc., while the other stretches southwards from the enormous volcano of "*Kirishima*" passing through "*Sakurajima*," the terrible monster in the Bay of Kagoshima, to the Kawabe islands and Formosa. There are many active volcanoes in both zones, and connected with which we have a terrible history of catastrophes.

Among the numerous bathing places in Kyūshū, "*Beppu*" and "*Unzen*" stand most prominent. The former lies at the foot of the volcanic group of "*Yufu*" on the shore of the picturesque Bay of Beppu, an indentation of the charming Inland Sea. Many springs come out in a novel and strange manner, some of them make ponds of boiling water, while the others dissipate themselves into the sand of the coast, and here the sand-bath taken on the roomy and clear beach is the most interesting and unique one of its kind found in this country. The volcano "*Unzen*" is a beautiful mountain on the peninsula of Shimabara and commands a fine panorama of land and sea. Its great eruption in 1792 still forms a theme of awe and topic of a conversation among the countrymen. Many thermal springs, especially "*Obama*" are well known for the bathing, and form excellent

summer resorts for natives as well as foreigners, and being situated only a few hours' steaming distance from Nagasaki, they are visited every summer both by the local foreign community and the numbers of Europeans and Americans living in China who seek for shelter and change from the summer heat and debilitating influences of that climate. The volcanoes "Aso" and "Kirishima," some of their peaks yet active, have also many bathing places. At the southern end of the Satsuma Peninsula there is the perfect cone of "*Kaimon*" and the crater harbour of "*Yamakawa*," and many springs are found in their vicinities, though the bathing houses are mostly a very simple and unassuming type.

As to hot springs in Formosa we cannot neglect to refer one to the famous one of "*Hokutō*," near Taihoku, the capital of the island. An abundant quantity of hot water is constantly flowing through a valley at the foot of the volcano "*Daiton*," and the deposit of the radio-active mineral "*Hokutōlite*" in that spring has recently attracted keen attention in the scientific world. The governing authorities undertook to plan a park and a splendid and magnificent public bath has been built which has no rival throughout the whole Empire of Japan. Other springs have been also discovered in the deep valleys of the central mountain chain, where the aborigines had previously built primitive bathing places by excavating the natural rock, further details of these being lacking we but record the fact.

Likewise of Korea we have not material enough yet to hand to describe any springs though there are many localities already known, and among which the much frequented bathing places of the following might be mentioned:—"*Tōrai* (Tong-nai) near the harbour of Fuzan and accessible by train, "*On-yō*" (On-yang) in Central Korea and not far from the railway station of Ten-an, and lastly "*Shu-otsu*" (Chyu-eur) in Northern Korea, this lying to the west of Ranan and Kyojō, where a sanatorium for the army is established.

Turning to the relationship between the mineral springs and volcanoes in Japan, it is of course obvious that the volcanic belts are usually accompanied by mineral springs, but we do not venture to assert that mineral springs are necessarily apart of or only confined to those districts. Some of them we find along the dislocation lines quite independent of volcanic zones, while others are noticed in the regions of old plutonic rocks. Where the volcanic regions is traversed by dislocation lines it is of course most favourable for the springs to find their way, as seen in the case of Shinano Highlands, Izu Peninsula, and the middle part of Kyūshū. From the accompanying map we note there are two sets of principal tectonic lines—longitudinal and transversal—which separate our islands into four quadrants. The above mentioned four regions are traversed by these tectonic lines as well as volcanic belts. Of the innumerable mineral springs in Northern Japan we find they are mostly limited to the north-western quadrant, while in the north-eastern quadrant there are only a few springs. The former is not only rich in volcanoes, but its younger beds have been influenced by local disturbances, and even at present are frequented by local earthquakes, while on the contrary the latter area has remained as plateau lands of older formation since remote geological ages without showing any peculiar disturbances but those due to denudation and erosion. Besides the hot springs, carbonated cold springs are found in the vicinity of Ōsaka, and the bottled liquids are in an ever increasing popular demand in the Far East as a table water under the names of "*Tansan*," and "*Hirano Water*." These springs are found along the line of dislocation on the border of the Tamba Plateau.

It is remarkable that the radio-active springs, which are highly esteemed for medical purposes, are mainly formed in the north-western quadrant, where the geology is represented by granite. The cold springs of *Masutomi* in the Province of Kai, and the *Misasa* hot springs in the Chūgoku district, are typical illustrations of this geologic feature.

Regarding the utilization of the products of mineral springs we have only a few things to say. Various kinds of sinters under the name of *Yu-no-hana* or "Flower of Spring" are sold in several bathing places for use in private or household baths. In Beppu, alum is prepared by the action of sulphureous steam upon alum-earth and in Kusatsu the incrustation of sulphur is collected. White clay, which is a bleaching product of volcanic rock, etc., produced by solfataric action at Beppu is used in the manufacture of paper. There are also calcareous and sulphur deposits of ancient hot springs, the former is found in the Prefecture of Akita in northern Honshū and is utilized as a flux in the smelting furnace, and the latter, which gives us the range and forms of marketable products of sulphur, is worked from many volcanoes and exported to various parts of the globe.

Chapter III

GEOLOGY OF THE JAPANESE EMPIRE.

The lowest stratified rocks composing the Japanese Empire are the Gneiss and Crystalline Schist Systems of the Archæan Group. Then follows Palæozoic formation, in which the earliest fossiliferous rocks in the empire are found. The Mesozoic formations were next deposited. During the sedimentation of the Palæozoic and the Mesozoic rocks, intrusions of granite, diorite, porphyry, gabbro, porphyrite, and diabase occurred. In the Cainozoic era, especially in the Tertiary age, volcanic rocks were erupted in different places and the sediments of volcanic ejectamenta formed the various kinds of tuffs.

ARCHÆAN GROUP.

1. *Gneiss System*:—The Gneiss System consists of biotite-gneiss, granite-gneiss, mica-schist, and amphibolite; and is frequently intercalated with saccharoidal limestone and granulite. Besides, two-mica-gneiss and pyroxene-gneiss are known in Korea, and in southern Manchuria. Various kinds of gneiss show in several cases the characters of the ortho-gneisses, while the mica-schist and limestone seem to be the para-gneisses metamorphosed by the contact action of eruptive rocks. Granites frequently intrude them in a confused manner, and these granites as well as the others which metamorphosed the Palæozoic rocks into gneisses and schists, are often treated together with the sedimentaries.

2. *Crystalline Schist System*:—The Crystalline Schist System consists of various schists of a phyllitic aspect with the characteristic components—sericite, epidote, and calcite. They are graphite-gneiss, epidote-gneiss, sericite-gneiss, graphite-schist, chlorite-schist, sericite-schist, piedmontite-schist, quartz-schist, and chlorite-amphibolite, often accompanied with eruptives, such as serpentine and gabbro, but not with granitic eruptives. Glaucophane-schist is known only in Shikoku and Kyūshū, and in the province of Kii. Crystalline schists are clearly distinguished from the rocks of the Gneiss System, except the amphibolites; but the distinction between them and the lowest series of the Palæozoic, consisting of amphibolites, pyroxenites, and phyllites, cannot always be clearly made.

PALÆOZOIC GROUP.

The Palæozoic of Japan consists of a series of formations of an enormous thickness, which may, in the order of superposition and lithological character, be divided into the Lower, Middle, and Upper Formations. The Upper Carboniferous or Permo-carboniferous fauna of stratigraphical importance is found only in narrow zones in the Middle and Upper Formations. The Lower Palæozoic is composed mainly of metamorphic volcanics or pyroclastic rocks, *i. e.*, pyroxenites and amphibolites, with subordinate layers of phyllites, crystalline limestone, and quartzite; and is often accompanied with serpentine and gabbro. The Middle and Upper Palæozoic consist of schalstein, sandstone, clayslate, quartzite, hornstone, radiolarian slate, adinole slate, limestone, and commonly rest conformably on the Lower Palæozoic. Among these rocks, crumpled quartzite and hornstone of various colours, adinole slate, schalstein with limestone, radiolarian slate, and *Fusulina* and crinoidal limestones, are easily recognizable and are indeed the marks of the correlation. Among the fossils, *Fusulina* and *Schwagerina* are most common. In Rikuzen there occurs a fossiliferous clayslate, containing a trilobite, brachiopods and bryozoans, and occupying an upper horizon.

In Korea, the lower series of the Palæozoic consists of metamorphic rocks, mainly of mica-schist, hornblende-schist, quartz-schist, and phyllite, a part of which seems to represent a certain horizon of v. Richthofen's Ta-ku-shan Series. It is found in scattered patches in North Ham-gyōng-dō, Kyōng-geui-dō, and Chōl-la-dō. The Upper or Korean Formation consists mainly of quartzite, clayslate, and sandstone in the lower part, and of limestone and marl in the upper, and seems to represent v. Richthofen's Sinian Formation. The low *Karst*-like plateau in south Phyōng-an-dō and Hoang-hai-dō consists of a thick complex of Carboniferous limestone, containing *Fusulina* and other foraminifera.

In the Liao-tung peninsula, the Palæozoic is divisible into the Cambro-silurian and the Carboniferous Systems. The lower series of the Cambro-silurian, *i. e.*, the Ta-ku-shan Series, consists mainly of quartzite with subordinate layers of quartzose sandstone and clayslate, sometimes conglomerate

and limestone. The upper formation, *i. e.*, the Sinian, consists mainly of an alternation of sandstone and claystone with subordinate layers of quartzite, conglomerate, and limestone in the lower part, and of limestone and marl with sandstone, claystone and quartzite in the upper. The limestone of the various horizons contains brachiopods, pteropods, cephalopods and trilobites. The Carboniferous is found only in small patches, as in the vicinities of Fu-chou and Chin-chou. The lower formation of the Carboniferous consists of an alternation of limestone and marl. The upper formation or Coal Measure consists mainly of sandstone and claystone with coal-seams. The claystone contains abundant plant-remains.

MESOZOIC GROUP.

1. *Triassic System*:—The Triassic System occupies small areas, and consists mainly of sandstone, and claystone or shale, sometimes intercalating tuff, limestone, and also anthracite. The Triassic in Rikuchū, Tosa, and Higo, yields marine fauna, such as *Ceratites* and *Pseudomonotis*, which are considered to be closely allied to those of the Californian Trias. The Trias of Nagato yields Rhætic, flora; in Bitchū the *Pseudomonotis* bed underlies a plant bed, which seems to be comparable to the Rhætic of Nagato.

2. *Jurassic System*:—The Jurassic of Japan is in small detached areas, consisting of claystone or shale, sandstone, and conglomerate; schalstein, crystalline limestone and also anthracite are found in the Jurassic of Nagato. The schalstein, which is widely distributed in Chūgoku and northern Kyushu and mapped as the Mesozoic of an unknown epoch, resembles the Liassic schalstein of Nagato. The Jurassic in Rikuzen, Echizen, and Nagato contains marine fauna; that in Echizen, Etchū, Echigo, Kaga, Mino, and Hida consists of brackish-water deposits with plant fossils. Its lower horizon often contains fresh-water shells.

In Korea, the Jurassic develops widely in the south-east, and consists of sandstone, conglomerate, claystone, and schalstein, intruded by dykes and sheets of porphyry and porphyrite. Plant fossils are imbedded in it. Other small patches of the Jurassic consist mainly of sandstone, conglomerate, and claystone or shale; and in some places schalstein and limestone occur.

3. *Cretaceous System*:—The Cretaceous System has the widest distribution of all the Mesozoic, and is rich in fauna and flora. The lower Cretaceous on the Pacific side of Honshū and Shikoku consists of series of shale and sandstone with the so-called Torinosu limestone. The limestone contains abundant fauna, consisting of foraminifera, corals, bryozoans, echinoids, bivalves and gastropods. The lower Cretaceous in the provinces of Rikuzen, Iwaki, Kōtsuke, Kii, Awa, and Tosa consists of conglomerate, sandstone and shale. Generally, a *Cyrena* bed occurs in the lower horizon, a plant bed a little higher, and a *Trigonia* sandstone in the uppermost. The upper Cretaceous, forming a long chain from the boundary between the provinces of Izumi and Kii, to Amakusa through the provinces of Awaji, Sanuki, Iyo, and Bungo, with strips in Kōtsuke and Shinano, consists chiefly of sandstone and shale with conglomerate, and contains marine fauna, and some flora. On Amakusa Island, several seams of anthracitic coal are found in the series. In southern Shikoku it consists of sandstone and shale with conglomerate and limestone. In Rikuzen the upper Cretaceous consists of sandstone and shale with limestone, quartzite and tuff. The Cretaceous of Hokkaidō, consisting of shale, sandstone, and conglomerate, yields abundant forms of ammonites, besides, shells and other forms. The Cretaceous of Karafuto consists of shale, sandstone, conglomerate, and marl, abundant fauna being imbedded in the strata.

4. *Doubtful Mesozoic*:—The Mesozoic consists mainly of shale and sandstone, sometimes with impure limestone. The Mesozoic of Taiwan consists mainly of claystone with sandstone.

CENOZOIC GROUP.

1. *Tertiary System*:—The Tertiary consists of sand, gravel, clay, tuff, sandstone, conglomerate, and shale, with intervening layers of limestone, marl, coal-seams, and diatom-earth. As a result of volcanic activity tuffs are very widely prevalent, sandstone and shale showing very frequently a tuffaceous character. The Eocene is known in Ogasawara-jima (the Bonin group) where the series yields the genus *Nummulites*, which has been also discovered in Okinawa Island, and the limestone of Nakaosaka in Kōtsuke contains *Orbitoides*. It is highly probable that there exists a horizon which represents a Miocene-Eocene Series. Whether the Miike coal field in Chikugo, where *Aturia ziczac* has been discovered, belongs to the Eocene or not, is now under discussion. Pliocene and

Miocene beds have been recognized, the former being indicated chiefly by shell remains and the latter by plants and a few foraminifera, though the definite boundary between them has not yet been traced except in a few cases. The various kinds of gasteropods, solenochonchæ, conchifera, brachiopods, echinoids, asteroids, foraminifera, etc., are abundantly found in the Tertiary, and those found in the Pliocene belong mostly to species now living in the Japanese and Chinese seas, only a few being extinct in the neighbouring waters. Sometimes the bones and teeth of mammals and rarely the remains of insects are found in the post-Miocene; shark's teeth specified as *Carcharodon megalodon* are found in several parts of Japan. The fossil flora is found most frequently in the pre-Pliocene or Miocene strata, and very often in the Pliocene. In the pre-Pliocene or Miocene flora, both European and Arctic elements are found; while most of the Pliocene flora is very intimately related to species now living especially in the mountain regions of Japan, and some to those now met with in other parts of the world. From these facts together with the evidence of fossil fauna it is supposed that the Pliocene period had a colder climate than that now prevailing in those regions.

In Korea the Tertiary consists of shale, sandstone, conglomerate and tuff, with coal, and is found in a few small scattered areas. In North Kyōng-syang-dō the shale and tuff of the series contain some fossil floras.

2. *Quaternary System*:—(a) *Diluvium*: The Diluvium consists of sand, gravel, clay, and pumice, often covered with a fine deposit of a volcanic nature, and forms low undulatory plateaus or elevated platforms as well as terraces. The Diluvial platform near the western coast of Taiwan is covered with laterite. (b) *Alluvium System*: The Alluvium consists of fluvial deposits of sand, gravel, and clay, as well as the sands of beaches and dunes.

3. *Raised Coral Reefs*:—Raised coral reefs are only found in Taiwan and Ryūkyū along the beach, or inland, where they form the low plateau.

IGNEOUS ROCKS.

Granite:—The granites consist of hornblende-granite, granitite, and hornblende-granitite with muscovite-granite and are the most widely extended of all the plutonic rocks. Contact phenomena of granite on the Palæozoic and Mesozoic sedimentaries have been observed in several places.

Porphyry:—The porphyries occur forming large masses in the inner zone, especially of South Japan, but mostly as small dykes throughout the mountains. The rocks belong chiefly to quartz-porphyry and often on one side approach granite, known as granite-porphyry, and on the other, liparite.

Diorite:—The diorite is generally quartz-diorite. The age of eruption is often uncertain. Structurally it is considered to be one of the older eruptives, but one mass in Sagami intrudes the Tertiary. Gabbro-diorite occurs in small areas, often accompanied by gabbro.

Gabbro, Peridotite, Serpentine, etc.:—The gabbro, peridotite, serpentine, etc., often occur together. The gabbro and serpentine are found mainly in the Crystalline Schist and the Palæozoic, but those in Chūgoku seem to belong to a later issue.

Diabase:—The diabase is found intercalated in the Palæozoic and the Mesozoic, and is accompanied with schalstein, showing a mighty eruption of it in the Palæozoic and the Mesozoic eras.

Porphyrite:—The porphyrite occurs in tolerably large areas in several places; but generally, it is found as dykes or sheets in the Palæozoic and the Mesozoic. In Chyōl-la-dō and Kyōng-syang-dō in Korea it occupies somewhat extensive mountain districts.

Liparite:—The liparite is of various kinds, and its tuffs resulting from the eruptions probably since the beginning of the Tertiary period, develop extensively with thick sediments especially in North Japan. Besides, the liparite occurs as dykes in several places, intruding the Tertiary as well as the older formations.

Andesite:—The andesites, including dacite, mica-hornblende-andesite, pyroxene-andesite, and olivine-pyroxene-andesite, have the widest distribution of all the igneous rocks. Among them, pyroxene-andesite and olivine-pyroxene-andesite are the most widely distributed and form huge volcanoes, especially those of North Japan, such as Fuji, Asama, Nasu, Zaō, Chōkai, Iwate, Iwaki, etc. Mica-hornblende-andesite builds the main volcanoes in South Japan, especially those along the coast of the Sea of Japan, such as Norikura, Hakusan, Daisen, Sambe, Kujū, Unzen, etc. The

dacite is limited in distribution, being found especially in the northern part of Honshū. Rhombic-pyroxene-andesite forms along the Inland Sea a volcanic series of small simple cones, such as Kabuto-yama in Settsu, Iino-yama in Sanuki, Kofuji in Iyo, etc. The andesite has been erupted extensively since the Tertiary period, being in most cases accompanied with tuff, agglomerate-tuff, lava-breccia, etc. Some andesites are also frequently found as sheets and dykes. The volcanoes of Japan are mostly composed of andesite, and are 165 in number, among which 63 active ones are enumerated. They are simple-coned, such as Fuji, Kaimon, etc.; double-coned as Asama, Sambe, etc.; or more complex, as Aso, Hakone, etc.

Basalt:—The basalt outcrops in rather small areas in Chūgoku, northern Kyūshū, and in northern Korea, forming plateaus and sometimes simple domes or cones, such as Kasa-yama in Nagato, Kannabe-yama in Harima, etc. Also it often occurs as intrusive sheets and dykes in these regions.

Volcanic Ash and Mud Lava:—Volcanic ash is widely distributed especially along the slopes of volcanoes, covering the mud lava, or sometime alternating with it. Mud lava always forms plateaus and the slopes of volcanoes. It occupies wide areas in Kyūshū, where it is considered to have been erupted from the volcanoes of Aso and Kirishima.

Chapter IV

CLIMATE OF JAPAN WITH SPECIAL REFERENCE TO HEALTH RESORTS IN JAPAN.

If there was no intense heat in August in Japan, who could be charmed by the autumnal glory of the coloured foliage in November? If there were no bitter winter monsoons from Siberia, who could expect the magnificent display of cherry blossoms all at once in April? We cannot incessantly endure a succession of good and fine days, so that Nature gives us summer to purify our blood by perspiring, and winter to brace our shattered nerves by its cold. The climate of Japan is moderate with two severe extremities both in time and in space.

Preceding the autumnal equinox, when the sun becomes somewhat lower in our middle north latitude, the high atmospheric pressure covering the North Pacific Ocean begins to send its force to the west, and the typhoons successively visiting Japan perform the great work of removing the action center of the atmosphere. The south-easterly monsoon gradually loses its intensity only to be replaced by the north-westerly monsoon, which threatens the inhabitants of the Far East. In Central Japan on some clear morning late in the October frost is seen whitening the stones and grasses even tinging the leaves. The mountain slopes are first seen variegated. After a few days the foliage in the valleys turn red or yellow and thence for about a month mountain slopes, hills, lakesides and wherever deciduous trees stand, present the autumnal glory of red and gold. No show window of a draper or mercer could ever be brighter and richer in colour than this. Some trees are as brilliant and as red as fire; others are of the colour of peach; some are as yellow as deep gamboge and still others are purple or white. Pine-trees are always mixed with these gaudily coloured trees and the deep blue of the sky over them gives a very pleasant contrast. Moreover we can see majestic mountain ranges whose shoulders are of ultramarine, but whose heads are as white as a sugar loaf. If we suppose a lake with clear water extending as a gigantic mirror between red and gold foliage; if we suppose a cascade hanging from the edge of white granite rocks, half-covered by coloured leaves and dark green pines, we have a fair picture of an autumnal day in Japan. Autumn scenes are very much appreciated in Japan, and Japanese sightseers never fail to visit some such places as *Arashiyama*, *Nikkō*, *Myōgi*, or *Mitake* during this season, to enjoy the autumnal tranquility amidst the sunny display of coloured nature. Such a pleasant season is but a prognostic of the coming penetrating winter.

From the middle of March there is a rapid change of climate in Japan. The high atmospheric pressure in Siberia begins to move to the Pacific Ocean and, in consequence, the severe, cold north-westerly monsoon gradually subsides and in its place the gentle beams of the spring sun warm the soil from which ice and snow have disappeared. We find that all moats, rivelets, ponds and marshes have become swollen with the water of spring, and the eggs of frogs are seen abundantly

in them. Plum trees in blossom from the previous month gradually give their places to soft green sprouts. At the end of March peach trees are in blossom and tadpoles are swimming about in the ponds. On the first or second week of April, cherry blossoms begin to open and are in their full glory all of a sudden. In this season nearly all the people in the great cities of Japan such as Tōkyō, and Kyōto, go out in full dress to play and dance under the blossoms. The weather is warm; swallows and butterflies are sometimes seen flying through the blossoms. Falling petals wafted on the breezes falls like snow flakes on the head and *kimono* (clothes) of the dancers. "*Which is the more beautiful, blossoms or girls?*" is repeated every year by sightseers. Such merry display of cherry blossoms and human flowers is but the reactional outburst against the solitary confinement of winter cold.

One of the most powerful factors in the variation of the climate in Japan is the **Monsoon**. Temperature, humidity, and other elements change very faithfully with it. The monsoon comes in winter from Siberia, and in summer from the Pacific Ocean. The empire of Japan extends from *Taiwan* (Formosa), whose southern half is in the tropics, to *Karafuto* (Saghalien) and *Chishima* (Kuriles) whose most northerly point is in Latitude N 52°. Hence the climate of Japan by no means uniform as already pointed out by many authorities, but it is certain that nearly all districts in Japan undergo the influence of the monsoons and hence the annual variation of the climate is very regularly repeated, and as far as any one of such districts as *Kyūshū*, *Naikai* (Inland Sea District) or *Ō-u* (north-eastern part of the Main Island) is concerned, the climate is uniform in the particular district. We can, thus, estimate the climate of a special place, if the general climate of the district in which the place is situated, is known, together with the orographical features of the place. This is illustrated by the following example. The temperature of *Hakone* must be lower than that of *Yokohama* by about 4.5° C, because Hakone is about 720 metres high above sea-level,—the average vertical gradient of the air temperature in mountainous districts in Japan is 0.6° C per 100 metres—while the temperature of *Miyanoshita* must be about 2° C higher than that of Hakone, because its height is about 420 metres. The observed mean temperatures at Hakone and Yokohama (from the report of Yokohama Met. Station) are shown in the following table, which proves the correctness of the above view.

Table I.

Average Temperature at Hakone and Yokohama. (° C).

| Name of Place | Jan. | Feb. | Mar. | Apr. | May | June | July | Aug. | Sept. | Oct. | Nov. | Dec. | Year |
|---------------|------|------|------|------|------|------|------|------|-------|------|------|------|------|
| Yokohama | 4.4 | 4.7 | 7.6 | 13.0 | 17.2 | 20.8 | 24.2 | 25.9 | 22.2 | 16.7 | 11.3 | 6.6 | 14.6 |
| Hakone | 0.0 | 0.6 | 3.5 | 8.2 | 12.3 | 16.9 | 20.1 | 21.1 | 17.0 | 12.3 | 6.4 | 1.7 | 10.0 |
| Diff. | 4.4 | 4.1 | 4.1 | 4.8 | 4.9 | 3.9 | 4.1 | 4.8 | 4.0 | 4.3 | 4.9 | 4.9 | 4.6 |

Thus Hakone and Miyanoshita must be cool in summer, but too cold in winter. *Odawara*, which is situated at the foot of *Hakone-yama* (Mt. Hakone) must be warmer than Yokohama in winter, because the west and the north side of this place are protected by mountain ranges which divert the raw cold of the north-westerly winter monsoons. The observed temperature of January is 9.8° C at Odawara and 8.5° C at Yokohama, the difference is 1.3° C; while that of August is 29.6° C at Odawara and 29.2° C at Yokohama; the difference is only 0.4° C. This fact coincides with the theory. In 1911, it was proposed to make climatological observations at mineral bath localities and health resorts. But there were many obstacles in the way, and since then data have been obtained only at about forty places. Some of these results are given in Table II, but the period over which observations extend is too short to discuss the climate of these places. Fortunately, however, meteorological stations of first and second order are so numerous and so closely situated in this country that for many bath localities and health resorts we can find some meteorological station standing in the close proximity to them, where the meteorological elements have been observed for many years. For example, we have Aomori Meteorological Station near *Asamushi* mineral bath resort (10 miles apart and both on the sea-shore), Utsunomiya Met. St. near *Nikkō*, Maebashi Met. St. near *Ikao*, Ōsaka Met. St. near *Hamadera*, Kōbe Met. St. near *Arima*, *Suma*, *Maiko* and *Akashi*, Matsuyama Met. St. near *Dōgo*, Oita Met. St. near *Beppu*, etc. The complete meteorological observations taken at all meteorological stations are given in the monthly and annual reports of the Central Meteorological Observatory of Japan, and an extract is given in the Japanese Almanac called

'*reki*.' We cannot here enter into minute descriptions of the climate observed at each of the meteorological stations. For our present purpose, Tables III—VI are prepared, in which the mean and the absolute extremes of air temperature, the mean and the absolute minimum humidity, the mean and the absolute maximum precipitation, with the number of rainy days and the number of the clear and fair days are given.

Table II.

Climatological Observations at some Health Resorts.

| | January | February | March | April | May | June | July | August | September | October | November | December | Year |
|---|---------|----------|-------|-------|------|------|------|--------|-----------|---------|----------|----------|-------|
| <i>Unzen</i> in Nagasaki Prefecture ; Observation, 1909–1912. | | | | | | | | | | | | | |
| Mean Temperature °C | 1,7 | 1,7 | 5,2 | 9,6 | 13,6 | 17,3 | 21,1 | 21,9 | 19,7 | 13,2 | 7,8 | 2,6 | 11,3 |
| Absolute Max. Temp. °C | | | | | | | | | | | | | 29,8 |
| Absolute Min. Temp. °C | | | | | | | | | | | | | -12,2 |
| Humidity % | 76 | 81 | 81 | 79 | 80 | 89 | 90 | 87 | 88 | 78 | 81 | 80 | 83 |
| Absolute Min. Humidity % | | | | | | | | | | | | | 27 |
| <i>Takeo</i> in Saga Prefecture ; Observation, 20 years. | | | | | | | | | | | | | |
| Mean Temperature °C | 5,3 | 4,7 | 7,9 | 13,0 | 17,3 | 20,5 | 24,8 | 27,1 | 23,0 | 16,9 | 11,9 | 7,3 | 15,0 |
| <i>Suma</i> in Hyōgo Prefecture ; Observation, 1898–1912. | | | | | | | | | | | | | |
| Mean Air Temperature °C | 6,8 | 7,1 | 8,5 | 13,2 | 17,3 | 21,2 | 24,7 | 27,5 | 22,0 | 16,4 | 10,7 | 7,3 | 15,0 |
| Mean of Daily Max. Temp. °C | 8,1 | 10,9 | 12,6 | 17,9 | 22,2 | 25,8 | 28,1 | 31,8 | 25,8 | 21,1 | 14,6 | 9,3 | 19,0 |
| Mean of Daily Min. Temp. °C | 0,5 | 3,4 | 4,4 | 8,5 | 12,5 | 16,6 | 21,3 | 23,3 | 18,3 | 11,8 | 6,8 | 5,3 | 11,1 |
| <i>Arima</i> in Hyōgo Prefecture ; Observation, 1898–1912. | | | | | | | | | | | | | |
| Mean Air Temperature °C | 2,5 | 2,0 | 5,1 | 11,1 | 15,6 | 19,6 | 23,2 | 24,3 | 20,8 | 14,7 | 8,8 | 4,2 | 12,7 |
| Mean of Daily Max. Temp. °C | 5,8 | 5,5 | 9,1 | 15,4 | 20,4 | 23,4 | 26,3 | 27,7 | 23,6 | 17,7 | 11,9 | 7,1 | 16,2 |
| Mean of Daily Min. Temp. °C | -0,7 | -1,4 | 1,2 | 6,9 | 10,9 | 15,9 | 20,1 | 20,9 | 18,1 | 11,8 | 5,9 | 1,4 | 9,2 |
| <i>Yumoto</i> in Kanagawa Prefecture ; Observation, 1902–1913. | | | | | | | | | | | | | |
| Mean Temperature °C | 5,3 | 5,3 | 8,3 | 13,2 | 16,7 | 20,2 | 23,5 | 24,7 | 21,6 | 16,5 | 11,6 | 7,4 | 14,5 |
| Absolute Max. Temp. °C | | | | | | | | | | | | | 34,3 |
| Absolute Min. Temp. °C | | | | | | | | | | | | | -7,0 |
| <i>Hakone</i> in Kanagawa Prefecture ; Observation, 1911–1913. | | | | | | | | | | | | | |
| Mean Temperature °C | 0,0 | 0,6 | 3,5 | 8,2 | 12,3 | 16,9 | 20,1 | 21,1 | 17,6 | 12,3 | 6,4 | 1,7 | 10,0 |
| Absolute Max. Temp. °C | | | | | | | | | | | | | 31,2 |
| Absolute Min. Temp. °C | | | | | | | | | | | | | -13,5 |
| <i>Kami-Suwa</i> in Nagano Prefecture ; Observation, 1911–1912. | | | | | | | | | | | | | |
| Mean Temperature °C | -2,5 | 1,5 | 4,3 | 11,0 | 15,5 | 19,2 | 23,0 | 24,2 | 19,6 | 13,1 | 7,5 | 1,0 | 11,4 |
| Mean Humidity % | 77 | 63 | 66 | 57 | 57 | 66 | 69 | 68 | 73 | 70 | 69 | 72 | 67 |

| | January | February | March | April | May | June | July | August | September | October | November | December | Year |
|---|---------|----------|-------|-------|------|------|------|--------|-----------|---------|----------|----------|------|
| <i>Asama</i> (near Matsumoto) in Nagano Prefecture; Observation, 1911-1912. | | | | | | | | | | | | | |
| Mean Air Temperature °C | -1.0 | 2.5 | 6.0 | 12.2 | 17.2 | 20.6 | 24.7 | 25.5 | 21.7 | 14.1 | 8.3 | 2.5 | 12.9 |
| Mean Humidity % | 30 | 71 | 64 | 50 | 60 | 73 | 60 | 75 | 74 | 72 | 60 | 72 | 60 |
| <i>Bessho</i> in Nagano Prefecture; Observation, 1911-1912. | | | | | | | | | | | | | |
| Mean Temperature °C | -0.2 | 2.8 | 6.0 | 12.8 | 17.2 | 20.6 | 25.3 | 26.4 | 21.1 | 15.1 | 9.1 | 2.3 | 13.2 |
| Mean Humidity % | 37 | 69 | 60 | 58 | 59 | 74 | 79 | 70 | 83 | 66 | 69 | 72 | 70 |
| <i>Karuizawa</i> in Nagano Prefecture; Observation, 1911. | | | | | | | | | | | | | |
| Mean Temperature °C | 1.8 | 2.1 | 5.0 | — | — | 17.3 | 18.8 | 21.9 | 19.9 | 11.5 | 10.1 | 0.4 | — |
| Mean Humidity % | 65 | 59 | 61 | — | — | 73 | 83 | 82 | 72 | 75 | 60 | 61 | — |
| <i>Nikkō</i> in Tochigi Prefecture; Observation, 1908. | | | | | | | | | | | | | |
| Mean Air Temperature °C | 0.5 | -0.5 | 3.6 | 9.0 | 13.3 | 17.7 | 18.6 | 22.1 | 15.8 | 12.1 | 5.0 | 1.7 | 9.9 |
| Mean Daily Range of Temp. °C | 13.4 | 14.3 | 12.5 | 14.5 | 6.8 | 13.2 | 10.2 | 11.8 | 10.8 | 13.2 | 14.5 | 12.0 | 12.3 |
| Absolute Max. Temp. °C | | | | | | | | | | | | | 35.5 |
| Absolute Min. Temp. °C | | | | | | | | | | | | | 9.5 |
| Humidity % | 75 | 75 | 74 | 79 | 76 | 83 | 91 | 84 | 91 | 81 | 68 | 64 | 78 |
| Absolute Min. Humidity % | | | | | | | | | | | | | 27 |
| <i>Kaminoyama</i> in Yamagata Prefecture; Observation, 22 years. | | | | | | | | | | | | | |
| Mean Air Temperature °C | 0.7 | 1.5 | 4.8 | 12.3 | 17.7 | 21.5 | 24.8 | 26.5 | 21.6 | 15.2 | 9.0 | 2.9 | 13.2 |
| <i>Takayu</i> in Yamagata Prefecture; Observation, 9 years. | | | | | | | | | | | | | |
| Mean Temperature °C | 2.8 | -3.2 | 0.5 | 7.0 | 12.0 | 16.3 | 20.1 | 21.5 | 17.0 | 11.5 | 4.7 | 1.2 | 8.7 |


Table III.

Mean, Absolute Maximum and Absolute Minimum Temperature.

| Locality | Name of Place | Height above sea | Latitude N | Longitude E | Number of years of observation | Mean Temperatures | | | | | | | | | | | | Highest Temperature | Lowest Temperature | |
|------------------|---------------|------------------|------------|-------------|--------------------------------|-------------------|----------|-------|-------|------|------|------|--------|-----------|---------|----------|----------|---------------------|--------------------|------|
| | | | | | | January | February | March | April | May | June | July | August | September | October | November | December | | | Year |
| Taiwan (Formosa) | Tainan | m. 14.3 | 22°59' | 120°12' | 15 | 17.3 | 16.4 | 19.7 | 23.3 | 26.0 | 27.2 | 27.7 | 27.4 | 27.1 | 24.9 | 21.4 | 18.4 | 23.1 | 36.9 | 3.0 |
| | Taihoku | 9.3 | 25° 2' | 121° 31' | 15 | 15.6 | 14.3 | 17.9 | 20.5 | 23.9 | 26.6 | 27.9 | 27.7 | 26.2 | 23.1 | 19.6 | 16.8 | 21.1 | 37.0 | -0.2 |
| Kyushu | Kagoshima | 119.7 | 31°35' | 130° 33' | 29 | 7.4 | 7.2 | 10.9 | 15.7 | 18.7 | 22.2 | 25.8 | 26.0 | 24.2 | 19.1 | 13.8 | 8.9 | 16.7 | 36.2 | 9.1 |
| | Nagasaki | 133.0 | 32°41' | 129° 52' | 33 | 5.8 | 5.8 | 9.1 | 14.3 | 17.9 | 21.6 | 25.5 | 26.7 | 23.8 | 18.0 | 12.6 | 7.6 | 15.7 | 36.7 | -5.2 |
| Shikoku | Oita | 5.9 | 33°14' | 131°36' | 25 | 5.8 | 5.4 | 8.3 | 13.2 | 17.0 | 21.1 | 24.9 | 26.0 | 22.7 | 17.3 | 12.3 | 7.7 | 15.1 | 36.0 | -7.5 |
| | Matsuyama | 32.4 | 33° 50' | 132°45' | 22 | 4.9 | 4.6 | 7.8 | 12.9 | 16.8 | 21.3 | 25.2 | 26.1 | 22.7 | 16.5 | 11.3 | 6.9 | 14.8 | 36.0 | -7.0 |

| Locality | Name of Place | Height above sea | Latitude N | Longitude E | Number of years of observation | Mean Temperatures | | | | | | | | | | | | Highest Temperature | Lowest Temperature | | |
|--------------|---------------|------------------|------------|-------------|--------------------------------|-------------------|----------|-------|-------|------|------|------|--------|-----------|---------|----------|----------|---------------------|--------------------|------|-------|
| | | | | | | January | February | March | April | May | June | July | August | September | October | November | December | | | Year | |
| Japan Proper | Main Island | Sakai | 2.9 | 35°33' | 133°14' | 29 | 4.1 | 3.6 | 6.9 | 11.0 | 16.1 | 20.5 | 24.5 | 26.0 | 22.1 | 16.3 | 11.1 | 6.4 | 14.1 | 37.8 | -9.7 |
| | | Kōbe | 58.3 | 34°41' | 135°11' | 15 | 4.9 | 4.4 | 7.6 | 13.2 | 17.5 | 21.3 | 25.0 | 26.6 | 23.0 | 17.2 | 11.9 | 6.9 | 15.0 | 36.5 | -5.5 |
| | | Kyōto | 49.4 | 35° 1' | 135°46' | 31 | 2.6 | 2.7 | 6.1 | 12.2 | 16.4 | 21.0 | 25.0 | 26.1 | 22.2 | 15.5 | 9.4 | 4.3 | 13.6 | 37.2 | -11.9 |
| | | Nagoya | 15.2 | 35°10' | 136°55' | 21 | 3.5 | 3.6 | 7.3 | 13.2 | 17.4 | 21.5 | 25.5 | 26.5 | 22.9 | 16.4 | 10.6 | 5.3 | 14.5 | 36.8 | -9.5 |
| | | Tokyo | 21.3 | 35°41' | 139°45' | 36 | 2.9 | 3.5 | 6.8 | 12.4 | 16.5 | 20.4 | 24.0 | 25.4 | 21.8 | 15.8 | 10.1 | 5.2 | 13.7 | 36.6 | -9.2 |
| | | Niigata | 25.6 | 37°55' | 139° 3' | 30 | 1.4 | 1.2 | 4.4 | 10.3 | 14.8 | 19.2 | 23.5 | 25.4 | 21.4 | 15.2 | 9.4 | 4.0 | 12.5 | 39.1 | -9.7 |
| | | Matsumoto | 582.1 | 36°14' | 137°59' | 14 | -2.0 | -1.7 | 2.4 | 9.0 | 13.7 | 18.4 | 21.0 | 22.5 | 18.2 | 11.6 | 6.0 | 0.9 | 10.1 | 35.2 | -24.8 |
| | | Utsunomiya | 125.0 | 36°34' | 139°53' | 21 | 0.9 | 1.5 | 4.9 | 11.1 | 15.4 | 19.6 | 22.9 | 24.2 | 20.6 | 14.4 | 8.1 | 2.7 | 12.2 | 35.7 | -14.8 |
| | | Maebashi | 112.9 | 36°24' | 139° 4' | 15 | 2.5 | 2.5 | 5.8 | 11.5 | 15.8 | 19.8 | 23.2 | 24.4 | 20.5 | 14.6 | 9.3 | 4.5 | 12.0 | 35.8 | -0.1 |
| Hokkaidō | | Aomori | 4.3 | 40°51' | 140°45' | 30 | -2.8 | 2.5 | 0.5 | 6.9 | 11.7 | 16.1 | 20.4 | 22.6 | 18.3 | 11.8 | 5.6 | -0.1 | 0.0 | 35.0 | -19.0 |
| | | Sapporo | 16.9 | 43° 4' | 141°21' | 35 | -6.2 | -5.3 | -1.6 | 5.1 | 10.4 | 14.8 | 19.0 | 20.8 | 16.1 | 9.3 | 2.7 | -3.3 | 6.8 | 34.1 | -25.6 |
| | | Nemuro | 26.7 | 43°20' | 145°35' | 32 | -5.0 | -5.5 | -2.5 | 2.9 | 6.5 | 9.9 | 14.2 | 17.4 | 15.2 | 10.5 | 4.3 | -1.4 | 5.5 | 31.9 | -22.7 |
| | | Karafuto | | | | | | | | | | | | | | | | | | | |
| Chōsen | | Udomari | 37.3 | 46°39' | 142°46' | 6 | -12.5 | -10.8 | -5.7 | 1.5 | 5.6 | 9.7 | 14.2 | 16.8 | 13.2 | 6.9 | -0.8 | -7.3 | 2.6 | 27.9 | -32.7 |
| | | Jin-sen | 67.6 | 37°29' | 126°32' | 8 | -2.3 | -3.0 | 2.6 | 9.2 | 14.3 | 19.1 | 23.0 | 24.3 | 20.2 | 14.3 | 5.8 | -1.6 | 10.5 | 34.0 | -18.7 |

Fig. 1.
Temperature.



The air temperature in Japan, except Taiwan (Formosa), belongs to the *temperate-zone type*. Nearly all temperature curves for places in Japan Proper lie between the curves for Nagasaki and Sapporo. If we exclude those for Ō-u and Hokkaidō, then the curves coalesce at the curve for Kyōto. The Tōkyō curve nearly coincides with that of Kyōto, which actually does in the warmer period. In these regions—from Kyūshū to Kwantō and Hokuroku excluding the high land in Central Japan—the mean monthly temperature does not fall below 0° C. In Japan Proper it lies between 5° C and 20° C for about 6 months of the year, excepting Kyūshū where for about 7 or 8 months.


Table IV.

Humidity.

| Name of Place | January | February | March | April | May | June | July | August | September | October | November | December | Year | Absolute Min. |
|---------------|---------|----------|-------|-------|-----|------|------|--------|-----------|---------|----------|----------|------|---------------|
| Tainan | 70 | 78 | 79 | 78 | 79 | 83 | 82 | 84 | 80 | 78 | 77 | 78 | 80 | 26 |
| Taihoku | 84 | 84 | 85 | 83 | 81 | 80 | 77 | 78 | 79 | 81 | 81 | 83 | 81 | 31 |
| Kagoshima | 71 | 70 | 72 | 75 | 77 | 83 | 82 | 80 | 78 | 73 | 72 | 72 | 75 | 14 |
| Nagasaki | 72 | 70 | 71 | 75 | 76 | 83 | 82 | 78 | 76 | 71 | 72 | 72 | 75 | 17 |
| Oita | 75 | 73 | 77 | 80 | 81 | 85 | 86 | 85 | 86 | 80 | 77 | 73 | 80 | 22 |
| Matsuyama | 73 | 71 | 74 | 77 | 78 | 80 | 80 | 81 | 83 | 80 | 76 | 72 | 77 | 15 |
| Sakai | 79 | 77 | 75 | 77 | 77 | 81 | 83 | 82 | 82 | 79 | 78 | 78 | 79 | 18 |
| Kôbe | 69 | 65 | 68 | 69 | 70 | 77 | 79 | 75 | 75 | 72 | 70 | 67 | 71 | 17 |
| Kyôto | 79 | 76 | 74 | 74 | 73 | 77 | 78 | 77 | 80 | 80 | 81 | 79 | 77 | 10 |
| Nagoya | 75 | 70 | 69 | 73 | 73 | 77 | 79 | 79 | 81 | 77 | 76 | 75 | 75 | 21 |
| Tôkyô | 65 | 63 | 69 | 75 | 77 | 82 | 84 | 82 | 83 | 80 | 74 | 67 | 75 | 8 |
| Niigata | 82 | 81 | 76 | 76 | 76 | 81 | 83 | 81 | 81 | 79 | 79 | 81 | 80 | 21 |
| Matsumoto | 82 | 77 | 74 | 71 | 71 | 76 | 80 | 82 | 83 | 83 | 78 | 79 | 78 | 17 |
| Utsunomiya | 72 | 68 | 70 | 73 | 70 | 80 | 84 | 84 | 84 | 80 | 77 | 72 | 77 | 7 |
| Machiashi | 63 | 59 | 62 | 68 | 73 | 79 | 84 | 84 | 84 | 78 | 69 | 63 | 72 | 15 |
| Aomori | 81 | 79 | 75 | 72 | 73 | 79 | 82 | 81 | 79 | 76 | 70 | 79 | 78 | 18 |
| Sapporo | 79 | 79 | 77 | 73 | 74 | 81 | 84 | 83 | 83 | 81 | 78 | 79 | 79 | 8 |
| Nemuro | 72 | 75 | 78 | 81 | 83 | 90 | 91 | 91 | 86 | 78 | 72 | 70 | 80 | 13 |
| Odomari | 85 | 83 | 81 | 82 | 83 | 86 | 88 | 88 | 84 | 79 | 75 | 81 | 83 | 28 |
| Jinsen | 69 | 62 | 62 | 69 | 74 | 80 | 84 | 81 | 73 | 60 | 66 | 63 | 71 | 17 |

Fig. 2.

Humidity.



On the Pacific coast the relative humidity varies yearly as does the air temperature on the same coast, but on the Japan Sea coast it changes like the rainy days. In summer there is no striking difference between these two districts. In winter the coast of Japan Sea is wet, the mean monthly humidity exceeding 80 per cent. But the Pacific and Inland regions are dry, the mean monthly humidity sinking below 65 per cent. In summer, it often exceeds 90 per cent on mountain slopes and on some capes and islands in Japan Proper, but this is not the case in ordinary residential localities. The dullest and the wettest period in Japan Proper is experienced in the *bai-u* season when, for about two weeks—from the end of June to the beginning of July—the weather is cloudy, warm and damp indoors. In Karafuto (Sagalien) and in the eastern part of Hokkaidō 90 per cent is surpassed in winter, but Chōsen (Korea) is very dry in winter.

Table V.

Amount of Precipitation. (mm)


| Name of Place | January | February | March | April | May | June | July | August | September | October | November | December | Year | Absolute
Max. in
24 hrs. |
|---------------|---------|----------|-------|-------|-----|------|------|--------|-----------|---------|----------|----------|------|--------------------------------|
| Tainan | 29 | 35 | 39 | 56 | 146 | 358 | 353 | 457 | 138 | 38 | 21 | 8 | 1677 | 385 |
| Taihoku | 95 | 132 | 104 | 137 | 206 | 245 | 196 | 328 | 262 | 103 | 80 | 80 | 2030 | 287 |
| Kagoshima | 91 | 87 | 153 | 230 | 240 | 372 | 294 | 186 | 228 | 128 | 94 | 81 | 2184 | 207 |
| Nagasaki | 82 | 82 | 133 | 198 | 200 | 326 | 239 | 191 | 220 | 111 | 84 | 82 | 1947 | 345 |
| Ōita | 50 | 58 | 123 | 135 | 170 | 232 | 224 | 161 | 231 | 152 | 76 | 49 | 1662 | 444 |
| Matsuyama | 55 | 50 | 92 | 121 | 143 | 199 | 169 | 99 | 181 | 112 | 66 | 58 | 1343 | 157 |
| Sakai | 199 | 141 | 144 | 136 | 109 | 173 | 171 | 130 | 223 | 157 | 159 | 191 | 1934 | 290 |
| Kōbe | 59 | 39 | 98 | 136 | 113 | 217 | 174 | 117 | 199 | 116 | 58 | 39 | 1364 | 199 |
| Kyōto | 63 | 61 | 113 | 167 | 144 | 237 | 209 | 146 | 206 | 132 | 77 | 50 | 1604 | 161 |
| Nagoya | 61 | 56 | 133 | 172 | 162 | 235 | 212 | 187 | 262 | 147 | 83 | 51 | 1762 | 240 |
| Tōkyō | 59 | 69 | 113 | 128 | 150 | 170 | 144 | 140 | 213 | 183 | 100 | 53 | 1520 | 172 |
| Niigata | 189 | 123 | 107 | 109 | 90 | 133 | 157 | 133 | 187 | 151 | 191 | 231 | 1799 | 135 |
| Matsumoto | 54 | 38 | 80 | 102 | 101 | 168 | 149 | 131 | 174 | 100 | 45 | 44 | 1187 | 160 |
| Utsunomiya | 41 | 47 | 95 | 130 | 151 | 198 | 247 | 230 | 238 | 119 | 69 | 40 | 1605 | 153 |
| Maebashi | 34 | 28 | 59 | 89 | 108 | 157 | 195 | 219 | 228 | 105 | 40 | 29 | 1290 | 164 |
| Aomori | 127 | 106 | 79 | 65 | 74 | 80 | 135 | 113 | 139 | 111 | 136 | 161 | 1326 | 108 |
| Sapporo | 68 | 56 | 59 | 52 | 61 | 63 | 84 | 95 | 136 | 106 | 96 | 93 | 968 | 124 |
| Nemuro | 30 | 21 | 46 | 70 | 95 | 88 | 86 | 98 | 134 | 93 | 81 | 60 | 901 | 122 |
| Ōdomari | 23 | 14 | 36 | 56 | 90 | 66 | 96 | 78 | 83 | 84 | 56 | 48 | 728 | 51 |
| Jinsen | 15 | 9 | 22 | 56 | 84 | 86 | 193 | 179 | 139 | 39 | 41 | 22 | 884 | 153 |

The amount of precipitation is moderate in Japan. The yearly amount is 1520 mm at Tōkyō, 1799 mm at Niigata, 1949 mm at Nagasaki, 2030 mm at Taihoku, 1326 mm at Aomori and 963 mm at Sapporo. The general tendency is that it diminishes towards the north. The heaviest rain ever experienced at Tōkyō was 172 mm in 24 hours, while 444 mm occurred at Ōita. The rainfall is intense on slopes facing monsoons. Hence the Japan Sea coast is rich in rain in winter, and the Pacific coast in summer.

Table VI.
Number of Rainy Days.

| Name of Place | January | February | March | April | May | June | July | August | September | October | November | December | Year |
|---------------|---------|----------|-------|-------|-----|------|------|--------|-----------|---------|----------|----------|------|
| Tainan | 6 | 5 | 6 | 7 | 10 | 15 | 17 | 19 | 10 | 5 | 4 | 4 | 107 |
| Taihoku | 17 | 17 | 18 | 15 | 15 | 14 | 13 | 10 | 13 | 14 | 15 | 16 | 184 |
| Kagoshima | 13 | 12 | 16 | 15 | 15 | 18 | 16 | 14 | 14 | 11 | 10 | 12 | 168 |
| Nagasaki | 16 | 13 | 15 | 14 | 13 | 17 | 14 | 13 | 13 | 10 | 11 | 15 | 166 |
| Osaka | 9 | 8 | 14 | 13 | 13 | 14 | 14 | 12 | 15 | 10 | 8 | 7 | 137 |
| Matsuyama | 12 | 10 | 15 | 13 | 13 | 14 | 13 | 10 | 14 | 10 | 10 | 11 | 146 |
| Sakai | 26 | 21 | 19 | 14 | 12 | 13 | 14 | 11 | 16 | 15 | 20 | 25 | 207 |
| Kobe | 11 | 9 | 15 | 13 | 12 | 14 | 13 | 11 | 15 | 10 | 9 | 9 | 140 |
| Kyōto | 13 | 12 | 15 | 14 | 14 | 15 | 15 | 12 | 15 | 11 | 11 | 12 | 162 |
| Nagoya | 9 | 8 | 13 | 13 | 12 | 15 | 15 | 12 | 16 | 11 | 10 | 9 | 144 |
| Tōkyō | 8 | 8 | 13 | 14 | 13 | 15 | 15 | 12 | 16 | 14 | 9 | 6 | 145 |
| Niigata | 27 | 23 | 22 | 15 | 14 | 15 | 15 | 12 | 17 | 19 | 22 | 27 | 228 |
| Matsumoto | 12 | 8 | 13 | 12 | 12 | 15 | 17 | 13 | 15 | 12 | 9 | 9 | 147 |
| Utsunomiya | 7 | 7 | 13 | 14 | 15 | 17 | 21 | 18 | 17 | 13 | 9 | 5 | 156 |
| Maebashi | 8 | 5 | 11 | 13 | 14 | 18 | 20 | 18 | 18 | 12 | 7 | 5 | 148 |
| Aomori | 20 | 22 | 21 | 13 | 13 | 13 | 14 | 12 | 16 | 17 | 22 | 27 | 210 |
| Sapporo | 19 | 17 | 17 | 12 | 13 | 13 | 12 | 13 | 16 | 17 | 19 | 20 | 189 |
| Nemuro | 11 | 10 | 12 | 12 | 13 | 15 | 15 | 14 | 15 | 13 | 13 | 12 | 155 |
| Odumari | 12 | 12 | 15 | 12 | 13 | 12 | 14 | 14 | 15 | 15 | 17 | 16 | 167 |
| Yinsen | 8 | 5 | 5 | 7 | 8 | 10 | 14 | 12 | 9 | 7 | 8 | 9 | 103 |

Fig. 3.
Number of Rainy Days.



By a rainy day is meant a day on which not less than 0.1 mm. of any form of precipitation is measured. This amount really is too small to be regarded as a day of rain; hence the number of rainy days thus defined is 10 to 20 per cent greater than that which we understand a rainy day to be in ordinary conversation. The average number of rainy days is 145 in Tōkyō and 228 in Niigata. In Taiwan and in Chōsen it is comparatively less.


Table VII.

Number of Clear and Fair Days.

| Name of Place | January | February | March | April | May | June | July | August | September | October | November | December | Year |
|---------------|---------|----------|-------|-------|-----|------|------|--------|-----------|---------|----------|----------|------|
| Tainan | 24 | 21 | 23 | 23 | 22 | 20 | 24 | 21 | 24 | 25 | 23 | 24 | 272 |
| Taihoku | 12 | 8 | 11 | 12 | 15 | 16 | 23 | 22 | 22 | 17 | 12 | 12 | 182 |
| Kagoshima | 21 | 18 | 18 | 16 | 16 | 10 | 18 | 24 | 19 | 21 | 23 | 25 | 229 |
| Nagasaki | 19 | 17 | 19 | 17 | 17 | 12 | 18 | 23 | 19 | 22 | 23 | 20 | 224 |
| Oita | 24 | 21 | 20 | 18 | 19 | 14 | 18 | 24 | 17 | 21 | 24 | 26 | 245 |
| Sakai | 8 | 8 | 14 | 17 | 18 | 13 | 16 | 22 | 15 | 19 | 17 | 12 | 179 |
| Kōbe | 24 | 22 | 20 | 18 | 17 | 12 | 15 | 20 | 16 | 20 | 24 | 26 | 235 |
| Kyōto | 21 | 18 | 18 | 16 | 17 | 12 | 15 | 20 | 15 | 20 | 22 | 24 | 218 |
| Nagoya | 26 | 24 | 22 | 18 | 18 | 12 | 16 | 21 | 16 | 20 | 24 | 27 | 243 |
| Tokyo | 25 | 21 | 20 | 16 | 16 | 12 | 15 | 20 | 14 | 18 | 22 | 27 | 226 |
| Niigata | 8 | 8 | 13 | 17 | 18 | 14 | 15 | 21 | 15 | 18 | 14 | 10 | 196 |
| Matsumoto | 21 | 21 | 19 | 17 | 17 | 12 | 12 | 17 | 13 | 18 | 23 | 24 | 212 |
| Utsunomiya | 26 | 24 | 21 | 18 | 18 | 12 | 12 | 17 | 14 | 19 | 24 | 28 | 234 |
| Maebashi | 26 | 24 | 21 | 17 | 16 | 10 | 10 | 14 | 11 | 19 | 24 | 28 | 219 |
| Aomori | 11 | 11 | 16 | 20 | 21 | 17 | 16 | 21 | 20 | 22 | 15 | 10 | 200 |
| Sapporo | 18 | 16 | 18 | 19 | 18 | 16 | 15 | 19 | 19 | 22 | 17 | 17 | 213 |
| Nemuro | 23 | 20 | 22 | 17 | 17 | 11 | 11 | 12 | 16 | 22 | 21 | 25 | 217 |
| Ōdomari | 24 | 21 | 21 | 17 | 12 | 13 | 12 | 14 | 20 | 22 | 19 | 19 | 213 |
| Jinsen | 24 | 24 | 22 | 21 | 21 | 18 | 14 | 20 | 22 | 26 | 25 | 27 | 262 |

Fig. 4.

Number of Clear and Fair Days.



The weather is fine for about two thirds of a year in Japan, excepting the Japan Sea coast, where about half a year is cloudy. The number of clear and fair days is 226 in Tōkyō, and 196 in Niigata. In Taiwan it is 272 and in Jinsen 262.

The climate of each district is as follows:—

Taiwan (Formosa). The southern half of Taiwan is in the tropics, hence the climate is tropical. The mean annual temperature at Tainan and at Taihoku are respectively 23.1° C and 21.6° C and

the absolute maximum temperatures are 36,0° C and 37,0° C; thus mean humidity is 80 and 81 per cent. Rain falls abundantly near Kiirun (Keelung); the yearly amount exceeding 5000 mm; but in Taihoku it is 2030 mm. The rainy season of Taiwan is June, July and August, during which thunder showers are a daily phenomenon. There is no snow in Taiwan, nor frost in the southern half of it; in the other half frost happens sometimes but very rarely. Rice crops are raised three times a year. The sugar-cane and the camphor-tree are two useful plants in Taiwan. Typhoons attack the island very frequently during the summer months. Sometimes it destroys all the sugar crops. The maximum wind velocity reached 56,8 m/s at Kōshun in Taiwan during the passage of an intense typhoon.

Kyūshū. Kyūshū is rather warm; the mean annual temperature at *Kagoshima* and at *Nagasaki* is respectively 16,7° C and 15,7° C. Snow and frost occur very seldom. In other climatological respects there is no conspicuous difference between Kyūshū and the Main Island.

Naikai (Inland Sea District). The characteristics of the climate of this region are dryness and a comparatively small amount of rainfall. The latter is 1343 mm and 1115 mm respectively at Matsuyama and Okayama. The number of fine days in this district is greater than that of the surrounding districts, because it is guarded by mountain ranges on all sides and the air currents come across these in a comparatively dry and warm state. The manufacture of common salt on the sea beach is advantageously carried out in this district.

San-in and **Nankai.** San-in is situated to the north of Naikai, and Nankai to the south of this, both separated from Naikai by mountain ranges. San-in has abundance of rain in winter, and Nankai in summer; the excessive rain in these districts corresponds to the deficiency of it in Naikai district. The annual amount of rainfall at *Sakai* in San-in is 1934 mm; at *Okayama* 1115 mm; and at *Kōchi* in Nankai 2727 mm. The number of rainy days in January is 26 in Sakai; 8 in Okayama and Kōchi.

Hokuroku. This district lies on the Japan Sea coast of the central part of the Main Island. The summer climate of this district is nearly the same as that of Kwantō, but the winter climate is too wet being under the influence of the north-westerly monsoon. The depth of snow at some places exceeds two metres. The temperature is not low comparatively speaking, because a warm sea current washes the coast. The mean temperature in January at *Kanazawa* in this district is 2,5° C, while at *Mito* on the Pacific coast and at a little low latitude it is 2,2° C.

Kinai and **Kwansai.** The climate of these districts approaches the average of Shikoku—Chūgoku (Nankai + Naikai + San-in) and Kwantō. The phenological season in this district is a little earlier than that in Kwantō.

Tōkaidō and **Kwantō.** The climate of this region is considered to be the representative of that of Japan, and shall be described later under the heading of "The yearly variation of the climate of Japan."

Central Japan. A little cooler than Kwantō, but with less rainfall.

Ō-u (North-eastern part of the Main Island). This region is much colder than Kwantō in winter, but not appreciably cooler in summer.

Hokkaidō. Though this district is very cold in winter—the January mean temperature at *Sapporo* being -6,2° C and at *Nemuro* -5,0° C—, it becomes warm very rapidly so that the summer air temperature reaches 20° C. The rice plant is cultivated in the western half of this island. The amount of rainfall is less; in Sapporo the annual amount is 968 mm. Hokkaidō is a pleasant place to live in summer, but somewhat uncomfortable place in winter.

Karafuto (Sagalien). This island is very cold and humid. Thunder occurs once or twice a year.

Chōsen (Korea). The temperature is very low in the northern part, but moderate in general. The winter climate is dry and fine. In some years there is no snow nor rain in January or December at all. The eastern coast is always warmer than the western coast.

Yearly Variation of Climate in Japan. As formerly mentioned the empire of Japan stretches from Latitude N 22° to N 52°, and hence the climate of the southern and northern extremities is quite different. In the southern half of Taiwan (Formosa) people never experience frost in the plains, and rice can be raised three times a year, while in Karafuto (Saghalien) and in Chishima (Kuriles) the ground is covered with ice and snow during more than half a year; and even in the maximum of their summer heat, we find braziers or hearths in every house.

As the Japan Islands form the boundary between the Asiatic Continent and the Pacific Ocean, the influence of monsoons and oceanic currents upon the climate is very remarkable. During the winter, the north-westerly monsoon brings moisture from Siberia and Japan Sea, and on reaching Japan, drops it abundantly on the districts bordering on the coast of Japan Sea, which, as a whole, is called the *Ura-Nippon* (Back Japan), hence the winter of this district is very gloomy and humid as formerly stated. The average number of rainy days at Niigata in January amounts to 27 *i.e.* 90 per cent of a month. The monsoon deposits its moisture on the north-westerly side of the central mountain range of Japan, and crosses the system in dry state. Hence on the Pacific coast the weather is very fine during the cold winter months, the number of clear and fair days in Tōkyō in December being 27 *i.e.* 90 per cent. Moreover the warm current called *kuro-shio* (Black Stream) washes the southern coast of Japan as far as *Chōshi*—coasts of Nankai, Tōkaidō and Kwantō—being very warm in winter. For example, the air temperature in January is 5.7° C at *Kōchi* in Nankai, 5.0° C at *Hamamatsu*, and 5.3° C at *Numazu* both in Tōkaidō. In these regions the camellia bears flowers throughout the winter; plum blossoms open in December; the palm flourishes in the open air and the ever green orange-tree is laden with golden fruit in winter. In the central high lands, the weather is very clear in winter, but the air temperature descends conspicuously. It is -2.0° C on the average of January at *Matsumoto*. *Lake Suwa* in this district is covered with thick ice, affording a reliable skating-ground. We can see silver thaw and diamond dust *i.e.* ice-crystals floating in the air, every cold morning at the lakeside.

The weather changes very regularly in the cold months in Japan. This depends on the periodic oscillation of the Siberian high atmospheric pressure. In Korea there is a proverb "*San-kan Shi-on*" that is to say, "three days cold, four days warm." In Japan Proper they say "Wind on every five days and rain on every ten days." The actual periods of this variation is not quite fixed, but fluctuates on an average every six or seven days. In the spring as the air temperature rises, the number of rainy days increases on the Pacific coast but decreases on the Japan Sea coast, which means the change in the monsoons. Rain during this season is known by the special name of "*haru-same*," whose characteristic is that it comes down very calmly and gently as if this silky threads fell soundless from the sky. Thus it scarcely harms the soft young buds but rather nourishes them. It becomes warm by stages with each rainfall. At the beginning of April a weather feature called "*hana-gumori*" (flower cloudy) prevails in the middle part of Japan, under which it is mild, cloudy and rather warm; and *cherry blossoms* begin to display their myriad magnificence. After the cherry we have *peonies*, *wistarias*, *azaleas* and other flowers successively until the end of May. Swallows come back from the south, and begin to build their nests in the interiors or entrances of dwelling-houses. April and May are most enjoyable months in Japan. Fields and woods are bright with fresh verdure and birds such as the nightingale, cuckoo, lark, bunting etc. sing their loveliest songs. In this season farmers prepare their rice-marsh. We see all the gray, withered field of March change into a sort of a lake in May in which thousands of frogs croak. At the beginning or middle of June farmers plant the young shoots of the rice-plant in these shallow marshes. This task is called "*ta-ue*." The wives and daughters of farmers tidily dressed are also in the water planting rice seedlings and singing "*ta-ue-uta*" (Rice planting songs).

The *bai-u* or plum rain begins at the end of June, and it continues about two weeks. In this season the sky is overcast and rain falls continuously. In Fig. 4. we see clearly the diminution of the number of clear and fair days in Tōkyō and Niigata in June. The *bai-u* does not occur in Hokkaidō. It cools June and July temperatures, and consequently the yearly maximum of temperature appears in August. From the middle of July the real Japanese summer begins. The air temperature exceeds 30° C on some days at almost all places in Japan Proper, except in the north-eastern part. During this season the city life is somewhat trying, but in the country it is not so, because since Japan is an island and a mountainous country, the sea, on the one hand, affords gentle breezes (land and sea breezes) and the hills and valleys, on the other, supplies the inhabitants with pleasant winds (mountain and valley winds). *Hakone*, *Nikkō*, *Karuizawa*, *Unzen* etc. are good summer resorts. During this season thunder of thermic origin occurs very frequently. Sometimes it is intense, but in general very refreshingly and clearing. The short night of summer is not sufficient to cool the heated earth and air in the day time. The burning sun projects its beams fiercely increasing from its rise. At about noon, we see many great towers of *cumulo-nimbus* above the mountains. They increase their heights; their heads attain the cirrus height, flattened at the top; they begin

to move from their original positions, and soon we hear the distant peals of thunder. After one hour or so, the sky is overcast, the thunder becomes louder and louder and at last the shower begins preceded by cool squally winds. This rain disposes of the heat and clears the dust away. All creatures are refreshed. It passes away after 20 or 30 minutes and the deep blue sky reappears. Thus thunder in Japan in summer is accompanied rather rarely by storm but most frequently by showers. From the end of August until the beginning of September there are frequent visits of Typhoons to Japan. September is the period of transition of the summer and winter monsoons. From the middle of September cool winds begin to blow. People gradually forget to use the fan. The number of rainy days increases a little during this month. October is another pleasant month in the Japan Proper. Fine weather continues till November on the Pacific coast but it begins to rain and snow on the Japan Sea coast. From the end of October until November farmers harvest their rice. The autumn tints in forest are beautiful. In villages we see red *kaki*-fruits still on the branches. *Chrysanthemums* are also in full bloom. We now feel the approach of winter by the cackles of emigrating wild-geese which come from the northern countries flying over the sea. Thus the yearly variation of the climate of Japan is about to repeated.

Chapter V

THE ORIGIN OF CLIMATE THERAPEUTICS IN JAPAN.

The origin of climate therapeutics in Japan is unknown and naturally is not recorded in history. But according to one story, Kōbō-Daishi, a famous and much revered Buddhist priest, opened for worship the eighty-eight temples in the island of Shikoku and the thirty-three temples in the western provinces, and appealing to the religious sentiments of the people, he advocated a circular pilgrimage to these temples in spring and early summer. This is now supposed to be a prominent example of combining climate therapeutics with religious propaganda. We cannot but esteem very highly the originality and far-sightedness of this sage priest, who is generally believed to have possessed a profound medical knowledge. The genuineness of this statement seems fully justified by the fact that these temples founded by him are not located on level land, too easy of access, but in mountainous districts where the air is pure and free from corruption.

Principal Forest Health Resorts. The best forest health resorts in Japan are in the Shinano Province, the Nasu Mountains, the Kiso Valley, and some parts of Hokkaidō. If some suitable grounds be chosen and properly taken care of, all of these places will doubtless present us with ideal climatic sanatoria.

see Plate 37

see Plate 28,
Plates 12-14

THE ORIGIN OF BALNEO-THERAPY IN JAPAN.

In looking through Japanese history we find in the record of the Age of the Gods, that people bathed in hot springs to recuperate. Following this history later tells us that the two Emperors, Kimmei and Kōtoku, resorted to hot spring baths for medical purposes. The *kojiki* ("Records of Ancient Events") gives evidence that the former Emperor proceeded to *Arima* Hot Spring in Settsu Province, and *Dōgo* Hot Spring in Iyo, to recuperate there. Gotō Konzan, a physician, is said to have utilized hot springs for medical treatment. Yamamura Tsūan, another physician, walked the round of various hot springs scattered in distant provinces, in order to test their composition and medical efficacy, and published the fact that *Kinosaki* in Tajima Province, and *Kusatsu* in Kozuke, etc., were of the foremost importance from a medical point of view.

see Plate 58
see Plates 63
and 64

see Plate 59
see Plate 60

THE ORIGIN OF SEA BATHING IN JAPAN.

Tracing back this subject we come across the frequent mention of "*shio-tōji*" (cure in the brine) in old books. The custom of sea bathing, as well as bathing in hot springs, is thus traceable back to the Age of the Gods. These two methods of bathing may be said to be equally of extremely ancient origin, but in mediæval ages hot spring baths alone were popular, while sea bathing dwindled into insignificance. Although of course no definite data are obtainable as regards the vicissitudes of

this form of ablution, it is not difficult to infer from the seagirt position of this country that the custom undoubtedly continued to be practised in different places.

Ancient Evidence of Method of Basking in the Sun. In the "*Owari Meisho-zue*" (Pictorial Records of Interesting Places in Owari Province) there is a picture showing sea bathing at Ōno, some five miles from Atsuta (near Nagoya). From this origin we gather sea bathing there dates back to some 700 years ago, or to be more accurate, 765 years ago people used to dip themselves in sea water until they felt cold, and then lie down on sun-burnt stones for warmth. This is nothing but a curious, ancient record of basking in the sun.

New Departure in Sea Bathing since the Restoration. In 1881 the then Governor Rempei Kunisada happened to visit this place during his prefectural tour. On learning the history of sea bathing there, he conferred with Mr. Shimpei Gotō, then the Chief of the Aichi Hospital, who was emphatically teaching the medical value of sea bathing, and had made a certain Shōzō Nijō set up a rest hut for changing the clothes. This erection of a bathing hut marked a new departure in the methods of sea bathing after the Meiji Restoration and gradually led to its present popularity.

It is not rare to find old instances of bathing in waterfalls as an invigorating tonic during the heat of summer. In some such waterfalls as in Mt. Takao, Musashi Province, Mt. Ō-iwa, Etchū Province, etc., popularly believed to be efficacious for curing mental derangement, there were constant bathers not only in summer but also in the colder seasons.

The Opening of Sea Bathing at Ōiso. As stated above, though evidence of sea and cool water bathing dates back to mythical ages, but it is only within the past thirty years and long after the introduction of this medical science in the west that the people here have realized or have been awakened to the therapeutic advantages of sea bathing, etc. In 1886 Dr. Jun Matsumoto for the first time drew the public attention to Ōiso as an excellent sea bathing place and thus brought about the local popularity of this method of therapeutics. Since then there have been established seaside sanatoria at Kamakura and Shichiri-ga-hama in Sagami Province, Suma in Harima, etc.

Sea Water Bath-Houses. Sea water bath-houses are still very rare in Japan and the Chū-eikwan outside the port of Yokohama in Kanagawa Prefecture is probably the only one of this kind now in existence.

Natural Sea Bathing Place. On the other hand, there is an endless number of seaside places naturally fit for bathing, the most important of which are dotted along the Pacific coasts of Tōkaidō, Nankaidō, and San-yōdō districts and also the islands of Shikoku and Kyūshū. Most of these places are easily accessible, endowed with a mild climate, and form excellent health resorts. The geographical nature of our islands makes the Pacific side extremely rich in suitable watering places, but no few are found also on the Japan Sea side.

Sand Baths. Sand baths have been and still are very popular on the sandy beaches in the neighbourhood of Beppu and Hamawaki Hot Springs, both of these places are naturally suitable for the purpose and are said to cure hæmorrhoids.

Hot Sand Baths. Hot sand baths have been practiced among the people for a considerable period. One such bath is located on the beach of Sōyu-no-hama, Ibusuki-gōri, Satsuma Province. Shōshi Tamiya, a physician, speaks of this place, as follows:—

"All patients lie down, naked or thinly clad, on the hot sand and the by-standers assist in covering the bathers bodies with the sand leaving only their faces exposed and using for the purpose large wooden spades. Within a short period the bodies are heated by degrees, the perspiration oozes out from every pore of the skin carrying away ill humours and impurities, thus lightening, toning-up, and invigorating the body. During the operation the patients feel as if they were in paradise, and various forms of illness are at once thrown off. Sand baths, as compared with ordinary hot spring baths, inspire a feeling in patients of mild comfort that is often preferable."

Vapour Baths. Vapour baths have existed in Japan for some centuries and were known as "*Yumushi-no-hō*" (the method of steaming). An old stone bath in Miyajima Island in Aki Province, was used as a kind of vapour bath and at one time patronized and held in high favour.

Sea Weed Baths. There is in the city of Takamatsu in Sanuki Province a bath known "*mushiburo*" (steam bath), which has existed for a considerable number of years. This is an instance of a weed bath, the sea weeds were collected and warmed, and people bathed in them to heal their illness, especially nephritis being claimed to be curable by perspiration in this heated substance. This peculiar method may also be classed as a variety or type of vapour bath.

Mineral Mud Baths. The practice of mineral mud baths is not yet generally known in Japan. In 1909 when Dr. Ishizu, an expert of the Tōkyō Hygienic Laboratory, visited the Beppu and Kamegawa Hot Springs to test their radio-activity, he had a bath prepared with sinter collected from "*Chino-ike Jigoku*" (Hell of Blood Pond). This name is given owing to the local red colour produced by a large proportion of ferric oxide. This was the first experiment with a sinter bath in Japan, but it has been followed by the inhabitants there in recognizing its benefits and since then has become quite popular.

Chapter VI

GENERAL REMARKS ABOUT HOT SPRINGS IN JAPAN.

The Situation of Hot Springs. Although Japan has numerous volcanoes, hot springs are mostly but not always located in volcanic districts. Hot springs gush out almost everywhere in this country; on the side of high mountains, on plains, in valleys and ravines, in rivers and lakes, and also on the sea shore and from the sea bottom. The character of the surroundings has, on one hand, much to do with their scenery and climate, but, on the other, with the difficulty or ease of access, supplies of variety of food and other daily necessities, at the resorts, water, drainage, purity of atmosphere, etc., on all of which their hygienic conditions greatly depend. The diversity of geographical features makes the range of choice for visitors very wide, and enables each to select for patronage any place they fancy most suitable for their individual requirements. Thus the division of Japanese hot springs, according to their geographical and scenic characteristics, is not only of special instructive interest to intending visitors but at the same time by pointing out the local hygienic conditions, it facilitates the simplicity and ease of forming a personal decision that is of mutual benefit or satisfaction.

1. Hot springs located in high mountains, and characterized by the so-called upland climate:—*Renge, Bandai, Tateyama*, etc.

2. Hot springs located on the sides of mountains and commanding a wide view:—*Ikao, Akakura, Nasu*, etc.

3. Hot springs located in thick forests, valleys, ravines, etc.:—*Shiobara* and other places.

4. Hot springs located on sea shores and combining the advantages of seaside therapeutics:—*Obama, Sedomo-Kanayama, Atami*, etc.

5. Hot springs close to the sea and thus combining facilities both forms of bathing:—*Beppu, Sedomo-Kanayama, Ibusuki Beach*, etc.

6. Hot springs located near lakes and having picturesque scenery:—*Yumoto* near Nikkō, *Ashi-no-yu, Suwa*, etc.

If we classify hot springs according to their altitude, the highest are the *Shibu* Hot Springs near Lake Suwa (6,950 ft. above sea-level), *Nakabusa* of Japan Alps (5,300 ft.), *Manza* (5,180 ft.), *Yumoto* near Nikkō (5,088 ft.), *Sandogoya* in Nasu (5,000 ft.), *Kamikōchi* (4,720 ft.), *Kusatsu* (4,500 ft.), *Goshiki* (3,000 ft.), etc. All these hot spring resorts have vigorous upland climate.

The Geology of the Spots where Hot Springs gush out. Most of the hot springs issue from andesite, and next in abundance are those from Tertiary deposits, but not a few springs issue from granite.

The Quantity of Flow. The flow of hot spring waters is various in character and quantity. The most active of springs throw up their jets to a height of from 60 to 100 ft. The "*O-yu*" of Atami is a famous intermittent spring and once sent up its perpendicular jets with energetic power, so much so that stone walls were built to break its force in order and curtail to avoid possible danger and it is thus deprived of the former grand appearance its unfettered play once inspired. Its activity when in full play is amazing still, and a never failing source of attraction. Another famous intermittent spring is that of *Onikōbe* in Rikuzen Province, which reaches a height of 20 ft. One of the hot springs in *Obama*, called "*Funtō-yu*," is also of intermittent nature. At *Senami* in Echigo Province boiling-hot water is being thrown up more than 90 ft. high, which may be said to form a wonderful sight. Some springs discharge less sensationally, abundant in their quantity, they simply form streams of hot water in the end, as is the case at *Noboribetsu, Beppu, Kamegawa, Onikōbe, Kirishima*, etc. At some places such streams make waterfalls which are used to turn mills, e.g., *Shigaku* Hot Springs in Iwami Province. They often accumulate in forming ponds of warm water

as at *Beppu* and *Noboribetsu*. As regards the origins of hot springs, they are again of diverse kinds. Some gush out of caves (*Ogawa* and *Futami* Hot Springs in Etchū Province, *Shigaku* Hot Springs, *Shin-Goshiki* Hot Springs in Uzen Province, "*Hashiri-yu*" Spring of Izusan, near Atami), in the middle of a river bed ("*Dokko-no-yu*" Spring of Shuzenji), from the bottom of a lake (*Tōgō* Hot Springs in Hōki Province), etc. Such extraordinary circumstances as the above examples are not rare. We may point out *Beppu* and *Itō* as the most prominent examples of these places where hot spring veins are numerous and abundant and their sources may easily be found by boring almost at any spot. The baths of the former hot springs number more than one thousand and those of the latter more than three hundred.

The Temperature of Hot Springs. The temperature of hot springs in Japan ranges very widely and numerous springs (*Naruko*, *O-yu* in Atami, *Senami*, etc.) are above one hundred degrees Centigrade, while such hot springs as *Noboribetsu*, *Nakabusa*, *Yumura*, *Owakidani* in Hakone, *Obama*, *Beppu*, *Kamegawa*, etc., show above ninety degrees Centigrade. Other hot springs of below eighty degrees are too numerous to be referred to in this brief work.

The Composition of Hot Springs. The composition of hot springs is of a complete variety, and it is no exaggeration to state that Japanese hot springs comprise all possible kinds of hot springs known to the world. As might be expected simple and salt springs are the most numerous, but apart from these, the greatest bulk of the remainder are sulphur springs, these being followed in number by the alkaline carbondioxated springs. Most springs of peculiar properties are those containing free mineral acids (*Kusatsu*, *Nasu*, *Noboribetsu*, *Kirishima*, etc.) and a large quantity of alumina and iron (*Kusatsu*, *Kannawa*, etc.), while not a few springs contain a small proportion of boric acid, iodine, bromine, lithium, manganese, etc.

The mineral springs of Japan, either hot or cold, may be chemically classified, as follows:—

Table VIII.
Classification of Mineral Springs in Japan.

| Classification † | Central Japan* | Eastern Japan* | Western Japan* | Northern Japan* | Hokkaidō | Kyūshū | Taiwan | Chōsen | Total |
|-----------------------------------|----------------|----------------|----------------|-----------------|----------|--------|--------|--------|-------|
| Simple cold springs | 64 | 50 | 10 | 10 | 1 | 20 | 0 | 6 | 161 |
| Simple thermals | 41 | 60 | 22 | 28 | 3 | 70 | 2 | 10 | 237 |
| Simple carbondioxated springs . . | 4 | 7 | 5 | 1 | 1 | 3 | 2 | 0 | 23 |
| Earthy carbondioxated springs . . | 1 | 5 | 0 | 6 | 1 | 3 | 2 | 0 | 18 |
| Alkaline carbondioxated springs . | 39 | 38 | 8 | 9 | 20 | 35 | 4 | 0 | 153 |
| Common salt springs. | 55 | 60 | 9 | 31 | 5 | 19 | 3 | 5 | 187 |
| Bitter springs | 6 | 27 | 4 | 21 | 4 | 16 | 1 | 0 | 79 |
| Iron springs | 12 | 14 | 0 | 3 | 1 | 2 | 1 | 3 | 36 |
| Sulphur springs | 23 | 26 | 10 | 36 | 14 | 18 | 6 | 9 | 142 |
| Acid hydrogen sulphide springs . | 0 | 6 | 0 | 4 | 0 | 1 | 0 | 0 | 11 |
| Acid vitriol springs | 2 | 3 | 0 | 0 | 1 | 0 | 1 | 0 | 7 |
| Alum vitriol springs | 2 | 5 | 0 | 0 | 0 | 1 | 3 | 0 | 11 |
| Not yet examined | 10 | 42 | 7 | 14 | 0 | 17 | 2 | 35 | 136 |
| Total | 268 | 244 | 75 | 103 | 51 | 205 | 27 | 68 | 1201 |

Bathing Equipments. Japan has the largest number of hot springs but they leave still much to be desired as regards bathing equipments. Only very few of them are decently equipped with private baths, so that foreign visitors may take baths at ease. This lack of privacy is due to the custom of common bathing to which Japanese have been accustomed for many centuries, distinctions being made only for both sexes but not for individuals. Also many of the most efficacious hot springs are found in remote districts and are chiefly patronized by the neighbouring inhabitants. At present in Japan, such poorer classes as farmers living among isolated mountain districts take

* See Map of Imperial Government Railways.

† See Chapter XIII.

advantage of the medical treatment with their local hot springs, more often than the better classes from towns and cities. As it is difficult for such poor people to find good doctoring, they economically patronize the natural methods of therapeutics so conveniently at hand. Successive generations of these people have been taught that hot springs are good for almost all diseases, and without any consideration of individual cases or ailments, in simple faith, they invariably go for prolonged stays to hot springs in the neighbourhood as the sole means for recuperation. The people of the better classes consider hot springs simple holiday resorts, and, to our great regret, do not often realize their importance for medical purposes. In recent years physical therapy has been more seriously considered among well-informed circles, some of whom now are trying to experiment on the therapeutic efficacy of such natural treatments as balneo-therapy, sun basking, air baths, etc. Under such transitional circumstances, we have to be contented to await development with the present lack of proper hygienic equipments at most hot spring resorts.

Baths. For bathing accommodation houses are built near the source of springs and water is conducted to baths therein by bamboo, wooden or iron conduit-pipes, etc., the water being thus kept always fresh. At some places baths are made directly over the source. The most comfortable are natural baths hollowed out of the rock formation with hot water flowing in from the rock crevices. The housing accommodation is frequently very simple, but from the modern scientific view-point of the emanation therapy, it should be found to be ideal. Ordinary baths are in most cases made of wood but also not seldom of marble and other natural rocks. Generally clear and pure water is conducted from hot springs and offered for drinking.

see Plates 6
and 42

see Plate 7

The Use of Mineral Springs. Besides the ordinary use of mineral springs for bathing the body, there are some places with combined vapour baths as well, these being prepared with the natural steam from hot springs (*Beppu, Shima, Nakabusa, Naruko*, etc.). A method of bathing peculiar to Japan is that of utilizing hot waterfalls, artificially made where the flow of hot spring water is abundant. These waterfalls are used for the so-called hot spring massage, bathers standing under them so that the falling water may strike their head, shoulder, etc. In the hot springs of *Noboribetsu, Kirishima, Beppu, Shiobara, Kusatsu, Takeo, Nakabusa, Shima, Naruko*, etc., there are these waterfalls of different forms, breadths, quantities, heights, temperatures, etc., to suit the individual requirements of bathers. Indeed this method is delightful beyond description and may well be said to be an ideal massage method. It is a custom of ancient origin in this country to drink mineral spring waters, so that long experience has taught the people the characteristic efficacy of various springs. At many hot spring resorts, the sediments deposited in reservoirs, conduit-pipes and other parts, are collected and sold as "*yu-no-hana*" (flower of spring), which are again used for preparing household baths. Large quantities of these sediments are produced at some places, e.g., *Kusatsu, Ikao, Beppu, Hakone, Shiobara, Arima*, etc. These substances contain very often sulphur, iron oxide and alumina. Concentrated common salt springs are evaporated and their residue is sold being similar and replacing the salt produced at *Karlsbad* (*Isobe* mineral spring salt). Iron springs are utilized for dyeing at *Arima, Ikao, Beppu*, etc., and textiles thus dyed are bought by visitors as mementoes. At various places small quantities of mineral water are used in the making of wafers, but the only kind, more or less effective, are those of *Isobe*, while others are in value simply nominal or of minor consequence.

see Plate 37

see Plates 7
and 67

see Plate 8,
fig. 1

At *Beppu* and *Kannawa* we observe that vegetables, cereals, and other daily food-stuffs, are cooked with the heat of steam rising from the ground. There are ovens at the road side in front of every house, on which pans and kettles are placed when necessary, the cooking being done within a few minutes. This is certainly a case of fully utilizing nature's resources and a curious sight. At *Yumura* in Tajima Province, the source of the hot springs forms a small pond, in which villagers dip and boil rice, wheat, beans, fresh vegetables, eggs, bean curds, etc. In that village only very few houses are provided with charcoal heated ovens. There is also a laundry at a spot on the river, a little below its junction with the hot water. The practice of cooking half-boiled eggs is witnessed at different hot springs, and the temperature of the springs at *Yamanaka* and *Yamashiro* in Kaga Province quite answers this purpose. When eggs are dipped in these springs for six or seven minutes, the yolk congeals and the white is boiled to a palatable nicety. At some places hot-houses are set up over hot springs (*Owani* Hot Springs), and at others the heat of springs is utilized to evaporate sea water (*Asamushi* Hot Springs). In the districts with subterranean hot spring water circulation we find the cultivation of vegetables is much quickened. (*Utsunomiya* Hot Springs in Hoki Province)

see Plate 5,
figs. 2 and 3

see Plate 8,
fig. 4

Although mineral spring waters are drunk at many places, comparatively few are bottled for sale or export to foreign countries. The principal ones exploited are the carbonated springs of *Arima*, *Takarazuka*, *Nunobiki*, *Kasagi*, *Isobe*, *Iizaka*, *Misumi*, etc. As Japan has mineral springs of all known kinds of chemical composition, it is not difficult to find good mineral waters that are equal, or sometimes superior, to the world-famous *Apollinaris*, *Vichy*, *Ems*, *Fachingen*, *Selters*, *Karlsbad*, etc. We greatly regret that the complete analyses of Japanese mineral springs have not yet been made known to the world and no enterprisers have undertaken their bottling on a large scale.

Public and Private Baths. There are two kinds of baths, viz., public and private. At the hot spring resorts where the gush is not abundant, it is not easy for every inn to have its own baths supplied from the spring. This difficulty is experienced particularly at those places that are much below the level of the origin of hot springs. The hot water may be conducted to make public baths but may not be of such a strong flow as to supply private baths of each inn. Seeing the better class people prefer only hot spring resorts having private baths available, naturally such resorts alone prosper. No charge is generally made for common baths which are mostly open at all hours to the public. The cure tax or the like as in Europe and America is not collected at Japanese hot springs.

As Japanese enjoy and make a habit of frequent bathing, not only health seekers and holiday makers but also ordinary travellers avoid towns for staying overnight, if hot spring places are at all conveniently near and such bathing to be had. This is well exemplified by the case of Beppu Hot Springs and the Town of Oita.

As a rule, hot spring resorts have ample water supply in the neighbourhood and there is no necessity of water-works. Though no special arrangements may be made for drainage, it is generally satisfactory. Such being the case, it is very rare indeed any epidemic touches hot spring resorts, in spite of the absence of these artificial arrangements. Some hot spring resorts are provided with small parks or gardens for short walks and also with amusement halls. Generally there are such historic monuments and remains as temples, shrines, old castles, etc., which are numerous all over this country, and these add greatly to the interest of walks in the neighbourhood and visitors are easily beguiled of any tedium of long stays.

The Means of Access. As hot springs are generally situated far away from large cities, the means of access leave still much to be desired. But the more famous ones have now the facility

of light railways, and electric and horse tramways, and at others omnibuses and *jinrikisha* are available. Across steep passes, where it is difficult to walk over, people can travel by "*kago*" (a kind of sedan chair) or on horse-back. Even if all these means fail, one generally finds full compensation in any personal effort walking through beautiful scenery on the way to the desired destination.

As for provisions, one may find some difficulty at most hot springs in securing luxurious food the daily fare being usually of a simple kind. When there are railway communications, daily one

can get fresh fish from the sea, while again often this is replaced by fish caught in rivers and lakes in the neighbourhood. Trout abounds in mountain lakes and constitute chief dishes at hot spring resorts in their vicinity. Fresh vegetables, chickens, and eggs are obtainable everywhere, but milk is rare except in the environs of cities and towns.

Inns. Hot spring inns are generally built on a thoughtfully selected site commanding a beautiful view, this at the same time being in close proximity to the origin of the springs, though sometimes there are separate well fit up buildings providing private baths for guests' use. We can count but a limited number of hotels, at hot spring resorts that accommodate foreigners adequately, but these are all highly recommendable and well managed, homely and comfortable in their appointments, giving a good table and attendance at a most reasonable rate. But foreign tourists with some little knowledge of Japanese manners and customs will find genuine native inns rather interesting, and if ready to put up with slight incidental inconveniences, this personal adaption affords the visitor such



jinrikisha

a wider latitude not alone in the choice of route but in a closer and more intimate enjoyment of the best the country has to offer.

The Methods of Bathing. As previously stated, the majority of Japanese hot springs are chiefly patronized by farmers and their families from the adjacent districts. They come either to recuperate (known as "*tōji*" or hot spring cure) or for a change and rest to be simply free from the cares and drudgery of daily tasks during the spring, when at short periods there is not much work on their farms. They take baths quite frequently and for great lengths of time, so much so that it is not seldom to find them bathing more than ten times a day and passing the whole night in baths, even in sleep. They sometimes bathe in rather hot water for one or two hours. Of course the length of time for bathing depends much upon the temperature of the spring. This long bathing is due to the possibly erroneous belief that the longer they bathe, the more effective it should be. But curious to say, this method, generally considered unhealthy, is not hurtful to them, probably on account of its being a habit. Even ignorant farmers strictly observe the rules for bathing set down for every spring and do not fail in propriety and inborn sense of politeness and good manners.

The Therapeutic Efficacy of Hot Springs. These farmers know from experience the therapeutic efficacy of hot springs and make full use of their knowledge. They have an implicit confidential faith in the tradition of each hot spring and choose the particular spring adapted, to their needs, with the result they are invariably successful in healing their weaknesses. If the desired cure is not entirely complete with one year's baths they know a stubborn case needs patience and thus they make a practice of coming for a certain period every year. The largest percentage of these patients are sufferers from wounds, burns, rheumatism, affections of stomach, skin diseases, syphilis, disorders of brain, uterus diseases, neuralgia, etc. There is a curious superstition quite popular among the lower class people that one can be exempted from diseases during that year, if one take spring baths on a certain day in the middle of June. On that particular day all hot springs are crowded with people of both sexes and all ages, coming from far and near. All bath-tubs are filled with bathers from the early dawn and the confusion defies description but possibly can be left to the readers' imagination.

The most curious of all methods of bathing is the so-called "time-bath" at *Kusatsu*, which is worth while describing in some details. Four baths daily at stated definite hours are the rule. Before entering the water, at the word of command of the bath-master all bathers take deal boards and stir up the water in the tub with a concerted rhythm of motion to cool it. This cooling operation is an amusingly novel sight to visitors. Then bathers kneel in rows along bath beams and each one pours one or two hundred dippers full of the hot water over their heads to prevent vertigo on entering the water, and it is said to clear the head and freshen the brain. Now by the direction of the bath-master they gradually enter the water. The time of a bath is not allowed to exceed three minutes, the temperature being about 52° C. The second row of bathers stand along the side of the tub waiting for the first bathers to leave the water. There are often a third, fourth, or even fifth row of bathers, and the more the number the later they enter the water, the last batch getting a less heat. The bath-master gives the following signals one after another: "two minutes remain"; "only one minute"; "put up with it a little longer"; "perseverance is necessary"; and "now then, get out." At the final order, who would remain behind! instantly all bathers jump out of the water. Bathers must submit to a sort of semi-military discipline and be obedient to the orders of the bath-master. This is certainly a curiosity of bathing procedure. Another oddity resembling this "time-bath" is the "numbering bath" of *Nasu* Hot Springs. Bathers count one thousand on their fingers, remaining motionless in the water during that time. On finishing one thousand they leave the water with a gasp. These methods are practiced to compel people to bathe for a certain length of time in water of a most uncomfortably high temperature. At *Yunogō* Hot Springs in Mimasaka Province, the temperature of the springs is equal to that of the human body and bathers are accustomed to bathings for great lengths of time. Even short baths are for two or three hours, during which time bathers remain in the water. Gradually they prolong the length of time and not a few bathe overnight. They say it is most pleasant in the winter to sleep in warm water. After all, the various seemingly strange practices and queer methods of bathing at remote hot springs have but been taught by the experience of many years, and it is little wonder that these are sometimes found to be quite in conformity with the scientific theories of the latest balneo-therapy.

The Chance Causes of Spring Discovery. The discovery of Japanese hot springs is very often traceable back to the mythological Age of the Gods, especially is this so with famous ones.

According to history and traditions, the instances of accidental discovery are numerous; some springs were found on seeing animals, especially deer, herons, etc., drink or heal at them; others were accidentally discovered by huntsmen, wood-cutters, itinerant priests, etc., when they were climbing mountains, etc. Hence we have such names as "spring of deer," "spring of herons," etc., which are very frequently met with. The locations of some hot springs are said to have been communicated to the discoverers in dreams. Several hot springs have been not seldom discovered on account of snow thawing too early or grasses growing too fast in the early spring on the grounds over them.

Chapter VII

JAPANESE MINERAL SPRINGS AND THEIR THERAPEUTICAL FACTORS.

Since the recent discovery of radium, the scientific and medical investigations as regards radio-activity have revealed the fact, that the radio-activity of mineral springs is one of the most effective therapeutic factors. Thus the hygienic authorities and hot spring managers began to pay serious attention to the matter. Last year the Department for Home Affairs ordered the Hygienic Laboratories to investigate the radio-activity of the principal mineral springs in Japan. This investigation is entirely in the charge of Dr. R. Ishizu, expert of the Tōkyō Hygienic Laboratory, and Mr. Y. Kinugasa and Mr. H. Kibezaki, both being assistant experts of the Hygienic Laboratory. Up to the date of writing more than 150 mineral springs of more than 1,000 origins have been experimented upon. The above do not cover the whole number of mineral springs now known in existence in the empire and even some famous hot springs are owing to the brief notice given left unrecorded. But the results of these experiments are sufficient to show the general nature and value of the radio-activity of Japanese mineral springs. This chapter, therefore, is devoted to the report thereof; and together with the table of analyses, we trust, will prove of some service for balneologists, physicians, patients, etc. in general.

According to the experiments so far made, the quantity of radium emanation in any mineral spring is closely connected with the geology, rather than altitude of the place, chemical composition and temperature of the water, etc.

The bulk of Japanese hot springs are located among volcanic rocks and these have very small quantities of radium emanation, e.g., *Noboribetsu*, *Nasu*, *Bandai*, *Kusatsu*, *Hakone*, *Beppu*, *Aso*, *Kirishima*, *Sakurajima*, etc.; all these are sulphur springs. Mineral springs with large quantities of radium emanation gush out only from granite formations, just as radio-active rocks are found exclusively in granite regions. *Masutomi* in Kai Province, *Naegi* and environs in Mino Province, *Misasa* in Hōki Province, *Murasugi* and *Tochiomata* in Echigo Province, etc., are prominent examples. *Masutomi* and *Takayama*, are cold mineral springs. The former is at the foot of Mt. Kimbu-zan to the north-west of Mitake; and there are found in the neighbourhood, scheelite, apatite, tourmaline, sulphide ores, etc. The latter named place is a granite region along the River Kiso, and affords tin sand, together with wolframite and fergusonite, besides monazite and naegite which both contain thorium. All hot springs of strong radio-activity are located in granite regions, but we cannot say that hot springs gushing out of granite formations are usually strong in radio-activity.

If we compare these mineral springs with those of the strongest radio-activity in Europe, *Masutomi* is inferior only to *Joachimsthal* and *Brambach* (both in Austria), but superior to *Gastein*, *Landeck*, *Baden-Baden*, etc., holding undoubtedly the third position in the world. *Misasa* is only next to *Ischia* (Italy) and almost equal to the most radio-active hot spring in *Gastein*, being superior to all other hot springs of the world in radio-activity.

As stated above, not only may Japan well be proud of the fact that it holds a prominent position in the world as regards the radio-activity of mineral springs, but also it holds a great healing blessing to our countrymen. These mineral springs cannot be said to lie in remote places, but their real value has not yet been fully realized by the general public. As yet opportunity is not still ripe for paving the way to a fuller access, protecting the origins of the springs, and equipping them properly so as to be used for therapeutic ends. Nothing is more regrettable than that we have to leave their management, in spite of their rich qualities, to the small means and limited exploitation of local enterprisers.

see Plates 35,
36 and 37

see Plates 11,
20, 22 and 23


ENVIRONS OF ENA MINERAL SPRINGS (strong radio-active) AND NAEGI



MASUTOMI MINERAL SPRINGS (of the strongest radio-activity in Japan) AND ITS ENVIRONS



EASTERN PART OF SAN-IN DISTRICT with most radio-active hot springs of Japan



From Left to Right:

| Hot spring | Geology | Radio-activity |
|------------|----------|----------------|
| Sekigane | Granite | 33.47 M. u. |
| Mosaka | " | 142.14 " |
| Asōu | Andesite | 4.33 " |

| Hot spring | Geology | Radio-activity |
|-----------------------|----------|----------------|
| Matsuzaki | Andesite | 5.87 M. u. |
| Kachimui and Hamamura | Granite | 8.57 " |
| Yoshioka | Tertiary | 4.36 " |
| Yoshikata | Alluvium | 5.83 " |

| Hot spring | Geology | Radio-activity |
|-------------------|----------------|----------------|
| Yudani (cold sp.) | Tertiary | 4.24 M. u. |
| Iwai | Older Tertiary | 4.17 " |
| Yumura | Granite | 1.76 " |
| Kinosaki | Tertiary | 8.41 " |

Fig. 1



Hot springs, issued from the gash of a stream at **Tamatsukuri**, near *Hakone*.
Can be used in various parts of the river bed.
Pl. I, p. 36, 37, 38, 39.

Fig. 3



Yamanaka or *Yamanaka*
Spring of *Yamanaka*, the
patron of traveller,
children, and women in
delicate condition.


Kusatsu Spa, rising
from fissures in volcanic
debris on a mountain
slope.

Yamanaka stands on the
ground floor, well
temple dedicated to the
dynasty. Pl. I, p. 36.

Fig. 2

Hot springs of *Yamanaka*, *Hakone*, *Hakone*
Pl. I, p. 36, 37, 38, 39.

Fig. 4



Shigaku Hot Springs,

gushing out of rocks, the gashway for the hot water
flows of hot water.
Pl. I, p. 36, 37, 38, 39.

Fig. 5



Hot springs of *Yamanaka*, *Hakone*, *Hakone*


Pl. I, p. 36, 37, 38, 39.

Fig. 1



A full view of the outflow of
Shigaku Hot Springs, Iwami Province
(Pl. I, p. 25; Pl. III, p. 39)

Fig. 3




Yumura Hot Springs, Lamo province
Here a small primitive bath is built direct upon the
source of springs, water issuing from
cracks in rocks.
(Pl. I, p. 27; Pl. III, p. 38)

Fig. 2



Dr. Ishizu and his assistant journeying on horse back to
Kusatsu Spa. (Pl. I, p. 28; Pl. III, p. 20)

Fig. 5



**Akakura Mountain
Spa**, 2,500 ft. above sea
level, situated on the
middle slope of Mt.
Myōō. It is a pleasant
summer resort with an
extensive view.
(Pl. I, p. 25; Pl. III, p. 40)

Dr. Ishizu trying an open air bath at the source of **Tsubame Hot
Springs**, near Akakura. This bath is believed to be more efficacious
than ordinary baths; water is conducted into it by bamboo pipes over
a distance of about 1.3 m. (Pl. I, p. 27; Pl. III, p. 44)

Fig. 1



Fig. 3



Fig. 4




Fig. 2



Hot Waterfalls for Massage, *Ph. I*, p. 27

1. *Yunokawa* at **Noboribetsu Spa**, paved walkway of hot waterfalls, the temperature varies from 13 to 40°C. *Ph. 1*, p. 27

2. Hot Water Bathing in *Yunokawa* at **Furukawa Hot Springs**, *Ph. 2*, p. 27

3. Hot Water Bathing in *Yunokawa* at **Furukawa Hot Springs**, *Ph. 3*, p. 27

4. Women bathing in *Yunokawa* at **Noboribetsu Spa**, *Ph. 4*, p. 27

Fig. 1



Yumura Hot Springs, or "Hot Water Field"

at **Kusatsu.**

Boiling water is conducted into wide, long, and shallow reservoirs from which sulphur deposits are collected.

(Pt. I, p. 27; Pt. III, p. 26)

Fig. 2




Villagers boiling rice and vegetables in a hot water pool at

Yumura Spa, Tajima Province.

(Pt. I, p. 27; Pt. III, p. 55)

Fig. 3




Cooking food in **Kan-**

nawa Spa, near Bippu.

Several houses along narrow, stony streets are provided with cement furnaces outdoors, steam rushing out through holes in the ground.

A kettle for boiling rice is placed on one furnace while vegetables are cooked on another covered with straw mats. (Pt. I, p. 27; Pt. III, p. 63)

Fig. 5



Carbonic acid gas produced from **Isobe Spring** is collected and conveyed to a factory for preparing carbonate of lime for tooth powder.

(Pt. I, p. 27; Pt. III, p. 22)

Yumura Hot Springs, Tajima Province

Villagers washing in the streams into which flowing hot water flows

(Pt. I, p. 27; Pt. III, p. 55)

Fig. 4



Fig. 1




Fig. 3




Fig. 2



Jikan-yu or "Time-Bath" at Kusatsu.

The Jikan-yu at Kusatsu is a very large bath, and is one of the most famous in Japan. It is situated in the town of Kusatsu, in the province of Suruga. The bath is a large, open-air structure, and is used by a large number of people. The water is very hot, and is said to be good for many ailments. The bath is a very popular place, and is a very important part of the life of the people of Kusatsu. The bath is a very large structure, and is used by a large number of people. The water is very hot, and is said to be good for many ailments. The bath is a very popular place, and is a very important part of the life of the people of Kusatsu.

Fig. 1



Kashima, a Lowenthal's Contactoscope placed in an old *Yatai*, a shrine temple dedicated to the God of Alchemy at the source of Ogawa Hot Springs, Echigo Province. (Pt. I, p. 37; Pt. III, p. 3)

Fig. 3



Collecting gas rising from *Chi no Umi*, "Island of a Blood Lake Hell," a boiling pool with odorous mineral-stained border, near *Yippu*. (Pt. I, p. 37; Pt. III, p. 63)

Fig. 2




Measuring the flow of water at the source of Nihonawa Cold Spring in **Masutomi**, Kai Province. (Pt. I, p. 37; Pt. III, p. 34)

Fig. 4



Schmidt's Electrometer.
The picture shows the apparatus set for practical test at **Masutomi**, Kai Province. (Pt. I, p. 37; Pt. III, p. 37)

Fig. 5



Dr. R. Ishizu and his assistant taking water to be examined for radioactivity, at the source of Akakura Hot Springs on *Mt. Mōko*, about 3.7 m. up the spa more than 4,000 ft. above the sea. (Pt. I, p. 37; Pt. III, p. 47)

Fig. 1



Fig. 1. View of the Misasa river, from which
the water is drawn.

Fig. 4



Fig. 4. A steep hill, wooded with pine trees, one
of the attractions of Misasa.

Fig. 3




Fig. 3. A view of the "Shrine" of Misasa, a temple
where the late war was fought. The
structure is old and in ruins. The
new and the old are
seen in the foreground.

Fig. 2




Fig. 2. A view of the Misasa river, from which
the water is drawn.

Fig. 5



Fig. 5. A view of the Misasa river, from which
the water is drawn.

Misasa Hot Spring Spa, Misasa, Japan


The Misasa Hot Spring Spa is a famous resort in the Misasa district, Japan. It is situated on the Misasa river, which is one of the main attractions of the area.

Fig. 1




Lake Shōji, a favorite summer retreat, at the N. W. foot of Mt. Fuji. (*Pt. II, p. 202*)

Fig. 1



Unzen Resort with hot springs. (*Pt. III, p. 666*)

Fig. 3



Haku-in-dō or
"White Cloud Grotto"
in **Shiobara**
Hot Springs.
(*Pt. III, p. 46*)

Fig. 2



Misaki Sacred Bridge in **Nikkō**. (*Pt. III, p. 18*)

Fig. 5



Miyunoshita Spa, Hakone. (*Pt. III, p. 27*)

Some well-informed men have of late realized their importance and are undertaking to improve their present conditions. If necessary arrangements be made and it become possible that their therapeutic efficacy is fully utilized for the benefit of patients in general, we believe this must be a great boon to the public at large. Especially considering this country is endowed with a mild climate and the natural beauties of scenery left undamaged by abuse of modern culture; and, moreover the prices of daily necessities being cheap, these place Japan in a position to ultimately become a health resort, unrivalled all the world over.

We find many hot springs of excellent composition, but there are very few where proper care is taken to protect the source of springs, conduct the water, build baths, and encourage bathing, so as to suit the theory of emanation therapy. In spite of the existence of gases with strong radio-activity at many places, no inhalatorium has yet been established to turn them to account. This deplorable state of affairs is ascribable to the fact, that emanation therapy has only been very recently introduced to Japan and very few hot spring managers have a thorough knowledge or idea of its theoretical and practical importance. But as the hygienic authorities have the matter under attention and are enforcing improvements, undoubtedly all the Japanese hot springs will be greatly improved in the not very distant future.

We are going to point out a few facts witnessed at various hot spring resorts in Japan that go to prove that radium emanation constitutes an important therapeutic factor of mineral springs.

1. Original springs are generally believed to be more efficacious, than the baths into which the water is conducted.
2. The baths prepared with cold mineral water, artificially heated, are generally inferior to natural hot springs in their therapeutic efficacy.
3. Baths are built as close as possible to the source of springs.
4. The efficacy of hot springs is fully experienced, when a bath is taken in the early morning before other bathers.
5. People bathe in those hot springs of which the temperature is somewhat equal to that of the human body, without heightening their heat artificially.
6. To wash the body with fresh water after bathing is generally believed to lessen the therapeutic efficacy of the springs.
7. People try, while taking baths, to be as near as possible to the source of springs, and apply the water fresh from the source to particular diseased parts.
8. There is a popular belief that it is best to drink at the source of springs.

Although of course there may be found more facts, by more attentive observation, the above are the most generally known and entirely conform with the theories of the modern emanation therapy.

Chapter VIII

SHORT REMARKS ABOUT EMANATION THERAPY.

The radio-activity of mineral waters has attracted much attention during the last decade, and it is now widely believed that the therapeutic value of many waters is largely due to their content of radio-active substances. Tests have shown that many of the slightly mineralized waters, which have proved to be efficacious for certain diseases, have various degrees of radio-activity, and consequently, as radium compounds have been found to cause marked physiologic reactions, the improvement in health following sojourn at springs has very naturally attributed to the radio-active properties of the waters. There are yet too few experimental data to settle this point definitely.

The emanation has been used therapeutically in various ways; it has been inhaled, injected, taken in draughts, and administered by means of baths. Some natural waters are radio-active, and their efficacy may in part be due to this. It is well known that a bottled mineral water drunk at home does not produce the same active benefit that it does when drunk fresh and nascent at the spring, and this may be due to the decay of its radio-activity. This will also explain the impossibility of successfully imitating a natural mineral water; it must be made radio-active as well as of the same chemical composition.

In the case of injection or other introduction directly into contact with the tissue the emanation tends to diffuse itself throughout the tissues; it has a predilection for the glands which form an internal secretion, and especially for the suprarenal capsules. It is eliminated by the lungs and the skin, and to a small extent by the kidneys. Injections of water or of vaseline impregnated with the emanation have been made by Radcliffe Crocker, Wickham and Degrais, in lupus, mycosis fungoides, and cancers, with benefit, and baths and drinks containing the emanation have been administered successfully by Strasser in cases of chronic articular rheumatism, sciatica, and neuritis. Gout is also favourably influenced by this treatment.

Löwenthal states that the emanation is chiefly absorbed by the respiratory passages, that it is of value in the treatment of chronic inflammatory processes and residues of the same, and that in therapeutic doses it is quite harmless. In order to subject the body to the permanent action of the emanation, it is necessary to apply the source of emanation several times a day, since the body gets rid of all traces through the urine and expired air within three or four hours. It is necessary in the case of baths so to construct the baths that the nose lies within the space of the bath itself, so that as much emanation may be absorbed as possible. He has found that the emanation is capable of activating body ferments; this action would mean that the absorbing power of the body would be increased, and in this way the beneficial action on inflammatory processes would be explained. Löwenthal comes to the conclusion that specific urate-dissolving powers are to be found in mineral waters; there may be an action whereby ferments which form and destroy uric acid may be activated by radium. It can be shown that by drinking the radium-containing waters of Baden-Baden the excretion of urea is increased by 34 per cent., while that of urates increased by 14 per cent., as compared to the excretion when the person is drinking the same quantity of ordinary water warmed to the same temperature.

Recently some valuable experiments to test the bactericidal action of radium rays have been carried out by Dr. Helen Chambers and S. Russ, and communicated by them to the Royal Society of Medicine. Their paper concludes as follows:

- “1. The emanation in concentration of less than a millicurie per cubic centimeter has a marked bactericidal action.
2. Agglutination of bacteria in distilled water is an early sequel to their irradiation.
3. Bacteria are more quickly destroyed by the emanation than are opsonine and leucocytes.”

Dr. Saubermann and Prof. Paul Lazarus have recently conducted a valuable research into the action and use of the emanation. The following remarks are taken from papers contributed by them to the meeting of the Röntgen Society on April 1st, 1913:

“Radium emanation can enter the organism in several ways, but in accordance with the character of an inert gas it totally and rapidly leaves the body. The main portion leaves through the lungs, a large part through the intestines, and some through the skin and kidneys. The mode of entry decides the degree and speed of the passage of the emanation through the organism. It lasts *seconds* by inhaling, *minutes* with an injection, *hours* when taken by the mouth and stomach or *per rectum*. Consequently we distinguish five ways of absorbing emanation:

1. Through the lungs.
2. Through the digestive organs.
3. Through the skin.
4. Through the medium of different forms of injections.
5. Through the employment of local external applications—on wounds, for instance.

Prof. Lazarus went on to speak of emanation administered by the mouth through the drinking of emanation water. This entered the organism in three ways: the first, by direct penetration of the stomach and intestines; the second, by diffusion into the capillaries of the lymph and portal vein systems, the emanation circulating through the liver; the third, a method which he termed ‘retrospiration,’ for his tests left no doubt that a very large portion of the imbibed emanation reached the arterial blood. By inhalation the emanation reached primarily the lungs, and secondly the intestines; but the opposite took place in the drink cure. The difference between the two resided principally in the speed of the passage of the emanation through the body. Though the emanation was absorbed more rapidly in the case of primary inhalation, it would be retained scarcely as long as it was being breathed; but in the case of the drink cure the emanation was gradually introduced into the circulation, and it left the system also relatively slowly. The important feature in emanation

treatment was the constant saturation of the body, which was best achieved by giving the patient a dose to drink every 20 minutes for a period of 3 hours. By administering 10 or more doses or sips from a bottle containing the total quantity, a constant saturation of the body was maintained in a manner that would have been impossible had the total quantity been administered at once. He called this form of application the 'sipping cure'."

In a concluding passage, after dealing with the established fact of the absorption of emanation through the intact skin, Prof. Lazarus was thus quoted by his colleague:

"Emanation injections, whether intravenous, subcutaneous, or intramuscular, must prove useless, because the emanation will escape per pulmones in a very short time. Effects can only be obtained if solutions of radium salts are employed, as though establishing a kind of radium depot within the body, which can produce emanation for some time. I have practiced such injections for the past six years in exsudations, in joints, or in serous cavities, but always locally, and I have never been able to record a damaging effect by permanently depositing emanation within the organism."

GENERAL PHYSIOLOGICAL RESULTS OBTAINED BY THE USE OF THE RADIUM EMANATION.

- "1. It promotes the growth of healthy cells while inimical to morbid cells.
2. It produces diuresis.
3. It stimulates the activity of the digestive tract, and also promotes digestion and relieves constipation.
4. It increases the excretion of uric acid.
5. It lowers the blood-pressure by dilating the capillaries and diminishing the viscosity of the blood.
6. It increases the sexual activity and power of generation. Some spas—e. g. Gastein and Baden-Baden—now known to be markedly radio-active have for a long time enjoyed a reputation as *rejuvenating* waters. Engelmann refers to Von Noorden's observations, and to the results of O. Hertwig's experiments, who ascertained a specific effect of radium rays on the nuclear substances of the two sexual cells, and to the observations of Halben, who succeeded in enhancing the secondary symptoms of sexual passion in water newts by radium emanation. Affections of the kidneys, tendency to breeding, and tendency to miscarriage, are regarded at present as contra-indications.
7. It modifies the constitution of the blood by diminishing the number of white corpuscles, and by increasing the number of red." (Saubermann, Archives of the Röntgen Ray, August, 1913).

DISEASES FOR WHICH EMANATION TREATMENT IS SUITABLE.

Gout, chronic articular rheumatism, gonorrhoeal rheumatism, rheumatoid arthritis, neuralgias of all kinds, certain diseases of women, high blood-pressure, premature old age.

There is a general consensus of medical opinion as to the efficacy of radium emanation treatment in gout (Engelmann, Von Noorden, Löwenthal, etc.). Treatment may combine drinking, inhalation, baths, compresses, but the most important is the *sipping cure*. The patient should every half hour sip a dose of water containing the emanation; the total daily dose should not exceed 1,000 Mache's units to begin with, but this may be gradually increased to as much as 10,000. The body will thus be kept constantly saturated with the emanation. After a few days a gouty reaction, accompanied by pains in the joints, sets in, to be followed by a slow but general improvement. Engelmann recommends a six weeks' course of treatment, which should be repeated once or twice yearly. The same general plan of treatment can be followed for the other diseases enumerated above. Engelmann, in recommending radium emanation treatment for neuritis and neuralgia, states that he has seen two extremely obstinate cases of herpes zoster in persons advanced in years, which had defied every other kind of treatment, quite surprisingly bettered. Even the lancinating pains of tabes are relieved. The author has successfully treated an obstinate case of post-herpetic neuralgia by the radiation of solid radium. Engelmann further recommends the emanation in bronchial asthma, arterio-sclerosis, and diseases of women, especially in cases in which it is important to assist the absorption of chronic exsudates and swellings; and in general adhesions, tissue thickenings of every kind cannot be otherwise than favourably influenced by a combined local and general emanation treatment. Advantageous results, he says, after radiation with radium preparations in cases of myoma and disturbances in

menstruation, have quite recently been reported from the Freiburg clinic for women. In these cases the radiation with radium rays took the place of radiation with X-rays. A histologically demonstrable influencing of the ovaries, and also occurrence of miscarriage after mere drinking of emanation water, have already been experimentally ascertained. The X-ray treatment of myoma might, in the authors' opinion, be replaced by radium radiations, for the latter possess two intrinsic advantages: the dose can be measured more accurately, and the gamma rays are more penetrating. Testimony as to the value of emanation treatment in arthritis deformans is to be found in the report of the Radium Institute (British Medical Journal, January 25, 1913):

"This extremely obstinate, progressive, and crippling malady is not infrequently strikingly benefited by the daily drinking of 250 cc of radium emanation solution of a strength of 1-2 millicuries per litre. The treatment must, however, be persisted in for quite a long time, and at least six weeks are likely to elapse before any change is noted. In a favourable case the articular and muscular pains are lessened or disappear, the movements of the affected joints become much freer, and are accompanied by less grating; the muscles controlling the joints regain much of their lost tone, and the general health of the patients is greatly improved. With the majority of the patients the emanation solution produces a definite diuresis, and with a few it acts slight laxative. Up to the present only 21 patients have been treated, but the results obtained are sufficient to warrant the hope that radium emanation solution will prove a most valuable addition routine medical treatment."

Cases of emanation treatment and its therapeutical effects, which are to be found in literature till 1911, are given in the following table:

Table IX.

Summary.

| No. | Name of Disease | Number of Treatment | of which | | in per cent | |
|-----|---|---------------------|----------|-----------|-------------|-----------|
| | | | cured | benefited | cured | benefited |
| 1 | Anaemia | 11 | 1 | 10 | 9,09 | 90,9 |
| 2 | Ankylosis | 1 | — | 1 | — | 100 |
| 3 | Acne | 1 | — | 1 | — | 100 |
| 4 | Asthma bronchiale | 1 | — | 1 | — | 100 |
| 5 | Anaesthesia and paraesthesia in the fifth nerve | 1 | — | 1 | — | 100 |
| 6 | Apoplexie | 11 | — | 10 | — | 90,9 |
| 7 | Apoplexia cerebri | 1 | — | 1 | — | 100 |
| 8 | Arteriosclerosis (stenocardie) | 5 | — | 4 | — | 80 |
| 9 | Arthritis urica | 125 | 38 | 63 | 30,5 | 50,4 |
| 10 | Arthritis chronica | 13 | 3 | 9 | 23,07 | 69,23 |
| 11 | Arthritis rheumat. chron. | 27 | 18 | 5 | 66,6 | 17,7 |
| 12 | Arthritis urica | 2 | — | 2 | — | 100 |
| 13 | Arthritis subacuta | 10 | 7 | 3 | 70 | 30 |
| 14 | Arthritis rheum. subacuta | 10 | 3 | 5 | 30 | 50 |
| 15 | Arthritis rheumatica | 42 | 18 | 10 | 42,86 | 38,09 |
| 16 | Arthritis deformans | 91 | 12 | 52 | 13,2 | 57,1 |
| 17 | Arthritis chron. deformans | 7 | — | 7 | — | 100 |
| 18 | Arthritis tbc. | 2 | — | 2 | — | 100 |
| 19 | Bronchial catarrh | 67 | — | 50 | — | 84 |
| 20 | Bronchitis chron. | 3 | 1 | 2 | 33,30 | 66,6 |
| 21 | Carc. mam. inop. | 4 | — | 1 | — | 25 |
| 22 | Carc. oesoph. inop. | 3 | — | 1 | — | 33 |
| 23 | Catarrh of frontal sinus and super. maxil. | 11 | 10 | 1 | 90,9 | 9,1 |
| 24 | Cephalaea | 1 | — | 1 | — | 100 |
| 25 | Cephalaea traumat. | 1 | — | 1 | — | 100 |
| 26 | Cephalaea rheumat., etc. | 1 | 1 | — | 100 | — |
| 27 | Cystitis chronica | 1 | 1 | — | 100 | — |
| 28 | Dacryocystitis | 1 | — | 1 | — | 100 |
| 29 | Ductect. in epulis | 8 | — | 1 | — | 12,5 |
| 30 | Distorsion (after treatment) | 2 | — | 2 | — | 100 |

| No. | Name of Disease | Number
of
Treatment | of which | | in per cent | |
|-----|---|---------------------------|----------|---------|-------------|---------|
| | | | cured | limited | cured | limited |
| 31 | Dupuytren's contraction | 1 | — | 1 | — | 100 |
| 32 | Edema | 9 | 1 | 5 | 11,1 | 55,5 |
| 33 | Exacerb. pains after affect. of acces. sinus | 1 | 1 | — | 100 | — |
| 34 | Exudate of joint | 21 | 9 | 11 | 42,86 | 52,4 |
| 35 | Gastroenteritis chron. | 11 | 1 | 9 | 9,09 | 81,81 |
| 36 | Gout | 11 | 4 | 5 | 36,4 | 45,5 |
| 37 | Glandular growth | 2 | 1 | 1 | 50 | 50 |
| 38 | Hemierania | 1 | — | 1 | — | 100 |
| 39 | Hydrosalpinx | 1 | 1 | — | 100 | — |
| 40 | Hypertrophy of prostate | 2 | — | 2 | — | 100 |
| 41 | Laryngitis | 1 | — | 1 | — | 100 |
| 42 | Lumbago | 21 | 11 | 9 | 57,1 | 28,6 |
| 43 | Lymphoma tbc. colli | 10 | — | 9 | — | 90 |
| 44 | Malignant swelling | 2 | — | 1 | — | 50 |
| 45 | Metritis chronica | 7 | — | 7 | — | 100 |
| 46 | Muscular rheumat. of head | 4 | 2 | 2 | 50 | 50 |
| 47 | Muscular rheumat. of shoulder | 10 | 5 | 4 | 50 | 40 |
| 48 | Muscular rheumat. of neck | 5 | 4 | 1 | 80 | 20 |
| 49 | Muscular rheumat. of breast | 3 | 2 | 1 | 66,7 | 33,3 |
| 50 | Myelitis chronica | 4 | — | 1 | — | 25 |
| 51 | Myocarditis | 35 | — | 7 | — | 77,2 |
| 52 | Myocarditis chronica | 5 | — | 4 | — | 80 |
| 53 | Myositis traumatica | 1 | — | 1 | — | 100 |
| 54 | Neuralgia of brachial plexus | 8 | 5 | 2 | 62,5 | 25 |
| 55 | Neuralgia in the 2 a. 3 branch. of the fifth nerve | 4 | — | 3 | — | 75 |
| 56 | Neuralgia and paraesthesia in ramus supraorb. after herpes zoster | 1 | — | 1 | — | 100 |
| 57 | Neuralgia (Neuritis) | 35 | 13 | 15 | 37,2 | 42,8 |
| 58 | Neuritis chron. and traum. | 2 | — | 2 | — | 100 |
| 59 | Neurasthenia | 58 | 21 | 29 | 36,2 | 50 |
| 60 | Omphalitis simpl. and dupl. | 8 | 4 | 2 | 50 | 25 |
| 61 | Osteomyelitis | 1 | — | 1 | — | 100 |
| 62 | Ovaria | 4 | — | 4 | — | 100 |
| 63 | Pain | 3 | — | 2 | — | 66,7 |
| 64 | Paralysis nervi facialis rheumatica | 1 | — | 1 | — | 100 |
| 65 | Paralysis infantum | 3 | — | 1 | — | 33,3 |
| 66 | Paralysis agtans | 5 | — | 1 | — | 20 |
| 67 | Parametritis and perimetritis | 15 | 5 | 10 | 33,3 | 66,7 |
| 68 | Peritonitis tbc. and chron. | 4 | 2 | 1 | 50 | 25 |
| 69 | Pyosalpinx | 1 | 1 | — | 100 | — |
| 70 | Polyarthrititis | 19 | 16 | — | 84,2 | — |
| 71 | Polyarthrititis rheumatica chron. | 25 | 10 | 7 | 40 | 28 |
| 72 | Rectum fistula tbc. | 2 | 1 | 1 | 50 | 50 |
| 73 | Rhynitis chronica | 38 | 19 | 8 | 50 | 21,1 |
| 74 | Rheumat. nasc. acut. | 10 | 4 | 4 | 40 | 40 |
| 75 | Rheumat. musc. sub. and chronica | 40 | 19 | 22 | 47,5 | 55 |
| 76 | Rheumat. artic. chron. | 250 | 93 | 27 | 37,2 | 50,8 |
| 77 | Rheumat. gonorrh. | 5 | — | 4 | — | 80 |
| 78 | Residua after fractur | 7 | 9 | — | 85,6 | — |
| 79 | Sciatica | 180 | 95 | 59 | 52,8 | 33 |
| 80 | Sclerodermie | 2 | — | 2 | — | 100 |
| 81 | Tales dorsalis | 66 | — | 33 | — | 50 |
| 82 | Tbc. palmum | 6 | — | 2 | — | 33,3 |
| 83 | Uterus myomatosus | 11 | — | 11 | — | 100 |
| 84 | Vaginitis | 1 | — | 1 | — | 100 |
| 85 | Vitium cordis | 8 | — | 3 | — | 37,5 |

THE THORIUM EMANATION.

The thorium emanation is given off more or less freely from the compounds of thorium, the best source, according to Prof. Soddy, being a preparation of radiothorium in a moist condition. Radiothorium is derived from thorium through two intermediate products named respectively methothorium No. 1 and methothorium No. 2. Methothorium No. 1 is produced as a by-product in the manufacture of gas-mantles; it resembles radium in its chemical nature, and cannot be separated from it; it disintegrates into methothorium No. 2. The latter radiates beta and gamma rays, and yields radiothorium. From the latter thorium X is produced, and from thorium X the thorium emanation. The emanation has a life of about 26 seconds. It radiates alpha rays, and leaves an active deposit called thorium A. This integrates into thorium B, this into thorium C, and this into thorium D. The important members of this group, from the medical point of view, are methothorium, thorium X, and its emanation.

The thorium emanation would also probably be found of use in the same class of cases in which the radium emanation has proved of value. The class of cases referred to include gout and rheumatism and other general diseases for which natural mineral waters are prescribed. It has been found that these waters are as a rule radio-active, and that with the smallest mineral content are the most radio-active. W. His states that radio-active baths and waters owe their therapeutic value to the emanation of radium. After three years' experience observations have been taken on over 100 cases of gout, and the large majority have been so obviously benefited that the results cannot have been due to chance. The emanation decomposes uric acid into urea and carbonic acid. Löwenthal states that the emanation stimulates the body ferments and increases the gaseous exchange.

Use of Radio-active Earths.—Reference has already been made to the production and use of the emanation. It can also be administered by the employment of radio-active earths. One of the best known of these is obtained in the process of uranium extraction; it contains certain oxides of iron, aluminium, uranium, and manganese, and traces of actinium, radium, and polonium. Of the latter, actinium is the most important. Pure actinium is estimated by some physicists (pure actinium has not been isolated, but it is believed that it would show an activity comparable with that of radium—Rutherford) to be ten times as radio-active as radium, the quantity present in the earth suffices to confer upon it a radio-activity of 0.15 as compared to 2,000,000 in the case of radium.

The following is a table of comparative radio-activities (Claude):

| | | | |
|----------|------------------|---------|------------|
| Actinium | about 20,000,000 | Uranium | about 1 |
| Radium | about 2,000,000 | Earth | about 0.15 |

One centigramme of the earth spread over one square centimeter yields a radiation about one-seventh the strength of that of uranium. The earth is an extremely weak radio-active preparation, which yields alpha, beta, and gamma rays, and an emanation. It is, however, relatively to radium, very cheap, a kilogramme costing thirteen shillings.

The earth has been employed for some time by Dr. Claude and others in the hospitals of Paris in the treatment of arthritis deformans, subacute and chronic rheumatism, and gonorrhœal rheumatism, also for neuralgia and certain cutaneous (pruritus) and gynæcological affections.

The affected joint having been cleansed, the moistened earth is spread in the form of a compress around it, and maintained in position by a piece of oil-silk and a bandage. It must not be allowed to become dry. No emanation would be given off if it became dry. The plaster can be left on all night, and applied on alternate nights for ten days.

It should always be remoistened and made quite plastic and soft before application. It does not lose its strength. In some cases it may be advisable to sterilize it by heat (open sores, introduction into the cavities of the body); this will not affect its activity.

Dr. Guyenot prepares radio-active earth or mud by mixing powdered pitchblende with twice its volume of hot water. This is applied direct to the part to be treated, and can be left on several hours under a piece of waterproof. This mud exercises a sedative effect, and is believed to promote the absorption of exsudations. It does not irritate the skin, its radio-activity is permanent; but when it has been used, it must be allowed to dry for a fortnight before it is used again.

Another method of employing the earth is to make the use of baths. 250 grm of the earth are mixed with the warm water of the bath; more may be used if desired, but this is the usual

strength. The patient can remain in the bath for half an hour, and have a series of a dozen in a fortnight or longer.

Good results are claimed to have been obtained by Claude and Teulière in arthralgia and subacute forms of arthritis. The pain, redness, and swelling, usually disappear with great rapidity, but affections of the lower limb appear to be more rebellious than those of the upper limb. The best results appear to have been obtained in subacute gonorrhœal rheumatism.

Mud baths have been for a long time in vogue at certain health resorts in Europe; it is possible that the beneficial results may have depended upon the presence of traces of radio-active substances.

Chapter IX

DETERMINATION OF RADIUM EMANATION.

For measuring the radio-activity of mineral waters, Schmidt's electrometer (made by Spindler and Hoyer, Göttingen) was employed. It has been standardized with a radium solution of known strength. In our tests the so-called bubbling method was employed. The water for examination was carefully poured into a vessel. After the circulating system was completed, air was bubbled through the water by means of a circulation rubber bulb, in order that the greater part of the emanation should be taken up by the air. The air was then freed from its humidity by a calcium chloride tube and pressed into the ionisation chamber. Great care was taken to have the emanation equally distributed in the whole circulation system. On the stop-cocks of the ionisation chamber being closed the quantity of emanation was determined in the usual way from the diminution in the charge after the maximum current had been reached. Here the leaf of the electrometer was always charged negatively, so as to concentrate the active deposit on the inner electrode.

As by this method only a part of the emanation present in the water was determined, the leak therefore was multiplied with a factor in order to get the leak of the total emanation contained in 1 litre of spring water.

The results of measurements of radium emanation have unfortunately been expressed in a great variety of units, and as a natural consequence it is impossible to compare some reports of radio-activity with others. In England and America it is customary to express radio-activity as an equivalent of radium or of uranium, a radio-active substance; on the Continent it is customary to express the results in electrostatic units. It is quite general, irrespective of the unit, to report the radio-activity per litre of water or per litre of the gas in the water. The quantity of emanation in equilibrium with 1 gram of radium is known as a curie of emanation, but as this is a very large unit estimates are expressed in fractional parts of a curie; thus, 516.87×10^{-10} curies per litre means that one litre of the water contains 516.87 ten-billionths of the quantity of emanation in equilibrium with 1 gram of radium. The Mache's unit, frequently encountered in reports of radio-activity, is defined as the saturation ionization current due to the radium emanation, free from decay products from a litre of water or gas, expressed in electrostatic units (*i*) multiplied by 1000; and 1 curie of emanation per litre equals a concentration of 2.75 billion Mache's units.

The Mache's unit being recognized and still in general use amongst medical circles, we have recalculated our result (as expressed in the international unit) by using the following factor:

The saturation current produced by the radiation from one curie (international unit) of emanation by itself is 2.75×10^6 e. s. units, (Rutherford's "Radio-active Substances and their Radiations," 1913, p. 502).

So we have

$$2.75 \times 10^6 \text{ Mache's units} = 1 \text{ curie,}$$

so that

$$1 \times 10^{-10} \text{ curies} = 0.275 \text{ Mache's units.}$$

For simplicity of manipulation, the original fontactoscope of C. Engler and H. Sieveking (made by Günther and Tegetmeyer, Braunschweig) or sometimes the same modified by Kohlrausch and

Löwenthal was employed in most cases of our determination instead of Schmidt's electrometer. The determination was carried on exactly according to the authors' original instructions. The corrections for the absorption of emanation in water, and for the absorption of the radiation by the walls of the ionisation chamber were made taking also the effect of active deposit into consideration. The gas evolving with the spring which was examined, was collected over the spring water by replacement as usual and was examined practically by the same method as adopted with the water. The volume of gas was reduced to the normal temperature, though the correction due to the atmospheric pressure was neglected as its observation were not made.

Those spring waters impregnated with considerable quantities of gas, for example, carbonic acid, hydrogen sulphide, etc., were previously treated with a sufficient quantity of caustic soda, whereat the eventual loss of emanation due to occlusion by precipitates was not taken into consideration.

As for the measurement of the volume of the circulation system, (excepting bottle for spring water to be examined) we have adopted the replacement method. The air in the system in question (ionisation chamber, circulation rubber bulb, calcium chloride tube, rubber tubes for connection, etc.) was first replaced with pure dried carbonic acid gas, the latter being introduced into a weighed potash bulb by a current of dried air free from carbonic acid, and then the quantity of carbonic acid absorbed was from this gravimetrically determined. From the quantity of carbonic acid gas the volume of the space which occupies the system can be easily calculated, taking into account variation of volume of gas due to temperature and atmospheric pressure.

The general opinion that most spring waters do not contain radium itself, but radium emanation derived from contact with radium-bearing minerals, is logically based on the fact that nearly all spring waters gradually lose their radio-activity after being removed from the source, whereas a solution of a radium salt, after having once reached equilibrium, continues to drive off radium emanation. There are, as far as we know, only a few cases of mineral waters, where the radio-activity is due to the presence of dissolved radium salt:—

The spring water of *Kreuznach* has proved to have rest-activity of radium. 8 litres of the thermal spring in *Gastein* caused after being removed of emanation yet a discharge of 0,5–0,8 volt per minute after 14 days, while in a test of *Carlsbad* waters a discharge of $1,065 \times 10^{-3}$ electrostatic units was observed after 25 days (Dora).

It is probable that the thorium emanation also has a share along with that of radium in radio-activity of mineral waters, but the former generally escapes our usual method of measurements. We know only that thorium emanation was found in some spring waters in *Homburg v. d. Höhe* (Kaiserbrunnen and Elisabethbrunnen).

Observation has as not yet been made upon the thorium emanation of the spring waters of Japan.

Table X.

Absorption of the Radium Emanation by Water (Boyle).

| Temperature | Coefficient of Absorption, α | Temperature | Coefficient of Absorption, α |
|-------------|-------------------------------------|-------------|-------------------------------------|
| 0° C | 0,507 | 20,3° C | 0,250 |
| 4,3° C | 0,424 | 26,8° C | 0,206 |
| 5,7° C | 0,398 | 31,0° C | 0,193 |
| 10,0° C | 0,340 | 34,8° C | 0,176 |
| 14,0° C | 0,303 | 39,1° C | 0,160 |
| 17,0° C | 0,280 | | |

Table XI.
Decay of Radium Emanation.

| Time <i>t</i> in hrs. | <i>e</i> ^{<i>t</i>} | Time <i>t</i> in days | <i>e</i> ^{<i>t</i>} | Time <i>t</i> in days | <i>e</i> ^{<i>t</i>} |
|-----------------------|------------------------------|-----------------------|------------------------------|-----------------------|------------------------------|
| 0 | 1.000 | 1 | 0.835 | 20 | 0.037 |
| 1 | 0.993 | 2 | 0.698 | 21 | 0.023 |
| 2 | 0.985 | 3 | 0.583 | 22 | 0.019 |
| 3 | 0.978 | 4 | 0.487 | 23 | 0.016 |
| 4 | 0.970 | 5 | 0.407 | 24 | 0.013 |
| 5 | 0.963 | 6 | 0.340 | 25 | 0.011 |
| 6 | 0.956 | 7 | 0.284 | 26 | 0.008 |
| 7 | 0.949 | 8 | 0.237 | 27 | (2) 775 |
| 8 | 0.942 | 9 | 0.198 | 28 | (2) 647 |
| 9 | 0.935 | 10 | 0.165 | 29 | 2 541 |
| 10 | 0.928 | 11 | 0.138 | 30 | (2) 452 |
| 11 | 0.921 | 12 | 0.115 | 30 | (3) 747 |
| 12 | 0.914 | 13 | 0.096 | 50 | 3 123 |
| 13 | 0.907 | 14 | 0.081 | 60 | (4) 204 |
| 14 | 0.900 | 15 | 0.067 | 70 | (5) 337 |
| 15 | 0.894 | 16 | 0.056 | 80 | 6 558 |
| 16 | 0.887 | 17 | 0.047 | 90 | 7 022 |
| 17 | 0.880 | 18 | 0.039 | 100 | 7 152 |
| 18 | 0.874 | 19 | 0.033 | | |
| 19 | 0.867 | | | | |
| 20 | 0.861 | | | | |
| 21 | 0.854 | | | | |
| 22 | 0.848 | | | | |
| 23 | 0.842 | | | | |

The numbers in brackets, e. g. (6) refer to the number of o's before the first significant figure.

DETERMINATION OF ELECTRIC CONDUCTIVITY.

As the physiological effect of mineral waters may be regarded as due to the action of ions contained in them, it is necessary to extend our investigation of the springs in this direction.

In mineral waters, which contain, as a rule, only a small amount of salts, we find that the greatest part of the dissolved substances is contained as ions and not as salt molecules. To obtain a gauge for degree of dissociation, we have to determine (1) electric conductivity, (2) depression of freezing point, or (3) elevation of boiling point. We have taken the first, as it can be easily measured even while travelling through the field. This determination was made with Pleissner's portable apparatus, made by Bosse & Co., Berlin. Observations were usually made at 18° C. as the normal temperature, necessary corrections were made for those at other temperatures.

Chapter X

THE SINTER DEPOSITS OF MINERAL SPRINGS.

A. On a small Sinter-Cone formed by the Geuser of Obama, Hizen.

Obama Hot Springs are reached by steamers in about four hours from Mogi, a hamlet in Hizen Province, well known for its pliocence plant fossils described by Nathorst. (Contributions à la flore fossile du Japon. Kongl. Svenska Vetenskaps-Akademiens Handlingar, Bandet 20, No. 2, 1883.)

* Denzō Satō. Beiträge zur Mineralogie von Japan, No. 4, p. 107, 1912.

Obama belongs to a region of hornblende-andesite, erupted from the isolated volcano *Unzen-dake*. One of the springs, called "*Funtō-yu*," is a geyser from a bore-hole protected by an iron pipe about 25 meters in length and 6 cm. in diameter. Here small sinter-cones are being formed at the bottom of a wooden tank placed above the bore-hole, from which the water is conducted by a pipe into the baths.

see Plate 4,
fig. 1

The cone, shown in Plate 4, fig. 1, is a good specimen of typical form, taken in 1900 and exhibited at present in the Mineral Museum of the Imperial Geological Survey (Tōkyō).

Denzō Satō who described the one of the cones writes as follows:—

see Plate 4,
fig. 2 a, 3

"The whole time required for its deposition seems to have been about five months, so that the daily precipitation amounts ca. 43 grm., which nearly coincides with that of the sinter-cones of Yuzawa. The water of this geyser is slightly alkaline and faintly salty in taste. The temperature is 100° C. (at air temperature 15°), flow of water about 39 hectolitres in 24 hours. The geyser is of intermittent nature and there is a single greater ejection in a number of smaller ones. The latter, which throws the water to a height of about 3 meters, occur every second, while the greater one, which sends it up to about 4.5 meters, takes place every five seconds.

Analysis made in the Geological Survey is as follows:—

Total residue per litre of water: 9.2. Specific gravity: 1.006 at 15.5° C.

| | | | |
|---|--------|-----------------------------|--------|
| NaCl | 5.6634 | CaCO ₃ | 0.3215 |
| KCl | 1.6339 | CaCl ₂ | 0.2754 |
| Na ₂ SO ₄ | 0.6358 | SiO ₂ | 0.2180 |
| MgCl ₂ | 0.5358 | FeCO ₃ | 0.0163 |

Thus the water has nearly the same composition as that of Yuzawa Hot Springs.

The cone from Obama is snow-white, and so loose that it can be scratched with the finger nail. The cone consists of:—

| | | | |
|----------------------------|---------|--|--------|
| CaO | 38.28 % | Cl | 1.03 % |
| CO ₂ | 23.28 " | Na ₂ O | 0.89 " |
| SiO ₂ | 16.59 " | Fe ₂ O ₃ | 0.41 " |
| Ign. loss | 10.46 " | K ₂ O | 0.20 " |
| MgO | 7.28 " | Al ₂ O ₃ | 0.18 " |
| MnO | 1.80 " | | |

The cone is flat and symmetrical, gently sloping, with a broad truncation, and a central orifice about 6 cm. in diameter. The slope is about 50° near the top, about 35° in the middle, and about 20° near the base. The peculiar surface sculpture consists of radial valleys, narrowed and deepened towards the bottom, instead of towards the top as is usually the case on volcanoes. Between the valleys there are rounded ridges cut across by broad grooves. The formation of those valleys is probably due to the erosive action of the hot water flowing down the slopes. The specimen, represented in the figure, weighs about 6.457 grm., and measures about 13.7 cm. in height and about 37 cm. in diameter at the base."

13. The Tufa Cones* formed at the Kuriyama Hot Spring in the Province of Shimozuke.

Regular cones of sinter are being produced at Yuzawa in Kuriyama-mura, Shimozuke Province, a place deep in the mountain and seldom visited by men. This locality lies about 19.5 m. N. W. of Nikkō, on an affluent of the upper course of the Kinu-gawa.

see Plate 4,
fig. 4

All the vicinity of Yuzawa is composed of liparite from which several hot springs gush out. The same rock standing on the western side of the Yuzawa waterfall has a long horizontal crevice, about 10 ft. above the river-level, with hot waters rushing out horizontally. From a cliff near by, almost on the same level, are found two other geysers, which are ejecting hot waters vertically to a height of about 10 ft. The horizontal as well as vertical geysers form cones around their jets, which grow naturally larger in the latter than in the former. These vertical sinter-cones consist of carbonate of lime, with a very small quantity of silica and sulphur. The inner part is of compact

* Prof. Wataru Watanabe—Nippon Kōgyō-Kwaishi (Journ. of the Mining Institute of Japan) No. 194, p. 865, 1909.

aragonite. The compact part has the hardness 3 and specific gravity 3,017 (a little higher than ordinary aragonite). The sinter cones are provided with a central canal, the wall of which is covered to unequal thickness with sharp-pointed crystals, slightly reddish by the presence of manganous carbonate. The canal becomes thus narrower, as the cone increases in height; and the emission of hot water decreases in force.

On account of the depression of pressure and temperature of the spring water, after gushing out from the ground, a part of calcium bicarbonate contained in solution loses carbonic acid and forms the neutral carbonate which forms a deposit on the jets of spring, increasing gradually in size and compactness. The sinter cone, shown in Plate 4, fig. 2, is the one, which has been formed during a period of about ten months (304 days) from Aug. 26th., 1907, to June 25th., 1908. and consisting of a white crystalline mass with a conical canal. Height: 2 ft. Diameter at the base: 1 ft. Weight: 3,7 *kwamme* (13,85 kilograms). Amount of daily precipitation: 12,1 *monme* (45,4 grams). The other one, fig. 3, has been formed during a period of about nine and half months (286 days) from Aug. 27th., 1908, to June 9th., 1909, at the same jet of geyser, showing a similarity to Mt. Fuji with the explosion-crater of Hō-eizan. This cone has two narrow canals opening at the top. The cone is white, but looks yellowish on the outside, by presence of floury sulphur. Height: about 8 inches. Diameter at the base: 1 ft. Weight: 3,25 *kwamme* (12,2 kilograms), Amount of daily precipitation: 11,3 *monme* (42,4 grams). The idea of preservation of scientifically interesting natural objects being quite rudimentary among ordinary sightseers, they damage these beautiful cones as soon as they grow up, apparently without any purpose of utilization. A cone, which has been grown to a height of more than 5 ft. and regarded as one of the largest ever found, was said to have been destroyed by an inundation of Sept., 1904.

The water of the geysers is transparent and colourless, weak salty in taste, and the temperature of spring measured at the orifice is 94° C. The flow of water amounts to 4,26 litres per minute. Analysis of water made by Prof. M. Kawakita, in the Engineering College of Tōkyō Imperial University, in 1909, gave the following results:— Specific gravity: 1,02.

In 1 kilogram of water are contained:

| | | | |
|---------------------|----------------|--------------------------|--------|
| Sodium | 0,5293 | Chlorine | 0,7065 |
| Potassium | 0,0238 | Sulphuric acid | 0,0883 |
| Calcium | 0,0640 | Silicic acid | 0,1266 |
| Magnesium | small quantity | Carbonic acid | 0,2076 |
| Aluminium | .. | Total | 1,7461 |
| Iron | .. | | |

Hypothetical form of combination.

| | | | |
|------------------------------|--------|-------------------------------|----------------|
| Sodium chloride | 1,1642 | Magnesium carbonate | small quantity |
| Calcium carbonate | 0,1600 | Ferrous carbonate | .. |
| Silica | 0,1266 | Manganous carbonate | .. |
| Sodium bicarbonate | 0,1241 | Alumina | .. |
| Sodium sulphate | 0,1134 | Total | 1,7414 |
| Potassium sulphate | 0,0531 | | |

C. Sinter-coated Leaves, Sinter Balls, and other Sinter Deposits.

Some of the mineral springs, especially the earthy carbondioxated springs, produce more or less abundant sinter deposits at the source and on the creek bed on which they flow, besides in the conduit and at the bathing-place. The deposits vary in form, structure, texture, colour, etc., according to the composition and temperature of the water, as well as the velocity of flow and other conditions. All these determine also the rate of growth of different sinters. Some of the sinter shows a considerable growth in a short time, that the wooden conduits ought to be replaced by new ones, every four or five years, unless the crust formed on the inner wall is removed from time to time.

Plate 1, fig. 2 shows a piece of the regularly stratified calcite sinter, 5 cm. in thickness, formed at the Shigaku Hot Springs, Shimane Prefecture, in a wooden conduit running along the slope of Mt. Sambe-yama, from the source lying about 0,6 m. off the bathing resort. The breadth of the surface in contact with hot water is 11,0 cm. and that of the face in contact with the wood is 18,5 cm. The whole thickness of this specimen, (5 cm.), is told to be the growth in less than eight years.

see Plate 4,
fig. 2

see Plate 4,
fig. 3

see Plate 1

The colour is yellowish-brown, varying in intensity in different layers, which differ from one another also in compactness.

see Plate 1,
fig. 3

Fig. 3 is a pair of weights made of the same sinter, for the hanging picture in Japanese rooms, just for the object of *souvenir* sold to visitors. Each of the pieces, about 45 grm. in weight, is in the form of a short hexagonal prism, 45 mm. in length, and 30 mm. in diameter, with a longitudinal canal right through the middle.

see Plate 1,
fig. 1

Plate 1, fig. 1 shows the same kind of the grayish-white sinter, deposited in a wooden conduit of the *Akakura* Hot Springs, Echigo Province. The hot water flows down over a distance of more than 4 miles, along a steep mountain path on the middle slope of Mt. Myōkō. The upper figure (a) shows the lower surface of a fragment of the deposit, 24,3–28,0 cm. in length and 20,0–22,5 cm. in breadth, weighing 7,65 kilograms and showing the impression of the grain of wood. (b) is a fracture surface of the same, measuring 7,3 cm. in breadth. This specimen was produced in a conduit with a comparatively low gradient, and shows such a loose texture as to be easily scratched with the finger nail. Those found in one of a steeper gradient is generally thinner, but harder and more compact.

see Plate 2,
fig. 1

Sinter-coated leaves occur on the creek-bed, near the *Masutomi* Mineral Springs in Kai Province, with the water of "*Kinsen-tō*" flowing in. They are represented in natural sizes in Plate 2, fig. 1. The largest and most complete leaf measures about 12,0 cm. in length and 5,5 cm. in the broadest part, and 3,0–3,5 mm. in thickness. The deposits are yellowish-brown in colour, but are sometimes coated with green algae.

see Plate 2,
fig. 2

The sinter, fig. 2, was collected in the neighbourhood of the hot waterfall at the source of the *Ogawa* Hot Springs, Etchū Province. We find there root of plants covered with sinter, consisting chiefly of very minute crystal individuals of calcite lying one upon another.

see Plate 2,
fig. 3

The sinter-balls of calcite, fig. 3, are collected from the basin of the hot waterfall of *Ogawa*. Some of them are almost perfectly spherical, but all the others are more or less irregular in shape. In some specimens, the surface is quite smooth and lustrous but in others it is slightly drusy. The colour of the balls is yellowish-brown, as in the other sinter deposits above mentioned. The largest specimen in the figure is about 22 mm. in diameter, and about 13,8 grm. in weight, while the smallest one is 10 mm. in diameter and about 3,5 grm. in weight.

Chapter XI

A FEW WORDS ABOUT THE RADIO-ACTIVE MINERALS IN JAPAN.

The occurrence of radio-active minerals is already known in Japan, but none of them affords good quantity. The following species have up to the present been found.

1. HOKUTŌLITE.

Hokutōlite is the new radio-active mineral, found in streamlets of the *Hokutō* Spring in the island of Taiwan (Formosa) and at the *Shibukuro* Spring on the *Yake-yama* Volcano in the Province of Ugo. This mineral name was given by Prof. Kōtōra Jimbō. (See List of New Minerals, in *Mineralogical Magazine*, London, Vol. XVI, p. 362, 1913.) The mineral is a mixture of BaSO_4 and PbSO_4 , crystallized in minute rhombic platy crystals of a yellowish-gray or brown colour and a resinous luster.

Hokutō, Taiwan. This well known place with its hotels and baths is situated on the south-western side of Mt. Shichisei-zan, which forms a part of the *Daiton* Volcano Group in Northern Taiwan. There are a number of solfataras with a considerable quantity of sulphur deposits and an explosion-crater having a perfectly flat bottom. Hot springs of 80° to 90° C. issue out of the bottom of this crater, and unite in forming a creek, here not more than half a meter in depth, and only about three meters in width. Almost all of the hotels use this water for bathing purpose, an exception being one large modern bath which is fed from a different source. The explosion-crater as is usual shows rocks bleached and decomposed to earthy matter, carrying efflorescence of a white soluble substance, accompanied by a small quantity of flourey sulphur. The following analytical result

was obtained with water from this creek, taken at the seat of a common bath now out of use, where the creek bed is already broad and suddenly descends, making a low waterfall, as if cut across by a fault line.

| | | | |
|---|-------|--|-------|
| Na ₂ SO ₄ | 19,55 | Al ₂ (SO ₄) ₃ | 2,00 |
| NaCl | 9,70 | Al ₂ (HIO ₆) ₃ | 0,02 |
| KCl | 7,10 | BaSO ₄ | 0,07 |
| Al ₂ (SO ₄) ₃ | 6,58 | PbSO ₄ | trace |
| NH ₄ Cl | 0,73 | H ₂ SiO ₃ | 2,07 |
| CaSO ₄ | 4,78 | HBO ₂ | 0,71 |
| FeSO ₄ | trace | Free acid (HCl) | 19,59 |
| Fe ₂ (SO ₄) ₃ | 5,42 | | |

This water is colourless and almost free from suspended matter. The taste is strongly acid and the smell of sulphuretted hydrogen is noticeable. The above analysis perfectly coincides with that of the water obtained at the spot where the streamlets on the crater bottom finally unite together on the western side, and therefore it is most probable that the waters at the original spring have the same composition as this. It must be emphasized that there is observed no deposition of *hokutōlite* at the point of origin, the temperature of water being possibly too high. The gravel in the creek is mostly of andesite, more or less decomposed on the surface, sometimes one meter in diameter, but usually not larger than a man's head. These blocks are cemented together by the river sand, and a yellow or brown ferruginous substance. The gravels, as well as the consolidated sand, are covered with a thin fibrous crust of *hokutōlite*, the outer part of which forms druses of rhombic tabular crystals. In the following tables are given (I) the analysis of the crystalline crust, and (II) that of its well crystallized external part:—

| | I | II | | I | II |
|--|-------|-------|---|-------|--------|
| PbO | 21,96 | 19,38 | K ₂ O | 0,00 | 0,14 |
| BaO | 32,04 | 42,27 | Na ₂ O | 0,53 | 1,53 |
| SrO | 0,93 | trace | H ₂ O | 2,53 | — |
| CaO | 0,51 | 0,17 | Ign. loss | — | 2,74 |
| SO ₃ | 30,81 | 31,70 | P ₂ O ₅ | 0,01 | 0,00 |
| Fe ₂ O ₃ | 3,93 | 0,43 | SiO ₂ | 1,27 | 0,97 |
| Al ₂ O ₃ | 0,88 | 0,48 | Total | 96,44 | 100,00 |
| MgO | 1,04 | 0,28 | | | |

Fluorine is present in I, but its quantity was not determined. The mineral contains, moreover, three radio-active elements, ionium, polonium and radium, besides 2 per cent of CeO₂ and La₂O₃, taken together. It is most noticeable that this mineral contains no uranium. It is also remarkable that the deposition of *hokutōlite* is almost entirely restricted to those parts of stones now covered with hot water. Considering that BaSO₄ and PbSO₄ exist only in an extremely small quantity in the water, it is doubtful whether this mineral is really deposited by the present spring water or not.

At the water line, a dirty gray compact crust is formed with no crystal formation and no radio-activity, and of following composition, quite different from *hokutōlite*:—

| | | | |
|--|-------|---|--------|
| SiO ₂ | 77,10 | Na ₂ O | 0,73 |
| SO ₃ | 2,48 | K ₂ O | 0,14 |
| Al ₂ O ₃ | 2,27 | TiO ₂ | trace |
| Fe ₂ O ₃ | 2,11 | P ₂ O ₅ | trace |
| CaO | 0,13 | Ign. loss | 14,94 |
| MgO | 0,30 | Total | 100,20 |

The colour of *hokutōlite* varies from brown or brownish-yellow to almost white. The crystals are in minute thick rhombic plates, about 2,5 mm. in diameter and half as much in thickness; and there are besides the prism and the base, minute pyramidal faces. The plates mostly form a subparallel growth on the basal pinacoid, which is the direction of distinct cleavage. Hardness is about 3,5 and specific gravity nearly 6.

Radio-activity is well noticeable.

Shibukuro, Ugo. On the volcano Yake-yama, a hot spring water is issuing, distinguished by its exceedingly thick deposit of floury sulphur, collected and sold for use in medical baths. The bottom

so this hot water shows scanty deposits of *hokutōlite*, but the mineral forms a deposit of about 6 cm. in thickness at a distance of about 2,640 ft. (800 meters) lower down the creek, where the water is not very high in temperature. Close to the bathing hut, there is a deposit of *hokutōlite* in an old dried-up bed of a creek formerly supplied with hot water. The mineral from the Shibukuro Spring is apparently identical with that of *Hokutō*, but the gray-coloured deposit found on the water line is missing at *Shibukuro*.

Radio-activity is less distinct than that of hokutōlite from Hokutō.

Ryōichi Ōhashi, lecturer at the Akita Mining College, in 1913, using a small piece of *Hokutōlite* found at the *Shibukuro* Hot Springs, Ugo Province, observed the radio-activity of the mineral by the photographic effect.

The specimen is a radial-concentric crust with the nucleus of a conglomerate, with alternate zones of white opaque and dull yellow; and blow-pipe tests indicate the presence of a comparatively large quantity of lead in the former, while in the latter a very small amount of it. The test specimen was cut and polished perpendicular to the concentric shells. After having been sufficiently dried a sensible photographic plate (Lion-special rapid), with the film upwards, was placed on the polished surface of the mineral, a sheet of paraffin paper interposed between the plate and the object.




Fig. 1 shows a photograph of this kind obtained after exposure for three weeks, giving a different impression of different layers of the crust. The photograph of the same section of the specimen, is shown below the radiograph of the same for comparison (Fig. 2).

It may be easily seen from the figure that the outer part of the crust affects the plate generally more sensibly than the inner part, and the white layer presents stronger radio-activity than that of the dull yellow colour. It is noticeable that the mineral has only a weak power of penetration, as the rays emitted are relatively very




Fig 1.

Hokutōlite containing radio-active elements, photographed by its own rays.




Fig. 2.

The same in natural size and state

easily stopped, a single sheet of paraffin paper intercepting them.

If the mineral be applied directly to the plate without any screen, it will be observed that the exposure may be shortened to two weeks in order to obtain the radiograph of the same intensity as shown in the figure, while a piece of red paraffin paper is sufficient to stop their rays almost completely.

Care must be taken for taking radiograph of the mineral to desiccate it well, as even a faint trace of water weakens considerably the sensibility of the plate for radiation.

2. RADIOBARITE.

Shibukuro, Ugo. The white to reddish white *radiobarite* of the Shibukuro Spring is collected only from the heap of rock dug out of the hot water creek with *hakatite*. Crystals are thick rhombic tabular, parallel to the basal pinacoid and generally 2–3 mm. in the longest diameter. The faces observed are the basal, pinacoid, the unit prism and the unit pyramid. Hardness is about 3.

Radio-activity is weak.

3. NAËGITE.

Naëgite is a mineral nearly identical with zircon. The name was given by Tsunashirō Wada after its locality, **Naegi**, in the Province of Mino, where it is collected in tin placers together with *beryl*, *tourmaline*, *quartz*, etc. *Naëgite* is greenish or brownish in colour and forms imperfect globular aggregates, sometimes more than 2.5 cm. in diameter, upon whose surface the terminal faces of crystal individuals are distinguishable. Single crystals varying from 3 to 5 mm. in length are not rare. The fresh crystals are generally highly vitreous in lustre. The crystal habit is often pyramidal but generally prismatic; the faces are the protoprism and a proto-pyramid, besides a very subordinate face of the other prism. Hardness is about 7.5 and specific gravity 4.09. The crystals are not uniform in colour throughout, the terminal portion being often turned to reddish-brown, while the inner part is grass-green. Extinction is straight. Refraction is strong but double refraction is weak. The result of chemical analysis is as follows:—

| | | | |
|----------------------------|-------|--|--------|
| ZrO ₂ | 55.30 | Nb ₂ O ₅ | 7.10 |
| ThO ₂ | 5.01 | Ta ₂ O ₅ | 9.12 |
| SiO ₂ | 20.52 | Yt ₂ O ₃ | 100.73 |
| UO ₃ | 3.03 | Total | |

Radio-activity is strong.

4. ZIRCON.

Zircon in good macroscopic crystals are of rare occurrence in Japan, being found in well developed but minute crystals in the gold placers near Kiirun (Keelung) in Taiwan (Formosa); and as large crystals in a graphite deposit of Chōsen (Korea).

Placers near Kiirun. *Zircon* is found there in minute columnar crystals, not more than 1 mm. in length, and shows the tetragonal prism, and ditetragonal and tetragonal pyramids. Crystals are sometimes beautifully red, but also yellow, brown, gray, violet or colourless. The red colour of the mineral fades away on ignition. Specific gravity is 4.7 and hardness is 7.5. *Distinctly radio-active.*

Shimpyō-ri, Chōsen (Korea). The zircon crystals, which are found at this place, are dark violet-red and translucent, and prismatic in habit, with two kinds of tetragonal pyramids and the unit prism. The individuals attain 7 cm. in length and 2.5 cm. in width. The chemical analysis is as follows:—

| | | | |
|--|-------|----------------------------|--------|
| SiO ₂ | 33.90 | TiO ₂ | 0.00 |
| ZrO ₂ | 63.32 | Residue | 0.20 |
| Fe ₂ O ₃ | 3.33 | Total | 100.00 |

Radio-activity is indistinct.

5. MONAZITE.

Monazite occurs in gold placers and in the mother-rock pegmatite of **Ishikawa-yama** in the Province of Iwaki, as yellow to brownish, semi-transparent crystals of a thick tabular form showing the faces of the orthopinacoid, two domes, the unit prism and the base. The individuals are minute and rarely exceed half a centimeter in the longest diameter. *Radio-activity is well noticeable.*

A red crystal badly developed within a topaz crystal of **Tanokami-yama** in the Province of Ōmi, shows strong similarity to *monazite*. It runs up to 6 mm. in the longest diameter.

6. COLUMBITE.

Columbite occurs in pegmatite as well as in diluvial sand deposits, near Ishikawa-yama in the Province of Iwaki, and in pegmatite at Yamano-o in the Province of Hitachi.

Ishikawa-yama, Iwaki. The black crystals, sometimes attaining the length of 9 cm. and the width of 5 cm., show the two vertical pinacoids prevailing and well developed, besides the base, prisms and pyramids. Cleavage is parallel to the macropinacoid. The mineral is there found embraced by *beryl*, *quartz* or *feldspar*. *Radio-activity is distinct*.

Yamano-o, Hitachi. Some small crystals of this mineral were collected there. The chemical composition is as follows:—

| | | | |
|--|-------|----------------------------|-------|
| Nb ₂ O ₅ | 57.95 | MgO | 2.26 |
| Ta ₂ O ₅ | 22.69 | SnO ₂ | 0.49 |
| FeO | 10.81 | WO ₃ | trace |
| MnO | 7.06 | Total | 99.89 |
| CaO | 1.13 | | |

Radio-activity is indistinct.

7. FERGUSONITE.

The mineral is found in a scanty amount, in the river sand of the environs of Naegi in the Province of Mino.

Naegi. The crystal individual is irregularly columnar, tapering on one end, showing hemihedral faces (the pyramid, prism and the base), usually not more than half a centimeter in length. Colour of the mineral varies from dark-gray to light-gray. The faces are rough and curved, except the pyramid. Specific gravity is 4.3. The chemical analysis gave the following result:—

| | | | |
|--|-------|--|-------|
| SiO ₂ | 2.10 | Yt ₂ O ₃ | 35.60 |
| Nb ₂ O ₅ | 48.39 | Er ₂ O ₃ | 1.04 |
| Ta ₂ O ₅ | 2.95 | Fe ₂ O ₃ | 4.12 |
| UO ₂ | 5.69 | H ₂ O | 99.89 |
| CeO ₂ | | Total | |

Conspicuously radio-active.

8. ORTHITE.

Macroscopically visible crystals of *orthite* are found in the biotite granite of **Mt. Hiei-zan** near Kyōto. They are minute, columnar, almost black, and with a vitreous luster. The observed faces are the base, the orthopinacoid, pyramids and the unit prism. The crystals are as usual elongated in *b* axis, and developed on both ends; they measure sometimes nearly 1 cm. in length and a few mm. in width. Zonal structure is almost wanting, while pleochroism is very strong. *Radio-activity is distinct*.

LITERATURE.

Tsunashirō Wada: "Minerals of Japan", 1904.

Tadasu Hiki: "On the Allanite Crystal from the Hiei Mountain", (Memoirs of the College of Science and Engineering, Kyoto Imperial University, 1903).

Matsutarō Hayakawa and Tomonori Nakano: "Die Radioaktiven Bestandteile des Quellsedimentes der Tennen von Holuto, Taiwan". Zeitschrift für Anorganische Chemie, Bd. 78, 1912.

Hiroshi Yoshida: "Bunte, etc." MS., 1914.

Journal of the Geological Society of Tokyo (Japanese).

Wada's Beiträge zur Mineralogie von Japan.

Chapter XII

ON THE GEYSER IN ATAMI.

BY K. HONDA AND T. TERADA.

PHYSICAL LABORATORY, IMPERIAL UNIVERSITY, TOKYO, JAPAN, 1903.

Different from other geysers, the geyser of Atami in the Province of Izu is characterised by the regularity of the eruption, which consists in alternate projections of hot water and steam, usually five times in succession. The water projected does not return to the orifice as it does in several other geysers. The orifice, which originally opened vertically upward, has been covered by a heap of stones to prevent the dangers caused by the eruption. At present three orifices are exposed, among which one is distinguished as the principal opening. Besides, there is another mouth hidden underground; the water projected by these orifices is distributed to several bath-houses by conduits. The water is of a strong saline taste, containing about $\frac{1}{2}$ per cent. of sodium chloride, *i. e.*, about one fifth of that contained in the sea water. The mouth of the geyser is not far from the sea coast and about 22 meters above the sea-level.

The eruption occurs usually five times in a day and night. During the time of repose, we see only a small quantity of steam rising from the mouth. As the time of the eruption approaches, a rumbling sound is heard underneath. The boiling water appears just inside the mouth. It soon retires and again appears. This state continues for about three quarters of an hour. Next a small quantity of hot water flows out intermittently. This is followed by an intermittent stream of moderate quantity with a longer period. The activity soon attains its maximum. A torrent of hot water gradually increasing in force is torn into a violent splash and projected with a great velocity by steam which gradually increases with the diminishing water. When the roaring sound of steam reaches the maximum, the water almost disappears. The steam now diminishes and is soon followed by the second gush of water. After these states have been repeated five or six times, the activity ends with the last steam which gradually subsides into an inconsiderable amount as seen at the beginning. It takes above two hours from the beginning to the last stage of the eruption. The time of repose is a little less than three hours on the average. These regular recurrences are often interrupted by an abnormal outburst, called *nagawaki*, at which the water and steam come out incessantly for above twelve hours and after which a long repose follows as a rule. In years rich in this anomaly it occurred almost monthly, whereas in the last few years only once or twice.

Our excursion to Atami was undertaken to make detailed observations as to the manner of eruption and if possible to get an insight into the internal mechanism of the geyser. Arrangements were contrived to keep records of the manner of each successive eruption.

A pendulum consisting of a brass rod and a heavy lead ball was hung before the principal orifice. The water and steam, projected nearly horizontally, deviate the pendulum by the impulsive pressure. The motion of the pendulum was transmitted by a cord and pulley to a recording pen guided by two vertical pillars with grooves, in which roll two friction-wheels attached to the penholder. The vertical displacement of the pen was recorded on a cylinder making one revolution every two hours. The pendulum was afterward removed to a position where the impulsive action of vapor and splash was shielded off and the flow of water only could be recorded.

To record the manner in which the steam is ejected, an arrangement was used which was nothing more than an air-thermograph. A small cylinder of sheet zinc was introduced into a mouth neighboring to the principal orifice, where it was possible to find a position such that the bulb was exposed to the heating of the steam only. This orifice, being a smaller branch of the main one, could be considered as representing the main one on a reduced scale. The bulb was connected by a fine copper capillary tube to one of the arms of a U-tube containing mercury. The motion of the mercury meniscus in another arm was recorded on a cylinder by a pen mounted on a float on the mercury and guided by two vertical pillars with grooves and two friction-wheels attached to the penholder. To determine the temperature of the water and steam a maximum thermometer was used.

To take continuous records of the exact time of eruption, and also the general manner of each eruption, a mercury tide-gauge, constructed after Mr. Nakamura's design,* was used. The lead pipe of the instrument was inserted to the neck of the geyser. The pressure of the projected water at the neck during each eruption was recorded on a cylinder revolving once every 24 hours. The automatic records of daily eruptions have been taken during the last two years.

In the following lines, a brief account of our results of observation is given.

I. ORDINARY ERUPTION.

An ordinary eruption consists of three distinct series, termed conventionally the first, the second and the third, differing in period and force and succeeding one another very regularly. Fig. 1 is a reproduction of one of the records obtained by our pendulum recorder. The abscissa represents the time, and the ordinate the deflection of the pendulum by the pressure of water. The




Fig. 1.

dotted line shows the velocity of the steam obtained by our air-thermograph. The first series, which begins the outburst, consists of a small quantity of water appearing with an average period of 1 minute and 40 seconds. After this intermittence has been repeated a score of times the second series follows. A moderate quantity




Fig. 2.

of water comes out three or four times with a mean period of six minutes. The water increases in quantity and force till at last the third or principal series sets in. On the first outburst of the third series we see very often the superposition of the last one of the second series. The third series is to be distinguished from the previous series by its violence and the quantity of water and steam. The sequence of the water and steam occurs usually five or six times with a mean period of about 11 minutes. Fig. 2 is a record obtained by the mercury tide-gauge, and shows regular periodic sequence of eruptions with a mean period of 24/5 hours.

II. ABNORMAL ERUPTION; NAGAWAKI.

The first *nagawaki* recorded by our arrangement began at 4:30 A.M. on January 14, 1905, from the fourth eruption of the third series. During two or three days before the *nagawaki*, the period of successive eruptions seems to have been slightly diminished; but in such a degree that may be found not seldom in our records without leading to either *nagawaki*, or anything extraordinary. The *nagawaki* began, as it were, almost suddenly in the midst of an ordinary eruption. The flow of hot water continued without interruption, gradually decreasing in quantity and mixed with steam. At 7:40 P.M., it came to a sudden repose; at 2:40 A.M. on the 15th, an intermittent flow of hot water resembling the second series of an ordinary eruption began and continued for about three hours. After a repose of four hours ordinary eruptions at last set in, but with the period remarkably shortened and the activity strikingly reduced. The number of eruptions per day was ten, a remarkable contrast to the ordinary frequency of five. The frequency decreased afterward very slowly with the time, and recovered its original value after the lapse of about a month.

The second *nagawaki*, which occurred on May 26, was quite similar to the previous one in its general aspects, though it took place in conjunction with the extraordinary decrease of the general activity.

It is an interesting coincidence that the two *nagawakis* of January and May began at the same phase of the ordinary eruption at nearly the same hour of day, and that a center of low atmospheric pressure was approaching from the Pacific in each case.

III. INFLUENCE OF ATMOSPHERIC PRESSURE.

Examining our records of long observation, we found a remarkable fact that low atmospheric pressure *retards* and high atmospheric pressure *accelerates* the eruption of the geyser. This curious fact may be seen from Fig. 3. Times of the successive eruptions are plotted as the ordinates corresponding to each day which is laid off as abscissa; corresponding points for successive days are connected into five broken curves (b)-(f). Curve (a) represents the daily change of the mean atmospheric pressure. The probable cause of the strange coincidence would be given later.




Fig. 3.

IV. TEMPERATURE OF HOT WATER DURING ERUPTION.

A maximum thermometer was placed about 1.5 meters inside the orifice and the temperature has been read daily since last April. It was found that the temperature at this depth was almost invariably 103° – 104° C. At the orifice, however, it was about 100° C., indicating a rapid cooling of the hot water. Hence it may easily be conjectured how hot the underground water would be at a depth of some ten meters.

V. VELOCITIES OF WATER AND STEAM.

The pressures of water and steam upon the bob of a special pendulum arrangement were recorded on the cylinder, and estimated afterward by the substitution of known weights. The total pressure upon a sphere is known to be approximately equal to one half of that upon a circular disk of the same diameter. Hence

$$p = \frac{\pi r^2 \rho v^2}{2}, \quad \text{or} \quad v^2 = \frac{2p}{\pi r^2 \rho},$$

where p is the pressure, r the radius of the bob, ρ the density and v the velocity of water or steam.

For the water we obtained, at the mouth,

$$v = 1.5\text{--}2.0 \text{ meters per second,}$$

and for the steam

$$v = 18\text{--}24 \text{ meters per second,}$$

the velocities varying within the limits according to the phase of each eruption. In this calculation, we took for the value of ρ the density of saturated vapor under ordinary atmospheric pressure.

VI. QUANTITIES OF WATER AND STEAM.

For the rough estimation of the quantity of water thrown out in an eruption, we measured directly the amount supplied to a number of tanks and calculated the total from the sections of several conduits. The quantity thus estimated amounts to 45 cubic meters.

The rough estimation of the quantity of steam was carried out in the following way:

Let the quantity of steam be denoted by Q . Then

$$Q = \int S_v v dt,$$

when S is the area of the orifice. If V be the ordinate in the curve representing the relation of velocity of steam to time, obtained by our special thermograph above referred to, we may put

$$v = kV, \quad \therefore \quad Q = S_v k \int V dt$$

Since v was determined by the pendulum experiment, and the corresponding V from the thermograph curve, k can easily be known; by calculation, it was found to be 500. S was estimated to be 200 cm.^2 ; for an instance, we obtained

$$\int V dt = 6500, \quad \therefore Q \doteq 500 \text{ kg},$$

the number must be considered as giving only the order of magnitude.

VII. EXTRAORDINARY DECREASE OF ACTIVITY.

During the course of the last few years, several wells have been bored in this district. Most of them give a moderate quantity of hot water only by pumping. Since the last year, the number of wells was greatly increased, amounting to about twenty in all. Sawaguchi's well, bored on March 27 of this year, burst out with great violence, throwing up a column of hot water about 8 meters high. On May 22, another one, Yonekura's well, of greater activity, has been opened within a few hundred meters of the geyser, giving hot water at a rate of 310 cubic meters per day. Two days afterwards another one, Higuchi's well, of not much less activity, was bored. After the boring of the Sawaguchi's a slight decrease in the frequency of the geyser was observed; on May 20 it was reduced to 4.4, though the force of each eruption presented no appreciable change. After the boring of the other two the frequency remarkably decreased; it was 3.6 on May 26. Moreover, the first and

second series of each eruption became considerably longer and the principal series was considerably lessened in force. After the *nagawaki* of May 27, the frequency was temporarily increased; but on June 11, it fell again to 3.2. As shown in Fig. 4, the first and second series lasted for three and a half hours and the third was




Fig. 4.

reduced to only three weak eruptions. Consequent decrease in the quantity of hot water caused trouble to several bath-houses supplied by the geyser, and therefore the above three wells were all stopped—Yonekura's on June 12, Sawaguchi's on the next day and Higuchi's on July 12. The frequency of eruption has gradually increased since then, and in the middle of August, it attained 4.5 which is yet somewhat short of the original value. As for the mode of each eruption, it has quite recovered in force.

VIII. LEVEL CHANGE AND TEMPERATURE OF WELLS.

Level change of two wells near the geyser were recorded by means of Honda's limnimeter. Nomura's well, which is within 200 meters of the geyser and quite high above the sea-level, shows a regular up and down motion corresponding to the eruption and repose of the neighboring geyser. Effects of the tidal and atmospheric pressure are also recognizable, but not very remarkably. (Fig. 5.) The temperature of the water of this well is 50°–60° C. In Suzuki's well, which is more remote




Fig. 5.



Fig. 6.

from the geyser than Nomura's, the effect of the geyser is not observed, but the level rises and falls with the tide. (Fig. 6.) The head of the water is 22 meters above the sea-level. Temperature measurement with a maximum thermometer was made in Abo's well, which is situated in the midway between the geyser and Yonekura's well. The results are tabulated as follows:

| | |
|--------------------------|----------|
| At the surface of water, | 62° 0 C. |
| 23.0 meters below, | 95° 5 |
| 28.5 " | 98° 3 |
| 29.7 " | 104° 7 |
| 31.0 " | 118° 2 |

Thus, a remarkably high temperature was found to exist in the surface layer of the district.

IX. EXPLANATION OF THE PHENOMENA.

Existing theories on geysers fail to explain the exact manner of eruption of the geyser of Atami. After a series of experiments with several forms of models, we arrived at a theory which we hope may fairly explain the phenomena of the geyser in question.

Referring to fig. 7, *A* is a cavity lying in a considerable depth; *a* is the vertical pipe and *b* a canal which supplies the water to *A*. We conceive a side canal *c* intermediate between *A* and *a*, which leads to a second cavity *C*, not shown in the figure. The temperature of the water in *a* and *c* is supposed to be lower than the corresponding boiling point. Water in *A* is heated by the wall of the cavity, the temperature of which is supposed to be decidedly higher than the boiling point at the depth. The source of the heat is probably to be attributed to the hot water and steam running through numerous veins and canals extending in the depth of the district. When the tension of the vapor in the cavity attains a critical value, the water is thrown off and then the steam follows.

When a certain amount of steam is given off, the pressure in the neck is reduced to such a degree that the water flows in from the side canal and stops the eruption momentarily. Soon the downward pressure of the water column is overcome by the tension of vapor and the second gush follows. These eruptions are repeated several times, till the vapor pressure is so reduced as to admit the comparatively colder water from the feed canal *b* and also from *c*. Thus the activity is quenched for a while till the next eruption begins. A model (Fig. 8) constructed according to this view, worked very satisfactorily. The manner of eruption was imitated in many details. The number of intermittences in an eruption increases with the temperature of the water in *c*.




Fig. 7.




Fig. 8.

The phenomena of *nagawaki* may be explained partly by the supposition that the underground temperature is raised above its normal value and so the temperature of the cavity *C* becomes higher than the ordinary case. If the temperature of the cavity *C* in our model be raised to a certain value, the eruption corresponding to *nagawaki* begins. It resembles the actual one even in some details. The cause of this occasional change of temperature is probably the change of the subterranean volcanic activity which keeps the underground temperature in this district considerably above the boiling point of water.

The fact that the frequency of eruptions immediately after *nagawaki*, is nearly doubled, may partly be explained, if we consider that the temperature of the heating cavity was raised during the course of the *nagawaki* by the incessant flowing of superheated water from a great depth. It may be added as a very suggestive fact that if in our model, a quantity of air be blown into the heating cavity, the frequency of eruption increases at first remarkably and gradually decreases with the gradual expulsion of air by successive eruptions; even the weakness of activity in the actual case is imitated with great faithfulness. During a few hours after *nagawaki*, the cavities as well as the canals leading to the orifice remain drained out, so that it is possible that air or other gases may enter into the cavities and cause the increased frequency of the eruption.

It is a common fact of observation that the temperature of some ordinary hot springs rises with low atmospheric pressure. This is undoubtedly due to the increase of flow due to the enhanced circulation of water caused by the reduction of the pressure. If in the supposed heating cavity of the geyser, the interchange of water due to the slow circulation through numerous veins and fissures (not shown in the figure), be accelerated by some cause, the time required for the sufficient heating for eruption must necessarily be prolonged. This consideration seems to explain partly the influence

of atmospheric pressure on the period of eruption above mentioned. Again, the possible influence of well-boring on the geyser, may be explained on the same basis. Such a well may increase the circulation of underground water in its vicinity and result in the retardation of the eruption of the geyser in a similar manner. Moreover, it is quite natural that the hot water should find its easier vent through a new passage opened with a less resistance, at the expense of the quantity originally given out by the old one alone. The prolongation of the first and second series of an eruption, suggests the slowness with which the pressure in the heating cavity approaches the critical value. The careful investigation of the variation of the frequency in connection with the boring and stopping of the wells, leads us to the strong belief that the striking coincidence of the well-boring and the decrease of the activity of the geyser, is a necessary, and not an accidental one. If the frequency of eruption does not yet quite attain its former value long after the stopping of the wells, we need not wonder at all, since some irreversible change in the subterranean mechanism might have happened during the period of the anomaly.

Chapter XIII

A. CLASSIFICATION OF MINERAL SPRINGS

As for the classification of the mineral springs, we have adopted the new system, which was proposed lately by the German chemists, Prof. Dr. E. Hintz and Dr. L. Grünhut in Wiesbaden. The classification is made according to their chemical constitution, based on the modern physico-chemical theory of solution.

According to the system the springs are classified as below: (The quantity of dissolved solid constituents given in this article is always that contained in 1 kilogram of the spring water).

1. **Simple (indifferent) thermals.** Simple (indifferent) thermals may be defined as those which issue from the earth at a higher temperature than 37° C. throughout the whole year, and at the same time are poor in free carbon dioxide and also in dissolved solid constituents. The quantity of free carbon dioxide should be less than 1 gram.

The *simple earthy thermals* may also be included in this division.

2. **Simple carbondioxated springs.** Simple carbondioxated springs contain more than 1 gram of free carbon dioxide, but less than 1 gram of the dissolved solid constituents.

Should springs contain less than 10 milligram-equivalents of ferrous or ferric ions and their presence still give the water a pharmacodynamic value, they may also be included under this heading, as far as the above condition applies.

3. **Earthy carbondioxated springs.** Earthy carbondioxated springs contain more than 1 gram of both free carbon dioxide and the dissolved solid constituents, as anions hydrocarbonate ions predominating and as cations calcium and magnesium ions. (Therefore calcium bicarbonate and magnesium bicarbonate appear as predominating constituents besides free carbon dioxide in the salt calculation).

4. **Alkaline springs.** Alkaline springs contain more than 1 gram of dissolved solid constituents, as anions hydrocarbonate ions predominating and as cations alkaline ions. When these spring waters are boiled, the carbonic acid gas evolves and the hydrocarbonate ions are changed into carbonate ions, which combine partly with the alkaline earth metal ions to form precipitates, while the greater part of the hydrocarbonate ions remains unchanged. Its hydrolysis causes the alkaline reaction of the water. When the quantity of free carbon dioxide exceeds 1 gram, we call the springs *alkaline carbondioxated springs*.

Should chlorine, sulphate, or alkaline earth metal ions be present in a predominating quantity, we call the springs *alkaline muriated*, *alkaline saline*, *alkaline muriated saline*, or *alkaline earthy springs* respectively, instead of *pure alkaline springs*.

5. **Common salt springs.** Common salt springs (*muriated springs*) contain more than 1 gram of dissolved solid constituents, chlorine ions being the predominating anions, and sodium ions the main cations. (Thus common salt appears to be present in the salt calculation as the most predominating constituent).

These springs may be called *simple earthy common salt springs*, when the quantity of chlorine ions as well as that of sodium ions is less than 200 milligram equivalents (or 15 gram of common salt in the salt calculation), but *concentrated common salt springs* when of larger quantities.

When the quantity of free carbon dioxide exceeds 1 gram, we call the springs *carbondioxated common salt springs*. When hydrocarbonate or sulphate ions are present in a predominating quantity, the common salt springs are not called real ones, but *alkaline, saline, or alkaline saline common salt springs* respectively. But when alkaline earth metal ions, or these and hydrocarbonate ions, or alkaline earth metal and sulphate ions predominate, the springs are called *earth-muriated, earthy or sulphated common salt springs* respectively.

6. **Bitter springs.** Bitter springs contain more than 1 gram of dissolved solid constituents, the sulphate ions predominating among anions. Should the sodium, calcium or magnesium ions be the principal cations, the springs are classified as *saline bitter springs, sulphated bitter springs, or real bitter springs*, as the case may be. If chlorine ions be present in addition to one of these three cations, the springs are named *muriated saline, muriated sulphated or muriated real bitter springs* respectively.

7. **Iron carbonate springs.** Iron carbonate springs contain generally more than 0,01 gram of ferrous ions with hydrocarbonate ions.

8. **Vitriol springs.** Vitriol springs contain generally more than 0,01 gram of ferrous or ferric ions with sulphate ions.

9. **Alum vitriol springs.** Alum vitriol springs contain generally more than 0,01 gram of ferrous or ferric ions with sulphate ions. The quantity of aluminium ions exceeds 10 milligram-equivalents.

10. **Acid vitriol springs.** Acid vitriol springs contain generally more than 0,01 gram of ferrous or ferric ions besides sulphate and hydrogen ions.

11. **Acid alum springs.** Acid alum springs may be defined as those which contain sulphate and hydrogen ions, and in which the quantity of aluminium ions exceeds 10 milligram-equivalents.

12. **Acid alum vitriol springs.** Acid alum vitriol springs contain generally more than 0,01 gram of ferrous or ferric ions, with sulphate and hydrogen ions, and more than 10 milligram-equivalents of aluminium ions.

13. **Sulphur springs.** Sulphur springs are those which contain hydrosulphide ions and sometimes free hydrogen sulphide in addition. Whether the springs contain free carbon dioxide, and consequently also free hydrogen sulphide, or not, we call them *hydrogen sulphide springs or sulphur springs* in the strict sense.

14. **Acid hydrogen sulphide springs.** Acid hydrogen sulphide springs contain free hydrogen sulphide with hydrogen ions.

B. KINDS OF MEDICINAL SPRINGS AND THEIR INDICATIONS.*

1. Simple Cold Springs.

Internal use of cold water stimulates peristaltic motion of the stomach and intestines, which causes as a result promotion of stool (evacuation of the bowels), especially when somewhat larger quantities of cold water are taken to the empty stomach in the morning. It is necessary for assisting rapid evacuation of the bowels to favour the action by adequate bodily exercise.

If one aims at the local specific action of waters, they should be taken warm.

The spas with cold mineral springs are therefore recommended to patients, who are disposed to take cold, also reconvalescents on the one hand, and those who suffer nervous, neurasthenic, hypochonderic complaints with congestion, but without any advanced organic disturbance, on the other hand.

Cold baths may be regarded also as one of the fat lowering systems. Adequate diet and the exercise cure are both essential in assisting the action.

Thus they are contra-indicated for the patients, when temperature of the body and fat deposit must be economized, such as is the case with nervous subjects and reconvalescents.

* The content of this section is a summary of the principles of the treatment of diseases by mineral waters, as given in the "Principles of the Treatment of Diseases by Mineral Waters" by Dr. J. J. Fox, 1916.

The actual effects of cold baths must obviously depend on their duration and temperature in each case.

2. Simple Thermals.

The healing property of this class of springs is based only on the action of water and its temperature, it being simply the local influence upon or through the outer surface of the skin.

3. Simple Carbondioxated Springs.

The therapeutical effect of the waters of this class is based on the hyperbolic action of water, as well as carbon dioxide, especially in the sense of local stimulation. The waters are used both internally or externally, but for external use as baths far the more frequently. Some of these waters are exported exclusively as table water, not being employed locally in any way for medical purposes.

Internal use: (1). If taken cold, the water promotes peristaltic action of the stomach and consequently cleansing and clearing it of its contents. (2). Local stimulation by carbon dioxide causes the dilatation of vessels of the mucous membrane and thus leads to an increased secretion and accelerated reabsorption of water thus influencing the amount of nutriment taken up; therefore if drunk in moderate quantity with or preferably after a meal, it promotes nutritive digestive action of the stomach and also drives off oppressive flatulency or sensation of fullness found sometimes after a generous meal.

When taken too frequently or in too large quantities, the gastric juice is diluted too strongly on the one hand, and, also as carbon dioxide leads to importunity, when it causes the stomach to dilate, disturbs its function on the other hand.

For the same reason it is unsuitable to use table water which is impregnated with an excess of carbon dioxide, as is used with artificial carbonic acid waters, unless it is drunk after dilution. When taken moderately, the water is beneficial in alleviating slight disturbance of digestion, atony, dyspepsia, catarrhs of stomach and the upper intestines, and also for patients suffering from urinary, kidney and bladder diseases, as well as disturbance of assimilation, gout, uric acid diathesis, and poisonings, etc. As for the action of elimination it surpasses the simple waters, both cold and warm, owing to its influence in accelerating the absorption of water. It is advisable to associate the milk cure in the first place with the treatment, taking some milk suitably warmed with the spring water.

4. Earthy Carbondioxated Springs.

These waters can be used as carbonic acid baths, owing to their content of free carbon dioxide, like the simple carbondioxated springs, when they are protected against loss of carbonic acid at heating. The presence of earth-alkaline metals invests them also with a feeble astringent property, which proves to be useful for the treatment of some diseases of the skin.

In the case of internal use also the action of carbonate of calcium and magnesium equally joins in those of water and carbonic acid, just as with the simple carbodioxated springs. They are recommended for chronic diarrhoea, gout, urine concrement, catarrh of the bladder and urinary passage.

5. Alkaline Springs.

The nature of the action of these baths is not well known or understood up to the present.

When the water contains a considerable quantity of carbonic acid, it stimulates the skin, as is the case with simple carbondioxated springs, and suitable temperature of bath is often beneficial for the catarrhic state of the female genital organs. The drink cure is contra-indicated for loss of vitality. The use of alkaline beverages is generally injurious when a little amount of acid is present in the gastric juice and the disposition of the stomach and intestines to catarrh is observed, such as in atonic dyspepsia of anaemia, chlorose and convalescence. Against phthisis pulmonum, malicious tumors, and every kind of invalidity, the use of stronger alkaline waters should absolutely be avoided. It is stated that the continual use of such mineral waters in large quantities may cause emaciation, atonic dyspepsia and at last cachexia even upon healthy person. The habit of many gouty patients to drink strongly alkaline waters for a long period as a diatetic beverage is therefore dangerous.

Consequently the quantity of water which is to be taken, and the duration of cure should be controlled by spa physician. On the contrary, weak alkaline waters containing carbonic acid gas can be used as a daily beverage for healthy person.

It may be, however, not hygienically advantageous, when such mineral water is impregnated with too large quantities of carbonic acid gas for such to be used as a table water. The volume of gas needs not exceed one and a half to one of water, the best waters of this class containing this amount of carbonic acid in the natural state; if three to four volumes of carbonic acid be added to the water, such a product must be regarded as injurious. The stomach of the drinker becomes swollen and by habitual use permanent dilatation of the stomach may be the result.

6. Common Salt Springs.

Baths of common salt springs of 0.5 to 25 per cent common salt content may be used with good result for: mal-nutrition, diseases of the blood and lymph, diseases of the bones and medulla, swelling of the spleen and the liver, exsudate formation, compensatory disturbance of the circulatory system, habitual abortus, gout, rheumatism, general adipositas, diabetes mellitus, a large number of diseases of the skin and scar formation.

For the drink cure also secondary constituents such as lithium, iron, Glauber's salt, etc., must be taken into consideration. The common salt springs, which are suitable for drinking (of about 25 per cent common salt content), have stimulating influence on the secretion of gastric juice and act in accelerating the proteid digestion. In the same manner also the motility of the stomach is stimulated. Moreover the common salt can be easily absorbed and acts then diuretically, while waters containing common salt have the further action of promoting peristaltic of the intestines and seem to act with stimulation upon the bile secretion, thus proving to be advantageous for bile stasis in the liver.

Hence the common salt springs are recommendable for cases of mal-nutrition, disturbance of the motility and secretion of the stomach, retarded function of the intestines, further for a large number of metabolism-anomalies, for which also other secondary constituents are probably efficacious. Besides these, the drink cure may be used for such afflictions as swelling of the liver and the spleen, inflammatory diseases of pancreas, and in the first place for exsudate formation in the body cavity.

Another curative method adopted at common salt spring resorts is by inhalation. It is recommendable for every form of chronic catarrh of respiratory organs, chronic bronchitis, catarrh of larynx, secondary diseases after catarrh of lung, callosity formation, atelectasis, and lastly also for scrofulous diseases of conjunctiva and chronic diseases of the nose and pharynx (rhinitis, ozaena). For the last mentioned diseases the nose douche, and gargling are also advisable.

7. Bitter Springs.

By the use of waters of this class the evacuation of the excrement increases and the dark colour of the latter proves a strong secretion of bile. When the drink cure is stopped, the evacuation of the bowels usually decreases very rapidly, so that constipation follows usually for a few days after the use of the water, while the increase of urine, which takes place during the use of bitter waters, also has its reaction lasting a few days, after the taking of the bitter waters ceases. The secretion of urea and sodium chloride increase, but that of uric acid on the contrary decreases. The influence of the water upon the mucous membranes of the stomach and intestines, *i. e.* the stimulation for their secretion, may be voluntarily regulated by the quantity of water; if taken in smaller quantities (250 grm.), the action is weakly diuretic, but in larger quantities (500 grm.), very strongly. Larger quantities of the water, if taken for a long duration, may cause sometimes heavy dyspepsia, or catarrhic phenomena, while smaller quantities do not disturb the function of the stomach, but may produce even an accelerating influence on its mobility and absorption, but results are somewhat different from the above when the waters are taken in much smaller quantities.

In the use of less than about 50 grm., though taken a few times daily, the bitter waters can be absorbed, but sometimes cause constipation even in the case of already existing catarrh of the intestines. A part of the sulphates decompose in its passage into sulphides and hydrogen sulphide. The recommendation of the waters to suit each particular ailment differs according to the contents being sodium, calcium or magnesium ion, or chlorine ion present in predominant quantities. As a

general rule the taking of the bitter waters are advisable for habitual constipation of a strong and well nourished person, nervous stages of abdomen and bowels with their secondary condition, and for excessive fat formation; equally a number of nervous disturbances which accompany congestion may be favourably treated. Also diabetes and gout may be benefited by the drink cure. In the case of these diseases, especially diabetes, the defective power of the organs for oxidation finds considerable assistance, evidently due to the influence of the sulphates. Smaller quantities of water can be used with greater frequency over an extended period in cases of stasis of liver, and for laziness of intestines during pregnancy when organic heart trouble is present.

8. Iron Springs.

The therapeutically most important constituent of the iron springs is, as the name denotes, of course iron. It is almost equivalent in its action in organism, whether it is contained in the form of ferric or ferrous ion in water. Iron spring waters are now regarded as one of the most suitable forms of introducing iron into the system, as we find by practical experience that it is able to supply the requisite iron to the blood by means of natural spring water for a longer duration without any interference of digestion or disturbance on the part of the stomach, and thus preferable to any other form of iron tonic remedy. Besides the iron the presence of free acids in these waters must be taken into consideration as highly valuable and efficacious constituents. These are hydrocarbonate ion and free carbonic acid, in the case of vitriol springs sulphate ion and in some few cases besides these, free sulphuric acid.

Iron carbonate and vitriol springs are used therapeutically for both drinking and bathing.

The drink cure is advocated in the first place for various kinds of anaemia and chlorosis, furthermore the iron waters containing arsenic are used for chronic nervous diseases and functional neurosis: hysteria, neurasthenia; the local action of the waters upon the mucous membrane of the stomach and intestines being beneficially utilized for chronic dyspepsia, and the waters of vitriol springs are specially applicable in cases of chronic diarrhoea. As in the bath cure the absorption of iron (or arsenic) into the body through the skin is naturally out of question, only the carbonic acid free or set free is under consideration as an active agent. For this reason the water to be used for bathing purpose should be carefully warmed. They are equally as with baths containing carbonic acid recommended for cases of functional neurosis, *viz.* neurosis of the heart, affections of the female genital organs, especially for irregularity or period disturbance, chronic inflammation, disposition to abortus, functional sterility and affections of the masculine genital organs: Spermatorrhoea, pollution, impotence caused by weakness.

The bath cure with vitriol springs containing no carbonic acid is applicable only for treatment of diseases of the skin when the waters contain a large quantity of arsenic.

9. Sulphur Springs.

All sulphur waters are easily decomposable. Though clear greenish or bluish when pouring forth from springs they quickly become cloudy on atmospheric contact and then sooner or later appear milky owing to deposited sulphur produced by the oxidation of hydrogen sulphide or sulphides. Even in closed bottles oxidation follows in a course of time. In consequence of this quick decomposition sulphur waters should be drunk at the spring, or taken only when fresh bottled.

In most sulphur springs the sulphur discharge varies in its proportion. The cause of this unstableness may be due to the variation of the pressure of the atmosphere. Under a higher pressure of air more hydrogen sulphide gas remains in the water than under a lower.

The origin of sulphur springs is various and in its details still only a little explained. In contrast with the sulphur content in thermal springs from younger formations, that in hot sulphur springs issuing from the crystalline schists, and between these and the transition formation, *i. e.* from granite, schists or crystalline limestones, may be derived from the decomposition of metal sulphides by hot water. Hydrogen sulphide in sulphur springs, both cold or warm, from younger formations, with more or less gypsum beds containing fossil remains of extinct plants or animals, is probably formed by decomposition of sulphates by organic matter in the wet way. Especially, when water containing calcium sulphate comes into contact with organic remains for a long time do we find organic

substances are oxidized at the expense of oxygen in gypsum and changed into carbonic acid. On the other hand, the sulphate is reduced to calcium sulphide by the carbon and hydrogen in organic substances. One part of calcium sulphide is, however, decomposed in the presence of the carbonic acid and water; and calcium carbonate is produced and the hydrogen sulphide liberated.

As all these decomposed products actually occur in sulphur springs from younger formations rich in fossil remains, we accept the afore mentioned explanation for the formation of hydrogen sulphide.

Sulphur waters are for drink cures, baths, douches, and also as a supplement of mud baths and inhalation.

The physiological action of sulphur waters by internal or external use is not yet sufficiently investigated; with regard to this subject we are directed for the most part to supposition and hypothesis.

The indications of sulphur springs concerned has been found quite empirically. In the drink cure with a sulphur spring water it seems to be of no importance whether sulphur is contained in the form of free hydrogen sulphide or of sulphides. In both cases sulphur is absorbed from the stomach and intestines. In the use of water for baths it is assumed that hydrogen sulphide enters into the body through the skin. In the blood, iron sulphide will be, it is assumed, formed through the iron of the blood and consequently a normal reformation of blood globules, formation taking place constantly, be quickened and the assimilation stimulated. At the same time a strong influence takes place upon the liver, the bile secretion being greatly increased. One part of the sulphur compounds, which are entered into the blood, is oxidized by oxygen which is present in the blood in abundance and this forms sulphates thrown off later in the urine.

Not all hydrogen sulphide absorbed is decomposed in the body, a small part of it departs unchanged through the skin and lungs. The evident influence on the respiration and the blood circulation is only partly attributed to the local action of gas upon the lungs. It seems that hydrogen sulphide acts by the mediation of the vagus, on the nerve centres of respiration and blood circulation.

With excessive application of strong sulphur waters, anaemia and weakening of the heart action may be the result. The therapeutical action of the drink cure with strong sulphur waters for furunculose is believed to be probably based on its antiseptic action. By the secondary constituents of sulphur waters (calcium, silicic acid, etc.), mineral nutriment can be given to the system.

One drinks from 100 to 1,000 ccm. of water daily this being divided into morning and afternoon potions and taken cold or warm, sometimes with warm milk. One should drink slowly and with long pauses and also take some form of physical exercise. One soon gets accustomed to the taste and smell of the water.

The baths are taken at the temperature of 33° to 36° C. and of different duration, between 10 and 40 minutes. Prolonged baths for three hours as are usual in some health resorts in Switzerland are not given in Germany, although it is said that they have an antimicrobial effect and so the prolonged sulphur baths have had high reputation for ages as a valuable remedy for wounds and ulcus, but we would not recommend prolonged baths in this water. After every sulphur bath, one hour's rest in bed is advisable.

One bathing course consists of 21 to 28 baths; but as patients always take a rest after several bathing days, 4 to 5 weeks are calculated for a cure.

For douching it is best to use a movable pipe, by which one can make the sulphur water act upon any part of body one may choose and under any desired pressure (as at Aachen, Nenndorf and other baths).

The douches are given in various degrees, moderate or weak, and at the same time lukewarm, warm or hot, and usually combined with sulphur plunge baths.

In the sulphur water inhalation it is not necessary to spray fluid upon the body, but merely to take in the liberated gases thrown into the room. Therefore the apparatus seen in celebrated sulphur gas institutions is an arrangement consisting of a flat fountain basin with a sulphur water spring in the centre, rising up in many fine sprays; the agitated particles falling giving off gas to the atmosphere. In such an inhalation house the patients remain usually from a quarter to one hour.

According to York Schwartz, who analyzed the air of the sulphur gas inhalatorium at Nenndorf (*Archiv der Pharmazie*, Bd. 26, S. 761, 1888) the sulphur is not, as one formerly thought, contained

in form of hydrogen sulphide, but as thiosulphuric acid. Schwartz explained the chemical process of the decomposition of sulphur waters as follows :

The main part of hydrogen sulphide will be oxidized to thiosulphuric acid, probably because hydrogen superoxide, in consequence of the strong evaporation of water produced in the inhalation room, acts as oxidizing agent. Schwartz found on an average 1.5 gm. thiosulphuric acid in 1 cubic metre of inhalation air, but only a small quantity hydrogen sulphide on the contrary. When sulphur water was sprayed around, the characteristic smell disappeared in the inhalation room, all hydrogen sulphide becoming quickly transformed into water and sulphur in consequence of this fine dispersion, without producing any intermediate product.

The sedative effect of inhalation, especially for coughs and dyspnoea, is attributed to the other spring gases, such as carbonic acid, nitrogen, etc., but its antibacterial action is due to the thio-sulphuric acid produced from hydrogen sulphide. Therefore we hold the inhalation of sulphur water especially good for infectious catarrhs of the respiratory organs.

For sulphur gas baths, small rooms heated to 28° C., are filled with sulphur gas by means of the hydrogen sulphide fountain, in the same manner as above mentioned. The nude patient sits on a cane chair, so that the gas may operate freely on all parts of the body surface. Such baths are especially prescribed for itching cutaneous diseases, sensibility neurosis, chronic catarrh of the respiratory organs and asthma nervosum.

In some sulphur thermals, sweating boxes and inhalation rooms are provided, being filled with vapour direct from springs. Gargling and nose douching with sulphur water is of great value for several chronic diseases of the mouth, pharynx, and nose.

The high credit of sulphur waters as a remedy is founded on old experience.

For a combined drink and bath cure, under circumstances in connection with sulphur gas inhalation or sulphur gas plunge baths come into consideration :

Gout (arthritis uratica) in every stage and in all forms, as Hufeland has stated, "perhaps the good effect lies here in the influence of sulphur water on the liver," which may be considered as the place of the uric acid formation.

Congestive hyperaemie of the liver, where swelling of the affected organs can easily and quickly be reduced by a sulphur water drink cure.

Formation of biliary calculus, as a much richer and diluter bile flows during and after the drink cure.

Haemorrhoids and abdominal plethora.

Diseases of respiratory organs, especially old cases of catarrh of the nose, pharynx, larynx, trachea, and bronchus, besides bronchial asthma, and even the first stage of phthisis pulmonum.

Subacute and chronic rheumatism of the joints and muscles, for which the sulphur mud bath is especially effective.

Arthritis deformans.

Neuralgia of various nervous stems and branches (ischias, lumbago, intercostal neuralgia, prosopalgia, etc.).

Neurosis, as chorea minor (chorea St. Viti) of rheumatic origin, neurasthenia and hysteria.

Paralysis of a rheumatic, goutic, and traumatic nature, also paralysis deriving from mercurialism ; but not such, where organic destruction in the brain or spinal cord underlies. In the first stage of tabes dorsalis sulphur mud baths are of value in alleviating lancinating pains, but they cannot affect the course of the illness or prevent its development.

Blood dyscrasia. Several cases of chlorose, especially those where the iron treatment proves to be ineffective, are beneficially treated with the sulphur water drink cure.

Chronic metal poisonings of mercury or lead.

Diseases of the skin : Sulphur baths have stood for years in high reputation for these diseases. The best result is to be obtained especially for chronic eczema (wet and dry lichen), acne vulgaris (face pimples), acne rosacea (copper pimples, copper nose), psoriasis vulgaris, pitiosis versicolor, furunculosis (disposition to furunkel and carbunkel formation), atonic ulcer crusis. On prurigo, pruritus cutaneus (skin itching) and ichthyosis an improvement is noticeable.

Diseases of women : Especially for catarrhs of uterus and vagina, disturbances or irregularities, perimetritic and parametritic exsudates, chronic ovarian inflammation, metritis and endometritis,

adhesive process in Fallopian tube and ovary, and also for complaints arising during klimakterium, sitz-baths and plunge-baths of sulphur water and sulphur mud baths are of great benefit.

Complaints which arise after wounds of soft parts of bones through shot, bruise or sting, suppurated wounds, bone pains after fracture, painful scars, etc.

Bone diseases.

Syphilis and their sequela. In these cases sulphur cures are combined with inunction treatment with mercury.

10. Alum Springs.

It is quite noticeable that hot springs containing comparatively large quantities of alum are not seldom met with in Japan, especially in volcanic districts. In most of these waters are present also ferric or ferrous compounds, but some of them contain only a trace of iron.

Alum waters are used internally for acute and chronic catarrhs and haemorrhage of the stomach and intestines, anaemia, neuralgia, haemorrhoids, etc. They are indicated as baths for catarrhs of mucous membrane of the female genital organs, mal-nutrition, rheumatism, and convalescence after various diseases.

Baths of alum waters are given at *Kusatsu*, *Myōban-Onsen* near Beppu, etc., in Japan.

11. Acid Springs.

Certain thermal waters, particularly the geyser springs of New Zealand and America, are distinguished by the presence of free acids. Free hydrochloric acid is present in considerable quantities in the Lemonade Spring, in California, at Tuscarora in Sour Spring, and in some of Canada.

Japan is rich in acid springs which contain considerable quantities of free mineral acids, with or without iron compounds. Those thermals, which issue in volcanic regions, belong to this class in most cases. *Kusatsu*, *Nasu*, *Gōra* in Hakone, *Shibu*, *Noboribetsu*, *Kirishima*, *Unzen*, *Aso*, *Hokutō*, etc., are good examples of acid waters.

One of the most remarkable of acid springs is to be found among the Oak Orchard Springs, in New York State, which is said to contain free sulphuric acid in the proportion of 40 per cent. of its total mineral content. The same acid exists, with or without hydrochloric acid, in the "acid sulphur waters" of Rotorua (New Zealand), to the amount of from 4 to 22 gr. per gallon. These waters and geyser springs of the Yellow Stone National Park (America), and of Iceland, contain certain large quantities of silicic acid. There is in Europe nothing comparable to this class of waters. They are not employed internally. In the bath their action is stated to be powerfully rubefacient and decongestive.

C. KINDS OF BATHS AND THEIR USE.

1. Subthermal Baths.

The temperature of the baths is somewhat near the point of thermal indifference 32.2° to 34° C. (90° – 93° F.), ranging between 36.6° and 26.7° C. (98° – 80° F.), or a little lower. Moreover the climatic character, the traditions and genius of the spa, promote and intensify the sedative action of the bath. *Schlangenbad*, *Néris*, *Bains-les-Bains*, *Baden-Weiler*, *Ragatz*, *Plombières*, *Buxton*, are stations of this kind.

Subthermal baths are both sedative and stimulating, and have a very large place in spa treatment. They withdraw heat to a moderate extent, and by reducing peripheral tone invite the blood into the extremities and the skin. The cardiac action is increased in force and markedly slowed; nervous excitation, whether peripheral or central, is diminished by the massive and equable impression of the water. These baths are, therefore, indicated in irritability of the vascular and nervous system, as well as in general debility and motor weakness, and in peripheral and insipient and reflex paralyses, in hyperaesthesia, nervous insomnia and neuralgias. They favour nutrition, moderate senile changes, and are much used in degenerative arthritis, atonic gout and rheumatism, and nervous exhaustion and convalescence.

Prolonged Subthermal Baths.—Immersion in waters at the point of thermal indifference may be prolonged for many hours or even day without discomfort. At the sub-alpine spa Loèche-les-Bains (4600 ft.) baths at 33.7° to 23.9° C. (93° – 75° F.) are commonly prolonged for from one to six hours; and similar baths are taken at Nérís (1150 ft.), Schinznach (1140 ft.), Bagnères de Bigorre (1800 ft.), at La Bourboule (2780 ft.), at certain subthermal spas in Japan such as *Yunogō*, *Tochiomata*, *Shimobe*, etc., and elsewhere. Their effect is both sedative and intensive.

Besides their value for some intractable skin affections, they are useful in a variety of nervous disorders, visceral, and especially entero-colic neuralgias, sciatica, spastic arterial hypertension, and occasionally in exophthalmic goitre.

2. Hyperthermal Baths.

Such baths range in temperature from 40° to 54.4° C. (104° – 130° F.). They are of necessity so limited in duration that the late and enervating effects of heat should not be developed.

Hyperthermal baths freely open the surface arterial system, accelerate the pulse and, after an initial rise, diminish the blood pressure. Always taken with very short immersions, one, two or three minutes, they constitute a very useful means of treatment for many cases. The hyperthermal baths supplies the powerful primary stimulation of heat, without its dangerous reactions.

Japan and New Zealand furnish typical examples of natural hyperthermal baths. At *Kusatsu* the temperature of the water is from 54.4° to 71° C. (130° – 160° F.). It is reduced by splashing to 48.9° C. (120° F.). Hot water is poured over the head before entering the bath, and the bather remains in the water for three or four minutes. The bodily temperature rises to 40° C. (104° F.) in six or eight minutes, and returns to normal in half an hour. Cold affusions are used after the bath. Hundreds of syphilitic and leprous patients are submitted to the treatment.

At Teplitz, Hammân-Meskoutine, and elsewhere various degrees of high temperature treatment are employed. It should be applied in practice with much care. The brief intensive and stimulating effect of these baths is helpful in many of the conditions enumerated in the last section, particularly in torpid skin and circulation. The hyperthermal footbath (*pediluvium*) is a valuable form of derivative treatment in many chronic ailments, especially in poor circulation with weak heart, in neuralgia, catarrhs, and constipation; and is insufficiently employed at the British spas. These and other hyperthermal applications induce locally arterial hyperaemia and raise the temperature of the part. They stimulate the oxidizing processes in chronic disease and may be used with much benefit where, as in arthritis, the arterial circulation is defective.

3. Vapour Baths (Ital. *Stufe*; Fr. *Étuve*).

Steam baths, natural or artificial, have been used in medicine from ancient times. The Romans utilized the vapour of the hot springs in their *thermae*; and there are many natural vapour baths, like the cave at Monsummano (temperature 33.3° to 35° C. (92° – 95° F.)). In the island of Ischia there are hotter *stufes*, such as San Lorenzo 51° C. (124° F.) and Castiglione 56° C. (133° F.), and a similar one at the Bagni di Nerone, near Pozzuoli.

The ancient Roman hyperthermal vapour bath is still in use at Plombières. Warm aqueous vapour softens the skin, induces perspiration and increases the fullness and frequency of the pulse. Both the cutaneous and pulmonary capillaries are dilated. Hot vapour baths soften and stimulate the harsh, hard, thickened, and inactive skin. In sensitive subjects the regulation of temperature is essential. When a sedative and tranquillising influence is sought, the moisture must predominate over the heat (as at Monsummano). It may be stated as a general law that the heat of every thermal application should bear an inverse proportion to the febrile excitement or nervous sensibility of the patient.

Assolini used vapour baths below blood-heat in inflammatory affections of the throat and respiratory passages. In chronic catarrhs a more stimulating vapour—Plombières, Mont Dore, or Ems—can be employed.

Natural, or artificial, vapours may also be applied locally, as at Aix-les-Bains and Luchon (the “Berthollet” bath) and at Bath and elsewhere. In Japan such baths are found at *Beppu*, *Atami*, *Shima*, *Nakabusa*, *Naruko*, etc. In addition to the indications already mentioned, vapour baths are

helpful in painful rheumatic and gouty affections (lumbago, sciatica, and neuralgia), and generally in the disorders for which the more stimulating and less sedative hot air bath is employed.

4. Sea Baths.

The sea bath comprises several factors. There is, first, the proper effect of the salt water at the natural temperature, varying according to season and the coast. Then the element of movement in the water.

This again varies from the gentlest undulation to strong mechanical stimulation, equivalent to a rapid succession of douches. The constant movement of water, the impact of the waves and the wind are all to be taken into account. It has been ascertained that the loss of heat in water moving at half a meter per second is one-third more than in a bath of still water. On the other hand, the deposit of salt on the skin from a hypertonic bath inhibits the evaporation of sweat *after the bath*, and this, in addition to the salt stimulus of peripheral nerves, makes the risk of "catching cold" much less after a marine than after a fresh-water bath. Lastly, the local climatic influences modify the effect, in the sense of augmenting or diminishing the stimulus. The sea bath causes an energetic primary contraction of the skin vessels with the "first chill." The circulatory and thermic reactions are more rapid than with ordinary water. Hayem has well said that the reaction is the "therapeutic event most desired and sought for." It is, therefore, important for the bather to leave the water before the "second chill," indicating the failure of reaction, occurs. Sea baths are believed to diminish the elimination of uric acid but to increase that of urea. All the vital actions are stimulated; body weight is increased. It follows that marine baths have their chief medical opportunity in lymphatic and scrofulous affections, local tuberculosis of glands, bones, joints, and in whatever conditions tonic salt baths are indicated. They are inadmissible in cardiac and rheumatic and gouty cases, in renal disease, bronchitis, emphysema, asthma, and in haemorrhagic phthisis and in other congestive states.

5. Effervescent Baths (Carbonated or Gaseous Baths, Luftperlbäder).

Natural effervescing waters owe their acidulous quality to varying proportions of carbonic acid gas. This gas, dissolved under a certain pressure, is uniformly diffused through the water, and exists partly in the dissociated form as ions. When the pressure is reduced, as in the bath, the gas is liberated and attaches itself in minute bubbles to the skin.

Among the thermal waters of this class are: Nauheim, in the Grand Duchy of Hesse 27,8° to 35° C. (82°–95° F.), which contains 2 to 3 per cent. of chloride of sodium; Oeynhausen, in Westphalia 25° to 32,8° C. (77°–91° F.), possessing a muriated water very rich in carbonic acid gas; Soden in the Taunus, and Salins-Moutier, at 35,6° C. (96° F.), with 1,3 per cent. of chloride of sodium. At Royat, in the Auvergne mountains, the Saint Mart and César Springs furnish a valuable thermal effervescing bath, slightly alkaline and salt. Châtel-Guyon, in the same district, has an effervescent alkaline muriated water at 31,7° C. (89° F.), which is employed for "flowing baths."

Among the cold waters there are effervescing chalybeate springs at Spa in Belgium, at Schwalbach, and at St. Moritz; and "gaseous iron baths" are to be had at these more or less tonic stations at subthermal temperatures. Also at Kissingen in Bavaria, Tarasp-Schuls in the Engadine, Marienbad in Bohemia, and elsewhere. *Yunogō* in Japan is one of the baths of this kind.

The peculiar effect of effervescent baths is due to the contained carbonic acid gas, and is to be explained by the fact that the specific heat of this gas is one-fifth that of water whilst its conductivity for heat is only one-fiftieth that of water. The point of thermal indifference for the human skin which is about 33,9° C. (93° F.) for water, is only 23,9° C. (75° F.) for carbonic acid gas, and in an effervescing bath the indifferent point must, therefore, lie somewhere between these two figures.

As a rule, the cooler the water the more gas it will contain, and it is owing to the stimulation of the gas that such baths can be taken much cooler than ordinary baths. In an effervescing bath of say 32,2° C. (90° F.), the body is exposed to thermic stimuli both of cold and heat, coexisting side by side on innumerable minute contiguous areas of the skin and constantly alternating upon the same area, the cold stimulus taking effect where the surface is in contact with the water and

the hot stimulus from the bubbles of gas. Some of the carbonic acid gas is absorbed; and, acting upon the sensory nerve endings, and directly or indirectly upon the blood vessels, alters and increases the circulation. On entering an effervescing bath below skin temperature there is, therefore, a momentary chilliness, soon succeeded by a sense of prickling or warmth, due to the collection of innumerable bubbles on the skin. The increasing feeling of warmth soon overcomes the feeling of cold, because the relative heat of the carbonic acid gas [Indifferent point 23.9°C . (75°F .)] more than neutralises the relative coolness of the water. The vaso-constrictor action of the water is an antagonised by the vaso-dilator action of the gas, the arterioles relax, and there is a more or less vivid reddening of the surface. The activity of the surface action is shown by the comparative pallor of those parts of the body not immersed in the bath.

Following the direct effect upon the skin, there is an indirect result upon the distribution of the blood, known as "decongestion of the viscera," in obedience to the law of inverse relation between the circulation of deep and superficial parts. The heart's action becomes less frequent to the existence of perhaps four to six or ten beats less per minute, the systole stronger, and the pulse more ample, from dilatation of the artery. The effect of the blood pressure varies. It is reduced in baths above the skin temperature, where the vaso-dilator action of the gas has overcome the vaso-constriction of cold, and in all cases where the peripheral vascular dilatation is marked. At lower temperatures, any below 31°C . (88°F .), the blood pressure may be raised, but the action is not the same in all persons and depends partly on the duration of the bath. Many observers have noted by percussion or by the X-rays, a diminution in the cardiac area, especially on the right side, after effervescing baths.

The total effect of these baths depends not only on their temperature and duration but on their saline and gaseous contents. A brief bath is usually purely stimulant whilst a prolonged bath is sedative and tonic. Diaphoresis and diuresis are reflexly encouraged. The effect on the nervous system is that of a mild, continuous subthermal stimulation, passing into sedation. Owing to the thermic stimulation of the gas these baths appear to be warmer than they really are. Consequently their action may be said to be twofold: (1) they favour an increased loss of heat, corresponding to their actual temperature and so induce the direct and indirect results of cooling baths; (2) they produce an active dilatation of the surface vessels such as results from thermal baths, but without application of positive heat or any elevation of the temperature of the body. The merit of effervescing baths, therefore, is that they combine the tonic and antipyretic effect of cool temperatures with an active peripheral stimulation. The presence of sodium chloride with the gas accelerates and enhances the peripheral effect. In addition to cardiac affections effervescing baths are employed for the sedative and tonic action in many conditions of nervous weakness, in the convalescence from influenza and other debilitating illness, in Graves's disease, also in anaemia, amenorrhoea, and congestive dysmenorrhoea. They are contra-indicated in advanced aortic disease and cardiac dyspnoea, but may be used with good results in many cases of angina pectoris.

Peat Baths (Moor- und Schlamm-bäder).

Peat, or moor, baths are prepared from the brown organic deposit underlying, often to the considerable depth, the more fibrous "fuel peat" in many northern peat mosses or bogs. This material consists largely of humus derived from the decomposition of sphagnum moss, and contains resins, silica, free sulphuric acid, and generally carbonate of iron and various salts. When exposed to the action of the heat, formic and other volatile acids are produced. Peats are usually classed as ferruginous, saline, or sulphuretted; and at many spas are fortified by mixture with the mineral water or with "mother lye."

During the last century the use of these baths has been especially developed in Bohemia. They were first used at Franzensbad, and employed there by Dr. Pöschmann in 1810 to 1815. Not long afterwards they were introduced at Marienbad. They are now given extensively in Germany, at Pyrmont, Schwalbach and other chalybeate baths, at Spa in Belgium and at Ronneby in Sweden. They were first employed in Great Britain in 1889 at Strathpeffer, and afterwards at Harrogate and Buxton.

The area of the peat deposits at Franzensbad is near a little extinct volcano outside the town. The moor is traversed by mineral springs containing iron. When first cut in the winter time it has

a greyish lardaceous appearance with an odour of sulphuretted hydrogen. After cutting it is exposed to the air for several months, during which time the sulphides of iron are converted into sulphates and the moor acquires an acid reaction. The "iron mineral-moor" of Franzensbad is said to contain as much as 32 per cent. of protosulphate of iron. It is friable, of a dark brown colour with a bituminous odour and an inky taste. When dry it is ground in a mill, treated with hot mineral water and stirred by hand. The temperature is adjusted by injecting steam into the bath. A single bath is said to contain nearly one pound of formic acid and other volatile substances.

Dr. Paul Cartellieri, the first writer on peat baths (1841) proved that the heat-conducting power of peat was less than that of water. For this reason the peat bath given at a temperature above the point of thermal indifference 33.9°C . (93°F .) does not feel so hot as water, and if given below that point does not abstract heat to the same extent as water does. The specific heat of peat is much lower than that of water. Upon entering the bath the layer of peat nearest to the skin rapidly cools to about the skin temperature. In the case of both mud and peat baths there is no constant changing of the particles which are in contact with a body. The layer next to the skin maintains a more or less constant temperature during the whole period of the bath, and the presence of this comparatively cool and non-conducting layer moderates and equalises the effects of heat. The point of thermal indifference for the peat bath is considerably higher than that of water, and has been placed by Kisch as high as 38.9°C . (102°F .). To the thermal effects must be added the stimulation due to constant chemical action taking place at the surface of the body, the astringent effect of the iron compounds and the antiseptic action of some of the acids. Lastly, the weight and comparative immobility of the peat or mud bath and the friction of the semi-solid poultice-like mass exert certain pressure on the surface circulation and impede muscular movements, especially those of respiration.

In suitable cases peat baths may be given at temperatures ranging from 37.8° to 44.4°C . (100° – 112°F .), and the higher temperatures are well borne if the bath is followed by cold affusion, as at Strathpeffer. When the heart's action or breathing are at all embarrassed, and always in baths of high temperature, the peat should not cover the front of the chest. A cold compress should be always applied to the head. Half baths or three-quarter baths are in these circumstances to be preferred. Local applications of peat, at from 46.1° to 48.9°C . (115° – 120°F .) are often used, principally as a poultice to the abdomen or as foot baths.

They furnish a convenient form of thermal treatment with marked cutaneous stimulation. They are prescribed for torpidity of the skin and circulation, in neuralgia and neuritis, especially of the peripheral nerves, in disorders of common sensation, in chronic rheumatic affections, and to promote the absorption of morbid deposits, as in pelvic peritonitis. In conjunction with chalybeate waters they are used for dysmenorrhoea and amenorrhoea in anaemic subjects, as at Schwalbach and Strathpeffer. They are also employed locally at high temperatures to the feet and legs, the arms or abdomen, either for their derivative effect, or to set up active hyperaemia in chronic arthritis, etc.

Mud Baths (Bains de Boue, Fango).

At many thermal sulphur spas soft, "butter-like" deposits are obtained from the vicinity of the springs, which are used as an application, often at high temperatures. At Pöstyén and Mehadia, in Hungary, and at the northern Italian spas (Vinadio, Acqui, Battaglia, Abano), as well as at Baden, near Vienna, a sulphuretted mud, or fango, is extensively employed. It is also imported into England for local applications, as at Matlock.

The mud bath may be given at 40°C . (104°F .), and at much higher temperatures 48.9° to 50°C . (120° – 122°F .) applied locally. These deposits, besides containing various mineral and organic matter, often exhibit a high degree of radio-activity. Sulphuretted mud baths are extensively used at Saint Amand in France and a similar mud is obtained from river deposits at Dax, near Bordeaux. In Germany sulphur mud baths are used at Eilsen and Nenndorf, and in Sweden at Loka, in combination with massage.

In Japan the mineral mud bath can be had only at *Kamegawa*, near Beppu.

A second variety of mud baths is prepared from sea, or salt, mud, such as those of the Baltic provinces, of the "Limans" on the Black Sea coast, and of Saki and others in the Crimea. The salt mud is heated by the sun, or artificially, to a high temperature, and plastered over the body, either wholly or partially. Similar salt muds are used at Laurvik, Modum, and Sandefjord, in

Norway, at Ischl and at Arensburg, in Livonia. Mud baths are essentially a hyperthermal treatment. The actions and indications are otherwise much the same as for peat baths, but the strong salt muds, like those of the Limans are especially valued for chronic rheumatism, for scrofulous conditions, and as an accessory treatment for syphilis. The pressure exerted by the mud has been regarded as helpful for the relief of varicose veins (as at Saint Amand).

Sand Baths (Arenation).

This ancient method of thermal treatment consists in immersing the body, wholly or in part, in sand, heated—by the sun or otherwise—to from 43.3° to 54.4° C. (110°–130° F.). Profuse perspiration ensues; the skin becomes reddened and encrusted with sand; and, the normal loss of heat being reduced, the body temperature may rise three or more degrees. These baths have been used since Roman times in the island of Ischia, and recently at Lavey, in Switzerland, and Koestritz, in Germany. They are employed sometimes, together with sun baths, for their special thermal effects in chronic diseases of the joints, especially degenerative arthritis. Cold, moist sand baths are also used as a vascular and nervous tonic in neurasthenia.

D. NOTABLE MINERAL SPRINGS ARRANGED ACCORDING TO CHEMICAL COMPOSITION.

1. Simple cold springs.

| No. | Spa | Prefecture | Altitude in ft. | Temperature in C. | Total residue in gm. |
|-----|---------------------------|---------------|-----------------|-------------------|----------------------|
| 1 | Komono 菰野 | Mie-ken | 2500 | 29° | 0.14 |
| 2 | Murasugi 村杉 | Niigata-ken | 400 | 25.6° | 0.35 |
| 3 | Shimobe 下部 | Yamanashi-ken | — | 34°–36° | 0.46 |
| 4 | Takayama 高山 | Gifu ken | 1200 | 10°–13° | 0.12 |
| 5 | Nekonaki 猫啼 | Fukushima-ken | — | 22° | — |
| 6 | Fukuroda 袋田 | Ibaraki-ken | — | 34° | 0.3 |
| 7 | Koshiki-iwa-shinden 越木岩新田 | Hyōgo-ken | — | 11°–19° | — |
| 8 | Sekine-Yunosawa 関根湯澤 | Yamagata-ken | — | 28° | — |
| 9 | Kaidani 栢谷 | Okayama-ken | — | 14.5° | — |

2. Simple thermals.

| No. | Spa | Prefecture | Altitude in ft. | Temperature in C. | Total residue in gm. |
|-----|--------------------------|---------------|-----------------|-------------------|----------------------|
| 1 | Yumoto—Tonosawa 湯本—塔之澤 | Kanagawa-ken | 150 | 42°–47.3° | 0.50–0.81 |
| 2 | Utsunomiya 姥子 | „ | 2877 | 40° | 0.65 |
| 3 | Dogo 道後 | Ehime-ken | 35 | 42°–47° | 0.78–0.83 |
| 4 | Beppu 別府 | Ōita-ken | 50 | 40°–66° | 0.68–0.93 |
| 5 | Itō 伊東 | Shizuoka-ken | — | 46°–47.8° | 0.98 |
| 6 | Takeo 武雄 | Saga-ken | 100 | 49° | 0.77 |
| 7 | Kami and Shimo-Suwa 上下諏訪 | Nagano-ken | 2500 | 47.5°–83° | — |
| 8 | Iizaka 飯坂 | Fukushima-ken | 550 | 50°–70° | 0.92–0.95 |
| 9 | Nasu 那須 | Tochigi-ken | 4000–5000 | 38°–71° | 0.58–0.85 |
| 10 | Nagaoka 長岡 | Shizuoka-ken | — | 44°–53° | 0.70 |
| 11 | Asama 淺間 | Nagano-ken | 1600 | 36.5°–53° | 0.44–0.45 |
| 12 | Andai 安代 | „ | 1043 | 55°–56° | 0.95 |
| 13 | On-yo 温陽 | Chōsen | — | 38°–48° | 0.21 |
| 14 | Goshiki 五色 | Yamagata-ken | 3000 | 38.5°–44.5° | 0.78 |
| 15 | Ao-ne 青根 | Miyagi-ken | 1800 | 43°–52° | 0.54 |

| No. | Spa | Prefecture | Altitude in ft. | Temperature in C. | Total residue in gm. |
|-----|---------------------|---------------|-----------------|-------------------|----------------------|
| 16 | Tōgō 東郷 | Tottori-ken | — | 40-49 | 0,93 |
| 17 | Tochiomata 栃尾又 | Niigata-ken | 920 | 38-39 | 0,28 |
| 18 | Iiatage 畑毛 | Shizuoka-ken | — | 38°-40° | 0,64-0,90 |
| 19 | Ko-na 古奈 | " | — | 52° | — |
| 20 | Tawarayama 俵山 | Yamaguchi-ken | 1000 | 40-42 | 0,08 |
| 21 | Hokutō 北投 | Taiwan | — | 43°-68,5° | 0,21-1,09 |
| 22 | Arifuku 有福 | Shimane-ken | 900 | 44,5-49 | 0,29 |
| 23 | Misasa 三朝 | Tottori-ken | 50 | 67°-71,5° | 0,86-0,87 |
| 24 | Kashi 甲子 | Fukushima-ken | 3000 | 45-51 | — |
| 25 | Innai-Yunosawa 院内湯澤 | Akita-ken | 679 | 39,5°-41° | 0,13 |
| 26 | Kamikōchi 上高地 | Nagano-ken | 4725 | 53,5° | — |
| 27 | Ōyu 大湯 | Niigata-ken | 900 | 53°-57° | 0,38 |
| 28 | Karurusu カルス | Hokkai-dō | 1120 | 48°-60° | 0,00 |
| 29 | Yumura 湯村 | Shimane-ken | 600 | 43° | 0,32 |
| 30 | Shuotsu 朱乙 | Chōsen | 4000 | 50 | 0,18 |

3. Simple carbondioxated springs.

| No. | Spa | Prefecture | Altitude in ft. | Temperature in C. | Total residue in gm. | Therapeutically essential constituents
free CO ₂ in gm. |
|-----|-------------------------|---------------|-----------------|-------------------|----------------------|---|
| 1 | Beppu 別府 | Ōita-ken | 50 | 48,5°-67° | 0,00-0,07 | 1,12-2,51 |
| 2 | Arima—Jigokudani 有馬—地獄谷 | Hyōgo-ken | 1155 | 16,5°-17,4 | 0,16-0,31 | 1,12-1,20 |
| 3 | Takarazuka 寶塚 | " | 82 | 14,7°-18,5° | 0,58 | 1,54 |
| 4 | Funagoya 船小屋 | Fukuoka-ken | — | 17,5°-21,0° | 0,17 | 1,94 |
| 5 | Kōbe—Zyareyama 神戸—砂連山 | Hyōgo-ken | — | 21,5° | 0,75 | 1,14 |
| 6 | Su-ō 蘇澳 | Taiwan | — | 23 | 0,14 | 1,57 |
| 7 | Oshio 大鹽 | Fukushima-ken | — | cold | 0,08 | 3,07 |
| 8 | Amabe 餘戸 | Gifu-ken | — | 15 | 0,43 | 1,00 |
| 9 | Gutehan 倶知安 | Hokkai-dō | — | cold | 0,43 | 1,31 |

4. Earthy carbondioxated springs.

| No. | Spa | Prefecture | Altitude in ft. | Temperature in C. | Total residue in gm. | Therapeutically essential constituents: | | |
|-----|--------------|------------|-----------------|-------------------|----------------------|---|---|----------------------------|
| | | | | | | free CO ₂ in gm. | Ca(HCO ₃) ₂ in gm. | Mg HCO ₃ in gm. |
| 1 | Shiraya 白矢 | Nara-ken | — | cold | 1,74 | 1,42 | 1,44 | 0,12 |
| 2 | Sawaguchi 澤口 | Akita-ken | — | " | 4,00 | 1,40 | 1,71 | 0,00 |
| 3 | Shōseidō 椒井洞 | Chōsen | — | " | 1,20 | 2,04 | 0,93 | 0,57 |

5. Alkaline springs.

| No. | Spa | Prefecture | Altitude in ft. | Temperature in C. | Total residue in gm. | Therapeutically essential constituents: | |
|----------------------|-------------|-------------|-----------------|-------------------|----------------------|---|---------------------------|
| | | | | | | NaHCO ₃ in gm. | Other constituents in gm. |
| a. Alkaline springs. | | | | | | | |
| 1 | Shiobara 鹽原 | Tochigi-ken | 1150 | 51,5°-60° | 0,97-1,10 | 0,49-0,70 | — |
| 2 | Ōtari 小谷 | Nagano-ken | 3785 | 46,1°-59° | 1,82-2,42 | 2,42-2,84 | — |

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in °C | Total residue
in grm. | Therapeutically essential constituents : | |
|-------------------------------------|-----------------------|--------------|--------------------|----------------------|--------------------------|--|--|
| | | | | | | NaHCO ₃
in grm. | Other constituents in grm. |
| b. Alkaline carbondioxated springs. | | | | | | | |
| 1 | Isobe 磯部 | Gumma-ken | 1200 | 13.5°-15.5° | 15.60-28.64 | 5.97-9.49 | free CO ₂ 0.97-1.18 |
| 2 | Sado no-Kanayama 瀬戸鉛山 | Wakayama-ken | - | 42°-60° | 4.09-5.07 | 2.57-4.17 | .. 0.94-1.56 |
| 3 | Inakamado 稲亀 | Kyōto-fu | - | cold | 4.83 | 4.00 | .. 1.87 |
| 4 | Yumura 湯村 | Hyōgo-ken | 150 | 90.5°-9.5° | 0.90 | 0.37 | .. 1.24 |
| c. Alkaline muriated springs. | | | | | | | |
| 1 | Shiobara 鹽原 | Tochigi-ken | 1150 | 15°-70° | 1.35-2.23 | 0.52-0.77 | NaCl 0.62-0.88 |
| 2 | Ureshino 嬉野 | Saga-ken | 400 | 95° | 1.55 | 1.30 | .. 0.38 |
| 3 | Kasagi 笠置 | Kyōto-fu | 205 | 13.6° | 5.08 | 4.28 | .. 1.61 |
| d. Alkaline earthy springs. | | | | | | | |
| 1 | Shirahone 白骨 | Nagano-ken | 3750 | 48°-52° | 1.35 | 0.29 | (Ca HCO ₃) ₂ 0.46
(Mg HCO ₃) ₂ 0.28 |

6. Common salt springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in °C | Total residue
in grm. | Therapeutically essential
constituents : | |
|--------------------------------|------------------|---------------|--------------------|----------------------|--------------------------|---|-------------------------------|
| | | | | | | NaCl
in grm. | Other constituents
in grm. |
| a. Weak common salt springs. | | | | | | | |
| 1 | Miyanoshita 宮ノ下 | Kanagawa-ken | 1123 | 36°-81° | 1.08-2.35 | 0.71-1.05 | |
| 2 | Sakokuri 底倉 | " | 1123 | 64°-76° | 1.83-2.19 | 1.15-1.61 | |
| 3 | Dogashima 堂ヶ島 | " | 790 | 46° | 1.46-1.79 | 0.97-1.39 | |
| 4 | Beppu 別府 | Oita-ken | 50 | 56°-60° | 2.57-3.33 | 1.72-2.26 | |
| 5 | Shiobara 鹽原 | Tochigi-ken | 1150 | 42°-71.5° | 1.41-3.84 | 0.69-2.33 | |
| 6 | Yuzawara 湯河原 | Kanagawa-ken | 350 | 41°-88.5° | 1.38-2.42 | 0.85-1.45 | |
| 7 | Owani 大鰐 | Aomori-ken | 248 | 62°-80° | 2.93-3.33 | 1.78-2.56 | |
| 8 | Shima 四萬 | Gumma-ken | 3000 | 55°-84° | 2.48 | 1.49 | |
| 9 | Kuradate 蔵持 | Aomori-ken | 248 | 56°-78° | 2.32-3.14 | 1.27-2.13 | |
| 10 | Ikariga-seki 碓ヶ関 | " | 680 | 54°-62° | 1.25-1.62 | 0.75-1.30 | |
| 11 | Oyu 大湯 | Akita-ken | 572 | 46°-70.5° | 1.62-2.10 | 1.12-1.54 | |
| 12 | Kamegawa 亀川 | Oita-ken | 50 | 57° | 1.22 | 0.68 | |
| 13 | Tamatukuri 玉造 | Shimane-ken | — | 64° | 1.58 | 0.91 | |
| 14 | Misasa 三朝 | Tottori-ken | 50 | 72° | 1.16 | 0.71 | |
| 15 | Shigaku 志學 | Shimane-ken | 1500 | 46.5° | 2.39 | 1.52 | |
| 16 | Masutomi 増富 | Yamanashi-ken | 3000 | 16°-21.5° | 2.18 | 1.40 | |
| b. Simple common salt springs. | | | | | | | |
| 1 | Katayamazu 片山津 | Ishikawa-ken | | 61°-79° | 15.79 | 7.28 | |
| 2 | Senami 瀬波 | Niigata-ken | | 102° | 4.19 | 3.38 | |
| 3 | Torai 東萊 | Chōsen | | 50°-53° | 5.50 | 2.96 | |
| 4 | Noboribetsu 登別 | Hokkai-dō | 600 | 76°-94° | 4.51 | 3.01 | |
| 5 | Yoshida 吉田 | Miyazaki-ken | | 42° | 4.90 | | |

| No. | Spa | Prefecture | Altitude
m. ft. | Temperature
in. C. | Total residue
in grm. | Therapeutically essential
constituents: | |
|--|-----------------|---------------|--------------------|-----------------------|--------------------------|--|--|
| | | | | | | NaCl
in grm. | Other constituents
in grm. |
| c. Concentrated common salt springs. | | | | | | | |
| 1 | Arima 有馬 | Hyōgo-ken | 1155 | 37.5-53.4 | 19,56-65,70 | 14,72-43,21 | |
| 2 | Isobe 磯部 | Gumma-ken | 1200 | 15.5-17.2 | 20,91-29,75 | 19,43-21,05 | |
| 3 | Yashio 八鹽 | " | | cold | 19,95-30,28 | 14,53-22,84 | |
| 4 | Oshio 大鹽 | Fukushima-ken | | " | 20,40 | 15,80 | |
| 5 | Kashio 鹿鹽 | Nagano-ken | | " | 28,57 | 25,85 | |
| 6 | Haraichi 原市 | Gumma-ken | | " | 27,60-27,86 | 19,25-19,87 | |
| 7 | Mukosan 武庫山 | Hyōgo-ken | | 12.8 | 22,7 | 17,06 | |
| d. Carbonic-acid common salt springs. | | | | | | | |
| 1 | Takarazuka 寶塚 | Hyōgo-ken | 82 | 18,5° | 13,70 | 11,13 | free CO ₂ 0,93 |
| 2 | Isobe 磯部 | Gumma-ken | 1200 | 15,5 | 15,60-28,64 | 11,01-28,64 | " 0,97-1,18 |
| 3 | Hirano 平野 | Hyōgo-ken | | 27 | 4,59-7,10 | 2,57-4,31 | " 1,17-1,19 |
| 4 | Yashio 八鹽 | Gumma-ken | | cold | 8,90-30,28 | 6,19-22,84 | " 1,00-1,40 |
| 5 | Haraichi 原市 | " | | " | 27,86 | 19,87 | " 1,51 |
| e. Alkaline common salt springs. | | | | | | | |
| 1 | Isobe 磯部 | Gumma-ken | 1200 | 15.5-17.2 | 15,60-29,75 | 11,01-21,05 | NaHCO ₃ 5,97-11,88 |
| 2 | Hirano 平野 | Hyōgo-ken | | 27° | 2,59-7,10 | 1,36-4,31 | " 0,94-1,74 |
| 3 | Ogawa 小川 | Toyama-ken | | 49°-60° | 1,08-1,22 | 0,59-0,67 | " 0,25-0,31 |
| 4 | Kwanshirei 關仔嶺 | Taiwan | | 44°-80° | 7,91-8,51 | 3,62-4,18 | " 3,58-3,78 |
| 5 | Yashio 八鹽 | Gumma-ken | | cold | 5,80-30,28 | 3,69-22,84 | " 1,58-5,00 |
| 6 | Haraichi 原市 | " | | " | 27,60-27,86 | 19,25-19,87 | " 6,03-11,19 |
| f. Saline common salt springs. | | | | | | | |
| 1 | Shuzenji 修善寺 | Shizuoka-ken | 250 | 55°-77° | 1,08-1,21 | 0,46-0,58 | Na ₂ SO ₄ 0,22-0,45 |
| 2 | Kamasaki 鎌先 | Miyagi-ken | 600 | 37-48 | 5,03-5,19 | 2,41-2,47 | " 1,50-1,69 |
| g. Earth-muriated common salt springs. | | | | | | | |
| 1 | Atami 熱海 | Shizuoka-ken | | 77°-108° | 8,1-9,87 | 4,79-5,66 | CaCl ₂ 2,57-3,19 |
| 2 | Arima 有馬 | Hyōgo-ken | 1155 | 37.8-38.3 | 19,56-19,66 | 14,72-43,21 | " 2,90-11,30 |
| 3 | Kinosaki 城崎 | " | | 44.2-60.3 | 5,32-5,91 | 3,02 | " 1,50 |
| 4 | Wakura 和倉 | Ishikawa-ken | | 70.5-93 | 20,93-21,55 | 11,03-18,67 | " 8,61-8,94 |
| 5 | Obama 小濱 | Nagasaki-ken | | 50-101 | 6,47-8,84 | 4,00-5,43 | { CaCl ₂ 1,01-1,26
MgCl ₂ 0,60-0,91 |
| 6 | Awara 蘆原 | Fukui-ken | | 53°-76° | 10,03-10,33 | 5,90-6,13 | CaCl ₂ 3,09-3,29 |
| 7 | Akayu 赤湯 | Yamagata-ken | 700 | 42°-58° | 2,81-3,97 | 1,90-2,06 | " 0,50-0,55 |
| 8 | Onogawa 小野川 | " | 1000 | 65.5°-73.5° | 5,17 | 3,60 | " 1,10 |
| 9 | Yunogo 湯郷 | Okayama-ken | 240 | 37,7°-38° | 2,25-2,27 | 1,17-1,18 | " 0,94-0,95 |
| 10 | Yunokawa 湯川 | Hokkai-dō | | 50 | 7,03 | 4,41 | |
| 11 | Matsunoyama 松之山 | Niigata-ken | 1000 | 44°-73° | 13,87-15,08 | 7,93-8,68 | CaCl ₂ 5,05-5,33 |
| 12 | Yunohama 湯野濱 | Yamagata-ken | | 43°-47,2° | 4,71-5,09 | 2,79 | " 1,51 |
| 13 | Atsushio 熱鹽 | Fukushima-ken | 1200 | 40 | 11,50 | 7,85 | " 2,24 |
| 14 | Morigasaki 森ヶ崎 | Tōkyō-fu | | 17 | 9,08 | 3,83 | { CaCl ₂ 0,90-1,00
MgCl ₂ 0,03 |
| 15 | Yunomoto 湯ノ本 | Nagasaki-ken | | 43-47 | 16,39-21,64 | 11,56-14,51 | { CaCl ₂ 0,82-1,74
MgCl ₂ 0,82-1,29 |
| 16 | Oshio 大鹽 | Fukushima-ken | | cold | 20,40 | 13,71 | { CaCl ₂ 1,00-1,00
MgCl ₂ 1,00-1,00 |

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in grm. | Therapeutically essential
constituents: | |
|--|----------------------|---------------|--------------------|----------------------|--------------------------|--|---|
| | | | | | | NaCl
in grm. | Other constituents
in grm. |
| h. Earthy common salt springs. | | | | | | | |
| 1 | Yunokawa 湯川 | Hokkai-dō | — | 40.5°-50° | — | — | — |
| 2 | Atsushio 熱鹽 | Fukushima-ken | 1200 | 35°-78° | 4.55- 5.11 | 2.65- 3.26 | {Ca(HCO ₃) ₂ 0.48-1.21
{Mg(HCO ₃) ₂ 0.17 0.26 |
| 3 | Akagi-Nashiki 赤城梨木 | Gumma-ken | 1490 | 20° | 2.25 | 1.08 | Mg(HCO ₃) ₂ 0.82 |
| 4 | Aoyama 青山 | Hokkai-dō | — | 42°-44° | 2.97 | 1.72 | {Ca(HCO ₃) ₂ 0.39
{Mg(HCO ₃) ₂ 0.29 |
| 5 | Masutomi 増富 | Yamanashi-ken | 3000 | 15°-24.5° | 5.98- 9.86 | 3.84- 6.49 | {Ca(HCO ₃) ₂ 0.96-1.29
{Mg(HCO ₃) ₂ trace-0.49 |
| i. Sulphated common salt springs. | | | | | | | |
| 1 | Shibu [Hirao] 澁 [平穩] | Nagano-ken | 6950 | 45°-76° | 1.21 | 0.53 | CaSO ₄ 0.36 |
| 2 | Shima 四萬 | Gumma-ken | 3000 | 62° | 1.47 | 0.72 | .. 0.45 |
| 3 | Atsumi 溫海 | Yamagata-ken | 50 | 45°-70° | 3.51 | 2.35 | .. 0.81 |
| 4 | Yunotsu 溫泉津 | Shimane-ken | — | 46°-50° | 7.06 | 4.79 | .. 1.10 |
| 5 | Yuwaku 湯涌 | Ishikawa-ken | 1300 | 41° | 3.31 | 2.15 | .. 0.71 |
| 6 | Omaki 大牧 | Toyama-ken | 850 | 49° | 3.29 | 1.86 | .. 0.82 |
| j. Hydrogen sulphide common salt springs. | | | | | | | |
| 1 | Ōyuzawa 大湯澤 | Akita-ken | 700 | 33.5°-45.5° | 11.31-14.78 | 7.00-11.03 | H ₂ S 0.23 |
| k. Common salt springs containing bromine or iodine. | | | | | | | |
| 1 | Arima 有馬 | Hyōgo-ken | 1155 | 38.3°-47° | 19.56-65.70 | 14.72-43.21 | Br 0.0116-0.0854 |
| 2 | Kinosaki 城崎 | " | — | 58.3° | 5.32 | 3.02 | .. 0.0005 |
| 3 | Isobe 磯部 | Gumma-ken | 1200 | 13.5°-15.5° | 26.91-29.75 | 19.43-21.05 | {Br 0.0113-0.0298
{I 0.0020-0.0047 |
| 4 | Mukosan 武庫山 | Hyōgo-ken | — | cold | 7.40 | 5.97 | I 0.0026 |
| 5 | Shikanoda 鹿野田 | Miyazaki-ken | — | .. | 10.04 | 14.43 | {Br 0.0784
{I 0.0294 |
| 6 | Ōseda 大瀬戸 | " | — | .. | 3.69 | 2.22 | {Br 0.0155
{I 0.0075 |
| 7 | Nanatsu-ido 七井戸 | Chiba-ken | — | .. | 16.01 | 13.76 | I 0.0476 |
| 8 | Shita 志太 | Shizuoka-ken | — | .. | 11.80 | 9.36 | .. 0.0127 |
| 9 | Miyagaki-uchi 宮垣内 | Wakayama-ken | — | .. | 12.03 | 8.29 | {Br 0.0030
{I 0.0008 |
| l. Common salt springs containing boric acid. | | | | | | | |
| 1 | Isobe 磯部 | Gumma-ken | 1200 | 15.5° | 26.91 | 19.43 | H ₂ BO ₂ 1.00 |
| 2 | Haraichi 原市 | " | 1200 | cold | 27.86 | 19.87 | .. 5.54 |
| 3 | Jōhōji 浄法寺 | Aomori-ken | — | " | 16.83 | 10.17 | .. 3.03 |

7. Bitter springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in grm. | Therapeutically essential
constituents in grm. |
|--------------------|-----------------|--------------|--------------------|----------------------|--------------------------|---|
| a. Bitter springs. | | | | | | |
| 1 | Kaminoyama 上ノ山 | Yamagata-ken | 574 | 56°-61.5° | 2.58 | |
| 2 | Shido-daira 志戸平 | Iwate-ken | 700 | 76° | 1.20 | |
| 3 | Ushio 海潮 | Shimane-ken | 300 | 41.5° | 1.22 | |

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in grm. | Therapeutically essential
constituents in grm. | |
|---------------------------------------|----------------|---------------|--------------------|----------------------|--------------------------|---|------------------------|
| b. Saline bitter springs. | | | | | | | |
| 1 | Shiobara 鹽原 | Tochigi-ken | 1150 | 55°-57.5° | 1.49 | Na ₂ SO ₄ | 0.84 |
| 2 | Higashiyama 東山 | Fukushima-ken | 850 | 34°-61° | 1.95 | " | 0.69 |
| 3 | Yoshina 吉奈 | Shizuoka-ken | 300 | 41°-50° | 1.15-1.24 | " | 0.62 |
| 4 | Funabara 船原 | " | " | 35°-47° | 1.00 | " | 0.47 |
| 5 | Iwai 岩井 | Tottori-ken | 40 | 46°-60° | 1.96-2.00 | " | 1.07-1.10 |
| 6 | Yujiku 湯宿 | Gumma-ken | " | 37.2°-79° | 1.42 | " | 0.64 |
| c. Sulphated bitter springs. | | | | | | | |
| 1 | Ikao 伊香保 | Gumma-ken | 2500
2700 | 45°-47° | 0.96-1.34 | CaSO ₄ | 0.18-0.45 |
| 2 | Izusan 伊豆山 | Shizuoka-ken | 42 | 60° | 1.42 | " | 0.75 |
| 3 | Iwai 岩井 | Tottori-ken | 40 | 45°-58.5° | 1.83-1.91 | " | 0.96-0.98 |
| 4 | Asamushi 浅蟲 | Aomori-ken | - | 61.5°-79° | 1.15-1.34 | " | 0.31-0.63 |
| 5 | Yugashima 湯ヶ島 | Shizuoka-ken | 625 | 41°-64° | 1.7 | " | " |
| 6 | Tohi 上肥 | " | 30 | 36°-66° | 1.37-1.69 | CaSO ₄ | 0.89-1.11 |
| 7 | Tochinoki 栃木 | Kumamoto-ken | 450 | 39°-45° | 2.08 | " | " |
| d. Muriated saline bitter springs. | | | | | | | |
| 1 | Yoshikata 吉方 | Tottori-ken | " | 24.4°-47.5° | 4.03-4.62 | { NaCl
{ Na ₂ SO ₄ | 1.42-1.60
1.49-1.78 |
| 2 | Kachimi 勝見 | " | " | 51.5°-55° | 1.12-1.36 | { NaCl
{ Na ₂ SO ₄ | 0.10-0.49
0.39-0.65 |
| 3 | Hamamura 濱村 | " | " | 45°-49° | 1.07 | { NaCl
{ Na ₂ SO ₄ | 0.40
0.32 |
| e. Muriated sulphated bitter springs. | | | | | | | |
| 1 | Yudanaka 湯田中 | Nagano-ken | 1043 | 74°-76° | 1.53 | { NaCl
{ CaSO ₄ | 0.70
0.65 |

8. Iron carbonate springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in grm. | Therapeutically essential
constituents: | |
|---|----------------------------|---------------|--------------------|----------------------|--------------------------|---|---|
| | | | | | | Fe HCO ₃ ₂
in grm. | Other constituents
in grm. |
| a. Iron carbonate springs. | | | | | | | |
| 1 | Beppu 別府 | Oita-ken | 50 | 57.5 | 0.85 | 0.0383 | " |
| 2 | Kwankaiji 観海寺 | " | 200 | 57 | 0.77 | " | " |
| 3 | Shibasaki 芝石 | " | 200 | 69 | 1.59 | 0.2207 | " |
| b. Iron carbonate common salt springs. | | | | | | | |
| 1 | Arima 有馬 | Hyogo-ken | 1155 | 37.8°-47° | 19.66-65.70 | 0.0088-0.8480 | NaCl 14.72-43.19 |
| 2 | Arimura (Sakurajima) 有村 櫻島 | Kagoshima-ken | " | 38.0-45 | 5.38-5.70 | 0.0952-0.2286 | " 4.03-4.07 |
| 3 | Anamori 穴守 | Tokyo-fu | " | 17° | 0.38 | 0.445 | 7.48 |
| 4 | Koyabara 小屋原 | Shimane-ken | 900 | 38.2 | 5.28 | 0.0324 | 3.01 |
| c. Iron carbondioxated common salt springs. | | | | | | | |
| 1 | Hirano 平野 | Hyogo-ken | " | 27 | 4.50-5.50 | 0.0002-0.0342 | { NaCl 2.57-3.33
{ H ₂ CO ₃ 1.12 |
| 2 | Yashio 八潮 | Gumma-ken | " | cold | 8.90 | 0.1779 | { NaCl 0.41
{ H ₂ CO ₃ 1.45 |

9. Vitriol springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically
essential
constituent

FeSO ₄ in gm. |
|-----|----------------|---------------|--------------------|----------------------|-------------------------|---|
| 1 | Rokuyō 鹿野園 | Nara-ken | — | 20.5° | 3.79 | 2.923 |
| 2 | Michi-no-o 道之尾 | Nagasaki-ken | — | 24° | — | — |
| 3 | Hisomoe 砒霜燃 | Kagoshima-ken | 2706 | 78.9° | 0.36 | 0.0400 |

10. Alum vitriol springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically essential constituents: | | |
|-----|----------------|-----------------------------|--------------------|----------------------|-------------------------|---|---|--|
| | | | | | | FeSO ₄
in gm. | Fe ₂ (SO ₄) ₃
in gm. | Al ₂ (SO ₄) ₃ in gm. |
| 1 | Yurimoto 湯ノ本 | Nagasaki-ken
(Iki Prov.) | — | 43°-47° | 19,84-21,64 | 0.2882 0.2969 | — | 0.650-0.669 |
| 2 | Isobe 磯邊 | Toyama-ken | — | cold | 20,52-31,33 | 4.1420-12.1192 | 0-2.1063 | 13,596-16,461 |
| 3 | Yoshima 好間 | Fukushima-ken | — | " | 18,75 | 4.5196 | — | 9.784 |
| 4 | Uchiyama 内山 | Nagano-ken | — | " | 10,28 | 7.0320 | — | 2.048 |
| 5 | Iiyugayama 日向山 | " | — | " | 8,77 | 3.6602 | 0.3965 | 3.247 |

11. Acid vitriol springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically essential constituents: | | |
|-----|----------------|--------------|--------------------|----------------------|-------------------------|---|---|--|
| | | | | | | HCl (free)
in gm. | H ₂ SO ₄ (free)
in gm. | Fe(SO ₄)
in gm. |
| 1 | Kusatsu 草津 | Gumma-ken | 3740 | 60°-63.9° | 4.07-4.71 | 0.7461-0.8485 | 2.6186-2.2868 | 0.1663-0.2658 |
| 2 | Unzen 温泉 | Nagasaki-ken | 2145 | 65° | 1.20 | — | 1.2887 | 0.2539 |
| 3 | Kowakidani 小湧谷 | Kanagawa-ken | 2000 | 35.6° | 1.13 | — | 0.1920 | 0.3982 |
| 4 | Kannawa 鍛輪 | Ōita-ken | 150 | 89°-90.5° | 2.67-3.51 | 0.2214-0.7910 | — | 0.0451-0.3513 |
| 5 | Noboribetsu 登別 | Hokkai-dō | 600 | 71.5° | 0.81 | — | 0.2316 | 0.0650 |
| 6 | Naruko 鳴子 | Miyagi-ken | — | 62 | — | — | 0.3307 | Fe ₂ (SO ₄) ₃ 0.0878 |
| 7 | Renge 蓮華 | Niigata-ken | 4500 | 36°-39° | 0.80 | — | 0.0845 | 0.0524 |
| 8 | Noroshi 狼煙 | Ishikawa-ken | — | 15° | 1.16 | — | 0.2893 | 0.6582 |
| 9 | Shimoburo 下風呂 | Aomori-ken | — | 17° | 4.03 | — | 0.3074 | 0.0364 |

12. Acid alum springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically essential constituents: | | |
|-----|--------------|---------------|--------------------|----------------------|-------------------------|---|---|---|
| | | | | | | HCl free
in gm. | H ₂ SO ₄ free
in gm. | Al ₂ (SO ₄) ₃
in gm. |
| 1 | Takayu 高湯 | Fukushima-ken | 2040 | 45°-49° | 1.52-1.70 | 0.01012 | 0.1152-0.2722 | 0.5645-0.9702 |
| 2 | Togeshita 峠下 | Hokkai-dō | — | cold | 3.89 | — | 0.0112 | 1.8459 |

13. Acid alum vitriol springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Theoretically essential constituents: | | | |
|-----|------------------------|------------|--------------------|----------------------|-------------------------|---------------------------------------|---|-----------------------------|--|
| | | | | | | HCl (free)
in gm. | H ₂ SO ₄ (free)
in gm. | FeSO ₄
in gm. | Al ₂ SO ₄ in gm. |
| 1 | Kusatsu 草津 | Gunma-ken | 3740 | 43° 64.4° | 0.31-7.25 | 0.2664 | 14.8861 | 0.2177 | 1.5603 |
| 2 | Beppu 別府 | Oita-ken | 50 | | 6.65 | | | 1.1833 | 0.2130 |
| 3 | Myōdan 明礬 | " | 490 | 63°-97° | 2.52 | | | 1.0870 | 0.1590 |
| 4 | Hokuto 北投 | Taiwan | - | 48.5° 94° | 9.21 | | 2.3737 | | 0.3745 |
| 5 | Isobe 磯邊 | Toyama-ken | - | cold | 5.01-42.00 | 0.00499 | 0.15048 | 0.71660 | 0.290544 |
| 6 | Mikamuruma 御番沼 | Miyagi-ken | - | | 15.03 | 4.0390 | 0.2021 | 1.1800 | 7.390 |
| 7 | Shiino-kero 流黒 | Akita-ken | 2700 2800 | | 2.65 5.02 | 1.2681 | 2.5497 | 0.1254 | 0.4247 |
| 8 | Fukunoin 龍ノ入 | " | - | cold | 27.03 | 0.0157 | | 3.5791 | 0.6746 |
| 9 | Yamanoto-kamiyama 山ノ神澤 | " | - | " | 15.07 | | | 3.2454 | 13.7179 |
| 10 | Fuketsu-ka-mi 龍澤神 | " | - | " | 10.01 | 0.0779 | | 5.2010 | 7.6859 |
| | | | | | | | | | 10.7539 |
| | | | | | | | | | 0.720 4.749 |
| | | | | | | | | | 1.120 |
| | | | | | | | | | 0.863 |
| | | | | | | | | | 0.089 |
| | | | | | | | | | 1.780 11.667 |
| | | | | | | | | | 7.390 |
| | | | | | | | | | 0.549 0.668 |
| | | | | | | | | | 7.326 |
| | | | | | | | | | 4.589 |
| | | | | | | | | | 3.367 |

14. Sulphur springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Theoretically essential constituents: | |
|---------------------|-----|---------------|--------------------|----------------------|-------------------------|---------------------------------------|------------------------------|
| | | | | | | H ₂ S in gm. | other constituents
in gm. |
| a. Sulphur springs. | | | | | | | |
| 1 | 武蔵 | Fukuoka-ken | | 41° 40' | 0.72 | 0.000226 | |
| 2 | 堀田 | Oita-ken | 300 | 30 | | | |
| 3 | 阿蘇 | Fukuoka-ken | | 47° 58' | 2.23-2.24 | 0.0054 0.0073 | |
| 4 | 野碓 | Nagato-ken | 603 | 41° 82' | 0.70 0.68 | 0.0110 0.3230 | |
| 5 | 明礬 | Oita-ken | 400 | 68° | | | |
| 6 | 湯谷 | Iwate-ken | 2517 | 05 | 0.70 | | |
| 7 | 須賀 | Myagi-ken | | 40.5 | | | |
| 8 | 相馬 | Tottori-ken | 500 | 12° 40' | 0.52 0.56 | 0.0000 0.0023 | |
| 9 | 中房 | Nagato-ken | 5300 | 50.5 | 0.55 | 0.0000 | |
| 10 | 二見 | Toyama-ken | 650 | 64.5 95 | 0.50 0.61 | 0.0004 0.0028 | |
| 11 | 燕 | Nagato-ken | 3000 | 42° 48' | | | |
| 12 | 田中 | Kumamoto-ken | 850 | 57° 64' | | | |
| 13 | 酸湯 | Yamaguchi-ken | 4000 | 60 | 0.60 | | |

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically essential
constituents : | |
|-------------------------------|-------------------|---------------|--------------------|----------------------|-------------------------|---|--------------------------------------|
| | | | | | | H ₂ S in gm. | Other constituents
in gm. |
| b. Hydrogen sulphide springs. | | | | | | | |
| 1 | Nikko-Yumoto 日光湯本 | Tochigi-ken | 5088 | 22°-69° | 0,88-1,44 | 0,0268-0,0506 | |
| 2 | Nasu-Yumoto 那須湯本 | " | 3000 | 28° | 0,62 | 0,0292 | |
| 3 | Iwōdani 硫黄谷 | Kagoshima-ken | 3343 | 48,7°-59,5° | 0,47 | 0,0522 | |
| 4 | Ei-no-o 榮之尾 | " | 2410 | 34°-39° | 1,02 | 0,2374 | |
| 5 | Tateyama 立山 | Toyama-ken | 4627 | 63° | 1,39 | 0,0022 | |
| 6 | Shiono-e 鹽ノ江 | Kagawa-ken | 713 | 17° | 0,47 | 0,0021 | |
| 7 | Oyuzawa 大湯澤 | Akita-ken | 700 | 33,5°-45,5° | 11,31-14,78 | 0,2246 | NaCl 7,00-11,03 |
| c. Alkaline sulphur springs. | | | | | | | |
| 1 | Akakura 赤倉 | Niigata-ken | 3000 | 55,5°-62° | 1,19 | 0,0268 | |
| 2 | Naruko 鳴子 | Miyagi-ken | — | 103° | 3,52 | — | |
| 3 | Nakabusa 中房 | Nagano-ken | 5300 | 74°-96° | 0,46-1,14 | 0,0039-0,0070 | |
| d. Muriated sulphur springs. | | | | | | | |
| 1 | Takedao 武田尾 | Hyōgo-ken | 340 | 10,5°-23,5° | 1,13 | 0,0014 | NaCl 0,73 |
| 2 | Misasa 三朝 | Tottori-ken | 50 | 56,5° | 1,24 | 0,0023 | " 0,71 |
| 3 | Yumoto 湯本 | Fukushima-ken | — | 48,9° | 3,16-3,20 | 0,0164-0,0105 | " 2,05 |
| 4 | Sawatari 澤渡 | Gumma-ken | 2200 | 38,0°-52,8° | 1,23-2,71 | 0,0255 | " 1,23 |
| e. Saline sulphur springs. | | | | | | | |
| 1 | Yamashiro 山代 | Ishikawa-ken | — | 59°-71,5° | 1,66-1,74 | 0,00102-0,00545 | Na ₂ SO ₄ 0,69 |
| 2 | Yunomine 湯ノ峯 | Wakayama-ken | 630 | 87,5°-92° | 1,38 | 0,01014 | " 0,70 |
| f. Sulphated sulphur springs. | | | | | | | |
| 1 | Yamanaka 山中 | Ishikawa-ken | 240 | 49° | 1,65 | 0,0010 | CaSO ₄ 0,93 |
| 2 | Kawarayu 川原湯 | Gumma-ken | 2110 | 28,9°-70,7° | 1,52-1,72 | 0,0034-0,0091 | — |

15. Acid hydrogen sulphide springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically essential constituents : | | |
|-----|---------------------|---------------|--------------------|----------------------|-------------------------|--|---|-----------------------------------|
| | | | | | | HCl (free)
in gm. | H ₂ SO ₄ (free)
in gm. | H ₂ S (free)
in gm. |
| 1 | Kusatsu 草津 | Gumma-ken | 3740 | 58°-64,6° | 2,38-5,60 | 0,3085-0,8742 | 1,8334-2,1674 | 0,0041-0,0055 |
| 2 | Unzen 温泉 | Nagasaki-ken | 2145 | 38°-65° | 0,36-1,20 | small quantity | 0,1860-1,2887 | 0,0012-0,0041 |
| 3 | Nasu-Yumoto 那須湯本 | Tochigi-ken | 3000 | 43°-74,5° | 0,72-1,91 | 0,0838-0,2140 | 0,2171-0,4580 | 0,240-0,264 |
| 4 | Yumohanazawa 湯之花澤 | Kanagawa-ken | 3300 | 40° | 0,7 | — | 0,0180 | 0,1205 |
| 5 | Manza 萬座 | Gumma-ken | 5180 | 50,6°-81,7° | 0,05-1,54 | 0-0,0865 | 0,0990-0,4535 | 0,0527-0,3256 |
| 6 | Takayu 高湯 | Fukushima-ken | 2640 | 45°-49° | 1,52-1,77 | 0,1012-0,1114 | 0,1152-0,2294 | 0,0447-0,0840 |
| 7 | Numajiri 沼尻 | " | — | 63° | 1,94 | 0,5185 | 0,1698 | 0,1271 |
| 8 | Shibu [Suwa] 澁 [諏訪] | Nagano-ken | 6950 | 27° | 0,86 | — | 0,6692 | 0,0167 |

16. Acid springs.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically essential
constituents: | |
|-----|----------|--------------|--------------------|----------------------|-------------------------|--|--|
| | | | | | | HCl in
gm. | H ₂ SO ₄ in
gm. |
| 1 | Unzen 温泉 | Nagasaki ken | 2145 | 50-74 | 0.36-0.37 | trace | 0.735-0.1170 |
| 2 | Dake 岳 | Aomori ken | 2500 | 45-83.9 | 1.00 | | 0.0004 |

17. Boric acid spring.

| No. | Spa | Prefecture | Altitude
in ft. | Temperature
in C. | Total residue
in gm. | Therapeutically essential
constituent: |
|-----|--------------|------------|--------------------|----------------------|-------------------------|---|
| | | | | | | HBO ₂ in gm. |
| 1 | Takaizumi 高泉 | Gumma ken | — | — | 2.16 | 1.43 |

While the foregoing has been passing through the press, fresh data have come to hand, and these are appended here by way of supplement.

APPENDIX A.

ON THE GEYSER AT ONIKÔBE, MIYAGI PREFECTURE.

By DR. K. HONDA AND MR. T. SONE.

The Onikôbe Geyser, or "*Fuki-age*," as it is called among the natives, is situated at Onikôbe, Tamatsukuri-gôri, Miyagi Prefecture, and is one of the hot springs that are scattered along the eastern side of the Central Group of the Northern Mountain Range in Japan Proper. The geyser lies in the valley of the river Arao, a tributary to the river Kitakami, and this district is extremely irregular and undulating in topography, being intersected by steep mountain chains and deep gorges alternately. Hot springs are exceedingly numerous in the neighbourhood, and are indeed counted by hundreds, a sure evidence of the existence of some fissures in the earth crust running through this district. Visitors to this district will certainly be surprised to see columns of dense vapour rising everywhere and tepid water flowing down in torrents. It is also interesting to note that some springs spout hot water from the bottom of the stream, while others situated in cavities make noises resembling peals of distant thunder.

Turning to the left across a small bridge and then proceeding a short distance along the course of the river Arao, one can notice a cataract of some ten feet in height shooting over the precipice. To the right, one can also perceive clouds of dense vapour hanging over a mountain-side which indicates the site of the well-known "*Fuki-age*" Geyser. The upper part of the cataract forms a flat rock-bed, on which at the foot of an impending cliff on its western edge lies the basin of the geyser.

The orifice of the geyser opens at the south-western end of the basin and leads obliquely deep into the rock-bed. When the pool thoroughly dries up, as it does after an eruption, a full sight can be obtained of the mouth of the subterranean canal. The interior of the canal forms a curve and is completely shut out from view. The mouth of the orifice is five inches by one foot. There are two stones and a piece of timber lying in front of the orifice and are said to be a relic of the abnormal overflow of the streams in the vicinity on the occasion of a flood in 1910. Formerly there was another orifice near the south-eastern rim of the basin, but it has now been blocked, leaving vestiges of its mouth no longer in action. (Fig. 1 and 2)




Fig. 1



Fig. 2

According to tradition, the "*Fuki-age*" Geyser is said to have been known as far back as the era of the Emperor Ojin, a sovereign in the remote period of the history of Japan. Upwards of 1080 years ago, the north-eastern part of Japan was visited by tremendous earthquakes lasting for several days, when, we are told, a number of mountains crumbled down, and several hot springs made their appearance. Towards the beginning of the 19th century, a certain Ōba, a native of Akita Prefecture, settled down near the basin of the geyser, and later on, his descendants erected a bath-room for the convenience of bathers in general.

At that time there were two orifices, one called "*Kōbō*" and the other "*Fudō*." It was always "*Kōbō*" or the left orifice that first burst in eruption, and as soon as its activity ceased, the "*Fudō*" or the right orifice followed in action. This order was seldom disturbed. The "*Kōbō*" was copious in volume of the outflow, but it shot up only a little over ten feet in height, while the "*Fudō*" with smaller mouth shot up to a height of between 30 and 40 feet. Tradition goes that on one occasion the eruption blighted branches of some trees growing over the cliff. This tale taken as a fact may be explained by the supposition that the activity of the geyser must have shown a secular variation, allowing the trees to grow fully in the epoch of comparative inactivity and blasting them when in later years the eruption acquired its full intensity. It is also said that the source of the spring was lined with stone all around so as to collect the water into a pool which was partially dammed up by wooden boards, and that, at the request of visitors, the basin could be drained and the eruption thus excited artificially.

Down to fifty or sixty years ago, the two orifices remained in activity, repeating eruptions three times in a day and night *i. e.* about at 7 a. m., noon and 5 p. m. The ejection from the one orifice attained a height of over 10 feet and lasted in action for a minute or so, and after the lapse of half a minute the other commenced activity, shooting up to a height of between 24 and 25 feet with a terrific vigour for an interval of about two minutes. Later on, the geyser began to lose its force by degrees, and especially since, in May 1875, the Miyazawa spring in the vicinity has burst in a sudden explosion and killed 39 persons, even the height of the "*Fudō*" decreased to 14 or 15 feet.

When in 1894 Mr. Juzo Iwasaki made a visit to the region, he found the two orifices still in activity, the frequency of the eruption being 7 times in a day and night with the interval of 3 hours and 25 minutes on the average.

The observations of Mr. Iwasaki were subsequently published in the "Chishitsu-gaku Zasshi" (vide No. 15, Vol. II) which are quoted below with few corrections as regards the dimensions of the orifices.

"The two orifices, the larger and the smaller, are located along the major axis of the pool, the smaller one near the stream and the larger at some distance from it. The larger aperture shoot up columns of hot water as high as above 20 feet for a period of 3 minutes. After that the water vanishes from the basin, leaving heaps of stones and sands very hot to touch. After the lapse of about two minutes, a faint sound is heard far from the bottom of the smaller orifice, when all of a sudden water bursts out of it, rising to an altitude of 30 feet. The eruption lasts 2 or 3 minutes. Then the basin is again drained of its contents. Half an hour later, the water appears again inside the mouth of the smaller orifice, and fills up the basin as well as the larger orifice."

After 1898-9 the smaller orifice gradually declined in activity. In August 1910 the district was devastated by a flood of a very disastrous description, and the bath-house which had been run for generations by the Ōbas was washed away, causing the loss of 23 lives. Since then the geyser has been greatly lessened in force, its height diminishing to between 7 and 8 feet, though it increased a good deal in frequency.

The following is an extract of the observations made by Mr. Denzō Satō in July, 1913:—

"Directly after the eruption, the basin is found completely dried up, leaving only heaps of stones, but in a few moments the water gradually begins to fill up the orifice, and makes its appearance in the bottom of the basin till, after the lapse of half an hour, it brims over and finds its way out over the depressed part of the south-eastern rim. Just at this moment effervescence is observed to take place near the orifice and this is succeeded by ebullition, the overflow gaining in volume at the same time. Still the eruption is not yet set in. Meanwhile the ebullition grows in intensity and after a certain interval of time the outflow shows a sudden increase in volume until it overflows the southern rim, too, as if it were pushed by some unknown agency underneath. There is, however, no noise heard yet. At the same time the water in ebullition presents a state of extraordinary agitation, followed by a sudden eruption torn into violent splashes, rising up first to the height of 5 or 6 inches, then to 1 or 2 feet and ultimately reaching the climax of 4 or 5 feet. The splashes attain the height of even more than 7 feet. Taken altogether, it is to be remarked that the eruption has considerably decreased in altitude compared with previous years. After continuing in action for some forty seconds, the spouts become lower and lower and finally collapse, when, almost simultaneously, the water in the basin disappears in the orifice, leaving heaps of stones thoroughly dried up. As the water is being drained into the aperture, a faint rumbling sound is heard. It takes one minute and a half from the beginning to the last stage of the display. After the quiescence of one hour and five minutes the next eruption sets in."

The above may, in substance, be taken as a fairly accurate description of the present state of the geyser, and it entirely coincides with the results of our personal observations made last year.

To investigate the period of eruption and its variations corresponding to different external influences, we constructed a kind of self-recording air-thermometer, the bulb of which was fixed in front of the orifice. The apparatus proved very trustworthy in keeping record of the successive eruptions which were repeated 22 or 23 times in 24 hours with fairly regular intervals. A closer examination, however, showed that the periods of eruptions during the course of a day varied within a certain range.

Considering that the period of eruption may be affected by the pressure, we carried out experiments by raising the level of the water artificially. The embanking the basin all around so as to dam up the issuing water, the water level was raised by about 60 mm., and several observations were carried out.

It was ascertained that the period of the eruption which recorded about 70 minutes at the normal level of water, was prolonged to about 75 minutes when the level was raised by roughly 60 mm.

Next the relations between the variation in atmospheric pressure and the period of eruption demanded our attention. The results of our observations showed that a rise of about 4 mm. in the

mercury column which corresponds to about 60 mm. in the water level, brought about a prolongation of the period by approximately 10 minutes.

Now in the foregoing experiments relating to the artificial elevation of water level, a prolongation of only 5 minutes was brought about by an increase of 60 mm. in the water level, so that the explanation based on the assumption that the increase of the atmospheric pressure is simply equivalent to the elevation of water level and brings about the rise of boiling point, is inadequate to account for this phenomenon. The explanation must be sought elsewhere.

It is a well-known fact that the temperature of some ordinary hot springs rises with low atmospheric pressure. The explanation generally accepted is that the reduction of pressure facilitates the flow of the underground water and hence retards the cooling of water as it circulate through the surface layer of the earth crust. This phenomenon must be taken into account as affecting the period of eruption in the case of the Onikōbe geyser. Strange to observe, in the case of the Atami geyser, low pressure is noted to prolong the period of eruption, a phenomenon which is quite contrary to the case under notice. This apparent contradiction must assuredly be attributed to the difference of internal structure and arrangements between the two geysers, the appraisable assumption being that the subterranean cavity at Atami lies at a greater depth, while that at Onikōbe is nearer the surface of the earth. In support of this hypothesis, we may note the fact that in the latter the difference of water level affects the period of eruption in no small measure. If the cavity lies deep in the earth, the wall surrounding the cavity will generally maintain higher temperature than the subterranean water, and supplies heat to water flowing into it, while in the case of a cavity seated in a smaller depth where the earth crust is not high in temperature, the water in the cavity will, on the contrary, get heated by the flow of the subterranean water of relatively higher temperature. In short, the process that the water in the cavity is heated will be different according as the supply of heat comes from the surrounding wall or is derived from the superheated subterranean water flowing into it. Thus, at the time of low pressure which causes vigorous circulation of water, the heating of the water in the cavity will be slow in the former case and quick in the latter. This may suffice to explain why the low pressure brings about the prolongation of the period of eruption at Atami and to shorten it at Onikōbe. The conclusion we have arrived at is entirely different from the view set forth by Messrs. Iwasaki and Shimizu who regarded that the surrounding wall as heating the water in the cavity.


In order to estimate the quantity of water issuing from the orifice, we measured the overflow with a tub, by receiving in it the water as it flows out of the outlet in the embankment erected around the pool to that special end in view. The quantity thus obtained measured about 37 litre per minute at 20 minutes and 39 litre at between 2 and 3 minutes before eruption, indicating thereby that hot water always gushes out at practically fixed rates.

The temperature of the water at the mouth of the orifice stood at 97° C. both in autumn and winter, and as for its temperature in the canal, we measured it by inserting, at the time of repose, a maximum thermometer 2 or 3 metres deep into it, and found it reading 104.5° C. It was impossible to push the thermometer any further into the canal as it was found to curve inward, but it was sufficient to indicate that the cavity must generally maintain a pretty high temperature.

EXPLANATION OF THE PHENOMENA.

The results of our observations afore mentioned lead to the following consideration with regard to the construction of the geyser, by which the characteristic phenomena of the eruption may be explained :—

In the past period, in which two orifices have been still active, two cavities, one with the larger neck (i) and another with the smaller one (ii), must have been connected with the pool by the vertical channels C and D respectively, both opening side by side at the bottom of the pool. These cavities receive hot water heated in the deeper part of the earth crust flowing up from several narrow canals and the water ascends into the pool through the channels, heating in the mean time the water present in the cavities.



The position of the cavities may be considered to be either nearly at the same level under the surface of the ground or in the depths, somewhat different from one another, provided only that the amount of water supplied to the cavity (II) is always considerably smaller than that to (I).

It can be easily seen according to the above consideration that the waters in the cavities are heated chiefly by the subterranean water of high temperature coming forth from the underground, because the temperature of the wall of the cavities lying not very deep under the ground may not be sufficiently high, and the water in the cavity (I) will be as necessary consequence heated more rapidly than that in (II). Therefore, even when the cavity (I) has attained a definite high temperature and the gradually

increasing tension of vapor occupying the upper part of the cavity above the water level causes at last an eruption of hot water from the larger orifice, the vapor tension in the cavity (II) is not yet sufficient to urge the water to eruption. As soon as the water of the pool is sucked in by the cavity (I) after the eruption from the larger orifice, the ebullition of water in the smaller cavity is brought about owing to the decreases of pressure under which it has been subjected and the water vapor generated will accumulate in its upper portion (g).

As the latent heat necessary for the evaporation of water must be supplied from the hot subterranean water flowing into the cavity and the increase of vapor tension in the space requires a definite time in order to attain its full strength sufficient for eruption, it follows that the eruption from the smaller orifice takes place in a definite lapse of time after that from the larger. It seems that the interval between the eruption from the larger orifice and the beginning of that from the smaller one has varied according to ages to some extent, *i. e.* about 30 seconds to 2 minutes.

The fact that the water level at the larger orifice does not descend after the eruption, holding a certain height for some time, agrees well with the above explanation.

Moreover, the volume of the cavity (I) may be considered to be somewhat larger than that of (II), as the determination of the amount of ejected water actually shows. As for the difference of the heights of water column thrown up into the air from two orifices, we may explain it by the presence of some obstacles near the orifices.

As for the recent quiescence of eruption from the smaller orifice there are of course several causes to be considered, but it is most probable that the narrow canals leading to the cavity (II) have become gradually narrower by deposition of mineral sinter, so that the hot water cannot heat the cavity to a degree sufficient to cause eruption.

Also the period of eruption has a tendency to become shorter in the course of time, *i. e.* the period of about 5 hours which was observed 50–60 years ago, has decreased to 3,5 hours in 1895, and at present again to only 1 hour and 6 minutes.

The decrease of the amount of jets as well as the activity of eruption of this geyser may probably be explained if we assume that the gradual erosion of the projected parts of the neck of the cavities, A and B, has decreased the effective volume of the cavity available for eruption, for the activity of intermittent eruption does not depend on the total volume of the cavity, but only on that portion (g and h) occupying its upper part; the erosion of the neck must cause not only the decrease of the effective volume of the cavity, but also the decrease of the period as well as the activity of eruption.

If these considerations be correct, the intermittent spring of Onikōbe will gradually shorten its period of eruption in the course of time and at last be deprived of its intermittent character after several decades.

Belgium.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|---------------|---|--|----------------------|-----------------|--|---|
| 1 | Chaudfontaine | in the valley of Vesdre | indifferent thermals containing NaCl and CaCl ₂ gaseous chalybeate chalybeate | 35.6°C. (96°F.) | 1000 | rheumatism, neuralgia and hepatic disorders. | very complete. Bath. Radioactive. ment. irradiation, electrotherapy, massage, etc. as to medicinal properties. Hydrated sulphur. Indicated for rheumatic and non-rheumatic. |
| 2 | Spa | at the junction of the Rivers Weyl and Picherotte | | 10°C. (50°F.) | | | |
| 3 | Haarlem | | | | | | |

France.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|------------------------|-----------------------|----------------|---|-----------------------------|-----------------|---|--|
| 1. Indifferent waters. | | | | | | | |
| 1 | Plombières | Vosges | silicate, arsenic | 25°-68.3°C. (77°-155°F.) | 1300 | gastro-intestinal, enteritis, chronic diarrhoea, functional nervous disorders, and appendicitis. | radioactive. |
| 2 | St. Amand | near Lille | | 21.1°-26.1°C. (70°-79°F.) | 100 | rheumatism, diseases of the joints, neuralgia, and chronic skin affections. | silicious sulphurated mineral baths. |
| 3 | Néris | near Montluçon | soft, faintly alkaline (NaHCO ₃) ₂ and saline | 46.1°-52.2°C. (115°-126°F.) | 1150 | nervous excitation, gastritis, enteralgia, diseases of spinal cord, neuralgia, rheumatism, dysmenorrhoea, constipation, and diseases of uterus. | radioactive, sodative. |
| 4 | Mont Dore | Auvergne | weak alkaline with a trace of As ₂ O ₃ and Fe HCO ₃) ₂ | 40°-46.7°C. (104°-116°F.) | 3400 | bronchitis, asthma, laryngitis, rheumatism, and chlorosis. | mineral spa |
| 5 | Dax | near Bayonne | | 31.1°-63.9°C. (88°-147°F.) | 130 | chronic rheumatic and joint affections, gout, neuralgia. | an important resort, both for summer and winter. |
| 6 | Bagnères de Bigorre | Pyrenees | calcareous (CaSO ₄ , MgSO ₄ , Na ₂ SO ₄) | 21.1°-50.6°C. (77°-123°F.) | 1800 | rheumatism, anaemia, disorders of the nervous system, catarrhal conditions of alimentary, respiratory, and genito-urinary systems. | mild climate. |
| 2. Muriated waters. | | | | | | | |
| 1 | Château-Guyon | Auvergne | some of the springs highly chalybeate with CO ₂ -gas | 48.9°C. (120°F.) | 1300 | dyspepsia, chronic, gastric, and intestinal catarrh, chronic constipation, congestion of the liver, tropical diseases. | radioactive. |
| 2 | Bourbon l'Archambault | near Moulins | also Na ₂ CO ₃ , CaCl ₂ and FeCO ₃ | | 870 | rheumatism, scrofula, and nervous diseases. | |
| 3 | Bourbon Lancy | Le Morvan | some of the springs are saline | 27.8°-57.8°C. (82°-130°F.) | 1780 | chronic rheumatism, chronic heart disease, and nervous conditions. | radioactive. |

| | | | | | | |
|---|---------------------|------------------------------|--|----------------------------------|------|---|
| 4 | Boullonne-les-Bains | Plateau de Langres | also LiCl, CaSO ₄ , NaBr, and trace of Fe, Mn | 43.3° 65.6° C.
(110° 150° F.) | 900 | scorbuta, rheumatism, catarrh of the nose, septal conditions, and treatment of chronic rhinitis, scurvy, iron deficiency, and uterine affections. |
| 5 | La Motte-les-Bains | Isère | also CaSO ₄ , NaBr | 51.1° 58.3° C.
124° 137° F. | | scorbuta, rheumatism, catarrh of the nose, and acute polydipsia. |
| 6 | Balaine | l'Étang de Thau | strong mineral | 12.2° 47.8° C.
54° 118° F. | 1200 | scorbuta, lymphatic affection, and constitutional weakness in women and children. |
| 7 | Salins | Junas, on the River Furieuse | also MgCl ₂ , Na ₂ SO ₄ , K ₂ SO ₄ and little of KBr and KI | cold | | |

Also La Bourboule, Salins-Montivert, etc.

3. Alkaline waters.

| | | | | | | |
|---|-----------------|---------------------|---|-----------------------------|------|--|
| 1 | Vichy | by the River Allier | strong alkaline (NaHCO ₃ ; more than 4%) with trace of As, F and free CO ₂ , some of the springs contain Fe | cold and thermal | 750 | dyspepsia, liver and bladder diseases, gallstones, catarrhes, uric acid disorders, and anæmia. |
| 2 | Vals | on the Volane | much free CO ₂ and little Fe | cold | 700 | kidney and bladder diseases, hyperæmia and gastric catarrh, diabetes, and rheumatism. |
| 3 | Roizat | Amvergne | gaseous with NaCl, Ca salts, little Fe and trace of As | 20°-35° C.
(68°-95° F.) | 1480 | anæmia, gout, dyspepsia, liver diseases, hepatic, nervous diseases, catarrh of the stomach, and diseases of women. |
| 4 | St. Nectaire | " | gaseous | 10°-43.0° C.
50°-111° F. | 2500 | chronic rheumatism, neuritis, dyspepsia, anæmia, catarrh, and catarrh of the stomach. |
| 5 | La Bourboule | Ponlague, " | mineral alkaline with Na ₂ AsO ₄ | thermal | 2780 | respiratory diseases, scurvy, chronic rheumatism, catarrh of the stomach, and diseases of women. |
| 6 | Evian-les-Bains | on the L. Geneva | weak alkaline, contain CaCO ₃ | cold | 1240 | gout, diabetes, and affections of urinary organs. |

Also Pongues-les-Bains, Lamalou, etc.

4. Sulphated mineral waters.

| | | | | | | |
|---|-----------------------|------------------------|--|--------------------------------|------|---|
| 1 | Evian-les-Bains | in the valley of Doubs | sulphated saline | 35° C. 95° F. | 1800 | chronic catarrhes, hyperæmia, and affections of the urinary organs. |
| 2 | Montet | near Evian-les-Bains | saline, lime; charged with CO ₂ gas | 35.6° C.
(96° F.) | 1000 | chronic catarrhes, catarrh of the stomach, and diseases of women. |
| 3 | St. Gervais-les-Bains | near Mont Blanc | weak sulphated salt | 30.5°-42.3° C.
103°-108° F. | 2075 | chronic catarrhes, catarrh of the stomach, and diseases of women. |

5. Calcareous waters.

| | | | | | | |
|---|-----------------|------------------------------------|--|------|------|---|
| 1 | Evian-les-Bains | Pluton of Mont-Fauilles, Vosges | CaCO ₃ , CaHCO ₃ , trace of Fe, As | cold | 1150 | gout, diabetes, and affections of urinary organs. |
| 2 | Evian-les-Bains | by the River Lône, 7 m. from Nancy | Na ₂ CO ₃ , CaHCO ₃ , CaHCO ₃ , MgHCO ₃ , NaHCO ₃ and free CO ₂ | " | 650 | hyperæmia, catarrh of the stomach, and diseases of women. |

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|---|--------------------|------------------------------------|--|--------------------------------|-----------------|--|--|
| 3 | Cransac | near the Volcano le Montet | | | 980 | | thermo-mineral sulphur springs |
| 4 | Marigny-les-Bains | Plateau of Monts Faucilles, Vosges | CaSO ₄ with Li | | 1246 | gout, gravel, diabetes, and liver and kidney derangements. | |
| 5 | Vittel | | "Grande Source," alkaline chloride;
"Source Salée," laxative) | cold | 1100 | "Grande Source"—mineral waters, gout, sciatica, neuralgia, neuritis, rheumatism, sciatica, lumbago, and abdominal aches; "Source Salée"—constipation, congestion of the liver, gall stones, and bronchial catarrh. | 2 principal springs, "Grande Source" and "Source Salée." |
| 6. Chalybeate waters. | | | | | | | |
| 1 | Lanvalou | Cevennes | chalybeate and weak alkaline | 15°-43.3°C.
(59°-110°F.) | 620 | rheumatism, sciatica, neuralgia, locomotor ataxia, chlorosis, and anaemia. | |
| 2 | Remes-les-Bains | 6 m. from Couiza-Montazels | chalybeate and saline | thermal | 1000 | chronic rheumatism, anaemia, and scrofula. | |
| 3 | Prezza | Larne Island | gaseous | | | | |
| Also some springs in Châtel Guyon, etc. | | | | | | | |
| 7. Sulphur waters. | | | | | | | |
| 1 | Aix-les-Bains | near L. Bourget, Savoy | Two chief springs; one contains alum, the other is simple sulphur spring | 4.38°-44.5°C.
(109°-112°F.) | 860 | gout, rheumatism, arthritis, neuritis, skin diseases, and affections of the mucous membranes. | Alum sulphur water used externally and for the other for inhalation. Aix-les-Bains is celebrated for treating skin-pimples, arising from injuries of the skin disease. |
| 2 | Marlioz | near Aix-les-Bains, " | very strong sulphur | cold | | laryngeal and bronchial affections. | for inhalation. |
| 3 | Challes-les-Bains | 3 m. from Chambéry, Pyrénées | very strong sulphur; 5% Na ₂ S with Br, I | " | 800 | chronic catarrh of the ear, chronic bronchitis, scrofula. | for internal use. |
| 4 | Bagnères de Luchon | | also alkaline springs; Na ₂ S 0.01-00.2% | 39.5°-53.3°C.
(103°-128°F.) | 3200 | chronic laryngitis and bronchitis, phthisis, gastric catarrh, rheumatism, and uric derangements. | radio-active. |
| 5 | Cauterets | | weak sulphur | 21.1°-32.2°C.
(70°-90°F.) | 2460 | chronic bronchitis, laryngitis, and pulmonary catarrh. | radio-active. |
| 6 | Eaux Bonnes | | | 25°-26.1°C.
(77°-79°F.) | 2200 | scrofula, rheumatism, respiratory and nervous disorders. | radio-active. |
| 7 | Eaux Chaudes | | | 27.2°-43.0°C.
(81°-111°F.) | 4200 | osteomyelitis and diseases of the bones and joints, chronic suppurative conditions, scrofula, syphilis, and anaemia. | |
| 8 | Barèges | | | 33.9°C.
(93°F.) | 2500 | all diseases of women, and functional and organic nervous diseases. | |
| 9 | St. Sauveur | | | 17.2°-77.2°C.
(63°-171°F.) | 2350 | rheumatism, scrofula, syphilis, chronic dry cough, and other skin diseases. | |
| 10 | Aix-les-Thermes | | | | | | |

| | | | | | | | |
|----|------------------|---------|------------------|---------------------------------|------|---|---------------------------|
| 11 | Amélie-les-Bains | " | alkaline sulphur | 21.1° 62.8° C.
(70° 145° F.) | 920 | dermatomy, skin diseases, rheumatism, asthma, and chronic catarrh of the respiratory system | in all in water |
| 12 | Uriage | " | saline sulphur | 27.2° 55° C.
(81° 131° F.) | 1350 | chronic skin diseases, scurfiness, and rheumatic affections, and constitutional weakness | suitable for internal use |
| 13 | Vernat-les-Bains | " | " | 7.8° 73.0° C.
(46° 165° F.) | 2000 | affections of the throat and chest, rheumatism, gout, sciatica, dyspepsia, neuritis, neurasthenia | |
| 14 | Gangne | Corsica | | thermal | | | |
| 15 | Pietrapola | " | | " | | | |

Germany.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|------------------------|----------------|---------------------------------------|--|----------------------------------|-----------------|---|--------------------------|
| 1. Indifferent waters. | | | | | | | |
| 1 | Wildbad | Black Forest | rich in N gas | 32.8° 40° C.
(91° 104° F.) | 1410 | chronic rheumatism and gout, diseases of spinal cord and urinary tract, chronic gastritis, diseases of women and skin disease. | |
| 2 | Schlagenthalen | near Mainz | rich in gases N and O ₂ | 27.8° 32.2° C.
(82° 90° F.) | 950 | neuritis, neurism, neuritis, dyspepsia, rheumatism, diseases of women, and chronic skin eruptions, and all diseases of structure. | very good and effective. |
| 3 | Baden-Wöden | Black Forest | Na ₂ SO ₄ , CaO, O ₂ , lit l, N gas | 25° 26.7° C.
(77° 80° F.) | 1450 | affections of throat and lungs, convalescence, rheumatism, neuralgia, the foot. | |
| 4 | Embsay | Pinnsau-Salsia (Reichensteiner range) | Na ₂ SO ₄ , H ₂ S | 20° 28.3° C.
(68° 83° F.) | 1500 | chronic rheumatism, neuritis, affections of the foot | very good |
| 5 | Warmbrunn | " | | 25° 42.8° C.
(77° 109° F.) | 1140 | chronic rheumatism, skin diseases, and respiratory affections | very good |
| 2. Miniated waters. | | | | | | | |
| 1 | Bad | near Frankfort on the Main | effervescent; 1.3% salt, rich in CO ₂ | 27.8° 35.3° C.
(82° 95.5° F.) | 470 | chronic rheumatism, neuritis, skin diseases, and all diseases of the foot | very good |
| 2 | Bad | near Mainz | Na ₂ CO ₃ , CaO, O ₂ , lit l, N gas | 27.8° 35.3° C.
(82° 95.5° F.) | 340 | chronic rheumatism, neuritis, skin diseases, and all diseases of the foot | very good |
| 3 | Bad | near Frankfort on the Main | effervescent; 1.3% salt, rich in CO ₂ | 27.8° 35.3° C.
(82° 95.5° F.) | 630 | chronic rheumatism, neuritis, skin diseases, and all diseases of the foot | very good |
| 4 | Bad | near Mainz | Na ₂ CO ₃ , CaO, O ₂ , lit l, N gas | 27.8° 35.3° C.
(82° 95.5° F.) | 380 | chronic rheumatism, neuritis, skin diseases, and all diseases of the foot | very good |

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remark |
|-----------------------|---------------|------------------|---|---|-----------------|---|--------------------------------------|
| 5 | Kissingen | Bavaria | effervescent; CaCl_2 and little FeCO_3 | cold | 650 | plethora and general debility, catarrh of the liver, chronic catarrh of the bladder, gonorrhoea, and kidney diseases. | pleasant, healthy, and invigorating. |
| 6 | Baden-Baden | Black Forest | also LiCl and trace of As | $45^\circ\text{--}70^\circ\text{C.}$
$113^\circ\text{--}158^\circ\text{F.}$ | 650 | rheumatism, gout, gastritis, and chronic catarrh of the bladder. | pleasant, healthy, and invigorating. |
| 7 | Oeynhausen | Westphalia | strong effervescent | $25\text{--}33.1^\circ\text{C.}$
$77\text{--}91.5^\circ\text{F.}$ | 230 | chronic diseases of spinal cord, tetanus, chorea, rheumatism, and bone diseases. | pleasant, healthy, and invigorating. |
| 8 | Schwaighausen | Tannus | also saline springs with (CO_2gas) | subthermal | 460 | chronic catarrhal affections of the respiratory organs, emphysema, scrofula, dyspepsia, and chronic heart disease. | pleasant, healthy, and invigorating. |
| 9 | Imbs | Prussia | | | | see also alkaline waters | |
| 3. Alkaline waters. | | | | | | | |
| 1 | Neuenahr | Rhine | gaseous | $36^\circ\text{--}48.0^\circ\text{C.}$
$86^\circ\text{--}120^\circ\text{F.}$ | 260 | liver and kidney affections, Bright's disease, gout, hysteria, and chronic gastritis, especially diabetes and uric acid troubles. | pleasant, healthy, and invigorating. |
| 2 | Salzmann | Prussian Silesia | saline, highly charged with $(\text{CO}_2\text{gas}; \text{NaHCO}_3\text{ } 2\%)$ | 10°C. (50°F.) | 1320 | acute and chronic catarrh of the respiratory organs, emphysema, and asthma, diseases of the digestive organs, kidneys, and bladder; gout, diabetes, and other uricemia. | pleasant, healthy, and invigorating. |
| 3 | Imbs | Prussia | mineralized-alkaline | $26.7^\circ\text{--}48.0^\circ\text{C.}$
$(80^\circ\text{--}120^\circ\text{F.})$ | 260 | chronic catarrh, gout, rheumatism, diseases of the digestive system, and urinary organs, and uterine disorders. | pleasant, healthy, and invigorating. |
| 4 | Elster | Saxony | gaseous alkaline sulphated and chalybeate | cold | 1550 | anemia, and chlorosis, hepatic dyspepsia, and gout. | pleasant, healthy, and invigorating. |
| 4. Calcareous waters. | | | | | | | |
| 1 | Wildungen | Hessen | gaseous, with CaHCO_3 , MgHCO_3 , and FeHCO_3 | cold | 980 | catarrh of the bladder, eczema, gravel, gout, and chronic nephritis. | pleasant, healthy, and invigorating. |
| 2 | Lippstange | Westphalia | weak | 21.1°C.
(70°F.) | 450 | pulmonary and other catarrhal conditions and pleurisy. | pleasant, healthy, and invigorating. |
| 3 | Elisen | Saxony | (some of the springs are strong sulphuretted) | cold | 230 | see also sulphur waters. | pleasant, healthy, and invigorating. |
| 5. Chalybeate waters. | | | | | | | |
| 1 | Schwaighausen | Prussian Silesia | gaseous alkaline | cold | 950 | same as for Spa (Baden-Baden). | pleasant, healthy, and invigorating. |
| 2 | Kudowa | Westphalia | gaseous | | 1310 | | pleasant, healthy, and invigorating. |
| 3 | Drilburg | Black Forest | " | cold | 730 | nervous diseases, rheumatism, gout, dyspepsia, and anemia. | pleasant, healthy, and invigorating. |
| 4 | Rippoldsau | Black Forest | " | | 1870 | anemia, debility, chlorosis, scrofula, nervous affections and diseases of women. | pleasant, healthy, and invigorating. |
| 5 | Pirmas | Westphalia | " (FeHCO_3 , CaHCO_3 , CaSO_4 and MgSO_4) | " | 420 | | pleasant, healthy, and invigorating. |

6. Sulphur waters.

| | | | | | | |
|---|-----------------|-------------------|-----------------------------------|-----------------------------------|-----|--|
| 1 | Aix-la-Chapelle | Aachen | strongly impregnated with sulphur | with 45.6° 74.5°C.
114° 169°F. | | chronic rheumatism, dropsy, catarrh of the skin, gout, paralytic, syphilitic, catarrh, and monalgia. |
| 2 | Weilach | Tannus | weak alkaline | cold | 440 | initial congestion, haemorrhoids, gout, in cases of urinary tract, ure, and urinary affections. |
| 3 | Nemtschaf | Prussia | strong sulphuretted | | 230 | |
| 4 | Meinberg | Lippe, Westphalia | | cold | | |
| 5 | Lüben | Saxony | strong sulphuretted | w | 230 | gouty and skin affections. |

Switzerland.

| No. | Spr. | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|---|-----------------|---------------|--|-------------------------------|-----------------|---|---|
| 1. Indifferent waters. | | | | | | | |
| 1 | Kaiserbrunn | | rich in N-gas | 31.7°-33.9°C.
(89°-93°F.) | 2000 | chronic rheumatism, sciatica, interstitial neuritis, catarrh, and nervous debility, and in the chest-lungs. | Exhaustive. Proffered in Rheuma, catarrh of the bladder, Biliary water, leucorrhoea, piles, flux, Erysipelas, the scaly skin, Acute Gout. |
| 2 | Fischbach-Pains | Fischbach | CaSO ₄ | 33.9° 51.1°C.
(93° 124°F.) | 4000 | sciatica, catarrh, and other rheumatic affections, chronic rheumatism, and neuritis. | |
| 2. Muriated waters. | | | | | | | |
| 1 | Baden | | strong muriated (nearly saturated brine). | | 806 | gouty, sciatic, rheumatism, gonorrhoea, dropsy. | |
| 2 | Schwefelballe | | | | | | |
| 3 | Rehna | Rehna Valley | strong muriated | | 1100 | badly healed wounds, ulcers, and other diseases of the skin, rheumatism, and neuralgia. | |
| 3. Alkaline waters. | | | | | | | |
| A. Leopold-Seelenbrunn muriated waters. | | | | | | | |
| 1 | Leopold | | alkaline saline | | 2025 | chronic rheumatism, sciatica, interstitial neuritis, catarrh, and nervous debility, and in the chest-lungs. | |
| 2 | Leopold | Lower Leopold | muriated, converging chalybeate, strong alkaline | cold | 3000 | chronic rheumatism, sciatica, interstitial neuritis, catarrh, and nervous debility, and in the chest-lungs. | |

4. Chalybeate waters.

| | | | | | | | |
|---|----------------|--|--------------------------------------|------|------|---|------------------------------|
| 1 | St. Moritz | betw. Lake St. Moritz and Lake Campfer | impregnated with CO ₂ gas | cold | 5820 | debility, chlorosis, catarrh, nervous affections, and for all complaints. | purest, most frequent remedy |
| 2 | Aquafossa | | Fe, As, Li | | | gout, rheumatism, aneurism, aneurism, and other diseases of women. | |
| 3 | San Bernardino | | | | | | |

Some of the springs in Tarrap-Schuls belong to this class.

5. Sulphur waters.

| | | | | | | | |
|---|----------|--------------|---|---------------------------|------|--|---------------------------|
| 1 | Baden | | | 36.7 52.2 C.
98 120 F. | 1230 | chronic rheumatism, gout, catarrh, and all the diseases of rheumatism, and all the diseases of the urinary system. | Proven since Roman times. |
| 2 | Schönach | | strong sulphuretted | 35 C. 95 F. | 1140 | chronic rheumatism, gout, catarrh, and all the diseases of the urinary system. | |
| 3 | Lavay | Rhone Valley | weak sulphur | 33.3 47.8 C.
92 118 F. | 1350 | chronic rheumatism, gout, catarrh, and all the diseases of the urinary system. | fine and hot sand baths |
| 4 | Yverdon | | alkaline sulphur | 23.6 C.
75 F. | 1430 | gout, chronic rheumatism, pyelitis, neuritis, pulmonary catarrh, leucorrhoea, and other diseases of women. | |
| 5 | Leuk | | strong sulphuretted, (CaSO ₄) | cold | 3630 | | |

6. Calcareous water.

| | | | | | | | |
|---|------------|--|------|------------------|--|---|-------------------------|
| 1 | Wessobrunn | | weak | 20.1 C.
79 F. | | affections of the reproductive organs, and pulmonary tuberculosis (in early stage). | used for drinking only. |
|---|------------|--|------|------------------|--|---|-------------------------|

Austria.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|------------------------|--------------|----------|------------------------------------|-------------------------------|-----------------|--|---|
| I. Indifferent waters. | | | | | | | |
| 1 | Gastein | Bohemia | | 25.6 - 49.5 C.
78 - 121 F. | 3400 | nervous diseases, gout, rheumatism, general debility, typhoid fevers, and other functional derangements, also employed as an "after cure." | the most radio-active of all thermals in Austria. |
| 2 | Teplitz | " | | 28.3 - 45.6 C.
83 - 114 F. | 730 | chronic rheumatism and gout, sciatica, lumbago, neuralgia, functional nervous disorders, also chronic skin diseases and scabies, and after results of severe injuries. | radio-active. |
| 3 | Johanniskbad | " | some of the springs are chalybeate | 20.5 C.
68 F. | 2070 | nervous diseases and neurasthenia, convalescence. | |

Italy.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-------------------------------------|---------------------------------|------------------------------|---|--------------------------------|-----------------|---|--|
| 1. Indifferent waters. | | | | | | | |
| 1 | Bormio | near Stervio Pass | CaCl_2 , CaSO_4 , MgSO_4 | 32.5° 40.6°C.
(90° 105°F.) | 4500 | gout, rheumatism, nervous and skin diseases. | Alpine po. |
| 2 | Battaglia | near Padua, Prov. of Venetia | little quantity of salt | 60° 71.1°C.
140° 160°F. | 1900 | rheumatism, gout, lumbago, sciatica, and neuralgia. | radioactive, natural vapour and fango baths. |
| 3 | Montesummano
"Grotto Giusti" | in Val-di-Nievole | | subthermal | | | Grotto contains a large cave warmed by thermal water and forms a natural vapour bath itself. |
| 2. Muriated waters. | | | | | | | |
| 1 | Alano | near Padua | weak muriated | 37.8° 83.9°C.
(100° 183°F.) | 100 | chronic joint affections. | radioactive, an "ergon" fango bath. |
| 2 | Salsomaggiore | | brine (130%) water, rich in Br and I | cold | 520 | scrofula, gout, rheumatism, lumbago, and pulmonary affections, diseases of women, and nervous affections. | called "Fiducia" fango bath. |
| 3 | Montecatini | near Pistoia | weak | 21.1° 31.1°C.
70° 88°F. | 920 | diseases of stomach and intestines, dyspepsia, diabetes, obesity, and gout. | |
| 3. Chalybeate and arsenical waters. | | | | | | | |
| 1 | Fischietti | near Pozzuoli | arsenical and chalybeate | | 5290 | | Alpine po. fango bath. |
| 2 | Croscio Reale | Piedmont | " | | 260 | see also sulphur water. | |
| 3 | Sciaccia | Sicily | alkaline chalybeate | 27.8° 37.8°C.
(82° 100°F.) | | | |
| 4. Muriated alkaline waters. | | | | | | | |
| 1 | Pozzuoli | Puteoli (Bay of Naples) | weak muriated | 26.7°C.
(80°F.) | | | famous "Solfatara" (an arsenical water) and numerous fango-baths (ergon, fango) and natural vapour baths (fango). |
| 2 | Ischia | Ischia | | up to 65°C.
(149°F.) | | gout, rheumatoid arthritis, and chronic affections. | "Solfatara" (an arsenical alkaline) at Ischia. Numerous hot springs known at present as the most radioactive hot springs in the world. |
| 3 | Casamiciola | | | | | | sea-baths. |
| 4 | Castellumare di Stabia | Bay of Naples | (some of the springs are ferruginous) | cold | | | |
| 5. Calcareous waters. | | | | | | | |
| 1 | Lucca | | | | | | |
| 2 | Chianciano | | | | | | |

6. Sulphur waters.

| | | | | | | | |
|---|-------------|------------------------|--|--|------|--|--|
| 1 | Aequi | 20 m. from Alessandria | CaSO_4 and I | 75°C.
(167°F.) | 450 | articular, peripheral neuritis, rheumatism, scabids. | climate free from damp. |
| 2 | Amadio | Piedmont | | | | | |
| 3 | La Porretta | Bologna | NaCl with trace of I, Br and As, also sulphuretted and carburetted hydrogen-gas | 32.8° – 35°C.
(91° – 95°F.) | 1100 | internal: hemorrhoid, abdominal phlegm, extraneous diseases, syphilis, rheumatism. | |
| 4 | Acetale | Sicily | | | | | |
| 5 | Aque Alluie | Province of Rome | | | | | |
| 6 | Viterbo | " | | | | | |
| 7 | Statena | Sicily | saline sulphur | 50° – 51.7°C.
(122° – 125°F.) | 260 | | curious vapour baths near Monte San Calogero, 3 m. East. |

Spain and Portugal.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|--------------------------------------|-------------------|---------------------------------------|---------------------------------------|--|-----------------|---|---------|
| 1. Indifferent waters. | | | | | | | |
| 1 | Estero | Province of Navarra, Spain | practically indif., feebly sulphurous | 46.1°C.
(115°F.) | | | |
| 2 | Caldas de Guarez | Spain | | " | | | |
| 2. Muriated waters. | | | | | | | |
| 1 | Caldas de Montbay | Laracena, Spain | | 48.9° – 71.1°C.
(120° – 160°F.) | 720 | respiratory, nervous, circulatory, cutaneous diseases | |
| 3. Bitter sulphated waters. | | | | | | | |
| 1 | Castell de Ferro | near the boundary of France and Spain | | | | | |
| 4. Calcareous and chalybeate waters. | | | | | | | |
| 1 | Castell de Ferro | near the boundary of France and Spain | | thermal | | | |

5. Alkaline waters.

| | | | | | | |
|---|--------------------|-------------------|---|-------------------------------|------|--|
| 1 | Vidago | N. Portugal | strong alkaline | 25° 28.2°C.
(77° 84.5° F.) | 5600 | no also sulphur water. |
| 2 | Panticoia | Pyrenees, Spain | some sulphur | | | |
| 1 | Panticosa | Pyrenees, Spain | Na ₂ S 0.02%; (some of the springs are alkaline) | 25° 28.2°C.
(77° 84.5° F.) | 5600 | bronchopulmonary, tuberculous, and respiratory troubles, dyspepsia, and skin diseases. |
| 2 | Carrizosa | near Malaga, " | | thermal | 2600 | |
| 3 | Montemayor y Lejar | near Salamanca, " | | | 2600 | gout, rheumatism, sciatica, neuralgia, skin diseases, beriberi, and syphilis. |
| 4 | Ledesma | " | | | | |
| 5 | Caldas da Rainha | Portugal | | | | |

6. Sulphur waters.

Sweden, Norway and Iceland.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|-------------------|--------------------------|---|----------------------|-----------------|-------------|----------------|
| 1 | Parestrand Syssel | Iceland | indifferent | 103.3°
(218° F.) | | | |
| 2 | Medewi | Sweden | chalybeate | | | | mineral baths |
| 3 | Pörla Brunn | " | " | | | | " |
| 4 | Paneverd | near Upsala, Sweden | " | | | | |
| 5 | Gustafsberg | near Uddevalla, " | " | | | | |
| 6 | Modum (St. Olafs) | near Christiania, Norway | effervescent chalybeate sulphurous chalybeate | | | | |
| 7 | Saudefjord | " | " | | | | |
| 8 | Laurvick | Iceland | ferruginous | | | | mineral baths. |
| 9 | Randamel | on the L. Lersjö, Sweden | | | | | |
| 10 | Loka | | | | | | |

Russia.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|-----------------|----------|---|----------------------|-----------------|-------------|---------|
| 1 | Schekhesa-Worsk | Caucasus | alkaline ferruginous | | 1800 | | |
| 2 | Pjatigorsk | " | some springs sulphurous, others ferruginous or alkaline | thermal | | | |

| | | | | | | |
|----|-------------|------------------------------|------------------------------------|------|--|-----------------------------|
| 3 | Katschenowa | near Moscow | effervescent chalybeate | | | hot medicinal |
| 4 | Odessa | Black Sea coast | | | | therapeutic, for rheumatoid |
| 5 | Astrakhan | on the Caspian Sea | | | | and treated mud baths |
| 6 | Cesed | in Livonia on the Baltic Sea | | | | |
| 7 | Chechoineck | Poland | strongly mineralized | | | |
| 8 | Longun | Caucasus | strong sulphurous alkaline sulphur | | | |
| 9 | Kemmer | near Riga | saline sulphur | cold | | peat mud medicinal |
| 10 | Stoypin | on the Kusum | mineralized sulphur | | | |
| 11 | Fusk | near Czakow | | | | |

Turkey, Bulgaria, etc.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|----------|------------------------------|-----------------------------|----------------------|-----------------|------------------------------------|---------------------|
| 1 | Saloniki | Macedonia | sulphur | thermal | | | hot baths |
| 2 | Calicut | Palestine | " | " | | | " |
| 3 | Adana | " | " | | | | medicinal hot water |
| 4 | Erzincan | Roumelia | " | thermal | | | for rheumatoid, for |
| 5 | Ida | Anatolia | " | 88° C. (190.4° F.) | | | water |
| 6 | Saparevo | Bulgaria | sulphur with organic matter | subthermal | | acidic, proved for rheumatoid, for | water |
| 7 | Varshet | " | sulphurous and calcareous | thermal | | for rheumatoid, for | water |
| 8 | Varshet | " | | subthermal | | for rheumatoid, for | water |
| 9 | Varshet | " | | thermal | | for rheumatoid, for | water |
| 10 | Varshet | " | | subthermal | | for rheumatoid, for | water |
| 11 | Varshet | " | sulphurous | thermal | | for rheumatoid, for | water |
| 12 | Varshet | near Philippopolis, Bulgaria | | " | | | for rheumatoid, for |
| 13 | Varshet | near Zlatna, Bulgaria | | thermal | | | for rheumatoid, for |
| 14 | Varshet | near Zlatna, Bulgaria | | " | | | for rheumatoid, for |
| 15 | Varshet | near Constantinople, Turkey | sulphur | thermal | | | for rheumatoid, for |
| 16 | Varshet | near Constantinople, Turkey | mineralized and ferruginous | " | | | for rheumatoid, for |

Greece.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|-------------|--------------------------------|----------------------|----------------------|-----------------|-------------|---------|
| 1 | Epidaurus | Epirus
Is. of Euboea | bitter sulphuretted | thermal | | | |
| 2 | Ilellopie | | " | " | | | |
| 3 | Adipsos | | " | " | | | |
| 4 | Lesbos | | " | " | | | |
| 5 | Thermopylae | Lesbos
in Is. of Aegean Sea | saline sulphur | | | | |
| 6 | Lepanto | | " | | | | |
| 7 | Katharsion | | " | thermal | | | |
| 8 | Ikaria | | " | " | | | |
| 9 | Kythnos | | " | " | | | |

Africa.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|-------------------|--------------------|-------------------------------|----------------------|-----------------|--|--|
| 1 | Helouan-les-Bains | near Cairo | muriated and muriated sulphur | 32.2°C.
(90°F.) | 200 | chronic, scanty, rheumatic conditions, anaemia, and chlorosis. | the "Cimes," formed by the masses of CaCO ₃ are a unique feature of these baths, some of them over 40 ft. high. |
| 2 | Bains-de-la-Reine | near Oran, Algiers | muriated | 54.4°C.
(130°F.) | | | |
| 3 | Hammam Melouan | " | " | 54.4°C.
(130°F.) | | | |
| 4 | Hammam-r'Irha | " | calcareous and chalybeate | hyper-thermal | 1800 | scout, rheumatism. | |
| 5 | Hammam-Meskoutin | " | | 95°C. (203°F.) | 1000 | | |
| 6 | Mara | Abyssinia | bitter sulphur | warm | | | |
| 7 | Okme | Nubia | | | | | |

India.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|------------|----------|----------------------|----------------------|-----------------|-------------|---------|
| 1 | Salt Range | Punjab | muriated | | | | |
| 2 | Nazgonda | Simla | chalybeate sulphur | | | | |
| 3 | Danera | | | | | | |

| | | | | | | |
|---|--------------|-------------------------------|------------------------------|---------------------|------|---------|
| 4 | Sona | near Delhi | " | 97.2°C.
(207°F.) | 5587 | London. |
| 5 | Alumikarn | Jalandhar Doab | " (some sp. contain I or Br) | thermal | | |
| 6 | Jawala Mukki | Kangra District of the Punjab | " (some sp. contain I) | | | |
| 7 | Canica | Ceylon | " | | | |
| 8 | Malacca | | " | | | |

New Zealand.

| No. | Spa | Location | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|-----|-----------|---------------------------|--|--------------------------|-----------------|--|---|
| 1 | Te Aroha | Auckland, N. Island | sulphuretted, muriated alkaline, and effervescing carbonated | 76.7°C. (170°F.) | | | |
| 2 | Tarapo | " | the Crow's Nest Spring ¹) is muriated, the Rushing ²) contains free acid, the others ³) are sulphurous or alkaline silicious | 87.8°C. (190°F.) thermal | 1500 | | |
| 3 | Piscine | " | "Rachel's Bath" soft saline silicious | 95.6°C. (194°F.) | | | said as the finest bath in the S. hemisphere. |
| 4 | Rotorua | S. shore of L. Rotorua, " | sulphuretted silicious, of which, some are strongly alkaline, while the others contain free H ₂ SO ₄ | thermal | 902 | acid waters, richly sulphuretted and containing silicious derivatives in the form of silica. | |
| 5 | Ohinemutu | Rotorua, " | muriated, containing carburetted and sulphuretted hydrogen | | | | |
| 6 | Haamer | S. Island | | | | | |

United States of America.

| No. | Spa | Location (State) | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|---------------------|-------------|------------------|--------------------------|----------------------|-----------------|-------------|---------|
| I. Muriated waters. | | | | | | | |
| 1 | Hot Springs | California | weakly muriated alkaline | 37.8°C. 100 F. | | | |
| 2 | Hot Springs | New York | " | cold | | | |

| No. | Spa | Location (State) | Composition of water | Temperature of water | Altitude in ft. | Indications | Remarks |
|--------------------------------|-------------------------------------|------------------|-------------------------------------|----------------------|-----------------|-------------|--|
| 3 | Hawthorn | New York | Lime and magnesia | | | | |
| 4 | Blue Lick Spring | Kentucky | | | | | |
| 5 | St. Catharine's Well | Ontario | | | | | |
| 6 | Caledonia Springs | " | | | | | |
| 2. Sulphated or bitter waters. | | | | | | | |
| 1 | Crab Orchard | Kentucky | | | | | |
| 3. Sulphuretted waters. | | | | | | | |
| 1 | Ridgfield | New York | | | | | saladation is used. |
| 2 | The Salt Sulphur Spring | W. Virginia | | | | | |
| 3 | Paroquet Spring | Kentucky | | | | | |
| 4 | Sharon and White Sulphur Springs | New York | | | | | |
| 5 | Sandwich Springs | Ontario | | | | | |
| 4. Acid waters.* | | | | | | | |
| 1 | Oak Orchard Spring | New York | free H ₂ SO ₄ | hot | | | abs. acid, arsenic, carbonated, silicic acid, and arsenic. |
| 2 | Yellowstone National Park | Wyoming | free HCl | " | 7000 | | |
| 3 | California Geyser (Lemonade Spring) | California | " | " | | | |

* Texas has some strong acid springs, and also Tuscarora in Canada.

* Texas has some strong acid springs, and also Tuscarora in Canada.

PART II (A)

- (I) The Tables of Analyses, Radio-activity, Electric Conductivity, Flow of Water, etc., of Mineral Springs, arranged geographically.

NOBORIBETSU

Location.—Noboribetsu mtn., Horokanai, 100 miles from Tokyo.

The springs issue from the Tufaceous Clay.

The hot spring "Taki-no-yu"

Analysis (calculated from the original numbers).

Analysed by The Tōkyō Imperial Hygienic Laboratory. 1910.

Specific gravity: 1,0001 at 14° C.

Temperature: *71,5° C.

Total residue: ca. 0,81.

Flow of water: ca. 54000 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|----------------|------------|-----------------------|
| Hydrogen ion (H ⁺) | 0,00241 | 2,38703 | 2,38703 |
| Potassium ion (K ⁺) | 0,01302 | 0,33597 | 0,33597 |
| Sodium ion (Na ⁺) | 0,07791 | 3,38004 | 3,38004 |
| Ammonium ion (NH ₄ ⁺) | 0,00662 | 0,36615 | 0,36615 |
| Calcium ion (Ca ⁺⁺) | 0,02669 | 0,66725 | 1,33450 |
| Magnesium ion (Mg ⁺⁺) | 0,00682 | 0,27997 | 0,55994 |
| Ferric ion (Fe ⁺⁺⁺) | 0,01818 | 0,32464 | 0,97392 |
| Aluminium ion (Al ⁺⁺⁺) | 0,01420 | 0,52399 | 1,57197 |
| | | | 10,90952 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0,05730 | 1,61636 | 1,61636 |
| Hydrosulphate ion (HSO ₄ ⁻) | 0,23157 | 2,38560 | 2,38560 |
| Sulphate ion (SO ₄ ⁻) | 0,33177 | 3,45378 | 6,90756 |
| | | 0,78649 | 15,72078 |
| Boric acid (meta) (HBO ₂) | 0,00227 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,22671 | | |
| | | | 1,01547 |

| | Grams |
|---|----------|
| Potassium chloride (KCl) | 0,02481 |
| Sodium chloride (NaCl) | 0,05368 |
| Ammonium chloride (NH ₄ Cl) | 0,01960 |
| Sodium sulphate (Na ₂ SO ₄) | 0,17502 |
| Calcium sulphate (CaSO ₄) | 0,09078 |
| Magnesium sulphate (MgSO ₄) | 0,03371 |
| Ferric sulphate [Fe ₂ (SO ₄) ₃] | 0,06495 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0,09001 |
| Free sulphuric acid (H ₂ SO ₄) | 0,23157 |
| Boric acid (meta) (HBO ₂) | 0,00227 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,22671 |
| | 1,004311 |

The spring thus may be classified as "acid vitriol spring".

Radio-activity. 0,18 Mache's units in 1 litre of water at 14,5° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, Oct. 16, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 14,58$.

The temperature of the water was measured at the uppermost end of the bore, about 100 m. from the bath.

The hot spring "Ōkawa-no-yu"

Analysis (calculated from the original numbers).

Analysed by The Tōkyō Imperial Hygienic Laboratory. 1910.

Specific gravity: 1,0011 at 13° C.

Temperature: 61° C.

Total residue: ca. 0,88.

Flow of water: ca. 450 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K ⁺) | 0,01719 | 0,43908 | 0,43908 |
| Sodium ion (Na ⁺) | 0,10500 | 4,59002 | 4,59002 |
| Ammonium ion (NH ₄ ⁺) | 0,00037 | 0,02046 | 0,02046 |
| Calcium ion (Ca ⁺⁺) | 0,07534 | 1,88350 | 3,76700 |
| Magnesium ion (Mg ⁺⁺) | 0,00717 | 0,29434 | 0,58868 |
| Ferrous ion (Fe ⁺⁺) | 0,00028 | 0,00500 | 0,01000 |
| | | | 10,41524 |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0,03276 |
| Sodium chloride (NaCl) | 0,23231 |
| Ammonium chloride (NH ₄ Cl) | 0,00110 |
| Sodium sulphate (Na ₂ SO ₄) | 0,04400 |
| Calcium sulphate (CaSO ₄) | 0,17445 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,09741 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,04300 |

Anions.

| | | | |
|--|---------------------|----------|---------|
| Chlorine ion (Cl^-) | 0.15708 | 4.43103 | 4.43103 |
| Sulphate ion (SO_4^{2-}) | 0.15280 | 1.50161 | 3.18322 |
| Hydrocarbonate ion (HCO_3^-) | 0.10988 | 1.80099 | 1.80099 |
| | 0.02600 | 15.05603 | 9.41524 |
| Boric acid (meta) (HBO_2) | 0.00057 | | |
| Silicic acid (meta) (H_2SiO_3) | 0.25523 | | |
| | 0.88180 | | |
| Free carbon dioxide (CO_2) | 0.00153 . . 0.03477 | | |
| | 0.88333 | | |

| | |
|---|---------|
| Ferrous bicarbonate [$\text{Fe}(\text{HCO}_3)_2$] | 0.00089 |
| Boric acid (meta) (HBO_2) | 0.00057 |
| Silicic acid (meta) (H_2SiO_3) | 0.25523 |
| | 0.88181 |
| Free carbon dioxide (CO_2) | 0.00153 |
| | 0.88334 |

The spring thus may be classified as “simple thermal”.

Radio-activity. 0.30 Mache's units in 1 litre of water at 15.6° C.

(Kohl.-Löw, fontact.; Y. Kinugasa, Oct. 16, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 7.78$.

The hot spring “Manju-no-yu” No. 1.

Analysis (calculated from the original numbers).

Analysed by The Tōkyō Imperial Hygienic Laboratory. 1910.

Specific gravity: 1.0009 at 14.5° C. Temperature: 50° C.

Total residue: ca. 0.84. Flow of water: ca. 540 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of “Ōkawa-no-yu”, may be classified as “simple thermal”.

Radio-activity. 0.39 Mache's units in 1 litre of water at 16° C.

(Kohl.-Löw, fontact.; Y. Kinugasa, Oct. 16, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 7.44$.

The hot spring “Shio-no-yu (Shimo-no-yu)”

Analysis (calculated from the original numbers).

Analysed by The Tōkyō Imperial Hygienic Laboratory. 1910.

Specific gravity: 1.0041 at 15° C. Temperature: 76° C.

Total residue: ca. 4.51. Flow of water: ca. 630 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-e quivalents |
|--|---------|------------|------------------------|
| Potassium ion (K^+) | 0.09535 | 2.43550 | 2.43550 |
| Sodium ion (Na^+) | 1.18642 | 51.47158 | 51.47158 |
| Ammonium ion (NH_4^+) | 0.00177 | 0.09790 | 0.09790 |
| Calcium ion (Ca^{++}) | 0.32809 | 8.20225 | 16.40450 |
| Magnesium ion (Mg^{++}) | 0.02135 | 0.87644 | 1.75288 |
| Ferrous ion (Fe^{++}) | 0.00028 | 0.00500 | 0.01000 |
| | | | 72.17236 |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.18169 |
| Sodium chloride (NaCl) | 3.01122 |
| Ammonium chloride (NH_4Cl) | 0.00524 |
| Calcium chloride (CaCl_2) | 0.81065 |
| Calcium bicarbonate [$\text{Ca}(\text{HCO}_3)_2$] | 0.14455 |
| Magnesium bicarbonate [$\text{Mg}(\text{HCO}_3)_2$] | 0.12831 |
| Ferrous bicarbonate [$\text{Fe}(\text{HCO}_3)_2$] | 0.00086 |
| Boric acid (meta) (HBO_2) | 0.00114 |
| Silicic acid (meta) (H_2SiO_3) | 0.25605 |
| | 4.53971 |
| Free carbon dioxide (CO_2) | 0.00804 |
| | 4.54775 |

Anions.

| | | | |
|--|---------------------|-----------|----------|
| Chlorine ion (Cl^-) | 2.43286 | 68.62793 | 68.62793 |
| Hydrocarbonate ion (HCO_3^-) | 0.21625 | 3.54443 | 3.54443 |
| | 4.28237 | 135.26103 | 72.17236 |
| Boric acid (meta) (HBO_2) | 0.00114 | | |
| Silicic acid (meta) (H_2SiO_3) | 0.25605 | | |
| | 4.53951 | | |
| Free carbon dioxide (CO_2) | 0.00804 . . 0.18273 | | |
| | 4.54760 | | |

Besides these trace of sulphate ion.

The spring thus may be classified as “common salt spring”.

Radio-activity. 0.46 Mache's units in 1 litre of water at 13.8° C.

(Kohl.-Löw, fontact.; Y. Kinugasa, Oct. 16, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 49.80$.

Table 1.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparent specific gravity | Electric
conductivity
% to 1 | Date when the
water was taken | Examined by |
|-----|----------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|---------------------------|------------------------------------|----------------------------------|-------------|
| | | | | | in
Miche's unit | at
(temp. in C.) | | | | |
| 1 | Taki-no-yu | acid vitriol | 71.5° | ca.
54.00 | 0.18 | 14.5° | K&L | 14.58 | Oct. 16, 1913 | Y. Kitagawa |
| 2 | Okawa-no-yu | simple | 61.0° | 450 | 0.30 | 15.0 | " | 7.78 | " | " |
| 3 | Manju-no-yu (No. 1) | " | 50.0° | 540 | 0.39 | 10.0 | " | 7.44 | " | " |
| 4 | " (No. 2) | sulphur | 48.0° | " | 0.21 | 15.0 | " | 6.07 | " | " |
| 5 | Shio-no-yu (<i>Kami-no-yu</i>) | common salt | 94.0° | 180 | 0.68 | 14.5° | " | 4.27 | " | " |
| 6 | " (<i>Shimo-no-yu</i>) | " | 76.0° | 630 | 0.46 | 13.5° | " | 49.80 | " | " |
| 7 | Me-no-yu | vitriol | 97.0° | 324 | " | " | " | " | " | " |

* These springs are for the public use.

KARURUSU

Location.—Noboribetsu-mura, Horobetsu-gun, Province Iburi, Hokkaidō.

The hot spring “Tsuru-no-yu”

Analysis (calculated from the salt table).

Analysed by The Tōkyō Imperial Hygienic Laboratory. 1900.

Total residue: ca. 0.96.

In 1 kilogram of the mineral water are contained:

Cations.

| | Grams | Milli-mols | Milligram-equivalents |
|---|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0107 | 0.2734 | 0.2734 |
| Sodium ion (Na ⁺) | 0.1722 | 7.4707 | 7.4707 |
| Calcium ion (Ca ⁺⁺) | 0.0718 | 1.7905 | 3.5810 |
| Magnesium ion (Mg ⁺⁺) | 0.0012 | 0.0493 | 0.0986 |
| | | | 11.4237 |

Anions.

| | | | |
|--|--------|---------|---------|
| Chlorine ion (Cl ⁻) | 0.0996 | 2.8096 | 2.8096 |
| Sulphate ion (SO ₄ ^{''}) | 0.3091 | 3.2188 | 6.4376 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.1330 | 2.1800 | 2.1800 |
| | 0.7976 | 17.7923 | 11.4272 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2038 | | |
| | 1.0014 | | |
| Free carbon dioxide (CO ₂) | 0.1525 | 3.4650 | |
| | 1.1539 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 0.1644 |
| Potassium sulphate (K ₂ SO ₄) | 0.0238 |
| Sodium sulphate (Na ₂ SO ₄) | 0.3312 |
| Calcium sulphate (CaSO ₄) | 0.1024 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.1684 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0074 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2038 |
| Free carbon dioxide (CO ₂) | 0.1525 |
| | 1.1539 |

Besides these a small quantity of ferrous, aluminium ion and trace of hydrophosphate ion.

The spring thus may be classified as “simple thermal”.

Table 2.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|--------------------------------|----------------|--------------------------|---|---------------------------------|-------|--------------------|--|----------------------------------|-------------|
| 1 | Tsuru-no-yu <i>Font. No. 1</i> | simple | 60.0° | ca. 210 | 0.38 | 15.2° | K.&L. | 11.10 | Oct. 15, 1913 | Y. Kinugasa |
| 2 | " " <i>No. 2</i> | " | 58.0° | 210 | 0.57 | 13.1° | " | 11.10 | " | " |
| 3 | Matsu-no-yu | " | 59.0° | " | 0.47 | 14.4° | " | 9.69 | " | " |
| 4 | Kame-no-yu | " | 52.0° | 108 | 0.37 | 13.6° | " | 10.60 | " | " |
| 5 | Tsuru-no-yu <i>Font. No. 3</i> | " | 48.0° | — | — | — | — | 10.73 | " | " |

JŌZANKEI

Location.—Hiragishi-mura, Toyohira-machi, Sapporo-gun, Province Ishikari, Hokkaidō.
The springs issue from the Liparite.

The hot spring "Moto-no-yu"

Analysis.

Analysed by The Sapporo Hospital. 1889.

Temperature: 91° C.

Total residue: ca. 3.21.

Flow of water: ca. 378 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

Hypothetical form of combination.

| | Grams | | Grams |
|--|-------|---|-------|
| Potassium (K) | 0.031 | Sodium chloride (NaCl) | 2.570 |
| Sodium (Na) | 1.064 | Sodium sulphate (Na ₂ SO ₄) | 0.067 |
| Calcium (Ca) | 0.150 | Potassium carbonate (K ₂ CO ₃) | 0.050 |
| Magnesium (Mg) | 0.013 | Sodium carbonate (Na ₂ CO ₃) | 0.070 |
| Iron (Fe) | 0.006 | Sodium silicate (Na ₄ SiO ₄) | 0.068 |
| Chlorine (Cl) | 1.560 | Calcium carbonate (CaCO ₃) | 0.371 |
| Sulphuric anhydride (SO ₃) | 0.054 | Magnesium carbonate (MgCO ₃) | 0.071 |
| Silica (SiO ₂) | 0.052 | | 3.267 |

The spring may be classified as "common salt spring".

Radio-activity. 0.92 Mache's units in 1 litre of water at 14.1° C.

(Kohl-Löw. fontact.; Y. Kinugasa, Oct. 12, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 50.67$.

Table 3.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|-------------|----------------|--------------------------|---|---------------------------------|-------|--------------------|--|----------------------------------|-------------|
| 1 | Moto-no-yu | common salt | 91.0° | ca. 378 | 0.92 | 14.1° | K.&L. | 50.67 | Oct. 12, 1913 | Y. Kinugasa |
| 2 | Naka-no-yu | simple | 80.0° | 810 | 0.77 | 14.8° | " | 36.27 | " | " |
| 3 | Shika-no-yu | " | 82.0° | 1458 | — | — | — | — | — | " |

AOYAMA

Location. Minamishimbetsu-mura, Isoyama, Shimane Prefecture, Honshu, Japan.

The hot spring "Furōkaku-no-yu" No. 1.

Analysis.

Analysed by K. Saki, apothecary, Itoyama, Shimane Prefecture.

Specific gravity: 1.00248 at 15° C. Temperature: 44° C.
Total residue: ca. 2.97. Flow of water: ca. 2772 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|---------|--|----------------|
| Potassium chloride (KCl) | 0.18593 | Formic acid and albumen (C ₂ H ₃ O ₂ + N ₂) | 0.0005 |
| Sodium chloride (NaCl) | 1.72372 | Silica (SiO ₂) | 0.1458 |
| Sodium bicarbonate (NaHCO ₃) | 0.30642 | | <u>3.28569</u> |
| Calcium sulphate (CaSO ₄) | 0.23215 | Free carbon dioxide (CO ₂) | <u>0.2474</u> |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.38653 | | <u>3.52933</u> |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.29244 | | |

The spring may be classified as "earthy common salt spring".

Radio-activity. 1.39 Mache's units in 1 litre of water at 11.5° C.

(Kohl-Low, contact. Y. Kimugasa, Oct. 18, 1913.)

Electric conductivity. $218 \times 10^{-4} = 36.17$.

Table 4.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | Apparatus employed | Electric conductivity 218×10^{-4} | Date when the water was taken | Examined by |
|-----|----------------------------------|-------------------|-----------------------|---|-------------------------------------|-------|--------------------|--|---------------------------------|-------------|
| 1 | Furōkaku-no-yu A. 1 | earthy comon salt | 44.0° | ca. 2772 | 1.39 | 11.5 | K. & L. | 36.17 | Oct. 18, 1913 | Y. Kimugasa |
| 2 | " A. 2 | " | 42.0° | — | 1.93 | 10.5 | " | 25.03 | " 19, " | " |
| 3 | Miyagawa-no-yu | " | 42.0° | 1530 | 0.54 | 11.5° | " | 24.45 | " 18, " | " |
| | Gas evolving from Miyagawa-no-yu | | | | Emanation per litre of gas at 0° C. | | E. & S. | | Date when the gas was collected | |
| | | | | | 3.54 | | | | | |
| | | | | | 4.08 | | | | | |
| | | | | | 4.16 | | | | " 19, " | " |

Determined according to the authors' original direction.

† Correction for the absorption of the radiation by the wall of the ionisation chamber was taken.

‡ Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made.

YUNOKAWA

Location. Yunokawa-mura, Kamekura, Prefecture Oshima, Hokkaido.

The springs issue from the Liparite overlaid by Alluvial Sand.

The hot spring "Taki-no-yu"

Analysis.

Analysed by Prof. Dr. T. Shimoyama, R. Kimura, and Y. Kimugasa.

Specific gravity: 1.0051 at 15° C. Temperature: 49° C.

Total residue: ca. 7.03. Flow of water: ca. 200 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained.

| | Grams | | Grams |
|--------------------------|-------|--|-------|
| Potassium (K) | 0.064 | Sulphuric anhydride (SO ₃) | 0.761 |
| Sodium (Na) | 1.735 | Silica (SiO ₂) | 0.054 |
| Calcium (Ca) | 0.644 | Organic substances | 0.065 |
| Magnesium (Mg) | 0.177 | Free carbon dioxide (CO ₂) | 0.615 |
| Chlorine (Cl) | 3.409 | | |

Besides these trace of iron and phosphoric acid.

The spring may be classified as "earth-muriated common salt spring".

Radio-activity. 7.35 Mache's units in 1 litre of water at 16.9° C.

(Kohl-Töw. fontact.; Y. Kinugasa, Oct. 8, 1913).

82.31 $\frac{1}{2}$ Mache's units in 1 litre of gas at 0° C.

(Eng.-Siev. fontact.; Y. Kinugasa, Oct. 10, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 83.38$.

Table 5.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres. | Emanation per
litre of water
in Mache's units
at (temp. in C.) | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|---|-------------------------------|--------------------------|--|---|-------|--------------------|--|---------------------------------------|-------------|
| 1 | Taki-no-yu
(<i>Rinchō-kwan</i>) | earth-muriated
common salt | 50.0° | ca. 216 | 7.35 | 16.9° | K.&L. | 83.38 | Oct. 8, 1913 | Y. Kinugasa |
| 2 | Shin-yu No. 1
(<i>Rinchō-kwan</i>) | earthy common
salt | 48.0° | — | 8.48 | 13.9° | " | 56.07 | " 9, " | " |
| 3 | " No. 2
(<i>Rinchō-kwan</i>) | " | 50.0° | — | 6.51 | 9.0° | " | 70.63 | " " | " |
| 4 | Chōju-yu (<i>Rinchō-kwan</i>) | " | 49.0° | — | 8.21 | 18.2° | " | 74.25 | " 8, " | " |
| 5 | Senshin-kwan-no-yu
No. 1 | " | 40.5° | — | 11.24 | 15.0° | " | 45.11 | " 10, " | " |
| 6 | " No. 2 | " | 48.0° | — | 13.21 | 14.6° | " | 50.02 | " 11, " | " |
| 7 | Hōmei-kwan-no-yu | " | 46.0° | 216 | 11.32 | 15.2° | " | 69.67 | " " | " |
| 8 | Toyokawa-no-yu | " | 42.5° | 216 | 7.96 | 14.5° | " | 42.04 | " 10, " | " |
| 9 | Tōyō-kwan-no-yu | " | 45.0° | 216 | 3.02 | 18.5° | " | 59.21 | " 8, " | " |
| 10 | Mura-yu | " | 50.0° | 216 | 7.47 | 12.7° | " | 76.87 | " 10, " | " |
| 11 | Rikuzō-no-yu $\frac{1}{2}$ | " | 66.0° | — | 5.54 | 18.4° | " | 83.28 | " 8, " | " |
| 12 | Taisho-kwan-no-yu $\frac{1}{2}$ | " | 43.5° | — | 3.55 | 17.4° | " | 70.35 | " " | " |
| 13 | Yoshikawa-no-yu $\frac{1}{2}$ | " | 63.5° | — | — | — | — | 84.83 | " " | " |
| | Gas evolving from: | | | | Emanation per litre
of gas at 0° C. | | | | Date when the
gas was
collected | |
| 1 | Taki-no-yu
(<i>Rinchō-kwan</i>) | } | | | 70.04 * | | E.&S. | | | |
| | | | | | 80.76 † | | | | " 10, " | " |
| | | | | | 82.31 ‡ | | | | | |
| 2 | Shin-yu No. 2
(<i>Rinchō-kwan</i>) | } | | | 63.14 * | | " | | | |
| | | | | | 72.78 † | | | | " " | " |
| | | | | | 74.17 ‡ | | | | | |
| 3 | Rikuzō-no-yu | } | | | 85.97 * | | " | | | |
| | | | | | 99.14 † | | | | " 9, " | " |
| | | | | | 101.03 ‡ | | | | | |

* Determined according to the authors' original direction.

† Correction for the absorption of the radiation by the wall of the ionisation chamber was made.

‡ Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made.

These springs are located on the sea beach of Nezaki, about 12 m. distant from Yunokawa.

ASAMUSHI

Location.—Nonai mura, Higashitsuru-ri, Ito-gun, Aomori-ken.

The springs issue from the Andesite overlaid by Alluvium.

The hot spring “Hadaka-no-yu”

Analysis (calculated from the salt table).

Analysed by The Medical Department of the Army.

Temperature: 61.5° C.

Flow of water: ca. 360 hectolitres in 24 hours.

Total residue: ca. 1.34.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K') | 0.0082 | 0.2097 | 0.2097 |
| Sodium ion (Na') | 0.1895 | 8.2213 | 8.2213 |
| Calcium ion (Ca'') | 0.1844 | 4.5985 | 9.1970 |
| Magnesium ion (Mg'') | 0.0050 | 0.2053 | 0.4106 |
| | | | 18.0386 |
| Anions. | | | |
| Chlorine ion (Cl') | 0.1988 | 5.6079 | 5.6079 |
| Sulphate ion (SO ₄ '') | 0.5914 | 6.1565 | 12.3130 |
| Hydrocarbonate ion (HCO ₃ ') | 0.0058 | 0.0951 | 0.0951 |
| | 1.1831 | 25.0943 | 18.0160 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0792 | | |
| | 1.2623 | | |

| | Grams |
|---|--------|
| Sodium chloride (NaCl). | 0.3280 |
| Potassium sulphate (K ₂ SO ₄) | 0.0100 |
| Sodium sulphate (Na ₂ SO ₄) | 0.1860 |
| Calcium sulphate (CaSO ₄) | 0.6250 |
| Magnesium sulphate (MgSO ₄) | 0.0189 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0070 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0792 |
| | 1.2623 |

Besides these trace of ferrous, hydrophosphate ion and boric acid.

The spring thus may be classified as “**sulphated bitter spring**”.

Radio-activity. 0.77 Mache's units in 1 litre of water at 19.7°C.

(Kohl.-Löw. fontact.; Y. Kinugasa, Oct. 6, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 13.72$.

The hot spring “Ō-yu”

Analysis.

Analysed by R. Minami, apothecary of Aomori. 1909.

Temperature: 66° C.

Flow of water: ca. 270 hectolitres in 24 hours.

Total residue: ca. 1.15.

The spring, being in its composition nearly the same as that of “Hadaka-no-yu”, may be classified as “**bitter spring**”.

Radio-activity. 1.73 Mache's units in 1 litre of water at 16.9° C

(Kohl.-Löw. fontact.; Y. Kinugasa, Oct. 6, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 11.01$.

The hot spring “Tsubaki-no-yu” No. 1.

Analysis.

Analysed by The Hirosaki Garrison Hospital. 1912.

Specific gravity: 1.00442 at 15° C.

Temperature: 79° C.

Total residue: ca. 1.23.

The spring, being in its composition nearly the same as that of “Hadaka-no-yu”, may be classified as “**sulphated bitter spring**”.

Radio-activity. 1.13 Mache's units in 1 litre of water at 16° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, Oct. 6, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 13.25$.

The hot spring “Yanagi-no-yu”

Analysis.

Analysed by The Hirosaki Garrison Hospital. 1912.
 Specific gravity: 1.00418 at 15° C. Temperature: 63° C.
 Total residue: ca. 1.25. Flow of water: ca. 360 hectolitres in 24 hours.
 The spring, being in its composition nearly the same as that of “Hadaka-no-yu”, may be classified as
 “sulphated bitter spring”.

Radio-activity. 0.47 Mache's units in 1 litre of water at 15.8° C.
 (Kohl-Löw. fontact.; Y. Kinugasa, Oct. 6, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 13.82$.

Table 6.

| No. | Spring | Classification | Temp. of spring
in °C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
temp. in °C. | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by | |
|-----|------------------------------------|------------------|---------------------------|---|--|--------------------|--|----------------------------------|--------------|-------------|
| 1 | Hadaka-no-yu
<i>Public bath</i> | sulphated bitter | 61.5° | ca. 300 | 0.77 | 10.7° | K. & L. | 13.72 | Oct. 6, 1913 | Y. Kinugasa |
| 2 | O-yu <i>Public bath</i> | bitter | 66.0° | 270 | 1.73 | 16.9° | .. | 11.01 | .. | .. |
| 3 | Tsubaki-no-yu <i>No. 1</i> | sulphated bitter | 79.0° | 519 | 1.13 | 16.0° | .. | 13.25 | .. | .. |
| 4 | .. <i>No. 2</i> | .. | 68.5° | | — | — | .. | 11.94 | .. | .. |
| 5 | Yanagi-no-yu | .. | 63.0° | 300 | 0.47 | 15.8° | K. & L. | 13.82 | .. | .. |
| 6 | Udō-no-yu | .. | 72.0° | — | 1.00 | 16.9° | .. | 14.48 | .. 7. .. | .. |
| 7 | Namuyano-yu <i>Sain-yu</i> | .. | 68.0° | — | 1.15 | 18.0° | .. | 14.36 | .. | .. |
| 8 | Botan-yu | .. | 70.0° | 180 | — | — | .. | — | — | — |

ŌWANI

Location.—Ōwani-mura, Minami-tsugaru-gun, Province Mutsu, Aomori-ken.
 The springs issue from the Liparite Dyke and Alluvium.

The hot spring “Umeka-no-yu”

Analysis.

Analyst: Unknown. Temperature: 62° C. Flow of water: ca. 126 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|---------|---|---------|
| Potassium chloride (KCl) | 0.02280 | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.07990 |
| Sodium chloride (NaCl) | 2.12000 | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.35000 |
| Ammonium chloride (NH ₄ Cl) | 0.03000 | Aluminium phosphate (AlPO ₄) | 0.00266 |
| Sodium sulphate (Na ₂ SO ₄) | 0.06320 | Ferric oxide (Fe ₂ O ₃) | 0.00130 |
| Sodium bicarbonate (NaHCO ₃) | 0.01500 | Silica (SiO ₂) | 0.00380 |
| Calcium sulphate (CaSO ₄) | 0.43850 | | 3.12710 |

The spring may be classified as “common salt spring”.

Radio-activity. 10.28 Mache's units in 1 litre of water at 20.1° C.
 (Kohl-Löw. fontact.; Y. Kinugasa, Oct. 20, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 45.42$.

The hot spring "Yamabuki-no-yu"

Analysis.

Analyst: Unknown. Temperature: 67° C. Flow of water: ca. 108 hectolitres in 24 hrs.

The spring, being in its composition nearly the same as the one at Yamabuki, may be classified as "common salt spring".

Radio-activity. 7.50 Mache's units in 1 litre of water at 18.4° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 20, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 44.38$.

Table 7.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emission per
litre of water
in
Mache's units
at
(temp. in C.) | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by | |
|-----|--|----------------|--------------------------|---|--|--------------------|--|----------------------------------|---------------|---------------------------|
| 1 | Umeke-no-yu
(Public bath) | common salt | 62.0° | ca.
126 | 10.28 | 20.1° | K & L | 45.42 | Oct. 20, 1913 | Y. Kinugasa
& T. Saitō |
| 2 | Yamabuki-no-yu
(Public bath) | " | 67.0 | 108 | 7.50 | 18.4 | " | 44.38 | " | " |
| 3 | Wakamatsu-no-yu
(Votum-no-yu Public bath) | salt | 62.5° | 144 | 8.77 | 20.7° | " | 40.06 | " 21, " | " |
| 4 | " (Hie-no-yu) | " | 63.0° | 144 | 9.69 | 17.3° | " | 43.00 | " 20, " | " |
| 5 | Kawara-no-yu
(Public bath) | " | 76.0° | — | 0.74 [#] | 17.1° | " | 34.73 | " | " |
| 6 | Kagasuke-no-yu | " | 69.0° | 126 | 9.27 | 20.0° | " | 40.96 | " 21, " | " |
| 7 | Aoyagi-no-yu
(Public bath) | " | 77.0° | 171 | — | — | — | — | — | — |

* As the source of the spring did not admit to take sample of water for examination without loss of emanation, the real quantity must be much larger than observed.

KURADATE

Location.—Kuradate-mura, Minami-tsugaru-gun, Province Mutsu, Aomori-ken.

The springs issue from the Alluvium.

The hot spring "Shimo-no-yu"

Analysis.

Analysed by E. Ōkawa, apothecary of Hirosaki. 1909.

Temperature: 72.5° C.

Flow of water: ca. 450 hectolitres in 24 hours.

Total residue: ca. 2.38.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|--|--------|
| Potassium chloride (KCl) | 0.1354 | Ferric oxide (Fe ₂ O ₃) | 0.0030 |
| Sodium chloride (NaCl) | 1.2650 | Boric acid (H ₃ BO ₃) | trace |
| Calcium sulphate (CaSO ₄) | 0.0304 | Silica (SiO ₂) | 1.147 |
| Magnesium sulphate (MgSO ₄) | 0.0816 | | 2.2916 |
| Aluminium phosphate (AlPO ₄) | 0.0350 | | |

The spring may be classified as "common salt spring".

Radio-activity. 2.86 Mache's units in 1 litre of water at 18° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 21, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 29.60$.

The hot spring “ Ō-yu ”

Analysis.

Analyst: Unknown.

Temperature: 72° C.

Total residue: ca. 2,32.

Flow of water: ca. 540 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of “ Shimo-no-yu ”, may be classified as “ common salt spring ”.

Radio-activity. 2.60 Mache's units in 1 litre of water at 19.4° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 21 1913).

Electric conductivity. $\times 10^{-4} = 28.22$.

The hot spring “ Hoyōen-no-yu ”

Analysis.

Analysed by E. Ōkawa, apothecary of Hirosaki. 1912.

Temperature: 78° C.

Specific gravity: 1.0023 at 15° C.

Total residue: ca. 2,39.

The spring, being in its composition nearly the same as that of “ Shimo-no-yu ”, may be classified as “ common salt spring ”.

Radio-activity. 1.87 Mache's units in 1 litre of water at 12.2° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 21, 1913).

Electric conductivity. $\times 10^{-4} = 29.63$.

Table 8.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|------------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|--------------------|--|----------------------------------|---------------------------|
| | | | | | in
Mache's units | at
(temp. in C.) | | | | |
| 1 | Shimo-no-yu (<i>Public bath</i>) | common salt | 72.5° | ca. 450 | 2.86 | 18.0° | K & L. | 29.60 | Oct. 21, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Ō-yu (<i>Public bath</i>) | .. | 72.0° | 540 | 2.60 | 19.4° | .. | 28.22 | .. | |
| 3 | Kami-no-yu (<i>Public bath</i>) | .. | 69.0° | — | 4.38 | 17.5° | .. | 27.72 | .. | |
| 4 | Hoyōen-no-yu | .. | 78.0° | — | 1.87 | 17.2° | .. | 29.63 | .. | |
| 5 | Kawabata-no-yu | .. | 56.0° | — | — | — | — | 28.48 | .. | |
| 6 | Netsu-no-yu | .. | 71.0° | — | — | — | — | 28.77 | .. | |

IKARIGASEKI

Location.—Ikariga-seki-mura, Minami-tsugaru-gun, Province Mutsu, Aomori-ken.

The springs issue from the Volcanic Ash.

The hot spring “ Netsu-no-yu ”

Analysis.

Analyst: Unknown.

Temperature: 61° C.

Total residue: ca. 1,25.

Flow of water: ca. 558 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|---------|---|---------|
| Potassium chloride (KCl) | 0,06952 | Ferrous bicarbonate $[\text{Fe}(\text{HCO}_3)_2]$ | 0,00225 |
| Sodium chloride (NaCl) | 1,01255 | Aluminium phosphate (AlPO_4) | 0,00250 |
| Potassium sulphate (K_2SO_4) | 0,00211 | Silica (SiO_2) | 0,12690 |
| Calcium sulphate (CaSO_4) | 0,07718 | | 1,32015 |
| Magnesium sulphate (MgSO_4) | 0,00381 | Carbon dioxide (free and in form of bicarbo- | |
| Ferrous sulphate (FeSO_4) | 0,00225 | nates) (CO_2) | 0,06900 |
| Calcium bicarbonate $[\text{Ca}(\text{HCO}_3)_2]$ | 0,02108 | | 1,38915 |

Besides these trace of bromine, iodine and boric acid.

The spring may be classified as "common salt spring".

Radio-activity. 1.20 Mache's units in 1 litre of water at 16,6° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 22, 1913).

Electric conductivity. $21,5 \times 10^{-4} = 18,37$.

The hot spring "Fuji-no-yu (Kuzuya-no-yu)"

Analysis.

Analysed by the Hirosaki Garrison Hospital. 1902.

Temperature: 55° C.

Total residue: ca. 1,30.

The spring, being in its composition nearly the same as that of "Netsu-no-yu", may be classified as "common salt spring".

Radio-activity. 0,92 Mache's units in 1 litre of water at 18,2° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 22, 1913).

Electric conductivity. $21,8 \times 10^{-4} = 16,73$.

The hot spring "Taki-no-yu"

Analysis.

Analysed by the Hirosaki Hospital. 1907.

Temperature: 61° C.

Total residue: ca. 1,62.

Flow of water: ca. 630 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Netsu-no-yu", may be classified as "common salt spring".

Radio-activity. 1,27 Mache's units in 1 litre of water at 18,5° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 22, 1913).

Electric conductivity. $21,8 \times 10^{-4} = 18,75$.

Table 9.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$21,5 \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|--|----------------|--------------------------|---|---------------------------------|-------------------|--------------------|---|----------------------------------|---------------------------|
| | | | | | in
Mache's units | at
temp. in C. | | | | |
| 1 | Netsu-no-yu (<i>Public bath</i>) | common salt | 61,0 | ca.
558 | 1,20 | 16,6° | K. & L. | 18,37 | Oct. 22, 1913 | Y. Kinugasa
(L. Saitō) |
| 2 | Hie-no-yu (<i>Onna-yu</i>)
<i>Public bath</i> | " | 54,0 | 300 | 0,63 | 17,3° | " | 10,06 | " | " |
| 3 | Fuji-no-yu (<i>Kuzuya-no-yu</i>) | " | 55,0 | " | 0,92 | 18,2 | " | 16,73 | " | " |
| 4 | " (<i>Bunkwan-no-yu</i>) | " | 62,0° | " | 1,39 | 14,4 | " | 18,15 | " | " |
| 5 | Taki-no-yu (<i>Shibata-no-yu</i>) | " | 61,0 | 630 | 1,27 | 18,5 | " | 18,75 | " | " |

DAI

Location.—Yumoto-mura, Hienuki-gun, Province Rikuchū, Iwate-ken.
The springs issue from the Diluvium.

Table 10.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|---|----------------|--------------------------|---|---------------------------------|---------------------|--------------------|--|----------------------------------|------------------------------|
| | | | | | "
Mache's units | at
(temp. in C.) | | | | |
| 1 | Tsuru-no-yu (<i>Public bath</i>) | simple | 67.5° | ca. 108 | 1.15 | 17.2° | K. & L. | 11.08 | Oct. 4, 1913 | Y. Kinugasa
{
T. Saitō |
| 2 | Suzume-no-yu (<i>Public bath</i>) | " | 54.0° | 72 | 0.53 | 15.9° | " | 10.37 | " | " |
| 3 | O-yu (<i>Public bath</i>) | acid | 58.0° | 126 | 0.43 | 17.4° | " | 10.33 | " | " |
| 4 | Yakushi-no-yu
(<i>Public bath</i>) | " | 53.0° | 108 | 0.18 | 16.5° | " | 9.65 | " | " |
| 5 | Matsu-no-yu (<i>Public bath</i>) | " | 78.0° | 102 | 0.95 | 15.9° | " | 13.40 | " | " |
| 6 | Kin-no-yu | salt | 68.0° | 135 | — | — | — | — | — | — |
| 7 | Gorō-no-yu | " | 62.0° | 144 | — | — | — | — | — | — |
| 8 | Taka-no-yu | " | 74.0° | 135 | — | — | — | — | — | — |
| 9 | Taki-no-yu | simple | 84.0° | 270 | — | — | — | — | — | — |
| 10 | Senshō-no-yu | salt | 62.0° | 108 | — | — | — | — | — | — |
| 11 | Shin-suzume-no-yu | " | 54.0° | 108 | — | — | — | — | — | — |
| 12 | Hotokezawa-no-yu | " | 62.0° | 90 | — | — | — | — | — | — |
| 13 | Me-suzume-no-yu | " | 58.0° | 90 | — | — | — | — | — | — |

SHIDODAIRA

Location.—Yuguchi-mura, Hienuki-gun, Province Rikuchū, Iwate-ken.
The spring issues from the Tertiary.

The hot spring "Warabi-no-yu"

Analysis.

Temperature: 76° C.

Total residue: 1.26.

Flow of water: ca. 389 hectolitres in 24 hours.

From the result of an old analysis made in 1881, the spring seems to be "bitter spring".

Radio-activity. 0.64 Mache's units in 1 litre of water at 18.8° C.

(Kohl.-Löw. contact.; Y. Kinugasa and T. Saitō, Oct. 3, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 14.54$.

ŌSAWA

Location.—Yuguchi-mura, Hienuki-gun, Province Rikuchū, Iwate-ken.
The springs issue from the Tertiary.

Table 11.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Total residue in
1 litre of water | | Apparent
viscosity
in poise | Electric
conductivity
in ohm ⁻¹ cm. | Date when the
water was taken | Examined by |
|-----|-----------------|----------------|--------------------------|---|--------------------------------------|-------------|-----------------------------------|--|----------------------------------|---------------------------|
| | | | | | Maché's units | Temp. in C. | | | | |
| 1 | Ō-yu Vent No. 1 | simple | 51° | ca. 544 | 0.72 | 18.2 | Kohl. | 7.34 | Oct. 3, 1913 | Y. Kinugasa
& T. Saitō |
| 2 | " Vent No. 2 | " | 51° | " | 0.43 | 18.2 | " | 7.34 | " | " |
| 3 | Taki-no-yu | " | 51° | 312 | " | " | " | " | " | " |

ŌYUZAWA

Location.—Yatate-mura, Kita-akita-gun, Province Ugo, Akita-ken.

The springs issue from the Diluvium.

The hot spring "No. 38 (Ō-yu)"

Analysis.

Analysed by the Hygienic Laboratory of Akita-ken. 1893.

Temperature: 45.5° C.

Flow of water: ca. 270 hectolitres in 24 hours.

Total residue: ca. 14.78.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|---------|---|---------|
| Sodium chloride (NaCl) | 11,0306 | Silica (SiO ₂) | 0.1060 |
| Magnesium sulphate (MgSO ₄) | 0.8880 | | 14.7748 |
| Ferrous sulphate (FeSO ₄) | 0.5377 | Free hydrogen sulphide (H ₂ S) | 0.2246 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | | | 14.0994 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 2.2125 | | |

The spring may be classified as "hydrogen sulphide common salt spring".

Radio-activity. 0.86 Maché's units in 1 litre of water at 10.9° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 24, 1913).

5.35 ‡ Maché's units in 1 litre of gas at 0° C.

(Eng.-Siev. fontact.; Y. Kinugasa and T. Saitō, Oct. 24, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 91.62$.

The cold spring "No. 39"

Analysis.

Analysed by the Hygienic Laboratory of Akita-ken. 1912.

Specific gravity: 1.0073 at 15° C.

Temperature: 33.5° C.

Total residue: ca. 11.31.

Flow of water: ca. 180 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "No. 38 (Ō-yu)", may be classified as

"hydrogen sulphide common salt spring".

Radio-activity. 0.98 Maché's units in 1 litre of water at 12.2° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 24 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 100.42$.

‡ Correction for the absorption of the radiation by the wall of the ionisation chamber and also for the absorption of the radiation by the ionisation cylinder in the ionisation chamber was made.

Table 12.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
(temp. in C.) | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|---|---------------------------------------|--------------------------|---|---|--------------------|--|--|---------------------------|
| 1 | No. 38 <i>Ô-yu</i> | hydrogen sul-
phide common
salt | 45.5° | ca. 270 | 0.86 10.9° | K. & L. | 91.62 | Oct. 24, 1913 | { Y. Kinugasa
T. Saito |
| 2 | No. 39 | " | 33.5° | 180 | 0.98 12.2° | " | 100.42 | " | " |
| 3 | Shimonaizawa-no-yu | simple | 34.0° | — | 1.24 10.5° | " | 8.58 | " 23, " | " |
| 4 | Aka-yu | iron carbonate | 32.0° | 650 | 0.90 9.9° | " | 145.03 | " | " |
| | Gas evolving from
No. 38 <i>Ô-yu</i> | | | | Emanation per
litre of gas at 0° C.
4.55 *
5.25 †
5.35 ‡ | E. & S. | | Date when the
gas was
collected
Oct. 24, 1913 | " |

* Determined according to the authors' original direction.
† Correction for the absorption of the radiation by the wall of the ionisation chamber was made.
‡ Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made

ÔYU

Location.—Ôyu-mura, Kazuno-gun, Province Rikuchû, Akita-ken.

The springs issue from the Pumice Layer in Quaternary System.

The hot spring "Shimo-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Akita-ken. 1913.

Specific gravity: 1.00139 at 15° C. Temperature: 69.5° C.

Total residue: ca. 2.04. Flow of water: ca. 414 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|---------------------|---|----------------|
| Potassium (K) | 0.01630 | Chlorine (Cl) | 0.92500 |
| Sodium (Na) | 0.66673 | Sulphuric anhydride (SO ₃) | 0.13551 |
| Calcium (Ca) | 0.09087 | Phosphoric anhydride (P ₂ O ₅) | trace |
| Magnesium (Mg) | very small quantity | Boric acid (H ₃ BO ₃) | small quantity |
| Ferric oxide (Fe ₂ O ₃) | " | Silica (SiO ₂) | 0.06975 |
| Alumina (Al ₂ O ₃) | 0.12475 | Carbon dioxide (CO ₂) | small quantity |

The spring may be classified as "common salt spring".

Radio-activity. 0.89 Mache's units in 1 litre of water at 15.4° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitô, Oct. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 28.65$.

The hot spring "Kawara-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Akita-ken. 1913.

Specific gravity: 1.00141 at 15° C. Temperature: 70.5° C.

Total residue: ca. 2.10. Flow of water: ca. 634 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Shimo-no-yu", may be classified as
"common salt spring".

Radio-activity. 0.66 Mache's units in 1 litre of water at 16.5° C.
 (Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 28.19$.

The hot spring "Kami-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Akita-ken. 1913.

Specific gravity: 1.001 at 15° C. Temperature: 58° C.

Total residue: ca. 1.62. Flow of water: ca. 630 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Shimo-no-yu", may be classified as
"common salt spring".

Radio-activity. 0.98 Mache's units in 1 litre of water at 16.8° C.
 (Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 19.85$.

The hot spring "Arase-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Akita-ken. 1913.

Specific gravity: 1.00888 at 15° C. Temperature: 46° C.

Total residue: ca. 2.03. Flow of water: ca. 619 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Shimo-no-yu", may be classified as
"common salt spring".

Radio-activity. 0.81 Mache's units in 1 litre of water at 15.9° C.
 (Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 16.61$.

Table 13.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
(temp. in C.) | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|-------------------------------------|----------------|--------------------------|---|---|--------------------|--|----------------------------------|-------------------------|
| 1 | Shimo-no-yu (<i>Public bath</i>) | common salt | 69.5° | ca.
414 | 0.89
15.4° | K. & L. | 28.65 | Oct. 25, 1913 | Y. Kinugasa
T. Saitō |
| 2 | Kawara-no-yu (<i>Public bath</i>) | " | 70.5° | 634 | 0.66
16.5° | " | 28.19 | " | " |
| 3 | Kami-no-yu | " | 58.0° | 630 | 0.98
16.8° | " | 19.85 | " | " |
| 4 | Arase-no-yu | " | 46.0° | 619 | 0.81
15.9° | " | 16.61 | " | " |

ŌDAKI

Location.—Jūnisho-machi, Kita-akita-gun, Province Ugo, Akita-ken.

The springs issue from the Soil in Quaternary System.

The hot spring "Tsuru-no-yu"

Analysis.

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881. Temperature: 61° C.

Flow of water: ca. 6012 hectolitres in 24 hours. Total residue: ca. 2.28.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|---|---------|--|---------|
| Potassium chloride (KCl) | 0,17572 | Silica (SiO ₂) | 0,04214 |
| Sodium chloride (NaCl) | 0,84384 | Boric acid (H ₃ BO ₃) | trace |
| Sodium sulphate (Na ₂ SO ₄) | 0,36612 | | 2,23956 |
| Sodium carbonate (Na ₂ CO ₃) | 0,03832 | Hydrogen sulphide (H ₂ S) | 0,00216 |
| Calcium sulphate (CaSO ₄) | 0,76541 | | 2,24172 |
| Magnesium sulphate (MgSO ₄) | 0,00801 | | |

The spring may be classified as "sulphur spring".

Radio-activity. 0.93 Mache's units in 1 litre of water at 12,0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 24, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 23,37$.

Table 14.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
(temp. in C.) | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|-------------------------------------|----------------|--------------------------|---|---|--------------------|--|----------------------------------|---------------------------|
| 1 | Tsuru-no-yu (<i>Public bath</i>) | sulphur | 61,0° | ca.
6012 | 0,93 12,0° | K. & L. | 23,37 | Oct. 24, 1913 | (Y. Kinugasa
T. Saitō) |
| 2 | Susuki-no-yu (<i>Public bath</i>) | bitter | 62,0° | — | 0,71 12,3° | .. | 21,62 | .. | .. |

SHIBUKURO

Location.—Tazawa-mura, Senhoku-gun, Province Ugo, Akita-ken.

The springs issue from the Augite Andesite.

The hot spring "Shibukurozawa-no-yu" *

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1912. Specific gravity: 1,0030 at 19° C.

In 1 kilogram of the mineral water are contained :

Cations.

| | Grams | Milli-
mols | Milligram-
equiv. valents |
|--|---------|----------------|------------------------------|
| Hydrogen ion (H ⁺) | 0,07500 | 74,25743 | 74,25743 |
| Potassium ion (K ⁺) | 0,02491 | 0,63627 | 0,63627 |
| Sodium ion (Na ⁺) | 0,04415 | 1,91540 | 1,91540 |
| Ammonium ion (NH ₄ ⁺) | 0,00011 | 0,00608 | 0,00608 |
| Calcium ion (Ca ⁺⁺) | 0,04870 | 1,21750 | 2,43500 |
| Magnesium ion (Mg ⁺⁺) | 0,01483 | 0,60878 | 1,21756 |
| Ferrous ion (Fe ⁺⁺) | 0,24846 | 4,43679 | 8,87358 |
| Aluminium ion (Al ⁺⁺⁺) | 0,09622 | 3,55055 | 10,65165 |
| | | | 99,99297 |

Anions.

| | | | |
|--|---------|----------|--------------------------------|
| Chlorine ion (Cl ⁻) | 2,69729 | 76,14358 | 76,14358 |
| Hydrosulphate ion (HSO ₄ ⁻) | 0,42034 | 4,33028 | 4,33028 |
| Sulphate ion (SO ₄ ⁼⁼) | 0,03776 | 0,76223 | 19,52446 |
| | | | 4,00777 177,86489 99,99832 |
| Boric acid (meta) (HBO ₂) | 0,04244 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,37022 | | |
| | | | 5,02043 |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0,0475 |
| Sodium chloride (NaCl) | 0,1121 |
| Ammonium chloride (NH ₄ Cl) | 0,0003 |
| Calcium chloride (CaCl ₂) | 0,1350 |
| Magnesium chloride (MgCl ₂) | 0,0580 |
| Ferrous sulphate (FeSO ₄) | 0,6746 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0,6078 |
| Boric acid (meta) (HBO ₂) | 0,0424 |
| Free hydrochloric acid (HCl) | 2,5497 |
| Free sulphuric acid (H ₂ SO ₄) | 0,4247 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,3702 |
| | 5,0223 |

Besides these a small quantity of organic substances.

The spring thus may be classified as "acid alum vitriol spring".

* The waters of "Suka-yu", an acid vitriol spring near by, contain trace of barium and lead ion.

INNAI-YUNOSAWA

Location.—Innai-machi, Okachi-gun, Province Ugo, Akita-ken.

The springs issue from the Tertiary.

The hot spring “Yunosawa-Onsen”

Analysis.

Analysed by Dr. R. Ishizu and Y. Kinugasa, 1913.

Specific gravity: 1.0004 at 15° C.

Temperature: 40.5° C.

Total residue: ca. 0.13.

Flow of water: ca. 24624 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | Hypothetical form of compound | Grams |
|---|----------|---|----------|
| Potassium oxide (K ₂ O) | 0.011203 | Potassium chloride (KCl) | 0.008950 |
| Sodium oxide (Na ₂ O) | 0.011700 | Sodium chloride (NaCl) | 0.011050 |
| Calcium oxide (CaO) | 0.025206 | Calcium sulphate (CaSO ₄) | 0.024055 |
| Magnesia (MgO) | 0.004424 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.002674 |
| Ferric oxide (Fe ₂ O ₃) | 0.002403 | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.044264 |
| Alumina (Al ₂ O ₃) | trace | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.016147 |
| Chlorine (Cl) | 0.010851 | Silica (SiO ₂) | 0.040103 |
| Nitric anhydride (N ₂ O ₅) | trace | | 0.147243 |
| Sulphuric anhydride (SO ₃) | 0.014150 | | |
| Silica (SiO ₂) | 0.040103 | | |
| Carbon dioxide (CO ₂) | 0.017532 | | |

The spring may be classified as “simple thermal”.

Radio-activity. 1.06 Mache's units in 1 litre of water at 13.6° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 27, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 1.24$.

Table 15.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water in Mache's units | Temp. in C. | Apparatus employed | Electric conductivity $\kappa_{18} \times 10^{-4}$ | Date when the water was taken | Examined by |
|-----|----------------|----------------|-----------------------|---|---|-------------|--------------------|--|-------------------------------|---------------------------|
| 1 | Yunosawa-Onsen | simple | 40.5° | ca. 24624 | 1.06 | 13.6° | K. & L. | 1.24 | Oct. 27, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Taki-no-yu | „ | 41.0° | — | .21 | 13.7° | „ | 1.25 | „ | „ |

ONIKŌBE

Location.—Onikōbe-mura, Tamatsukuri-gun, Province Rikuzen, Miyagi-ken.

The springs issue from the Tertiary.

The hot spring “Todoroki-no-yu”

Analysis.

Temperature: 53° C.

Total residue: ca. 0.35.

Flow of water: ca. 270 hectolitres in 24 hours.

From the result of an old analysis made in 1881, the spring seems to be “simple thermal”.

Radio-activity. 0.49 Mache's units in 1 litre of water at 20.3° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 1, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 4.99$.

TAMATSUKURI

Location.—Yuzumi-mura, Tamatsukuri-gun, Province Rikuzen, Miyagi-ken.
The springs of Naruko, Shin-kuruma, Moto-kuruma and Akayu issue from the Pyroxene Andesite, but that of Kawatabi from the Tertiary.

Table 16.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\frac{1}{4} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|--------------------------|------------------|--------------------------|---|---------------------------------|-------------------|--------------------|--|----------------------------------|---------------------------|
| | | | | | in
Maché's units | at
temp. in C. | | | | |
| 1 | Taki-no-yu No. 1 | sulphur | 40.5° | ca. — | 1.01 | 20.0° | K. & L. | 15.30 | Sept. 29, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | " No. 2 | " | 92.0° | — | 0.24 | 19.3° | " | 44.80 | Oct. 1, " | " |
| 3 | Unagi-no-yu | alkaline sulphur | 100.0° | — | 0.13 | 18.4° | " | 14.60 | Sept. 29, " | " |
| 4 | Shin-unagi-no-yu | alkaline | 93.0° | — | 0.17 | 20.2° | " | 21.32 | " 30, " | " |
| 5 | Genzō-no-yu | sulphur | 51.0° | 270 | 0.32 | 18.0° | " | 42.24 | " 20, " | " |
| 6 | Kawara-no-yu | salt | 48.0° | 54 | 0.30 | 20.3° | " | 15.29 | Oct. 1, " | " |
| 7 | Tamamono-no-yu | carbonated | 70.0° | 90 | — | — | — | — | — | — |
| 8 | Uba-no-yu | " | 70.0° | 81 | — | — | — | — | — | — |
| 9 | Moto-yu | salt | 67.5° | — | 0.56 | 19.2° | K. & L. | 11.31 | Sept. 30, 1913 | { Y. Kinugasa
T. Saitō |
| 10 | Mushi-yu | " | 70.5° | — | — | — | — | 0.09 | " | " |
| 11 | Takashige-no-yu | " | 54.0° | — | 0.54 | 20.2° | K. & L. | 13.77 | " | " |
| 12 | Kame-no-yu | " | 48.0° | 90 | — | — | — | — | — | — |
| 13 | Ubuchi-no-yu | " | 86.5° | — | 0.34 | 19.0° | K. & L. | 15.06 | Sept. 30, 1913 | { Y. Kinugasa
T. Saitō |
| 14 | Yamane-no-yu | " | 73.0° | — | — | — | — | 22.51 | " | " |
| 15 | Senshin-no-yu | " | 78.0° | 90 | — | — | — | — | — | — |
| 16 | Reisen-no-yu | " | 78.0° | 90 | — | — | — | — | — | — |
| 17 | Goten-no-yu | carbonated | 58.0° | — | 0.28 | 18.8° | K. & L. | 8.17 | Oct. 1, 1913 | { Y. Kinugasa
T. Saitō |
| 18 | Taki-no-yu (Public bath) | " | 58.0° | — | — | — | — | 6.47 | " | " |
| 19 | Ō-yu. | " | 50.0° | 648 | — | — | — | 8.13 | " | " |
| 20 | Suzuki-no-yu | " | 42.5° | — | 0.53 | 20.6° | K. & L. | 7.16 | Sept. 30, 1913 | " |
| 21 | Fuji-no-yu | — | 48.0° | — | — | — | — | 8.14 | " | " |
| 22 | Me-no-yu | — | 43.0° | — | — | — | — | 7.74 | " | " |
| 23 | Naka-no-yu (Public bath) | carbonated | 45.0° | 864 | — | — | — | 7.44 | Oct. 3, 1913 | " |
| 24 | Mayu-no-yu | " | 46.0° | 648 | — | — | — | 7.50 | " 1, " | " |

KAMASAKI

Location.—Fukuoka-mura, Katta-gun, Province Iwaki, Miyagi-ken.
The springs issue from the Pyroxene Andesite and its Agglomerate.

The hot spring "Ichijō-no-yu"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1909.

Specific gravity: 1.0046 at 14° C. Temperature: 48° C.

Total residue: ca. 5.03.

Flow of water: ca. 3150 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained

| Cations. | Grams | Mille-
moles | Mil-
equ. |
|--|---------|-----------------|--------------|
| Potassium ion (K ⁺) | 0,05218 | 1,33209 | 1,33209 |
| Sodium ion (Na ⁺) | 1,43355 | 62,19306 | 62,19306 |
| Ammonium ion (NH ₄ ⁺) | 0,00021 | 0,01162 | 0,01162 |
| Calcium ion (Ca ⁺⁺) | 0,22988 | 5,74700 | 11,49400 |
| Magnesium ion (Mg ⁺⁺) | 0,01193 | 0,48974 | 0,97948 |
| Aluminium ion (Al ⁺⁺⁺) | 0,00106 | 0,03911 | 0,11733 |
| | | 76,12755 | |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 1,50624 | 42,48114 | 42,48914 |
| Sulphate ion (SO ₄ ⁼⁼) | 1,35357 | 14,08880 | 28,17760 |
| Hydrophosphate ion (HPO ₄ ⁼⁼) | 0,00103 | 0,01073 | 0,02146 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0,33185 | 5,43935 | 5,43935 |
| | 4,02127 | 131,84061 | 76,12755 |
| Boric acid (meta) (HBO ₃) | 0,28250 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,00414 | | |
| | 5,20791 | | |
| Free carbon dioxide (CO ₂) | 0,06169 | 1,40199 | |
| | 5,26960 | | |

Besides these trace of ferrous ion.

The spring thus may be classified as “**saline common salt spring**”.

Radio-activity. 0,87 Mache's units in 1 litre of water at 16,4° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 26, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 51,43$.

The hot spring “Mogamiya-no-yu”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1912. Specific gravity: 1,0048 at 15° C.

Temperature: 38,5° C.

Total residue: ca. 5,19.

The spring, being in its composition nearly the same as that of “Ichijō-no-yu”, may be classified as “**saline common salt spring**”.

Radio-activity. 1,83 Mache's units in 1 litre of water at 18,4° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 26, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 44,20$.

The hot spring “Kimuraya-no-yu”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1912. Specific gravity: 1,0051 at 13° C.

Temperature: 43° C.

Total residue: ca. 5,15.

The spring, being in its composition nearly the same as that of “Ichijō-no-yu”, may be classified as “**saline common salt spring**”.

Radio-activity. 0,80 Mache's units in 1 litre of water at 19,9° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 27, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 48,79$.

Table 17.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
(temp. in C.) | Apparatus employed | Electric
conductivity
$\times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|----------------|-----------------------|--------------------------|---|---|--------------------|--|----------------------------------|---------------------------|
| 1 | Ichijō-no-yu | saline common
salt | 48,0° | ca.
3150 | 0,87 16,4° | K. & L. | 51,43 | Sept. 26, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Mogamiya-no-yu | " | 38,5° | — | 1,83 18,4° | " | 44,20 | " | " |
| 3 | " | " | 37,0° | " | — | — | 51,28 | " 27, " | " |
| 4 | Kimuraya-no-yu | " | 43,0° | " | 0,80 19,9° | K. & L. | 48,79 | " | " |

ATSUMI

Location.—Atsumi-mura, Nishi-tagawa-gun, Province Uzen, Yamagata-ken.

The springs issue from the Tertiary Conglomeratic Tuff.

The hot spring "Yu-atsumi-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1886.

Specific gravity: 1,003.

Total residue: ca. 3,51.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K') | 0,0477 | 1,2183 | 1,2183 |
| Sodium ion (Na') | 0,9251 | 40,1344 | 40,1344 |
| Calcium ion (Ca'') | 0,2612 | 6,5162 | 13,0324 |
| | | | 54,3851 |
| Anions. | | | |
| Chlorine ion (Cl') | 1,5052 | 42,4598 | 42,4598 |
| Sulphate ion (SO ₄ '') | 0,5726 | 5,0608 | 11,9216 |
| | 3,3118 | 96,2895 | 54,3814 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,1064 | | |
| | 3,4182 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|--|--------|
| Potassium chloride (KCl) | 0,0910 |
| Sodium chloride (NaCl) | 2,3480 |
| Calcium chloride (CaCl ₂) | 0,0610 |
| Calcium sulphate (CaSO ₄) | 0,8118 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,1064 |
| | 3,4182 |

Besides these trace of magnesium, hydrophosphate ion, hydrogen sulphide and boric acid.

The spring thus may be classified as "sulphated common salt spring".

Radio-activity. $4,04 \times 10^{-10}$ curies in 1 litre of water * at 24,2° C.

(Schmidt's electrom.; H. Kakehi, June 2, 1914).

1,01 Mache's units (recalculated).

* Measurement has been made with the sample brought to the laboratory by applicant, taking into account decay of radium emanation with the time.

KAMINOYAMA

Location.—Kani-no-yama-machi, Minami-nagasaki-gun, Fukuoka-ken, Yamaguchi-ken.

The springs issue from the Liparite.

Table 18.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|-------------------------------------|----------------|--------------------------|---|--|-------------------|--------------------|--|---------------------------------------|---------------------------|
| | | | | | in
Mach's units | at
temp. in C. | | | | |
| 1 | Atatamari-yu (<i>Public bath</i>) | bitter | 61.0° | ca.
3944 | 1.70 | 13.8 | K. & L. | 30.00 | Oct. 28, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Hie-no-yu (<i>Public bath</i>) | .. | 62.0° | 3944 | 1.83 | 13.8 | .. | 27.40 | .. | .. |
| 3 | Kawashimaya-no-yu No. 2 | .. | 61.5° | 488 | 2.17 | 13.5 | .. | 28.72 | .. | .. |
| 4 | .. No. 1 | .. | 56.0° | — | — | — | .. | 31.73 | .. | .. |
| | Gas evolving from
Hie-no-yu | | | | Emanation per litre
of gas at 0° C. | | | | Date when the
gas was
collected | |
| | | | | | 13.45* | | E. & S. | | Oct. 28, 1913 | .. |
| | | | | | 15.51† | | | | | |
| | | | | | 15.80‡ | | | | | |

* Determined according to the authors' original direction.
† Correction for the absorption of the radiation by the wall of the ionisation chamber was made.
‡ Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made.

AKAYU

Location.—Akayu-machi, Higashi-okitama-gun, Province Uzen, Yamagata-ken.

The springs issue from the Tertiary.

The hot spring “Tamba-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tokyo Imperial Hygienic Laboratory, 1903.

Specific gravity: 1.0015 at 13.5° C.

Temperature: 30° C.

Total residue: ca. 2.95.

Flow of water: ca. 207 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|---|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0252 | 0.6437 | 0.6437 |
| Sodium ion (Na ⁺) | 0.7766 | 34.5597 | 34.5597 |
| Calcium ion (Ca ⁺⁺) | 0.2836 | 7.0723 | 14.1446 |
| Magnesium ion (Mg ⁺⁺) | 0.0043 | 0.0574 | 0.109 |
| | | | 49.4548 |

| | |
|--|--------|
| Potassium chloride (KCl) | 0.0480 |
| Sodium chloride (NaCl) | — |
| Sodium bicarbonate (NaHCO ₃) | 0.1785 |
| Calcium chloride (CaCl ₂) | 0.117 |
| Magnesium chloride (MgCl ₂) | 0.0051 |

Anions.

| | | | | | |
|--|--------|---------|---------|--|--------|
| Chlorine ion (Cl ⁻) | 1,5230 | 42,9873 | 42,9873 | Calcium sulphate (CaSO ₄) | 0,2958 |
| Sulphate ion (SO ₄ ²⁻) | 0,2085 | 2,1705 | 4,3410 | Silicic acid (meta) (H ₂ SiO ₃) | 0,0519 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0,1206 | 2,1242 | 2,1242 | | 3,0206 |
| | 2,9687 | 89,6111 | 49,4525 | Free carbon dioxide (CO ₂) | 0,0208 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0519 | | | | 3,0414 |
| | 3,0206 | | | | |
| Free carbon dioxide (CO ₂) | 0,0208 | 0,4727 | | | |
| | 3,0414 | | | | |

Besides these trace of ferrous, aluminium, iodine, hydrophosphate ion and boric acid.

The spring thus may be classified as “earth-muriated common salt spring”.

Radio-activity. 1.33 Mache's units in 1 litre of water at 16,6° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 29, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 36,29$.

The hot spring “Ō-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1907.

Specific gravity: 1,0025 at 15° C.

Temperature: 58° C.

Total residue: ca. 3,07.

Flow of water: ca. 908 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of “Tamba-no-yu”, may be classified as “earth-muriated common salt spring”.

Radio-activity. 1.45 Mache's units in 1 litre of water at 16,6° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 29, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 44,40$.

The hot spring “Ama-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1903.

Specific gravity: 1,0015 at 13,5° C.

Temperature: 49° C.

Total residue: ca. 2,91.

Flow of water: ca. 179 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of “Tamba-no-yu”, may be classified as “earth-muriated common salt spring”.

Radio-activity. 1.38 Mache's units in 1 litre of water at 16,1° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Oct. 29, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 40,83$.

Table 19.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|------------------------------------|-------------------------------|--------------------------|---|---------------------------------|-------|--------------------|--|----------------------------------|---------------------------|
| 1 | Tamba-no-yu (<i>public bath</i>) | earth-muriated
common salt | 50,0° | ca. 207 | 1,33 | 16,6° | K. & L. | 36,29 | Oct. 29, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Ō-yu (<i>public bath</i>) | .. | 58,0° | 908 | 1,45 | 16,6° | .. | 44,40 | .. | |
| 3 | Ama-yu (<i>public bath</i>) | .. | 49,0° | 179 | 1,38 | 16,1° | .. | 40,83 | .. | |
| 4 | Mori-no-yu | .. | 42,0° | 195 | 0,78 | 16,4° | .. | 40,15 | .. | |

ONOGAWA

Location.—Misa-wa-mura, Minami-okitama-gun, Province Uzen, Yamagata-ken.
The springs issue from the Tertiary.

The hot spring “Onogawa-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1906.

Specific gravity: 1.0045 at 15° C. Temperature: 66° C. Total residue: ca. 5.67.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|---|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.1310 | 3.3461 | 3.3461 |
| Sodium ion (Na ⁺) | 1.5369 | 66.6767 | 66.6767 |
| Calcium ion (Ca ⁺⁺) | 0.4644 | 11.5801 | 23.1602 |
| Magnesium ion (Mg ⁺⁺) | 0.0110 | 0.4516 | 0.9032 |
| | | | 94.0862 |

| Anions. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Chlorine ion (Cl ⁻) | 3.2252 | 90.9788 | 90.9788 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.0929 | 0.9671 | 1.9342 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0.0714 | 1.1703 | 1.1703 |
| | 5.5328 | 175.1707 | 94.0833 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1229 | | |
| | 5.6557 | | |
| Free carbon dioxide (CO ₂) | 0.0156 | 0.3545 | |
| | 5.6713 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.2496 |
| Sodium chloride (NaCl) | 3.9007 |
| Calcium chloride (CaCl ₂) | 1.1629 |
| Calcium sulphate (CaSO ₄) | 0.1318 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0217 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0661 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1229 |
| | 5.6557 |
| Free carbon dioxide (CO ₂) | 0.0156 |
| | 5.6713 |

Besides these trace of iodine, hydrophosphate ion and boric acid.

The spring thus may be classified as “earth-muriated common salt spring”.

Table 20.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water in Mache's units at (temp. in C.) | Apparatus employed | Electric conductivity $\times 10^{-4}$ | Date when the water was taken | Examined by |
|-----|-----------------------------------|----------------------------|-----------------------|---|--|--------------------|--|-------------------------------|----------------|
| 1 | Taki-no-yu (<i>Public bath</i>) | earth-muriated common salt | 70.0° | — | 5.78 14.3° | K. & L. | 65.76 | Oct. 30, 1913 | Y. K. Nagasaka |
| 2 | Ama-no-yu (<i>Public bath</i>) | .. | 73.5° | — | 1.95 11.0° | .. | 71.47 | .. | .. |
| 3 | Ōgiya-no-yu No. 1 | .. | 65.5° | .. | 0.92 12.6° | .. | 49.31 | .. | .. |
| 4 | .. No. 2 | .. | 66.5° | — | 2.23 10.3° | .. | 49.71 | .. 31. .. | .. |
| 5 | Suzuki-no-yu | .. | 68.0° | — | 3.45 12.2° | .. | 73.00 | .. 30. .. | .. |

SEKINE-YUNOSAWA

Location.—Yamakami-mura, Minami-okitama-gun, Province Uzen, Yamagata-ken.
The spring issues from the Gneiss.

The cold spring “Yunosawa-Onsen”

Temperature: 28° C.

No accurate analysis has ever been made, but the spring seems to be “simple cold spring”.

Radio-activity. 2.64 Mache's units in 1 litre of water at 6.6° C.

(Kohl-Low contact), Y. Kato, et al. J. S. C. No. 1, 1910.

Electric conductivity. $\times 10^{-4} = 5.01$.

NAMEGAWA

Location.—Yamakami-mura, Minami-okitama-gun, Province Uzen, Yamagata-ken.
The spring issues from the Tertiary.

The hot spring "Namegawa-Onsen"

Analysis.

Analysed by the Hygienic Laboratory of Yamagata-ken. 1909.
Specific gravity: 1,00069. Temperature: 55° C. (air temp. 22° C.).
Total residue: ca. 1,43. Flow of water: ca. 340 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|--|--------|
| Potassium chloride (KCl) | 0,1109 | Magnesium carbonate (MgCO ₃) | 0,0730 |
| Sodium chloride (NaCl) | 0,5364 | Aluminium phosphate (AlPO ₄) | 0,0416 |
| Calcium chloride (CaCl ₂) | 0,5067 | Silica (SiO ₂) | 0,0450 |
| Calcium sulphate (CaSO ₄) | 0,1457 | | 1,7357 |
| Calcium carbonate (CaCO ₃) | 0,2764 | | |

The spring may be classified as "earth-muriated common salt spring".

GOSHIKI

Location.—Yamakami-mura, Minami-okitama-gun, Province Uzen, Yamagata-ken.
The springs issue from the Tertiary.

The hot spring "Goshiki-Onsen" No. 1

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1904.
Specific gravity: 1,001 at 14° C. Temperature: ca. 44,5° C.
Total residue: ca. 0,87. Flow of water: ca. 778 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents | | Grams |
|--|--------|------------|-----------------------|---|--------|
| Potassium ion (K ⁺) | 0,0316 | 0,8072 | 0,8072 | Potassium chloride (KCl) | 0,0602 |
| Sodium ion (Na ⁺) | 0,2329 | 10,1041 | 10,1041 | Sodium chloride (NaCl) | 0,2241 |
| Calcium ion (Ca ⁺⁺) | 0,0580 | 1,4464 | 2,8928 | Sodium bicarbonate (NaHCO ₃) | 0,5274 |
| Magnesium ion (Mg ⁺⁺) | 0,0020 | 0,0838 | 0,1676 | Calcium sulphate (CaSO ₄) | 0,0475 |
| | | | 13,9717 | Magnesium sulphate (MgSO ₄) | 0,0101 |
| Anions. | | | | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,1779 |
| Chlorine ion (Cl ⁻) | 0,1644 | 4,6375 | 4,6375 | Silicic acid (meta) (H ₂ SiO ₃) | 0,0805 |
| Sulphate ion (SO ₄ ^{''}) | 0,0416 | 0,4331 | 0,8662 | | 1,1277 |
| Hydrocarbonate ion (HCO ₃ [']) | 0,5166 | 8,4680 | 8,4680 | Free carbon dioxide (CO ₂) | 0,3762 |
| | 1,0471 | 25,9801 | 13,9717 | | 1,5039 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0805 | | | | |
| | 1,1276 | | | | |
| Free carbon dioxide (CO ₂) | 0,3762 | 8,5500 | | | |
| | 1,5038 | | | | |

Besides these trace of ferrous, aluminium, hydrophosphate ion and boric acid.

The spring thus may be classified as "alkaline spring".

Radio-activity. 1,25 Mache's units in 1 litre of water at 6,1° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 1, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 10,22$.

Table 21.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
temp. in C. | | Apparatus employed | Electric
conductivity
$\gamma_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|---------------------|----------------|--------------------------|---|---|------|--------------------|--|----------------------------------|---------------------------|
| 1 | Goshiki-Onsen No. 1 | alkaline | 44.5° | ca. 778 | 1.25 | 0.1 | K & L | 10.22 | Nov. 1, 1913 | Y. Kinugasa
(T. Saitō) |
| 2 | " No. 2 | " | 38.5° | — | 1.95 | 0.5 | " | 8.04 | " | " |
| 3 | Shin-goshiki-Onsen | simple | 40.5° | 346 | 0.82 | 6.5° | " | 10.17 | " | " |

YUNO

Location.—Yuno-mura, Date-gun, Province Iwashiro, Fukushima-ken.

The springs issue from the Tertiary.

The hot spring "Hashimoto-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1897. Temperature: 56° C.

Flow of water: ca. 2160 hectolitres in 24 hours. Total residue: ca. 0.90.

In 1 kilogram of the mineral water are contained:

Cations.

| | Grams | Milli-
mols | Milligram-
equivalents |
|--------------------------------|--------|----------------|---------------------------|
| Potassium ion (K') | 0.0038 | 0.0971 | 0.0971 |
| Sodium ion (Na') | 0.2183 | 9.4707 | 9.4707 |
| Calcium ion (Ca'') | 0.0332 | 0.8279 | 1.6558 |
| Magnesium ion (Mg'') | 0.0012 | 0.0493 | 0.0986 |
| | | | 11.3222 |

Anions.

| | | | |
|--|--------|---------|---------|
| Chlorine ion (Cl') | 0.1208 | 3.4076 | 3.4076 |
| Sulphate ion (SO ₄ '') | 0.3801 | 3.9569 | 7.9138 |
| | 0.7574 | 17.8005 | 11.3214 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1055 | | |
| | 0.8629 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|--|--------|
| Sodium chloride (NaCl) | 0.1993 |
| Potassium sulphate (K ₂ SO ₄) | 0.0085 |
| Sodium sulphate (Na ₂ SO ₄) | 0.4315 |
| Calcium sulphate (CaSO ₄) | 0.1124 |
| Magnesium sulphate (MgSO ₄) | 0.0057 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1055 |
| | 0.8629 |

Besides these a small quantity of ferrous and aluminium ion and trace of hydrophosphate ion and boric acid.

The spring thus may be classified as "simple thermal".

Radio-activity. 0.38 Mache's units in 1 litre of water at 19° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 10.51$.

The hot spring "Kiri-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1897. Temperature: 57° C.

Flow of water: ca. 432 hectolitres in 24 hours. Total residue: ca. 0.79.

The spring, being in its composition nearly the same as that of "Hashimoto-no-yu", may be classified as "simple thermal".

Radio-activity. 0.39 Mache's units in 1 litre of water at 18.1° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 10.84$.

The hot spring "Kitsune-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1898. Temperature: 49° C.
Flow of water: ca. 540 hectolitres in 24 hours. Total residue: ca. 0.64.

The spring, being in its composition nearly the same as that of "Hashimoto-no-yu", may be classified as "simple thermal".

Radio-activity. 0.36 Mache's units in 1 litre of water at 16.8° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 10.93$.

Table 22.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|----------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|--------------------|--|----------------------------------|---------------------------|
| | | | | | in
Mache's units | at
(temp. in C.) | | | | |
| 1 | Hashimoto-no-yu
(Public bath) | simple | 56.0° | ca. 2160 | 0.38 | 19.0° | K. & L. | 10.51 | Sept. 25, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Kiri-no-yu (Public bath) | " | 57.0° | 432 | 0.39 | 18.1° | " | 10.84 | " | " |
| 3 | Kitsune-yu (Public bath) | " | 49.0° | 540 | 0.36 | 16.8° | " | 10.93 | " | " |
| 4 | Anabara-no-yu (Public bath) | " | 58.5° | 432 | 0.92 | 17.4° | " | — | " 24, " | " |
| 5 | Inariya-no-yu | " | 55.0° | 432 | — | — | — | 10.37 | " 25, " | " |
| 6 | Izumiya-furu-yu | " | 68.5° | 108 | — | — | — | 11.24 | " | " |
| 7 | Izumiya-shin-yu | " | 56.5° | — | — | — | — | 10.98 | " | " |
| 8 | Wahaya-no-yu | " | 55.0° | 216 | — | — | — | 10.45 | " | " |
| 9 | Matsubaya-no-yu No. 1 | " | 48.0° | 432 | — | — | — | 9.23 | " | " |
| 10 | " No. 2 | " | 51.0° | — | — | — | — | 10.63 | " 24, " | " |

IIZAKA

Location.—Iizaka-machi, Shinobu-gun, Province Iwashiro, Fukushima-ken.

The springs issue from the Tertiary.

The hot spring "Hako-yu"

Analysis.

Analyst: Unknown. 1891.

Specific gravity: 1.006. Temperature: 56.5° C.

Total residue: ca. 0.95. Flow of water: ca. 108 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|---------|---|---------|
| Sodium chloride (NaCl) | 0.20695 | Sodium silicate (Na ₂ SiO ₄) | 0.12860 |
| Sodium sulphate (Na ₂ SO ₄) | 0.11150 | Calcium sulphate (CaSO ₄) | 0.46020 |
| Potassium silicate (K ₄ SiO ₄) | 0.04320 | | 0.95945 |

The spring may be classified as "simple thermal".

Radio-activity. 0.50 Mache's units in 1 litre of water 18.7° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 25, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 10.60$.

Table 23.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Radio-activity
in 100 gms. of water | | Apparent specific gravity | Electric conductivity
at 16°C. | Date when the
water was taken | Examined by |
|-----|---|----------------|--------------------------|---|--|-------------|---------------------------|-----------------------------------|----------------------------------|---------------------------|
| | | | | | Mache's units | temp. 16°C. | | | | |
| 1 | Hako-yu (<i>Public bath</i>) | Simple | 56.5° | 108 | 0.50 | 18.7 | 1.001 | 10.60 | Sept. 25, 1913 | Y. Kinugasa
(T. Saitō) |
| 2 | Taki-no-yu .. | " | 59.0° | 180 | 0.68 | 15.8 | " | 12.28 | " 24, " | " |
| 3 | Shinsabako-no-yu .. | " | 70.0° | — | 0.52 | 18.0 | " | 12.32 | " 25, " | " |
| 4 | Spring, Sanatorium of II
Division, Army) | " | 56.5° | — | 0.67 | 16.3 | " | 10.14 | " 24, " | " |
| 5 | Tennōji-yu (<i>Public bath</i>) | " | 50.0° | 144 | 0.71 | 16.2 | " | 11.58 | " | " |
| 6 | Kin-taki-no-yu | " | 56.0° | 108 | — | — | " | 10.00 | " 25, " | " |
| 7 | Akagawa-yu | " | — | 144 | — | — | " | — | — | — |
| 8 | Senshu-kaku-no-yu | " | 63.0° | — | — | — | " | 12.24 | " 25, " | Y. Kinugasa
(T. Saitō) |
| 9 | Kwasui kwan-no-yu
<i>Ryōgi-aki</i> | " | 55.5° | — | — | — | " | 12.08 | " | " |
| 10 | " <i>Long spring</i> | " | 53.5° | — | — | — | " | 12.11 | " | " |
| 11 | Horikiri-no-yu
<i>Kamagata-no-yu</i> | " | 68.0 | — | — | — | " | 11.05 | " | " |

HIGASHIYAMA

Location.—Higashiyama-mura, Kitasaku-gun, Province Iwashiro, Fukushima-ken.
The springs issue from the Tertiary or the Quartz Augite Andesite.

The hot spring "Sugi-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory, 1902.

Specific gravity 1.000 at 15° C. Temperature: 59° C. Total residue ca. 1.95.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0921 | 2.3525 | 2.3525 |
| Sodium ion (Na ⁺) | 0.3898 | 16.9111 | 16.9111 |
| Calcium ion (Ca ²⁺) | 0.1797 | 4.4813 | 8.9626 |
| | | | 28.2262 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.3407 | 7.6135 | 9.0135 |
| Sulphate ion (SO ₄ ⁻²) | 0.8937 | 2.3036 | 18.6072 |
| | 1.8961 | 42.6620 | 28.2207 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0357 | | |
| | 1.9318 | | |

| | |
|--|--------|
| Potassium chloride (KCl) | 0.1755 |
| Sodium chloride (NaCl) | 4.34 |
| Sodium sulphate (Na ₂ SO ₄) | 1.89 |
| Calcium sulphate (CaSO ₄) | 0.6093 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0357 |

Besides these trace of magnesium, hydrophosphate ion and boric acid.

The spring thus may be classified as "**saline bitter spring**".

Radio-activity. 3.14 Mache's units in 1 litre of water at 9.6° C.

(Kohl-Löw. contact.; Y. Kinugasa and T. Saitō, Nov. 3, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 20.94$.

Table 24.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|--------------------------|----------------|--------------------------|---|---------------------------------|---------------------|--------------------|--|----------------------------------|--------------------------|
| | | | | | in
Maché's units | at
(temp. in C.) | | | | |
| 1 | So-yu <i>Public bath</i> | saline bitter | 56,5° | ca. 180 | 3,23 | 8,7° | K. & L. | 21,14 | Nov. 3, 1913 | {Y. Kinugasa
T. Saito |
| 2 | Sugi-no-yu | " | 50,0° | — | 3,14 | 9,6° | " | 20,94 | " | " |
| 3 | Sazanami-no-yu | " | 58,0° | 180 | 3,02 | 9,2° | " | 22,15 | " | " |
| 4 | Saru-no-yu | " | 43,0° | 180 | 3,20 | 8,1° | " | — | " | " |
| 5 | Fudô-no-yu | " | 42,0° | — | 4,59 | 10,6° | " | 16,19 | " | " |
| 6 | Nuru-yu | " | 34,0° | — | 5,30 | 10,0° | " | 14,64 | " | " |
| 7 | Takino-yu | " | 48,0° | 360 | — | — | " | — | — | — |
| 8 | Suga-no-yu | " | 61,0° | 360 | — | — | " | — | — | — |
| 9 | Kitsune-no-yu | " | 38,0° | 180 | — | — | " | — | — | — |
| 10 | Mujina-no-yu | " | 43,0° | 180 | — | — | " | — | — | — |
| 11 | Sumi-no-yu | " | 58,0° | 180 | — | — | " | — | — | — |
| 12 | Me-arai-no-yu | " | 58,0° | 180 | — | — | " | — | — | — |
| 13 | Ana-yu | " | 58,0° | 180 | — | — | " | — | — | — |
| 14 | Koga-no-yu | " | 60,0° | 180 | — | — | " | — | — | — |

KASHI

Location.—Nishigô-mura, Nishi-shirakawa-gun, Province Iwaki, Fukushima-ken.

The springs issue from the Granite.

Table 25.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|-------------|----------------|--------------------------|---|---------------------------------|---------------------|--------------------|--|----------------------------------|--------------------------|
| | | | | | in
Maché's units | at
(temp. in C.) | | | | |
| 1 | Moto-yu | simple | 51,0° | — | 4,59 | 10,2° | K. & L. | 12,63 | Nov. 5, 1913 | {Y. Kinugasa
T. Saito |
| 2 | Tengu-no-yu | " | 48,5° | — | 2,86 | 10,6° | " | 12,28 | " | " |
| 3 | Yujin-no-yu | " | 50,0° | — | 3,38 | 8,9° | " | 11,37 | " | " |

NEKONAKI

Location.—Ishikawa-machi, Ishikawa-gun, Province Iwaki, Fukushima-ken.

The springs issue from the Gneiss.

Table 26.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|------------------------------------|
| | | | | | in
10 ⁻¹⁶ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Nekonaki-no-yu | simple | — | ca. 22 | 21.47 | 5.91 | 16.0° | Schm. | Sept. 24, 1914 | S. Hanzawa
and
H. Yanagisawa |
| 2 | Tomaki-no-yu | " | 13.0° | — | 3.21 | 0.88 | 16.0° | " | " 30, " | |
| 3 | Tomaki-furu-yu | " | — | — | — | 2.18 | 17.0° | K. & L. | " " | |
| 4 | Spring A (<i>Chônakusen</i>) | " | — | — | — | 4.08 | 16.5° | " | " " | |
| 5 | Spring B (<i>Chaya</i>) | " | — | — | — | 1.82 | 19.0° | " | " 25, " | |
| 6 | Spring C
(<i>Yakôji temple</i>) | " | — | — | — | 3.18 | 16.0° | " | " 20, " | |
| 7 | Spring D | " | — | — | — | 1.95 | 15.5° | " | " 24, " | |
| 8 | Spring E | " | — | — | — | 3.07 | 16.0° | " | " 25, " | |

BOBATA

Location.—Bobata-mura, Ishikawa-gun, Province Iwaki, Fukushima-ken.

The springs issue from the Gneiss.

Table 27.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|------------------------------------|
| | | | | | in
10 ⁻¹⁶ curies | in
Mache's units | at
(Temp. in C.) | | | |
| 1 | Shimo-no-kyû-yu | simple | — | ca. — | — | 0.49 | 21.0° | K. & L. | Sept. 28, 1914 | S. Hanzawa
and
H. Yanagisawa |
| 2 | Spring A (<i>Ch. Yabô</i>) | " | — | — | — | 1.52 | 15.7° | " | " " | |
| 3 | Shimo-no-yu No. 1 | " | 27.0° | 285 | 21.05 | 3.31 | 24.0° | Schm. | " " | |
| 4 | " No. 2 | " | — | — | — | 2.36 | 21.0° | K. & L. | " " | |
| 5 | Toi-no-kuchi | " | — | — | — | 1.89 | 18.5° | " | " 20, " | |
| 6 | Spring B (<i>K. Watanabe</i>) | " | — | — | — | 9.69 | 20.8° | " | " " | |
| 7 | Naka-no-yu No. 1 | " | — | — | 8.58 | 2.36 | 15.0° | Schm. | " 20, " | |
| 8 | " No. 2 | " | — | — | — | 2.94 | 19.0° | K. & L. | " 27, " | |
| 9 | Kami-no-moto-yu | sulphur | 14.0°
17.0° | 130 | 43.48 | 11.96 | 19.0° | Schm. | " 26, " | |
| 10 | Kami-no-shin-yu | simple | — | — | 11.41 | 3.14 | 16.0° | " | " " | |
| 11 | Spring C | " | — | — | — | 0.18 | 21.0° | K. & L. | " 28, " | |

SENAMI

Location.—Senami-machi, Iwafune-gun, Province Echigo, Niigata-ken.

The hot spring “Senami-Funtō”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1905.

Specific gravity: 1.0035 at 14° C. Temperature: 102° C.

Total residue: ca. 4.19. Flow of water: ca. 9000 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0786 | 2.0077 | 2.0077 |
| Sodium ion (Na ⁺) | 1.3333 | 57.8438 | 57.8438 |
| Calcium ion (Ca ⁺⁺) | 0.1323 | 3.2993 | 6.5996 |
| Magnesium ion (Mg ⁺⁺) | 0.0003 | 0.0127 | 0.0254 |
| | | | 66.4765 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 2.1162 | 59.6954 | 59.6954 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.2759 | 2.8722 | 5.7444 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0632 | 1.0359 | 1.0359 |
| | 3.9998 | 126.7670 | 66.4757 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1928 | | |
| | 4.1926 | | |
| Free carbon dioxide (CO ₂) | 0.1596 | | 3.6273 |
| | 4.3522 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.1497 |
| Sodium chloride (NaCl) | 3.3748 |
| Sodium bicarbonate (NaHCO ₃) | 0.0132 |
| Calcium sulphate (CaSO ₄) | 0.3910 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0695 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0016 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1928 |
| | 4.1926 |
| Free carbon dioxide (CO ₂) | 0.1596 |
| | 4.3522 |

Besides these a small quantity of ferrous and aluminium ion and trace of ammonium, nitrate, hydrophosphate ion and boric acid.

The spring thus may be classified as “common salt spring”.

Radio-activity. 0.22×10^{-10} curies in 1 litre of water at 28.0° C.

(Schmidt's electrom.; Y. Kinugasa and T. Saitō, Aug. 23, 1914).

0.06 Mache's units (recalculated).

ONNAGAWA

Location.—Onnagawa-mura, Iwafune-gun, Province Echigo, Niigata-ken.

The hot spring “Takanosu-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1904.

Specific gravity: 1.002 at 15° C. Temperature: 63° C. (air temp. 31° C.)

Total residue: ca. 2.45. Flow of water: ca. 979 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|---|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0414 | 1.0575 | 1.0575 |
| Sodium ion (Na ⁺) | 0.6836 | 29.6573 | 29.6573 |
| Calcium ion (Ca ⁺⁺) | 0.1376 | 3.4314 | 6.8628 |
| Magnesium ion (Mg ⁺⁺) | 0.0020 | 0.0821 | 0.1642 |
| | | | 37.7418 |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|--|--------|
| Potassium chloride (KCl) | 0.0788 |
| Sodium chloride (NaCl) | 1.3032 |
| Sodium sulphate (Na ₂ SO ₄) | 0.5250 |
| Calcium sulphate (CaSO ₄) | 0.2984 |

Anions.

| | | | |
|--|--------|---------|---------|
| Chlorine ion (Cl') | 0.8371 | 23.3315 | 23.3315 |
| Sulphate ion (SO ₄ '') | 0.5732 | 5.9671 | 11.9342 |
| Hydrocarbonate ion (HCO ₃ '') | 0.1511 | 2.4798 | 2.4798 |
| | 2.4160 | 66.0037 | 37.7425 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0980 | | |
| | 2.5140 | | |
| Free carbon dioxide (CO ₂) | 0.1418 | 3.2227 | |
| | 2.6558 | | |

| | |
|--|--------|
| Magnesium sulphate (MgSO ₄) | 0.0094 |
| Sodium bicarbonate (NaHCO ₃) | 0.2007 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0980 |
| | 2.5140 |
| Free carbon dioxide (CO ₂) | 0.1418 |
| | 2.6558 |

Besides these trace of hydrophosphate ion.

The spring thus may be classified as "common salt spring".

Radio-activity. 4.00 Mache's units in 1 litre of water at 27.0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 25, 1914).

Table 28.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|--------------------------------|----------------|-----------------------|---|------------------------------|------------------|------------------|--------------------|-------------------------------|-------------------------|
| | | | | | in 10 curies | in Mache's units | at (temp. in C.) | | | |
| 1 | Takanosu-Onsen | common salt | air temp. 03.0 | ca. 979 | — | 4.00 | 27.0° | K. & L. | Aug. 25, 1914 | Y. Kinugasa
T. Saitō |
| 2 | Takase-no-yu
(Kiyara-no-yu) | salt | 72.0° | 504 | 15.13 | 4.16 | 27.8° | Schm. | " | " |
| 3 | Moto-yu | " | 50.0° | 65 | 11.27 | 3.10 | 26.6° | " | " 24. " | " |
| 4 | Takahashi-kwan no-yu | " | 52.0° | — | — | 3.20 | 26.6° | K. & L. | " | " |
| 5 | Sugai-no-yu | " | 48.0° | — | — | 3.60 | 25.7° | " | " | " |

MURASUGI

Location.—Sasaoka-mura, Kita-kambara-gun, Province Echigo, Niigata-ken.

The springs issue from the Granite.

The cold spring "Murasugi-Kōsen" No. 1

Analysis (calculated from the original numbers).

Analysed by the Niigata Higher Medical School. 1914.

Specific gravity: 1.0034 at 15° C.

Temperature: 25.6° C. (air temp. 30° C.)

Total residue: ca. 0.35.

Flow of water: ca. 530 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|----------------------|---------|------------|-----------------------|
| Sodium ion (Na') | 0.10417 | 4.51931 | 4.51931 |
| Calcium ion (Ca'') | 0.00810 | 0.20200 | 0.40400 |
| Magnesium ion (Mg'') | 0.00177 | 0.07266 | 0.14532 |
| | | | 5.06863 |

| | |
|--|---------|
| Sodium chloride (NaCl) | 0.03525 |
| Sodium sulphate (Na ₂ SO ₄) | 0.25171 |
| Sodium bicarbonate (NaHCO ₃) | 0.02084 |
| Calcium sulphate (CaSO ₄) | 0.02754 |
| Magnesium sulphate (MgSO ₄) | 0.00885 |
| | 0.35221 |

Anions.

| | | | |
|--|---------|---------|---------|
| Chlorine ion (Cl') | 0.02127 | 0.60000 | 0.60000 |
| Sulphate ion (SO ₄ '') | 0.20209 | 2.10379 | 4.20758 |
| Hydrocarbonate ion (HCO ₃ '') | 0.01525 | 0.24996 | 0.24996 |
| | 0.35265 | 7.74772 | 5.05754 |

The spring thus may be classified as "simple cold spring".

Radio-activity. 180.41 × 10⁻¹⁰ curies in 1 litre of water at 28.9° C.

(Schmidt's electrom.; Y. Kinugasa and T. Saitō; Aug. 20, 1914).

49.61 Mache's units (recalculated).

Discharge of Radium Emanation.

| Spring | Temp. of spring in C. | Conduit | | Per minute | | | Per hour | | | Per day (24 hours) | | | Remarks |
|--------------------------------|-----------------------|-------------------------------------|----------------------------|----------------------------|------------------------------|---------------------|---------------------------------|------------------------------|---------------------|---------------------------------|---|-------------------------------------|-----------------------|
| | | Distance
(source—bath
in feet | Material | Flow of water
in litres | Emanation | | Flow of water
in hectolitres | Emanation | | Flow of water
in hectolitres | Emanation | | |
| | | | | | in
10 ⁸ curies | in
Mache's units | | in
10 ⁸ curies | in
Mache's units | | in
10 ⁶ curies
(microcuries) | in
10 ³ Mache's units | |
| Murasugi Kōsen
<i>No. 1</i> | 25.6° | ca.
48 | bamboo and
wooden pipes | 36.8 | 66 | 1826 | 22 | 3984 | 110 | 530 | 956 | 2629 | heated for
bathing |

Table 29.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|--|----------------|-----------------------|---|-------------------------------------|------------------|------------------|--------------------|---------------------------------|---------------------------|
| | | | | | in 10 ¹⁰ curies | in Mache's units | at (temp. in C.) | | | |
| 1 | Murasugi-Kōsen No. 1 | simple | air temp. 25.6° (30°) | ca. 530 | 180.41 | 49.61 | 28.9° | Schm. | Aug. 20, 1914 | Y. Kinugasa
(T. Saitō) |
| 2 | " No. 2 | " | 26.0° (31.5°) | — | 176.61 | 48.57 | 27.6° | " | " 21, " | " |
| 3 | " No. 3 | " | 13.5° (30°) | — | — | 40.28 | 26.0° | K. & L. | " | " |
| 4 | Shiroyama-Kōsen | " | 22.5° (30°) | — | — | 13.74 | 26.3° | " | " | " |
| 5 | Imaita-Kōsen No. 1 * | " | 18.7° (32°) | — | — | 7.45 | 26.9° | " | " 20, " | " |
| 6 | " No. 2 * | " | 16.0° 32 | — | — | 4.19 | 26.8° | " | " | " |
| 7 | Uba-yu * | " | 14.0° (28°) | — | — | 3.26 | 27.0° | " | " 22, " | " |
| | Gas evolving from Murasugi-Kōsen No. 2 | — | — | — | Emanation per litre of gas at 0- C. | | | " | Date when the gas was collected | " |
| | | | | | — | 169.98 | — | " | Aug. 21, 1914 | " |

The geological structure of the place, where the spring gushes out, is not yet examined.

DEYU

Location.—Sasaoka-mura, Kita-kambara-gun, Province Echigo, Niigata-ken.

The springs issue from the Granite.

Table 30.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Radio-activity per litre of water | | | Apparatus employed | Date when the water was analysed | Examined by |
|-----|-------------------------------------|----------------|-----------------------|---|-----------------------------------|------------------|---------------|--------------------|---|----------------|
| | | | | | % 10 ⁻¹⁰ curies | in Mache's units | (temp. in C.) | | | |
| 1 | Chosenkutsu-no-yu | carbonated | 39.5° (27°) | ca. 830 | 12.09 | 3,33 | 30.1 | Schub | Aug. 22, 1914 | J. Y. Kinugasa |
| 2 | Toshundai-no-yu (No. 1) | " | 33.0° (27°) | | | 11.57 | 26.1° | K. & L. | " | " |
| 3 | " (No. 2) | " | 31.0° (25°) | | | 11.31 | 26.4 | " | " | " |
| 4 | Shiranoya-no-yu | " | 33.5° (27°) | | — | 6.99 | 26.3° | " | " | " |
| | Gas evolving from Chosenkutsu-no-yu | | | | | 14.60 | | " | Date when the gas was analysed
Aug. 22, 1914 | " |

TOCHIOMATA

Location.—Yunotani-mura, Kita-uo-numa-gun, Province Echigo, Niigata-ken.

The springs issue from the Granite.

The hot spring "Tochiomata-no-yu" No. 1

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1896.

Total residue: ca. 0.28.

Flow of water: ca. 1800 hectolitres in 24 hours.

Temperature: 39° C. (air temp. 24.5° C.)

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

Cations.

| | Grams | Milli-grams | Milligram-equivalents |
|---------------------------------|--------|-------------|-----------------------|
| Potassium ion (K ⁺) | 0.0013 | 0.0332 | 0.0332 |
| Sodium ion (Na ⁺) | 0.0266 | 1,1540 | 1,1540 |
| Calcium ion (Ca ⁺⁺) | 0.0435 | 1,0847 | 2,1694 |
| | | | 3,3566 |

Anions.

| | Grams | Milli-grams | Milligram-equivalents |
|--|--------|-------------|-----------------------|
| Chlorine ion (Cl ⁻) | 0.0171 | 0,4541 | 0,4541 |
| Sulphate ion (SO ₄ '') | 0.1394 | 1,4511 | 2,9022 |
| | | 0,2267 | 4,1771 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0308 | | |
| | 0.2577 | | |

| | Grams |
|--|--------|
| Sodium chloride (NaCl) | 0.0013 |
| Potassium sulphate (K ₂ SO ₄) | 0.0029 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0501 |
| Calcium sulphate (CaSO ₄) | 0.1474 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0308 |
| | 0.2577 |

Besides these a small quantity of ferrous and aluminium ion and trace of nitrate and hydrophosphate ion.

The spring thus may be classified as "**simple thermal**".

Radio-activity. 94.03×10⁻¹⁰ curies in 1 litre of water at 25.7° C.

(Schmidt's electrom.; Y. Kinugasa and T. Saitō, Aug. 27, 1914).

25.86 Mache's units (recalculated).

Discharge of Radium Emanation.

| No. | Spring | Temp. of spring in C. | Conduit | | Per minute | | | Per hour | | | Per day (24 hours) | | | Remarks |
|-----|--------------------------|-----------------------|--------------------------------|------------|-------------------------|---------------------------|------------------|------------------------------|---------------------------|----------------------------------|------------------------------|---|----------------------------------|---------------------------------------|
| | | | Distance (source bath in feet) | Material | Flow of water in litres | in 10 ⁸ curies | in Mache's units | Flow of water in hectolitres | in 10 ⁸ curies | in 10 ³ Mache's units | Flow of water in hectolitres | in 10 ⁸ curies (microcuries) | in 10 ³ Mache's units | |
| 1 | Tochiomata-no-yu No. 1 | 39° | 0 | — | 125 | 118 | 32.33 | 75 | 7052 | 194 | 1800 | 1693 | 4655 | { bath-tank is direct upon the source |
| 2 | Jizai-kwan-uchi-yu No. 1 | 36° | 180 | iron pipes | 16.4 | 27 | 730 | 10 | 1590 | 44 | 236 | 382 | 1050 | |

Table 31.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|--------------------------|----------------|-------------------------|---|------------------------------|------------------|------------------|--------------------|-------------------------------|---------------------------|
| | | | | | in 10 ¹⁰ curies | in Mache's units | at (temp. in C.) | | | |
| 1 | Tochiomata-no-yu No. 1 | simple | 39.0° (air temp. 21.5°) | ca. 1800 | 94.03 | 25.86 | 25.7° | Schm. | Aug. 27, 1914 | { Y. Kinugasa
T. Saito |
| 2 | " No. 2 | " | 39.0° (26°) | — | — | 23.33 | 27.6° | K. & L. | " 29, " | " |
| 3 | " No. 3 | " | 35.5° (26°) | — | — | 27.18 | 26.2° | " | " 27, " | " |
| 4 | " No. 4 | " | 28.5° (24.5°) | — | — | 56.41 | 25.3° | " | " 29, " | " |
| 5 | Jizai-kwan-uchi-yu No. 1 | " | 36.0° (27.5°) | 236 | 161.74 | 44.48 | 26.5° | Schm. | " | " |
| 6 | " No. 2 | " | 38.0° (24.5°) | 236 | — | 24.05 | 28.1° | K. & L. | " | " |
| 7 | O-yu (Otoko-yu) | " | 57.0° (33°) | 1170 | 11.25 | 3.09 | 29.8° | Schm. | " 28, " | " |
| 8 | " (Onna-yu) | " | 56.0° (33°) | — | — | 3.52 | 27.5° | K. & L. | " | " |
| 9 | Sakashita-no-yu | " | 56.0° (33°) | — | — | 3.62 | 26.3° | " | " | " |
| 10 | Yakushi-no-yu | " | 53.0° (33°) | — | — | 3.43 | 24.2° | " | " | " |
| 11 | Kawagashi-no-yu No. 1 | " | 56.0° (33°) | — | — | 4.14 | 29.0° | " | " | " |
| 12 | " No. 2 | " | 56.0° (33°) | — | — | 3.83 | 23.7° | " | " | " |

YUZAWA

Location.—Yuzawa-mura, Minami-uo-numa-gun, Province Echigo, Niigata-ken.

The springs issue from the Misaka Series.

Table 32.

| No. | Spring | Classification | Temp.
of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------|----------------|-------------------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|----------------------------|
| | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Moto-yu | salt | 45.0 ² (air temp. 29.5°) | | 1.61 | 29.3 | K. & L. | Aug. 31, 1914 | (Y. Kanagata)
(T. Sait) |
| 2 | Chūbu-no-yu | .. | 42.0 ² (29.1°) | | 1.30 | 29.1 | .. | .. | .. |
| 3 | Tamago-no-yu | .. | 41.5 ¹ | | 1.23 | 26.7° | .. | .. 30. " | .. |
| 4 | Kami-no-yu | .. | 45.0 ⁶ (29.1°) | | 1.45 | 29.6° | .. | .. 31. " | .. |
| 5 | Shin-yu | .. | 43.0° (29.5°) | - | 1.23 | 29.3° | .. | .. | .. |
| 6 | Kiridōshi-no-yu | .. | 37.0 ⁷ | - | 2.73 | 27.8 | .. | .. | .. |

AKAKURA

Location.—Nakayama-mura, Nakakubiki-gun, Province Echigo, Niigata-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring "Akakura-Onsen"

Analysis.

Analysed by the Takata Hospital. 1896.

Total residue: ca. 1.19.

Temperature: 62° C. (air temp. 20° C.)

Flow of water: ca. 7200 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| Grams | | Hypothetical form of combination | |
|--|--------|---|--------|
| Potassium oxide (K ₂ O) | 0.0544 | Sodium chloride (NaCl) | 0.0468 |
| Sodium oxide (Na ₂ O) | 0.6560 | Potassium carbonate (K ₂ CO ₃) | 0.0071 |
| Calcium oxide (CaO) | 0.1431 | Sodium carbonate (Na ₂ CO ₃) | 0.1027 |
| Magnesia (MgO) | 0.0378 | Calcium sulphate (CaSO ₄) | 0.3688 |
| Ferric oxide (Fe ₂ O ₃) | 0.0034 | Ferrous carbonate (FeCO ₃) | 0.0070 |
| Chlorine (Cl) | 0.0284 | Calcium carbonate (CaCO ₃) | 0.0865 |
| Sulphuric anhydride (SO ₃) | 0.2603 | Magnesium carbonate (MgCO ₃) | 0.1323 |
| Combined carbon dioxide (CO ₂) | 0.2533 | Silica (SiO ₂) | 0.1678 |
| Silica (SiO ₂) | 0.1678 | Organic substances | 0.1785 |
| Organic substances | 0.1785 | | 1.1926 |
| Free carbon dioxide (CO ₂) | 0.5384 | Free carbon dioxide (CO ₂) | 0.5384 |
| Hydrogen sulphide (H ₂ S) | 0.0268 | Hydrogen sulphide (H ₂ S) | 0.0268 |
| | 2.3482 | | 1.1926 |

Besides these trace of aluminium and phosphoric acid.

The spring may be classified as "earthy alkaline sulphur spring".

Table 33.

| No. | Spring | Character | Temp. of spring in C. | Emanation per litre of water | | Apparatus employed | Date when the water was taken | Examined by |
|-----|---------------------|--------------------------|-----------------------|------------------------------|----------------|--------------------|-------------------------------|---------------|
| | | | | in Mechie's units | at temp. in C. | | | |
| 1 | Matara-Oson / / A. | earthly alkaline sulphur | 55.5° air temp. | 0.33 | 21.2 | K. & L. | July, 20, 1913 | Dr. R. Ishizu |
| 2 | " / / A. | " | 62.0 20 | 1.04 | 22.6° | " | Aug. 13, 1914 | Y. Kinugasa |
| 3 | " / / A. | " | 62.0 20 | 0.54 | 23.0 | " | " | " |
| 4 | Isakami-Oson / / A. | sulphur | 48.0 22.5 | 0.09 | 27.2° | " | July, 29, 1913 | Dr. R. Ishizu |
| 5 | " / / A. | " | 44.0 27.5 | 0.23 | 27.5 | " | " | " |
| 6 | " / / A. | " | 42.0 27.5 | 0.74 | 26.3 | " | " | " |

Fubine, located at *Sekiguchi-dera*, about 2.5 m. N. from Akakura, is grouped for convenience in this table.

MATSUNOYAMA

Location.—Matsunoyama-mura, Higashi-kubiki-gun, Province Echigo, Niigata-ken.

The springs issue from the Tertiary.

The hot spring "Netsu-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1895.

Temperature: 63° C.

Total residue: ca. 15.08.

in 1 kilogram of the mineral water are contained

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram,

| Cations. | Grams | Milli-moles | Milli-equivalents |
|--|---------|-------------|-------------------|
| Potassium chloride (KCl) | 0.1652 | 2.6181 | 2.6181 |
| Sodium chloride (NaCl) | 8.6831 | 148.4208 | 148.4208 |
| Calcium chloride (CaCl ₂) | 1.3287 | 18.0000 | 97.9600 |
| | | | 248.9989 |
| Anions. | | | |
| Chloride ion (Cl ⁻) | 7.531 | 24.7550 | 247.0550 |
| Sulphate ion (SO ₄ ⁻²) | | 27.5 | 1.9530 |
| | 14.5100 | 148.6504 | 249.0080 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0896 | | |
| | 14.4204 | | |

| | |
|--|---------|
| Potassium chloride (KCl) | 0.1652 |
| Sodium chloride (NaCl) | 8.6831 |
| Calcium chloride (CaCl ₂) | 1.3287 |
| Calcium sulphate (CaSO ₄) | 0.1334 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0896 |
| | 14.4204 |

Besides these trace of bromine, iodine, hydrophosphate ion, boric acid and organic substances.

The spring thus may be classified as "earth-muriated common salt spring".

NASU

Location: Nasu-ku, Nasu-shi, Tochigi-ken, Japan.

The spring issues from the Aogite Andesite.

YUMOTO

The hot spring "Shika-no-yu"

Analysis calculated from the salt table.

Analysed by the Tōkyō Imperial Hygienic Laboratory 1911.

Temperature: 74.5° C.

Total residue: ca. 1.91.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams. | Milli-mols. | Meq. residue-equivalent. | | |
|--|--------|-------------|--------------------------|---|--------|
| Hydrogen ion (H ⁺) | 0.0018 | 11.6832 | 11.6832 | Potassium sulphate (K ₂ SO ₄) | 0.137 |
| Potassium ion (K ⁺) | 0.0071 | 0.1814 | 0.1814 | Sodium sulphate (Na ₂ SO ₄) | 0.115 |
| Sodium ion (Na ⁺) | 0.0307 | 1.5922 | 1.5922 | Calcium sulphate (CaSO ₄) | 0.2072 |
| Calcium ion (Ca ⁺⁺) | 0.0611 | 1.5237 | 3.0474 | Magnesium sulphate (MgSO ₄) | 0.3045 |
| Magnesium ion (Mg ⁺⁺) | 0.0609 | 2.5000 | 5.0000 | Ferrous sulphate (FeSO ₄) | 0.0595 |
| Ferrous ion (Fe ⁺⁺) | 0.0219 | 0.3918 | 0.7836 | Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.3209 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0507 | 1.8708 | 5.6124 | Free hydrochloric acid (HCl) | 0.2140 |
| | | | 27.9002 | Free sulphuric acid (H ₂ SO ₄) | 0.5609 |
| Anions. | | | | Silicic acid (meta) (H ₂ SiO ₃) | 0.2530 |
| Chlorine ion (Cl ⁻) | 0.2080 | 5.8674 | 5.8674 | | 2.0491 |
| Hydrosulphate ion (HSO ₄ ⁻) | 0.5551 | 5.7186 | 5.7186 | Free hydrogen sulphide (H ₂ S) | 0.0240 |
| Sulphate ion (SO ₄ ^{''}) | 0.7828 | 8.1461 | 16.2982 | | 2.0731 |
| | 1.7961 | 39.4782 | 27.8842 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2530 | | | | |
| | 2.0491 | | | | |
| Free hydrogen sulphide (H ₂ S) | 0.0240 | | | | |
| | 2.0731 | | | | |

Besides these trace of hydrophosphate ion and organic substances.

The spring thus may be classified as "acid hydrogen sulphide vitriol spring".

Radio-activity. 0.41 Mache's units in 1 litre of water at 13.9° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 7, 1913)

Electric conductivity. $\kappa_{18} \times 10^{-4} = 37.57$.

The hot spring "Komatsu-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1912.

Specific gravity: 1.0013 at 15° C.

Temperature: 43° C.

Total residue: ca. 0.72.

The spring, being in its composition nearly the same as that of "Shika-no-yu", differing only in the smaller proportion of iron from the latter, may be classified as "acid hydrogen sulphide spring".

Radio-activity. 0.56 Mache's units in 1 litre of water at 13.8° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 7, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 23.47$.

The hot spring "Kiraku-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1912.

Specific gravity: 1.0013 at 15° C.

Temperature: 28° C.

Total residue: ca. 0.72.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|---|--------|--|--------|
| Sodium chloride (NaCl) | 0,0356 | Silica (SiO ₂). | 0,1303 |
| Potassium sulphate (K ₂ SO ₄). | 0,0294 | | 0,6248 |
| Sodium sulphate (Na ₂ SO ₄). | 0,0092 | Carbon dioxide (free and in form of bicar- | |
| Sodium carbonate (Na ₂ CO ₃). | 0,0740 | bonates) (CO ₂). | 0,1078 |
| Calcium sulphate (CaSO ₄). | 0,2308 | Free hydrogen sulphide (H ₂ S). | 0,0292 |
| Magnesium carbonate (MgCO ₃). | 0,1082 | | 0,7618 |
| Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃). | 0,0073 | | |

Besides these a very small quantity of phosphoric acid and trace of organic substances.

The spring may be classified as "**hydrogen sulphide spring**".

Radio-activity. 1.08 Mache's units in 1 litre of water at 13,0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 7, 1913)

Electric conductivity. $\kappa_{18} \times 10^{-4} = 6,31$.

KITA

The hot spring "Ai-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1912.

Specific gravity: 1,0010 at 15° C.

Temperature: 54° C.

Total residue: ca. 0,64.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|---|--------|---|--------|
| Potassium chloride (KCl) | 0,0111 | Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃). . . | 0,0024 |
| Potassium sulphate (K ₂ SO ₄). | 0,0026 | Silica (SiO ₂). | 0,1598 |
| Sodium sulphate (Na ₂ SO ₄). | 0,0511 | | 0,6144 |
| Sodium carbonate (Na ₂ CO ₃). | 0,1173 | Carbon dioxide (free and in form of bicar- | |
| Calcium sulphate (CaSO ₄). | 0,2188 | bonates) (CO ₂). | 0,3398 |
| Magnesium carbonate (MgCO ₃). | 0,0513 | | 0,9542 |

Beside these trace of iodine, phosphoric and boric acid.

The spring may be classified as "**simple thermal**".

Radio-activity. 1.15 Mache's units in 1 litre of water at 15,3° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 7, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 5,55$.

BENTEN

The hot spring "Kawa-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1912.

Specific gravity: 1,0010 at 16° C.

Temperature: 54° C.

Total residue: ca. 0,85.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|---|--------|--|--------|
| Sodium chloride (NaCl) | 0,0410 | Alumina (Al ₂ O ₃). | 0,0008 |
| Potassium sulphate (K ₂ SO ₄). | 0,0096 | Silica (SiO ₂). | 0,1589 |
| Sodium sulphate (Na ₂ SO ₄). | 0,1615 | | 0,9171 |
| Sodium bicarbonate (NaHCO ₃). | 0,1044 | Carbon dioxide (free and in form of bicarbo- | |
| Calcium sulphate (CaSO ₄). | 0,2770 | mates) (CO ₂). | 0,2705 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,1617 | | 1,1876 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0022 | | |

Besides these trace of phosphoric and boric acid.

The spring may be classified as "**simple thermal**".

Radio-activity. 2.07 Mache's units in 1 litre of water at 10,2° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 7, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 8,13$.

DAIMARU-ZUKA

The hot spring "Sakura-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1911. Specific gravity: 1.0009 at 15° C.
Temperature: 71° C. Total residue: ca. 0.75.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|--|--------|
| Potassium sulphate (K_2SO_4) | 0.0389 | Silica (SiO_2) | 0.2083 |
| Sodium sulphate (Na_2SO_4) | 0.1665 | | 0.7540 |
| Calcium sulphate ($CaSO_4$) | 0.2174 | Carbon dioxide (free and in form of bicar- | |
| Magnesium sulphate ($MgSO_4$) | 0.0740 | bonates) (CO_2) | 0.0190 |
| Magnesium carbonate ($MgCO_3$) | 0.0479 | | 0.7730 |
| Ferric oxide and alumina ($Fe_2O_3+Al_2O_3$) | 0.0010 | | |

Besides these trace of chlorine, iodine, nitric and phosphoric acid.

The spring may be classified as "simple thermal".

Radio-activity. 0.42 Mache's units in 1 litre of water at 13.4° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 7, 1913).

Electric conductivity. $\times_{18} \times 10^{-1} = 6.46$.

SANDO-GOYA

The hot spring "Sando-goya-Onsen"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1911.
Specific gravity: 1.0005 at 15° C. Temperature: 52° C. (air temp. 18° C.)
Total residue: ca. 0.59. Flow of water: ca. 1800 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|--|--------|
| Sodium chloride ($NaCl$) | 0.0031 | Alumina (Al_2O_3) | 0.0014 |
| Sodium sulphate (Na_2SO_4) | 0.1837 | Silica (SiO_2) | 0.3385 |
| Sodium carbonate (Na_2CO_3) | 0.0084 | | 0.5973 |
| Calcium sulphate ($CaSO_4$) | 0.0006 | Carbon dioxide (free and in form of bicar- | |
| Magnesium carbonate ($MgCO_3$) | 0.0578 | bonates) (CO_2) | 0.1449 |
| Ferrous carbonate ($FeCO_3$) | 0.0038 | | 0.7422 |

Besides these trace of phosphoric acid.

The spring may be classified as "simple thermal".

Composition of the sinter-deposit (analysed in 1912).

In 100 parts of the yellowish powder are found as main constituents:

| | | | |
|------------------------|-------|--------------------------------------|----------------|
| Sulphur | 92.10 | Iron oxide, alumina and silica . . . | small quantity |
| Ignition loss. | 1.34 | | |
| Water. | 4.20 | | |
| | 97.64 | | |

Table 34.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|------------------------------------|-----------------------------------|--------------------------|---|---------------------------------|-------------------|--------------------|--|----------------------------------|---------------------------|
| | | | | | in
Mache's units | at
temp. in C. | | | | |
| 1 | Shika-no-yu (<i>Public bath</i>) | acid hydrogen
sulphide vitriol | 74.5° | ca. | 0.41 | 13.9° | K. & L. | 37.57 | Nov. 7, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Komatsu-no-yu | acid hydrogen
sulphide | 43.0° | | 0.56 | 13.8° | " | 23.47 | " | |
| 3 | Kiraku-no-yu | hydrogen
sulphide | 28.0° | — | 1.08 | 13.0° | " | 6.31 | " | |
| 4 | Taki-no-yu (<i>Public bath</i>) | " ?) | 74.0° | | 0.73 | 13.8° | " | 37.28 | " | |
| 5 | Takaomata-no-yu | sulphur | 34.0 | | — | — | — | — | — | |
| 6 | Ai-no-yu | simple | 54.0° | — | 1.15 | 15.3° | K. & L. | 5.55 | Nov. 7, 1913 | { Y. Kinugasa
T. Saitō |
| 7 | Tengu-no-yu | " | 53.5° | — | — | — | — | — | — | |
| 8 | Izumi-no-yu | " | 51.0° | — | — | — | — | — | — | |
| 9 | Kawa-no-yu | " | 54.0° | — | 2.07 | 10.2° | K. & L. | 8.13 | Nov. 7, 1913 | { Y. Kinugasa
T. Saitō |
| 10 | Ai-no-yu | " | 48.0° | — | — | — | — | 8.05 | " | |
| 11 | Sakura-no-yu | " | 71.0° | — | 0.42 | 13.4° | K. & L. | 6.46 | " | |
| 12 | Ai-no-yu | " | 61.5° | — | — | — | — | 6.72 | " | |
| 13 | Sando-goya-Onsen | " | 52.0° | 1800 | — | — | — | — | — | |
| 14 | Itamuro-no-yu | " | 38.0° | 540 | — | — | — | — | — | |

SHIOBARA

Location —Shiobara-mura, Shiōya-gun, Province Shimozuke, Tochigi-ken.

ŌAMI

The springs issue from the Tertiary.

The hot spring “Ishi-ai-no-yu”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1912.

Temperature: 57.5° C.

Specific gravity: 1.0018 at 15° C.

Total residue ca. 1.49.

In 1 kilogram of the mineral water are contained

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram

| Cations. | Grams | Milli-
mols | Milligram-
equivalents | | Grams |
|--|---------|----------------|---------------------------|--|--------|
| Potassium ion (K') | 0.00838 | 0.21405 | 0.21405 | Potassium chloride (KCl). | 0.0160 |
| Sodium ion (Na') | 0.28609 | 12.41041 | 12.41041 | Sodium chloride (NaCl). | 0.0328 |
| Ammonium ion (NH ₄ ') | 0.00212 | 0.11726 | 0.11726 | Ammonium chloride (NH ₄ Cl). | 0.0063 |
| Calcium ion (Ca'') | 0.14571 | 3.64275 | 7.28550 | Sodium sulphate (Na ₂ SO ₄) | 0.8422 |
| | | | 20.02722 | Calcium sulphate (CaSO ₄) | 0.4411 |

Anions.

| | | | | | |
|--|---------|----------|----------|---|--------|
| Chlorine ion (Cl') | 0,03164 | 0,89252 | 0,89252 | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,0649 |
| Sulphate ion (SO ₄ '') | 0,88050 | 0,10015 | 18,33230 | Boric acid (meta) (HBO ₂) | 0,0085 |
| Hydrocarbonate ion (HCO ₃ ') | 0,04895 | 0,80233 | 0,80233 | Free carbon dioxide (CO ₂) | 1,5040 |
| | 1,40336 | 27,24547 | 20,02715 | | 0,0650 |
| Boric acid (meta) (HBO ₂) | 0,00851 | | | | 1,5690 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,09218 | | | | |
| | 1,50405 | | | | |
| Free carbon dioxide (CO ₂) | 0,06501 | 1,47750 | | | |
| | 1,56906 | | | | |

The spring thus may be classified as “saline bitter spring”.

Radio-activity. 4,15 Mache's units in 1 litre of water at 8,0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 10, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 16,19$.

FUKUWATA

The springs issue from the Tertiary Tuff.

The hot spring “Hadaka-no-yu”

Analysis.

Analysed by the Yokohama Imperial Hygienic Laboratory. 1887. Specific gravity: 1,0014 at 15° C.

Temperature: 50° C.

Total residue: ca. 1,41.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Sodium chloride (NaCl) | 0,7840 | Alumina (Al ₂ O ₃) | 0,0020 |
| Potassium sulphate (K ₂ SO ₄) | 0,0653 | Silica (SiO ₂) | 0,0720 |
| Sodium sulphate (Na ₂ SO ₄) | 0,1460 | | 1,3901 |
| Sodium carbonate (Na ₂ CO ₃) | 0,0326 | Carbon dioxide (free and in form of bicar- | |
| Calcium carbonate (CaCO ₃) | 0,2655 | bonates) (CO ₂) | 0,5544 |
| Magnesium carbonate (MgCO ₃) | 0,0212 | | 1,9445 |
| Ferrous carbonate Fe(CO ₃) | 0,0015 | | |

Besides these trace of lithium, strontium, manganese, bromine, iodine, and nitric, phosphoric and boric acid, and organic substances.

The spring may be classified as “common salt spring”.

Radio-activity. 5,76 Mache's units in 1 litre of water at 12,6° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 8, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 19,63$.

The hot spring “Hie-no-yu”

Analysis.

Analysed by the Yokohama Imperial Hygienic Laboratory. 1887. Specific gravity: 1,0019 at 15° C.

Temperature: 50° C.

Total residue: ca. 2,36.

The spring, being in its composition nearly the same as that of “Hadaka-no-yu”, may be classified as “common salt spring”.

Radio-activity. 1,42 Mache's units in 1 litre of water at 9,8° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 8, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 30,16$.

SHIONOYU

The springs issue from the Tertiary Tuff.

The hot spring "Naka-no-yu"

Analysis.

Analysed by the Yokohama Imperial Hygienic Laboratory. 1887. Specific gravity: 1.0029 at 15° C.
Temperature: 71.5° C. Total residue: ca. 3.84.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Potassium chloride (KCl) | 0.1114 | Alumina (Al ₂ O ₃) | 0.0048 |
| Sodium chloride (NaCl) | 2.3271 | Silica (SiO ₂) | 0.1306 |
| Potassium sulphate (K ₂ SO ₄) | 0.6645 | | 3.8059 |
| Calcium chloride (CaCl ₂) | 0.0997 | Carbon dioxide (free and in form of bicar- | |
| Calcium carbonate (CaCO ₃) | 0.4240 | bonates (CO ₂) | 0.4958 |
| Magnesium carbonate (MgCO ₃) | 0.0386 | | 4.3017 |
| Ferrous carbonate (FeCO ₃) | 0.0100 | | |

Besides these trace of lithium, strontium, manganese, bromine, iodine, and phosphoric and boric acid, and organic substances.

The spring may be classified as "common salt spring".

Radio-activity. 0.87 Mache's units in 1 litre of water at 12.5° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 9, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 51.00$.

HATAORI

The springs issue from the Tertiary.

Analysis of the spring water (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1901. Total residue: ca. 1.35.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams | Milli-mols | Milligram-equivalents | | Grams |
|--|--------|------------|-----------------------|---|--------|
| Cations. | | | | Sodium chloride (NaCl) | 0.6238 |
| Potassium ion (K ⁺) | 0.0260 | 0.6641 | 0.6641 | Potassium sulphate (K ₂ SO ₄) | 0.0580 |
| Sodium ion (Na ⁺) | 0.4028 | 17.4751 | 17.4751 | Sodium sulphate (Na ₂ SO ₄) | 0.0413 |
| Calcium ion (Ca ⁺⁺) | 0.0571 | 1.4240 | 2.8480 | Sodium bicarbonate (NaHCO ₃) | 0.5234 |
| Magnesium ion (Mg ⁺⁺) | 0.0144 | 0.5911 | 1.1824 | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.2312 |
| | | | 22.1694 | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0866 |
| Anions. | | | | Silicic acid (meta) (H ₂ SiO ₃) | 0.1716 |
| Chlorine ion (Cl ⁻) | 0.3780 | 10.6629 | 10.6629 | | 1.7359 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.0599 | 0.6236 | 1.2472 | Free carbon dioxide (CO ₂) | 0.5569 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.6261 | 10.2623 | 10.2623 | | 2.2928 |
| | 1.5643 | 41.7031 | 22.1724 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1716 | | | | |
| | 1.7359 | | | | |
| Free carbon dioxide (CO ₂) | 0.5569 | 12.6568 | | | |
| | 2.2928 | | | | |

Besides these trace of lithium, ferrous, manganous, aluminium, bromine, iodine and hydrophosphate ion, boric acid, and organic substances.

The spring thus may be classified as "muriated alkaline spring".

SUMAKI

The springs issue from the Tertiary.

The hot spring "Taki-no-yu"

Analysis.

Analysed by the Yokohama Imperial Hygienic Laboratory. 1887. Specific gravity: 1.0011 at 15° C.

Temperature: 62.5° C.

Total residue: ca. 0.78.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|--|--------|
| Potassium chloride (KCl) | 0.0097 | Alumina | 0.0012 |
| Sodium chloride (NaCl) | 0.2849 | Silica (Al ₂ O ₃) | 0.1660 |
| Potassium sulphate (K ₂ SO ₄) | 0.0575 | | 0.7900 |
| Sodium carbonate (Na ₂ CO ₃) | 0.0352 | Carbon dioxide (free and in form of bicar- | |
| Calcium carbonate (CaCO ₃) | 0.1763 | bonates) (CO ₂) | 0.3161 |
| Magnesium carbonate (MgCO ₃) | 0.0573 | | 1.1061 |
| Ferrous carbonate (FeCO ₃) | 0.0019 | | |

Besides these trace of lithium, strontium, manganese, bromine, iodine, phosphoric acid, boric acid and organic substances.

The spring may be classified as "simple thermal".

Radio-activity. 0.91 Mache's units in 1 litre of water at 9.6° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 10, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 11.44$.

MONZEN

The springs issue from the Tertiary.

The hot spring "Kawara-no-yu"

Analysis.

Analysed by the Yokohama Imperial Hygienic Laboratory. 1887. Specific gravity: 1.0012 at 15° C.

Temperature: 54° C.

Total residue: ca. 1.33.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Potassium chloride (KCl) | 0.0715 | Alumina (Al ₂ O ₃) | 0.0010 |
| Sodium chloride (NaCl) | 0.6745 | Silica (SiO ₂) | 0.2105 |
| Potassium sulphate (K ₂ SO ₄) | 0.0051 | | 1.4216 |
| Sodium carbonate (Na ₂ CO ₃) | 0.0149 | Carbon dioxide (free and in form of bicar- | |
| Calcium carbonate (CaCO ₃) | 0.2210 | bonates) (CO ₂) | 0.5102 |
| Magnesium carbonate (MgCO ₃) | 0.1221 | | 1.9318 |
| Ferrous carbonate (FeCO ₃) | 0.0024 | | |

Besides these trace of lithium, strontium, manganese, bromine, iodine, and nitric, phosphoric and boric acid, and organic substances.

The spring may be classified as "common salt spring".

Radio-activity. 0.79 Mache's units in 1 litre of water at 12.0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 9, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 21.01$.

The hot spring "Jirakubō-no-yu"

Analysis.

Analyst: Unknown.

Specific gravity: 1,0015 at 15° C.

Temperature: 50° C.

Total residue: ca. 148.

The spring, being in its composition nearly the same as that of "Kawara-no-yu", may be classified as "common salt spring".

Radio-activity. 1.03 Mache's units in 1 litre of water at 12,1° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 9, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 16,09$.

FURUMACHI

The springs issue from the Tertiary.

The hot spring "Gosho-no-yu"

Analysis.

Analysed by the Yokohama Imperial Hygienic Laboratory. 1887.

Specific gravity: 1,0013 at 15° C.

Temperature: 51,5° C.

Total residue: ca. 1,16.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Sodium chloride (NaCl) | 0,2564 | Alumina (Al ₂ O ₃) | 0,0006 |
| Potassium sulphate (K ₂ SO ₄) | 0,1037 | Silica (SiO ₂) | 0,1778 |
| Sodium sulphate (Na ₂ SO ₄) | 0,0167 | | 1,2021 |
| Sodium carbonate (Na ₂ CO ₃) | 0,4977 | Carbon dioxide (free and in form of bicar- | |
| Calcium carbonate (CaCO ₃) | 0,1023 | bonates (CO ₂) | 0,4080 |
| Magnesium carbonate (MgCO ₃) | 0,0466 | | 1,6101 |
| Ferrous carbonate (FeCO ₃) | 0,0003 | | |

Besides these trace of lithium, strontium, manganese, bromine, iodine, and nitric, phosphoric and boric acid, and organic substances.

The spring may be classified as "alkaline spring".

Radio-activity. 0.72 Mache's units in 1 litre of water at 12,1 °C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 9, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 14,54$.

Composition of the sinter-deposit (analysed in 1908).

Sample 1.

In 100 parts of the grayish coarse grain, slightly soluble in water with a strong acid reaction, are found as soluble constituents:

| | | | |
|----------------------------------|-------------|-----------------------------|----------------|
| Free sulphuric acid | 10,270 pts. | Calcium oxide | small quantity |
| Sulphates. | 0,789 " | Potassium oxide } | laint quantity |
| Iron oxide and alumina | 0,680 " | Sodium oxide } | |
| Ignition loss | 1,940 " | Magnesia | |
| | | Chlorine | trace |

Sample 2.

In 100 parts of the brownish coarse grain, almost soluble in water with a strong acid reaction, are found:

Insoluble part. 20,866 pts.

| | | | |
|-----------------------|-------------|--|------------|
| Soluble part. | 79,134 pts. | { CaO | 0,600 pts. |
| | | { Fe ₂ O ₃ | 4,320 " |
| | | { Al ₂ O ₃ | 6,480 " |
| | | { SO ₃ | 28,465 " |

Table 35.

| No. | Spring | Classification | Temp. of spring in C. | Emanation per litre of water | | Apparatus employed | Electric conductivity at 18° C. | Date when the water was taken | Examined by |
|-----|------------------|-------------------|-------------------------|------------------------------|----------------|--------------------|---------------------------------|-------------------------------|---|
| | | | | in March's mts. | at temp. in C. | | | | |
| 1 | Ishii | Ishi-ai-no-yu | saline bitter | 57.5° | 4.15 | 8.0 | K. & L. | 10.19 | Nov. 10, 1913 { Y. Kinugasa
T. Saitō |
| 2 | | Kawara-no-yu | " | 55.0 | 4.11 | 8.0 | " | 15.51 | |
| 3 | | Hadaka-no-yu | common salt | 50.0° | 5.76 | 12.6° | " | 19.63 | |
| 4 | Fukuwata | Hie-no-yu * | " | 50.0° | 1.42 | 9.8° | " | 30.16 | " 9, " " |
| 5 | | Awa-no-yu * | " | 44.5° | 2.36 | 12.5° | " | 25.98 | |
| 6 | | Iwa-no-yu * | " | 48.0° | 2.77 | 10.9° | " | 26.71 | |
| 7 | Shiogama | Fudō-no-yu | " | 42.0° | — | — | — | — | — |
| 8 | | Fukuwata-uchi-yu | " | — | — | — | — | — | |
| 9 | | Hashimoto-no-yu * | " | 65.0° | — | — | — | — | |
| 10 | Shimoyu | Naka-no-yu | " | 71.5° | 0.87 | 12.5 | K. & L. | 51.00 | Nov. 9, 1913 { Y. Kinugasa
T. Saitō |
| 11 | | Iwa-no-yu | " | 69.0° | 1.11 | 12.5° | " | 50.80 | |
| 12 | | Hie-no-yu | " | 45.0° | — | — | — | 41.16 | |
| 13 | Hataori | Moto-yu | muriated alkaline | 65.0° | — | — | — | 19.74 | " " " |
| 14 | | Hato-no-yu * | " | 57.0° | 1.52 | 11.5° | K. & L. | — | |
| 15 | | Mujina-no-yu * | " | 63.5° | 1.50 | 10.4° | " | 33.35 | |
| 16 | Menzen Sumaki | Hie-no-yu | " | 70.0° | — | — | — | — | — |
| 17 | | Kawara-no-yu | " | 55.0° | — | — | — | — | |
| 18 | | Taki-no-yu | simple | 62.5° | 0.91 | 9.6° | K. & L. | 11.44 | Nov. 10, 1913 { Y. Kinugasa
T. Saitō |
| 19 | Furumachi Yumoto | Kawara-no-yu * | common salt | 54.0° | 0.79 | 12.0° | " | 21.01 | |
| 20 | | Jirakubō-no-yu * | " | 50.0° | 1.03 | 12.1° | " | 16.09 | |
| 21 | | Hana-no-yu | earth-muriated alkaline | — | — | — | — | — | Nov. 9, 1913 { Y. Kinugasa
T. Saitō |
| 22 | Furumachi Yumoto | Gosho-no-yu * | alkaline | 51.5° | 0.72 | 12.1° | K. & L. | 14.54 | |
| 23 | | Asahi-no-yu * | — | 48.3° | 1.07 | 11.6 | " | 5.33 | |
| 24 | | Takara-no-yu | — | 42.0° | 0.77 | 12.1° | " | 7.62 | " " " |
| 25 | Furumachi Yumoto | Fudō-no-yu | alkaline | 60.0° | — | — | — | — | |
| 26 | | Kajiwara-no-yu | muriated alkaline | — | — | — | — | — | — |

* These springs are for public use.

NIKKŌ-YUMOTO

Location.—Nikkō-machi, Kami-tsuga-gun, Province Shimozuke, Tochigi-ken.

The springs issue from the Quartz Porphyry.

The hot spring "Gosho-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 66° C.

Total residue: ca. 1.20.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|----------------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0249 | 0.6871 | 0.6871 |
| Sodium ion (Na ⁺) | 0.1452 | 6.2993 | 6.2993 |
| Calcium ion (Ca ⁺⁺) | 0.1535 | 3.8271 | 7.6542 |
| Magnesium ion (Mg ⁺⁺) | 0.0061 | 0.2504 | 0.5008 |
| | | | 15.1430 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.0788 | 2.2228 | 2.2228 |
| Sulphate ion (SO ₄ ^{''}) | 0.5843 | 6.0827 | 12.1654 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0452 | 0.7409 | 0.7409 |
| | 1.0400 | 20.1111 | 15.1291 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1168 | | |
| | 1.1568 | | |
| Free carbon dioxide (CO ₂) | not determined | | |
| Free hydrogen sulphide (H ₂ S) | 0.0500 | | |
| | 1.2068 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|---|----------------|
| Sodium chloride (NaCl) | 0.1300 |
| Potassium sulphate (K ₂ SO ₄) | 0.0600 |
| Sodium sulphate (Na ₂ SO ₄) | 0.2900 |
| Calcium sulphate (CaSO ₄) | 0.4700 |
| Magnesium sulphate (MgSO ₄) | 0.0300 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0600 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1168 |
| | 1.1568 |
| Free carbon dioxide (CO ₂) | not determined |
| Free hydrogen sulphide (H ₂ S) | 0.0500 |
| | 1.2068 |

The spring thus may be classified as "hydrogen sulphide spring".

Radio-activity. 1.58 Mache's units in 1 litre of water at 7.0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 12, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 12.45$.

The hot spring "Ara-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1912.

Specific gravity: 1.0010 at 15° C.

Temperature: 69° C.

Total residue: ca. 1.16.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|--|--------|---|--------|
| Sodium chloride (NaCl) | 0.1160 | Silica (SiO ₂) | 0.1653 |
| Potassium sulphate (K ₂ SO ₄) | 0.0429 | | 1.1582 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0302 | Carbon dioxide (free and in form of bicar-
bonates) (CO ₂) | 0.1749 |
| Sodium carbonate (Na ₂ CO ₃) | 0.1111 | Free hydrogen sulphide (H ₂ S) | 0.0289 |
| Calcium sulphate (CaSO ₄) | 0.6552 | | 1.3620 |
| Magnesium carbonate (MgCO ₃) | 0.0317 | | |
| Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃) | 0.0058 | | |

Besides these a very small quantity of phosphoric acid, trace of iodine and organic substances.

The spring may be classified as "hydrogen sulphide spring".

Radio-activity. 0.88 Mache's units in 1 litre of water at 9.2° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 12, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 12.25$.

The hot spring "Tsuru-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1912.

Specific gravity: 1.0013 at 15° C.

Temperature: 62° C.

Total residue: ca. 0.88.

The spring, being in its composition nearly the same as that of "Ara-yu", may be classified as "hydrogen sulphide spring".

Radio-activity. 2.08 Mache's units in 1 litre of water at 8.1° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 12, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 11.87$.

The hot spring "Kawara-no-yu"

Analysis.

Analysed by the Hygienic Laboratory of Tochigi-ken. 1912. Specific gravity: 1.0017 at 15° C.
Temperature: 64° C. Total residue: ca. 1.44.

The spring, being in its composition nearly the same as that of "Ara-yu", may be classified as "hydrogen sulphide spring".

Radio-activity. 1.73 Mache's units in 1 litre of water at 8.4° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 12, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 12.72$.

Table 36.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water
in Mache's units
at (temp. in C.) | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|----------------|----------------------|--------------------------|---|-----|--------------------|--|----------------------------------|---------------------------|
| 1 | Gosho-no-yo * | hydrogen
sulphide | 60.0 | 1.58 | 7.0 | K. & L. | 12.45 | Nov. 12, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Ara-yu | " | 60.0 | 0.88 | 9.2 | " | 12.25 | " | " |
| 3 | Tsuru-no-yu * | " | 62.0 | 2.08 | 8.1 | " | 11.87 | " | " |
| 4 | Kawara-no-yu * | " | 64.0 | 1.73 | 8.4 | " | 12.72 | " | " |
| 5 | Donsu-no-yu * | " | 42.5 | 0.72 | 7.8 | " | 11.12 | " | " |
| 6 | Naka-no-yu * | " | 49.0 | - | - | " | - | - | - |
| 7 | Uba-no-yu | " | 65.0 | - | - | " | - | - | - |
| 8 | Taki-no-yu * | " | 22.0 | - | - | " | 9.20 | Nov. 12, 1913 | { Y. Kinugasa
T. Saitō |
| 9 | Tade-no-yu | " | 56.0 | - | - | " | - | - | - |
| 10 | Sasa-no-yu * | " | 62.7 | - | - | " | - | - | - |

* These springs are for public use.

KUSATSU

Location.—Kusatsu-machi, Agatsuma-gun, Province Kōzuke, Gumma-ken.

The springs issue from the Volcanic Detritus.

The hot spring "Yubatake-no-yu"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1913.

Specific gravity: 1.0032 at 18° C.

Temperature: 58° C.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|----------------|-----------------|---------------------------|
| Hydrogen ion (H ⁺) | 0.03086 | 30,55446 | 30,55446 |
| Potassium ion (K ⁺) | 0.01703 | 0.43499 | 0.43499 |
| Sodium ion (Na ⁺) | 0.03670 | 1,59210 | 1,59210 |
| Ammonium ion (NH ₄ ⁺) | 0.00106 | 0.05863 | 0.05863 |
| Calcium ion (Ca ⁺⁺) | 0.09967 | 2,49175 | 4,98350 |
| Magnesium ion (Mg ⁺⁺) | 0.03271 | 1,34278 | 2,68556 |
| Ferrous ion (Fe ⁺⁺) | 0.11444 | 2,04357 | 4,08714 |
| Aluminium ion (Al ⁺⁺⁺) | 0.17022 | 6,28118 | 18,84354 |
| | | | 63,24001 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.64573 | 18,21523 | 18,21523 |
| Hydrosulphate ion (HSO ₄ ⁻) | 2,14506 | 22,09807 | 22,09807 |
| Sulphate ion (SO ₄ ^{''}) | 1.09275 | 11,37570 | 22,75140 |
| Hydrophosphate ion (HPO ₄ ^{''}) | 0.00860 | 0.08957 | 0.17914 |
| | 4.39483 | 96,57812 | 63,24384 |
| Boric acid (meta) (HBO ₂) | 0.01502 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.24978 | | |
| | 4.65963 | | |
| Free hydrogen sulphide (H ₂ S) | 0.00546 | | |
| | 4.66509 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|---|---------------|
| Potassium chloride (KCl) | 0.0325 |
| Sodium chloride (NaCl) | 0.0932 |
| Ammonium chloride (NH ₄ Cl) | 0.0031 |
| Calcium chloride (CaCl ₂) | 0.2763 |
| Magnesium chloride (MgCl ₂) | 0.1279 |
| Ferrous sulphate (FeSO ₄) | 0.3107 |
| Aluminium phosphate [Al ₂ (HPO ₄) ₃] | 0.0102 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 1.0651 |
| Free hydrochloric acid (HCl) | 0.3085 |
| Free sulphuric acid (H ₂ SO ₄) | 2.1674 |
| Boric acid (meta) (HBO ₂) | 0.0150 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2498 |
| | 4.6597 |
| Free hydrogen sulphide (H ₂ S) | <u>0.0055</u> |
| | 4.6652 |

The spring thus may be classified as “acid hydrogen sulphide alum vitriol spring”.

Radio-activity. 0.13 Mache's units in 1 litre of water at 14° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, April 8, 1913).

0.83 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, April 5, 1913).

The hot spring “Takino-moto-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 64.4° C.

Total residue: ca. 5.60.

The spring, being in its composition nearly the same as that of “Yubatake-no-yu”, may be classified as “acid hydrogen sulphide alum vitriol spring”.

The hot spring “Kunshi-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1888.

Specific gravity: 1.0019 at 25° C.

Total residue: ca. 2.38.

The spring, being in its composition nearly the same as that of “Yubatake-no-yu”, may be classified as “acid hydrogen sulphide alum vitriol spring”.

The hot spring “Washi-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 60° C.

Total residue: ca. 4.09.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milli-gram-equivalents |
|--|---------------|------------|------------------------|
| Hydrogen ion (H ⁺) | 0.0444 | 43.9703 | 43.9603 |
| Sodium ion (Na ⁺) | 0.0778 | 3.3753 | 3.3753 |
| Calcium ion (Ca ⁺⁺) | 0.1885 | 4.7007 | 9.4014 |
| Magnesium ion (Mg ⁺⁺) | 0.0499 | 1.9130 | 3.8260 |
| Ferrous ion (Fe ⁺⁺) | 0.0989 | 1.7602 | 3.5324 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0034 | 0.1255 | 0.3765 |
| | | | (44.779) |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.7252 | 20.4513 | 20.4513 |
| Hydrosulphate ion (HSO ₄ ⁻) | 2.2633 | 23.3162 | 23.3162 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.0871 | 10.2759 | 20.5518 |
| | 4.4352 | 109.8874 | 64.3193 |

Besides these trace of hydrophosphate ion and organic substances.

The spring thus may be classified as "acid vitriol spring".

Radio-activity. 0.13 Mache's units in 1 litre of water at 17.5° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, April 9, 1913).

1.37 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, April 6, 1913).

The hot spring "Goza-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.
Temperature: 53.9° C.

Specific gravity: 1.001 at 15° C.
Total residue: ca. 4.71.

The spring, being in its composition nearly the same as that of "Washi-no-yu", may be classified as "acid vitriol spring".

The hot spring "Saino-kawara-no-yu"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1913.

Specific gravity: 1.0028 at 19° C.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milli-gram-equivalents |
|--|----------------|------------|------------------------|
| Hydrogen ion (H ⁺) | 0.02790 | 27.62376 | 27.62376 |
| Potassium ion (K ⁺) | 0.01254 | 0.32031 | 0.32031 |
| Sodium ion (Na ⁺) | 0.03459 | 1.50065 | 1.50065 |
| Ammonium ion (NH ₄ ⁺) | 0.00106 | 0.05863 | 0.05863 |
| Calcium ion (Ca ⁺⁺) | 0.08263 | 2.06575 | 4.13150 |
| Magnesium ion (Mg ⁺⁺) | 0.02923 | 1.19991 | 2.39982 |
| Ferrous ion (Fe ⁺⁺) | 0.11169 | 1.99446 | 3.98892 |
| Aluminium ion (Al ⁺⁺⁺) | 0.15019 | 5.54207 | 16.62621 |
| | | | 56.64980 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.55722 | 15.71848 | 15.71848 |
| Hydrosulphate ion (HSO ₄ ⁻) | 1.97131 | 20.30813 | 20.30813 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.98697 | 10.27452 | 20.54904 |
| Hydrophosphate ion (HPO ₄ ⁻) | 0.00343 | 0.03573 | 0.07146 |
| | 3.01852 | 36.33684 | 36.33684 |
| Boric acid (meta) (HBO ₂) | 0.01132 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.21493 | | |
| | 4.16501 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0239 |
| Sodium chloride (NaCl) | 0.0878 |
| Ammonium chloride (NH ₄ Cl) | 0.0031 |
| Calcium chloride (CaCl ₂) | 0.2291 |
| Magnesium chloride (MgCl ₂) | 0.1143 |
| Ferrous sulphate (FeSO ₄) | 0.3033 |
| Aluminium phosphate [Al ₂ (HPO ₄) ₃] | 0.0041 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.9449 |
| Free hydrochloric acid (HCl) | 0.2664 |
| Free sulphuric acid (H ₂ SO ₄) | 1.9918 |
| Boric acid (meta) (HBO ₂) | 0.0113 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2149 |
| | 4.1650 |

Besides these trace of hydrogen sulphide.

The spring thus may be classified as "acid alum vitriol spring".

The hot spring "Jizō-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 56.5° C. (air temp. 11.2° C.)

Total residue: ca. 4.72

The spring, being in its composition nearly the same as that of "Saino-kawara-no-yu", may be classified as "acid alum vitriol spring".

Radio-activity. 0.14 Mache's units in 1 litre of water at 15° C.

0.65 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, April 5, 1913).

The hot spring "Netsu-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 56° C. (air temp. 10.1° C.).

Total residue: ca. 4.57.

The spring, being in its composition nearly the same as that of "Saino-kawara-no-yu", may be classified as "acid alum vitriol spring".

Radio-activity. 0.16 Mache's units in 1 litre of water at 17.7° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, April 8, 1913).

0.70 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, April 6, 1913).

The hot spring "Shirane-san-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1891.

Specific gravity: 1.0109 at 15° C.

Total residue: ca. 7.25.

The spring, being in its composition nearly the same as that of "Saino-kawara-no-yu", may be classified as "acid alum vitriol spring".

Composition of the sinter-deposit (analysed in 1911).

100 parts of the yellowish sinter-deposit consist of:

| Soluble in water | | Insoluble in water | |
|--|------------|--|------------|
| Total soluble matters. | 0.430 pts. | Sulphur. | 94.42 pts. |
| of which | | Silica (SO ₂) | 2.34 " |
| Free acids (as H ₂ SO ₄). | 0.098 " | Fe ₂ O ₃ , Al ₂ O ₃ , MgO, CaO | 2.01 " |

Composition of the gas (analysed in 1913).

1000 c.c. of gas* evolving from the spring consist of:

| | Shirahata-no-yu | Netsu-no-yu | Jizō-no-yu. |
|---------------------------------|-----------------|----------------|----------------|
| CO ₂ | 730 c.c. | 550 c.c. | 670 c.c. |
| O ₂ | 20 " | 75 " | not determined |
| Hydrocarbons and nitrogen . . . | not determined | not determined | " |

* The analysis is not accurate: the numbers given are only the approximate value.

Table 37.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|--------------------|---------------------|--|--------------------------|--|-------------------|--------------------|---------------------------------------|---------------|
| | | | | in
Mach's units | at
temp. in C. | | | |
| 1 | Yubatake-no-yu | acid hydrogen sul-
phide alum vitriol | 58.0° (air temp. 10.2) | 0.13 | 14.0 | K. & L. | Apr. 8, 1913 | Dr. R. Ishizu |
| 2 | Takin-no-yu | " | 64.4 | — | — | — | — | — |
| 3 | Katsun-no-yu | " | — | — | — | — | — | — |
| 4 | Washi-no-yu | acid vitriol | 60.0° | 0.13 | 17.5 | K. & L. | Apr. 9, 1913 | Dr. R. Ishizu |
| 5 | Jizō-no-yu | acid alum vitriol | 50.5 (11.2) | 0.14 | 15.0 | " | " 5, " | " |
| 6 | Gozan-no-yu | acid vitriol | 63.9 | — | — | — | — | — |
| 7 | Sakurakawa-no-yu | acid alum vitriol | 43.0 (10.7) | 0.18 | 20.0 | K. & L. | Apr. 7, 1913 | Dr. R. Ishizu |
| 8 | " (Shirahata-no-yu) | " | 49.5 (14.0) | 0.20 | 20.0 | " | " " | " |
| 9 | Netsu-no-yu | " | 56.0° (10.15) | 0.16 | 17.7 | " | " 8, " | " |
| 10 | Shirane-san-no-yu | " | — | — | — | — | — | — |
| 11 | Shirahata-no-yu | " | 58.0 (10.9) | 0.22 | 18.0 | K. & L. | Apr. 9, 1913 | Dr. R. Ishizu |
| 12 | Chiyo-no-yu | " | 56.0° | 0.17 | 18.0 | " | " 6, " | " |
| 13 | Drinking water | " | — | 0.03 | 10.0 | " | " 5, " | " |
| Gas evolving from: | | | | | | | | |
| | | | | Emanation per
litre of gas at 0° C. | | | Date when the
gas was
collected | |
| 1 | Yubatake-no-yu | — | — | 0.83 | — | K. & L. | Apr. 5, 1913 | Dr. R. Ishizu |
| 2 | Washi-no-yu | — | — | 1.37 | — | " | " 6, " | " |
| 3 | Jizō-no-yu | — | — | 0.65 | — | " | " 5, " | " |
| 4 | Netsu-no-yu | — | — | 0.70 | — | " | " 6, " | " |
| 5 | Shirahata-no-yu | — | — | 0.66 | — | " | " 9, " | " |
| 6 | Chiyo-no-yu | — | — | 0.71 | — | " | " 7, " | " |

SHIMA

Location.—Sawada-mura, Agatsuma-gun, Province Kōzuke, Gumma-ken.

The springs issue from the Misaka Series.

The hot spring "Tokiwa-no-yu"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1913.

Specific gravity: 1.0014 at 15° C.

Temperature: 62° C.

Total residue: ca. 1.47.

Flow of water: ca. 635 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

Cations.

| | Grams | Milli-grams | Centi-grams |
|--|---------|-------------|-------------|
| Potassium ion (K ⁺) . . . | 0.01735 | 0.1735 | 0.41291 |
| Sodium ion (Na ⁺) . . . | 0.00001 | 0.0001 | 12.26073 |
| Ammonium ion (NH ₄ ⁺) . . . | 0.00026 | 0.0117 | 0.1117 |
| Calcium ion (Ca ⁺⁺) . . . | 0.0147 | 1.47 | 8.23850 |
| Magnesium ion (Mg ⁺⁺) . . . | 0.00262 | 0.10755 | 0.21510 |
| Ferrous ion (Fe ⁺⁺) | 0.00140 | 0.02500 | 0.05000 |
| | | | 21.22162 |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0331 |
| Sodium chloride (NaCl) | 0.0001 |
| Ammonium chloride (NH ₄ Cl) | 0.0001 |
| Calcium chloride (CaCl ₂) | 0.0787 |
| Calcium sulphate (CaSO ₄) | 0.4539 |
| Calcium carbonate (CaCO ₃) | 0.0117 |
| Magnesium sulphate (MgSO ₄) | 0.0001 |

Anions.

| | | | | | |
|--|---------|----------|-----------------|---|--------|
| Chlorine ion (Cl^-) | 0.50127 | 14,13879 | 14.13879 | Ferrous bicarbonate [$\text{Fe}(\text{HCO}_3)_2$] | 0.0045 |
| Sulphate ion (SO_4^{--}) | 0.32048 | 3,33625 | 6.67250 | Boric acid (meta) (HBO_2) | 0.0567 |
| Hydrocarbonate ion (HCO_3^-) | 0.02503 | 0,41026 | 0,41026 | Silicic acid (meta) (H_2SiO_3) | 0.1089 |
| | 1,31577 | 34,85512 | 21,22155 | | 1,4816 |
| Boric acid (meta) (HBO_2) | 0.05667 | | | Free carbon dioxide (CO_2) | 0.0500 |
| Silicic acid (meta) (H_2SiO_3) | 0.10891 | | | | 1,5316 |
| | 1,48135 | | | | |
| Free carbon dioxide (CO_2) | 0.04995 | 1,13523 | | | |
| | 1,53130 | | | | |

The spring thus may be classified as "sulphated common salt spring".

Radio-activity. 0.82 Mache's units in 1 litre of water at 10.5° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō; Nov. 15, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 18.66$.

The hot spring "Iwane-no-yu"

Analysis.

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881. Temperature: 83.5° C.

Flow of water: ca. 526 hectolitres in 24 hours. Total residue: ca. 2,48.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|--|--------|
| Potassium chloride (KCl) | 0,1349 | Calcium sulphate (CaSO_4) | 0,5865 |
| Sodium chloride (NaCl) | 1,4891 | Silica (SiO_2) | 0,1373 |
| Sodium sulphate (Na_2SO_4) | 0,1137 | | 2,4825 |
| Magnesium chloride (MgCl_2) | 0,0210 | | |

Besides these trace of iron.

The spring may be classified as "common salt spring".

Radio-activity. 1.27 Mache's units in 1 litre of water at 10.4° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 15, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 30.66$.

Table 38.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|--------------------------------|----------------------------|--------------------------|---|---------------------------------|-------------------|--------------------|--|----------------------------------|---------------------------|
| | | | | | in
Mache's units | at
temp. in C. | | | | |
| 1 | Tokiwa no-yu | sulphated com-
mon salt | 62.0° | ca. 635 | 0.82 | 10.5° | K. & L. | 18.66 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Shinkoku tō | salt | 55.0° | 311 | 0.87 | 10.5° | " | 19.74 | " | " |
| 3 | Tsubame no-yu | " | 50.0° | 311 | — | — | — | — | — | — |
| 4 | Otake no-yu | " | 54.0° | 234 | — | — | — | — | — | — |
| 5 | Shio no-yu | " | 58.5° | 94 | — | — | — | 20.00 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 6 | Meiji no-yu | " | 67.0° | 208 | — | — | — | — | — | — |
| 7 | Shio no-yu <i>Seizen-kyōan</i> | " | 74.0° | 524 | 0.82 | 11.7° | K. & L. | 28.47 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 8 | Furo sen | " | 74.0° | 389 | — | — | — | — | — | — |
| 9 | Iwane no-yu | common salt | 83.5 | 520 | 1.27 | 10.4 | K. & L. | 30.66 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 10 | Shio no-yu | salt | 77.0 | 138 | — | — | — | — | — | — |
| 11 | Ryugū no-yu | " | 84.0 | 635 | 0.64 | 10.9° | K. & L. | 34.05 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 12 | Kawara no-yu | " | 60.0 | 285 | — | — | — | — | — | — |
| 13 | Hinotani no-yu | " | 59.0 | 247.3 | 0.43 | 9.4 | K. & L. | 10.80 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 14 | Shimo no-yu | " | 55.0° | 156 | — | — | — | 10.22 | — | " |

Corrigendum to P. 53 :—

This table of analysis is to be exchanged with the corresponding table which is misprinted.

In 1 kilogram of the mineral water are contained:

Cations.

| | Grams | Milli-
moles | Milligram-
equivalents |
|--|---------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 1,43430 | 36,63600 | 36,63600 |
| Sodium ion (Na ⁺) | 9,82587 | 426,28503 | 426,28503 |
| Ammonium ion (NH ₄ ⁺) | 0,00254 | 0,14049 | 0,14049 |
| Calcium ion (Ca ⁺⁺) | 0,26901 | 6,72525 | 13,45050 |
| Magnesium ion (Mg ⁺⁺) | 0,05976 | 2,45320 | 4,90640 |
| Ferrous ion (Fe ⁺⁺) | 0,00410 | 0,07321 | 0,14642 |
| | | | <u>481,56484</u> |

Anions.

| | | | |
|--|----------------|-----------|------------------|
| Chlorine ion (Cl ⁻) | 13,07537 | 368,83970 | 368,83970 |
| Bromine ion (Br ⁻) | 0,01133 | 0,14170 | 0,14170 |
| Iodine ion (I ⁻) | 0,00200 | 0,01577 | 0,01577 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 6,86973 | 112,60067 | 112,60067 |
| | 31,55401 | 953,91102 | 481,59784 |
| Boric acid (meta) (HBO ₂) | 0,99695 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,02279 | | |
| | 32,57375 | | |
| Free carbon dioxide (CO ₂) | 0,96555 | 15,82609 | |
| | 33,53930 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|-----------------|
| Potassium chloride (KCl) | 2,73304 |
| Sodium chloride (NaCl) | 19,42517 |
| Sodium bromide (NaBr) | 0,01460 |
| Sodium iodide (NaI) | 0,00236 |
| Ammonium chloride (NH ₄ Cl) | 0,00752 |
| Sodium bicarbonate (NaHCO ₃) | 7,90974 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 1,08955 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,35913 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,01303 |
| Boric acid (meta) (HBO ₂) | 0,99695 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,02279 |
| | 32,57380 |
| Free carbon dioxide (CO ₂) | 0,96555 |
| | <u>33,53943</u> |

Anions.

| | | | | | |
|--|---------|----------|-----------------|---|--------|
| Chlorine ion (Cl') | 0,50122 | 14,13879 | 14,13879 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0045 |
| Sulphate ion (SO ₄ '') | 0,32048 | 3,33625 | 6,67250 | Boric acid (meta) (HBO ₂) | 0,0567 |
| Hydrocarbonate ion (HCO ₃ ') | 0,02503 | 0,41026 | 0,41026 | Silicic acid (meta) (H ₂ SiO ₃) | 0,1089 |
| | 1,31577 | 34,85512 | 21,22155 | | 1,4816 |
| Boric acid (meta) (HBO ₂) | 0,05667 | | | Free carbon dioxide (CO ₂) | 0,0500 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,10891 | | | | 1,5316 |
| | 1,48135 | | | | |
| Free carbon dioxide (CO ₂) | 0,04995 | 1,13523 | | | |
| | 1,53130 | | | | |

The spring thus may be classified as "sulphated common salt spring".

Radio-activity. 0,82 Mache's units in 1 litre of water at 10,5° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō; Nov. 15, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 18,66$.

| | | | Temp | Flow
in
lit | i
Mache' | a
(temp | Apparat | E
conc
at | Date
water | Exam |
|----|----------------------------------|----------------|-------|-------------------|-------------|------------|---------|-----------------|---------------|---------------------------|
| 1 | Tokiwa-no-yu | sulphated com- | 62,0° | ca.
635 | 0,82 | 10,5° | K. & L. | 18,66 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 2 | Shinkoku-tō | salt | 55,0° | 311 | 0,87 | 10,5° | " | 19,74 | " | " |
| 3 | Tsubame-no-yu | " | 59,0° | 311 | — | — | — | — | — | — |
| 4 | Otake-no-yu | " | 54,0° | 234 | — | — | — | — | — | — |
| 5 | Shio-no-yu | " | 58,5° | 94 | — | — | — | 20,00 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 6 | Meiji-no-yu | " | 67,0° | 208 | — | — | — | — | — | — |
| 7 | Shio-no-yu, <i>Sekisen-kawan</i> | " | 74,0° | 524 | 0,82 | 11,7 | K. & L. | 28,47 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 8 | Furō-sen | " | 74,0° | 380 | — | — | — | — | — | — |
| 9 | Iwane-no-yu | common salt | 83,5° | 526 | 1,27 | 10,4° | K. & L. | 30,00 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 10 | Shio-no-yu | salt | 77,0° | 138 | — | — | — | — | — | — |
| 11 | Ryūgo-no-yu | " | 84,0° | 635 | 0,64 | 10,9° | K. & L. | 34,05 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 12 | Kawara-no-yu | " | 60,0° | 285 | — | — | — | — | — | — |
| 13 | Hinatani-no-yu | " | 59,0° | 2473 | 0,43 | 9,4° | K. & L. | 10,80 | Nov. 15, 1913 | { Y. Kinugasa
T. Saitō |
| 14 | Shimo-no-yu | " | 55,0° | 150 | — | — | — | 10,22 | — | " |

ISOBE

Location.—Isobe-mura, Usui-gun, Province Kōzuke, Gumma-ken.

The springs issue from the Tertiary.

The cold spring “Isobe-Kōsen”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0249 at 10° C.

Temperature: 15.5° C. (air temp. 3° C.).

Total residue: ca. 26.91.

Flow of water: ca. 720 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|----------------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.14343 | 3.66360 | 3.66360 |
| Sodium ion (Na ⁺) | 9.82587 | 426.28503 | 426.28503 |
| Ammonium ion (NH ₄ ⁺) | 0.00254 | 0.14049 | 0.14049 |
| Calcium ion (Ca ⁺⁺) | 0.26901 | 6.72525 | 13.45050 |
| Magnesium ion (Mg ⁺⁺) | 0.05976 | 2.45320 | 4.90640 |
| Ferrous ion (Fe ⁺⁺) | 0.00410 | 0.07321 | 0.14642 |
| | | | 448.59244 |
| Anions. | Grams | Milli-mols | Milligram-equivalents |
| Chlorine ion (Cl ⁻) | 13.07537 | 368.83970 | 368.83970 |
| Bromine ion (Br ⁻) | 0.01133 | 0.14170 | 0.14170 |
| Iodine ion (I ⁻) | 0.00200 | 0.01577 | 0.01577 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 4.85816 | 79.62891 | 79.62891 |
| | 28.25157 | 887.96686 | 448.62608 |
| Boric acid (meta) (HBO ₂) | 1.40515 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.02279 | | |
| | 29.67951 | | |
| Free carbon dioxide (CO ₂) | 2.41581 | 54.90477 | |
| | 32.09532 | | |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.2735 |
| Sodium chloride (NaCl) | 21.3537 |
| Sodium bromide (NaBr) | 0.0146 |
| Sodium iodide (NaI) | 0.0024 |
| Ammonium chloride (NH ₄ Cl) | 0.0075 |
| Sodium bicarbonate (NaHCO ₃) | 3.7293 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 1.0896 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.3591 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0130 |
| Boric acid (meta) (HBO ₂) | 0.9970 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0228 |
| | 27.8625 |
| Free carbon dioxide (CO ₂) | 2.4158 |
| | 03.2783 |

Besides these trace of nitrate and sulphate ion.

The spring thus may be classified as “alkaline carbon dioxide concentrated common salt spring”.

Radio-activity. 0.47×10^{-10} curies in 1 litre of water at 15.5° C.

0.13 Mache's units (recalculated). (Schmidt's electrom.; S. Hanzawa, Jan. 30, 1914).

1.41×10^{-10} curies in 1 litre of gas at 0° C.

0.39 Mache's units (recalculated). (Schmidt's electrom.; S. Hanzawa, Jan. 31, 1914).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 245.12$.

The cold spring “Ji-no-yu”

Temperature: 9° C. (air temp. 2° C.).

Radio-activity. 3.78×10^{-10} curies in 1 litre of water at 9° C.

1.04 Mache's units (recalculated).

7.42×10^{-10} curies in 1 litre of gas at 0° C.

2.04 Mache's units (recalculated).

(Schmidt's electrom.; S. Hanzawa, Feb. 1, 1914).

Electric conductivity. $\kappa_{18} \times 10^{-4} = 34.83$.

Table 39.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|------------------------------------|---|--------------------------|---|--|-------------------------|---------------------|--------------------|---------------------------------------|--------------------------|
| | | | | | in
10 ¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Isobe-Kōsen | alkaline car-
bonated
concentrated
common salt | 15.5°
(30°) | ca.
7.20 | 0.47 | 0.13 | 15.5° | Schm. | Jan. 30, 1914 | S. Hanzawa |
| 2 | Ji-no-yu | | 9.0
(20) | | 3.78 | 1.04 | 9.0° | " | Feb. 1, " | " |
| 3 | Rokutanda-Kōsen | alkaline | 9.5°
(9.50) | 7 | — | 0.64*
0.74†
0.75‡ | 9.0° | E. & S. | Jan. 30, " | { S. Hanzawa
T. Saitō |
| 4 | Ushio-Kōsen | — | 8.0°
(10) | — | — | 0.91*
1.05†
1.07‡ | 8.5° | " | " | " |
| 5 | Yunosawa-Kōsen
(Kōngōji temple) | — | 7.5°
(80) | — | — | 0.82*
0.95†
0.97‡ | 10.0° | " | " | " |
| 1 | Gas evolving from
Isobe-Kōsen | — | — | — | Emanation per litre
of gas at 0° C. | | | Schm. | Date when the
gas was
collected | |
| 2 | Ji-no-yu | — | — | — | 1.41 | 0.39 | — | " | Jan. 31, 1914 | " |
| | | | | | 7.42 | 2.04 | — | " | Feb. 1, " | " |

* Determined according to the authors' original direction.
† Correction for the absorption of the radiation by the wall of the ionisation chamber was made.
‡ Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made.
§ These springs are located in *Matsuida*, 4.1 m. W. from Isobe.

IKAO

Location.—Ikao-machi, Gumma-gun, Province Kōzuke Gumma-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring “Nomi-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1903. Specific gravity: 1,0008 at 15° C.

Temperature: 46° C.

Total residue: ca. 0.96.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents | Grams |
|---|--------|----------------|---------------------------|--|
| Potassium ion (K ⁺) | 0.0069 | 0.2529 | 0.2529 | Sodium chloride (NaCl) 0.0468 |
| Sodium ion (Na ⁺) | 0.0892 | 3.8699 | 3.8699 | Potassium sulphate (K ₂ SO ₄) 0.0220 |
| Calcium ion (Ca ⁺⁺) | 0.1224 | 3.0549 | 6.1098 | Sodium sulphate (Na ₂ SO ₄) 0.1007 |
| Magnesium ion (Mg ⁺⁺) | 0.0274 | 1.1248 | 2.2496 | Sodium bicarbonate (NaHCO ₃) 0.1393 |
| Ferrous iron (Fe ⁺⁺) | 0.0076 | 0.1342 | 0.2684 | Magnesium chloride (MgCl ₂) 0.1036 |
| Manganous ion (Mn ⁺⁺) | 0.0017 | 0.0310 | 0.0620 | Calcium sulphate (CaSO ₄) 0.2769 |
| | | | 12.8127 | Calcium bicarbonate [Ca(HCO ₃) ₂] 0.1646 |

Anions.

| | | | |
|--|--------|--------|---------------|
| Chlorine ion (Cl^-) | 0.1955 | 2.776 | 2.776 |
| Sulphate ion (SO_4^{2-}) | 0.2754 | 2.877 | 5.7340 |
| Hydrocarbonate ion (HCO_3^-) | 0.2498 | 4.044 | 4.044 |
| | 0.8889 | 18.451 | 12.877 |
| Silicic acid (meta) (H_2SiO_3) | 0.2068 | | |
| | 1.0057 | | |
| Free carbon dioxide (CO_2) | 0.6757 | 15.350 | |
| | 1.7714 | | |

| | |
|---|--------|
| Magnesium bicarbonate [$\text{Mg}(\text{HCO}_3)_2$] | 0.0053 |
| Ferrous bicarbonate [$\text{Fe}(\text{HCO}_3)_2$] | 0.0242 |
| Manganous bicarbonate [$\text{Mn}(\text{HCO}_3)_2$] | 0.0055 |
| Silicic acid (ortho) (H_2SiO_4) | 0.2068 |
| | 1.0057 |
| Free carbon dioxide (CO_2) | 0.6757 |
| | 1.7714 |

Besides these trace of aluminium, bromine, iodine, hydrophosphate ion and organic substances.

The spring thus may be classified as “**sulphated bitter spring**”.

Radio-activity. **0.67** Mache's units in 1 litre of water at 10.1°C .

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 14, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = \mathbf{10.28}$.

The hot spring “Ohaguro-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1903.

Specific gravity: 1.001 at 15°C .

Temperature: 47°C .

Total residue: ca. 1.34.

The spring, being in its composition nearly the same as that of “Nomi-yu”, may be classified as “**sulphated bitter spring**”.

Radio-activity. **0.47** Mache's units in 1 litre of water at 11.0°C .

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 14, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = \mathbf{13.41}$.

The hot spring “Ōseki-no-yu *”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1903.

Specific gravity: 1.0008 at 15°C .

Temperature: 44.5°C .

Total residue: ca. 0.98.

Flow of water: ca. 25400 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of “Nomi-yu”, may be classified as “**sulphated bitter spring**”.

Radio-activity. **0.37** Mache's units in 1 litre of water at 10.1°C .

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Nov. 14, 1913).

Electric conductivity. $\kappa_{18} \times 10^{-4} = \mathbf{10.36}$.

Table 40.

| No. | Spring | Classification | Temp. of spring
in $^\circ \text{C}$ | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Electric
conductivity
$\kappa_{18} \times 10^{-4}$ | Date when the
water was taken | Examined by |
|-----|----------------|------------------|---|---|---------------------------------|----------------------------------|--------------------|--|----------------------------------|-----------------------------|
| | | | | | in
Mache's units | at
temp. in $^\circ \text{C}$ | | | | |
| 1 | Nomi-yu | sulphated bitter | 46.0° | — | 0.67 | 10.1 | K & I | 10.28 | Nov. 14, 1913 | (Y. Kinugasa
& T. Saitō) |
| 2 | Ohaguro-yu | .. | 47.0 | — | 0.47 | 11.0 | .. | 13.41 | .. | .. |
| 3 | Ōseki-no-yu | .. | 44.5 | 25400 | 0.37 | 10.1 | .. | 10.36 | .. | .. |
| 4 | Fukiage-no-yu | .. | 45.0 | — | 0.51 | 9.9° | .. | 10.53 | .. | .. |
| 5 | Kurozeki-no-yu | .. | 46.0° | — | — | — | .. | 11.03 | .. | .. |

* This is the name of hot water in a reservoir into which all the springs of Ika flow together.

YUJIKU

Location — Niharu-mura, Tone-gun, Province Kōzuke, Gumma-ken.

The springs issue from the Tertiary Tuff.

The hot spring “Sen-tō”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914. Specific gravity: 1.0007 at 19° C.
Temperature: 67° C. Total residue: ca. 1.42.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.01225 | 0.31289 | 0.31289 |
| Sodium ion (Na ⁺) | 0.28531 | 12.37786 | 12.37786 |
| Ammonium ion (NH ₄ ⁺) | 0.00042 | 0.02323 | 0.02323 |
| Calcium ion (Ca ⁺⁺) | 0.16417 | 4.10425 | 8.20850 |
| Magnesium ion (Mg ⁺⁺) | 0.00044 | 0.01806 | 0.03612 |
| Ferrous ion (Fe ⁺⁺) | 0.00035 | 0.00625 | 0.01250 |
| | | | 20.97110 |

| Anions. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Chlorine ion (Cl ⁻) | 0.13095 | 3.69395 | 3.69395 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.77710 | 8.08974 | 16.17948 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0.06697 | 1.00768 | 1.00768 |
| | 1.43796 | 29.72391 | 20.97111 |
| Boric acid (meta) (HBO ₂) | 0.00710 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.06098 | | |
| | 1.50604 | | |

Free carbon dioxide (CO₂) . . . small quantity

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.02334 |
| Sodium chloride (NaCl) | 0.19644 |
| Ammonium chloride (NH ₄ Cl) | 0.00124 |
| Sodium sulphate (Na ₂ SO ₄) | 0.64110 |
| Calcium sulphate (CaSO ₄) | 0.48712 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.08489 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.00264 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.00111 |
| Boric acid (meta) (HBO ₂) | 0.00710 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.06098 |
| | 1.50596 |

The spring thus may be classified as “saline bitter spring”.

Radio-activity. 2.30×10^{-10} curies in 1 litre of water at 18.0° C.

(Schmidt's electrom.; M. Komori, April 29, 1914).

0.63 Mache's units (recalculated).

33.36×10^{-10} curies in 1 litre of gas at 0° C.

9.18 Mache's units (recalculated).

(Schmidt's electrom.; M. Komori, April 30, 1914).

Table 41.

| No. | Spring | Classification | Temp. of spring
in °C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------------|----------------|---------------------------|---|--|---------------------|--------------------|--------------------|--|-------------|
| | | | | | in
10 ¹⁰ curies | in
Mache's units | at
temp. in °C. | | | |
| 1 | Sen-tō | saline bitter | air temp.
67° | ca. — | 2.30 | 0.63 | 18.0° | Schm | Apr. 29, 1914 | M. Komori |
| 2 | Kubo-yu | „ | 79° (29°) | 648 | 2.56 | 0.70 | 12.0° | „ | „ 30, „ | „ |
| 3 | Kawaburi-yu | „ | 37.2° | — | 1.50 | 0.41 | 13.5° | „ | May 2, „ | „ |
| 4 | Ko-daki | „ | 47° | 90 | 2.90 | 0.80 | 15.5° | „ | „ 3, „ | „ |
| 5 | Ō-daki | „ | 50° | 360 | 3.11 | 0.86 | 14.8° | „ | „ 4, „ | „ |
| | Gas evolving from
Sen-tō | | | | Emanation per litre
of gas at 0° C.
33.36 9.18 | | | „ | Date when the
gas was
collected
Apr. 30, 1914 | „ |

YUBISO—YUBARA

Location: Minakamigata, Tonegun, Province Kōzuke, Gamma-ken.

The springs issue from the Granite.

The hot spring "Yubiso-no-yu" No. 2

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0001 at 16° C.

Temperature: 88° C. (air temp. 26° C.).

Total residue: ca. 0.5.

Flow of water: ca. 974 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

Cations.

| | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K') | 0.00451 | 0.11520 | 0.11520 |
| Sodium ion (Na') | 0.09000 | 3.90455 | 3.90455 |
| Ammonium ion (NH ₄ ') | 0.00026 | 0.01438 | 0.01438 |
| Calcium ion (Ca'') | 0.05214 | 1.30250 | 2.60500 |
| Magnesium ion (Mg'') | 0.00022 | 0.00903 | 0.01806 |
| Ferrous ion (Fe'') | 0.00070 | 0.01250 | 0.02500 |
| | | | 6.68219 |

Anions.

| | | | |
|--|---------|----------|---------|
| Chlorine ion (Cl') | 0.14650 | 4.13258 | 4.13258 |
| Sulphate ion (SO ₄ '') | 0.06295 | 0.65532 | 1.31064 |
| Hydrocarbonate ion (HCO ₃ ') | 0.07568 | 1.24045 | 1.24045 |
| | 0.43296 | 11.38651 | 6.68367 |
| Boric acid (meta) (HBO ₂) | 0.02838 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.08829 | | |
| | 0.54963 | | |
| Free carbon dioxide (CO ₂) | 0.04542 | 1.03227 | |
| | 0.59505 | | |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.00859 |
| Sodium chloride (NaCl) | 0.22843 |
| Ammonium chloride (NH ₄ Cl) | 0.00077 |
| Calcium chloride (CaCl ₂) | 0.00546 |
| Calcium sulphate (CaSO ₄) | 0.08016 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.09700 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.00132 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.00223 |
| Boric acid (meta) (HBO ₂) | 0.02838 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.08829 |
| | 0.54963 |
| Free carbon dioxide (CO ₂) | 0.04542 |
| | 0.59505 |

The spring thus may be classified as "simple thermal".

Radio-activity. 1.97 Mache's units in 1 litre of water at 19.5° C.

(Kohl-Löw. fontact.; M. Komori and K. Matsubara, April 30, 1914)

Table 42.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water in Mache's units at temp. in C. | Apparatus employed | Date when the water was taken | Examined by |
|-----|----------------------|----------------|-----------------------|---|--|--------------------|-------------------------------|---------------------------|
| 1 | Yubiso | | | | | | | |
| | Yubiso-no-yu (No. 1) | simple | 81.0 (26) | ca. 97.2 | 1.32 19.0 | K. & L. | Apr. 30, 1913 | M. Komori
K. Matsubara |
| 2 | " (No. 2) | " | 88.0 (26) | | 1.97 19.5 | " | " | " |
| | Yubara | | | | | | | |
| 3 | Fujiya-no-yu | " | 57.0° (21.5°) | | 1.32 17.5 | " | " | " |
| 4 | Ubukata-no-yu | " | 51.5 | | 0.89 15.5 | " | " | " |

IRINOYU

Location — Sakamoto-machi, Usui-gun, Province Kōzuke, Gumma-ken.

The springs issue from the Pyroxene Andesite.

The hot spring "Iri-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1890. Specific gravity: 1.0011 at 15° C.
Temperature: 37° C. (air temp. 15° C.). Total residue: ca. 1.90.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0060 | 0.1533 | 0.1533 |
| Sodium ion (Na ⁺) | 0.6250 | 27.1150 | 27.1150 |
| Calcium ion (Ca ⁺⁺) | 0.0537 | 1.3392 | 2.6784 |
| Magnesium ion (Mg ⁺⁺) | 0.0094 | 0.3859 | 0.7718 |
| | | | 30.7185 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.9907 | 27.9464 | 27.9464 |
| Sulphate ion (SO ₄ ^{''}) | 0.1333 | 1.3877 | 2.7754 |
| | 1.8181 | 58.3275 | 30.7218 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0325 | | |
| | 1.8506 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|--|--------|
| Potassium chloride (KCl) | 0.0039 |
| Sodium chloride (NaCl) | 1.5864 |
| Potassium sulphate (K ₂ SO ₄) | 0.0089 |
| Magnesium chloride (MgCl ₂) | 0.0368 |
| Calcium sulphate (CaSO ₄) | 0.1821 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0325 |
| | 1.8506 |

Besides these trace of hydrophosphate ion.

The spring thus may be classified as "common salt spring".

Radio-activity. 0.77[†] Mache's units in 1 litre of water at 4.8° C.

(Eng.-Siev. fontact.; S. Hanzawa and T. Saitō, Feb. 1, 1914).

Table 43.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water
in
Mache's units
at
(temp. in C.) | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-------------|----------------|--------------------------|---|--------------------|----------------------------------|--------------------------|
| 1 | Iri-no-yu | common salt | 37.0° (air temp. 15°) | { 0.65 *
0.75 † 4.8°
0.77 ‡ | E. & S. | Feb. 1, 1914 | { S. Hanzawa
T. Saitō |
| 2 | Me-no-yu | " | 33.5° (15°) | { 0.61 *
0.71 † 4.1°
0.72 ‡ | " | " | " |
| 3 | Kaede-no-yu | " | 31.5° (15°) | { 0.63 *
0.73 † 4.1°
0.74 ‡ | " | " | " |

* Determined according to the authors' original direction.
† Correction for the absorption of the radiation by the wall of the ionisation chamber was made.
‡ Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made.

AKAGI-NASHIKI

Location.—Kurohone-mura, Seta-gun, Province Kōzuke, Gumma-ken.

The springs issue from the Volcanic Detritus.

The cold spring “Nashiki-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1908.

Specific gravity: 1.0022 at 15° C.

Temperature: 20° C.

Total residue: ca. 2.25.

Flow of water: ca. 195 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-grams | Milligram-equivalents |
|--|--------|-------------|-----------------------|
| Potassium ion (K ⁺) | 0.0525 | 1.3410 | 1.3410 |
| Sodium ion (Na ⁺) | 42.55 | 18,4599 | 18.4599 |
| Calcium ion (Ca ⁺⁺) | 0.1280 | 3.1920 | 6.3840 |
| Magnesium ion (Mg ⁺⁺) | 0.1392 | 5.7143 | 11.4286 |
| | | | 37.6135 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.9241 | 26.0677 | 26.0677 |
| Sulphate ion (SO ₄ ^{''}) | 0.0211 | 0.2197 | 0.4394 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.6801 | 11.1474 | 11.1474 |
| | 2.3705 | 66.1420 | 37.6545 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1298 | | |
| | 2.5003 | | |
| Free carbon dioxide (CO ₂) | 0.7901 | 17.9568 | |
| | 3.2904 | | |

| | |
|---|--------|
| Potassium chloride (KCl) | 0.1000 |
| Sodium chloride (NaCl) | 1.0000 |
| Calcium chloride (CaCl ₂) | 0.3300 |
| Magnesium chloride (MgCl ₂) | 0.0150 |
| Calcium sulphate (CaSO ₄) | 0.0300 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.8155 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1298 |
| | 2.5003 |
| Free carbon dioxide (CO ₂) | 0.7901 |
| | 3.2904 |

Besides these trace of ferrous and aluminium ion.

The spring thus may be classified as “earthy common salt spring”.

TOMOE

Location.—Tomoe-mura, Kashima-gun, Province Hitachi, Ibaraki-ken.

The springs issue from the Diluvium.

The cold spring “Tomoe-Kōsen” No. 1

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1911.

Specific gravity: 1.0012 at 21° C.

Temperature: 16.9° C.

Total residue: ca. 3.34.

Flow of water: ca. 270 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-grams | Milligram-equivalents |
|--|---------|-------------|-----------------------|
| Potassium ion (K ⁺) | 0.03707 | 0.94687 | 0.94687 |
| Sodium ion (Na ⁺) | 1.00735 | 46.26161 | 46.26161 |
| Ammonium ion (NH ₄ ⁺) | 0.00159 | 0.08794 | 0.08794 |
| Calcium ion (Ca ⁺⁺) | 0.03787 | 0.94650 | 1.89300 |
| Magnesium ion (Mg ⁺⁺) | 0.06191 | 2.54146 | 5.08292 |
| Ferrous ion (Fe ⁺⁺) | 0.00224 | 0.04000 | 0.08000 |
| | | | 54.35234 |

| | |
|---|---------|
| Potassium chloride (KCl) | 0.07064 |
| Sodium chloride (NaCl) | 2.66607 |
| Ammonium chloride (NH ₄ Cl) | 0.0471 |
| Sodium sulphate (Na ₂ SO ₄) | 0.04903 |
| Calcium sulphate (CaSO ₄) | 0.12878 |
| Magnesium sulphate (MgSO ₄) | 1.0003 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.24981 |

Anions.

| | | | |
|--|---------|-----------|----------|
| Chlorine ion (Cl') | 1,65233 | 46,61016 | 46,61016 |
| Sulphate ion (SO ₄ '') | 0,20408 | 2,12451 | 4,24902 |
| Hydrocarbonate ion (HCO ₃ '') | 0,21312 | 3,49320 | 3,49200 |
| | 3,27653 | 103,05225 | 54,35238 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,06544 | | |
| | 3,34197 | | |
| Free carbon dioxide (CO ₂) | 0,02135 | 0,48523 | |
| | 3,36332 | | |

| | |
|---|---------|
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,00712 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,06544 |
| | 3,34193 |
| Free carbon dioxide (CO ₂) | 0,02135 |
| | 3,36328 |

Besides these a small quantity of boric acid.

The spring thus may be classified as "common salt spring".

Radio-activity. 4.72×10^{-10} curies in 1 litre of water at 19.8° C.

(Schmidt's electrom.; M. Komori, April 26, 1914).

1.30 Mache's units (recalculated).

TSUKUBA

Location.—Ishioka-machi, Niiharu-gun, Province Hitachi, Ibaraki-ken.

The springs issue from the Diluvium.

Table 44.

| No. | Spring | Classification | Temp. of spring in °C. | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|---------------------------|----------------|------------------------|------------------------------|------------------|-----------------|--------------------|-------------------------------|----------------------------------|
| | | | | in 10 ¹⁰ curies | in Mache's units | at temp. in °C. | | | |
| 1 | Well K. H. (H. H. H. H.) | — | 10.0° | ca. | 1.10 | 13.0° | K. & L. | Jan. 25, 1914 | { Dr. K. Ishizu
H. Yanagisawa |
| 2 | " F. H. (H. H. H. H.) | — | 17.0° | — | 0.49 | 13.0° | " | " | " |
| 3 | " (O. Nakazawa) | — | 14.0° | — | 0.55 | 13.5° | " | " | " |
| 4 | Spring A* (K. H. H. H.) | simple | 14.0° | — | 0.28 | 13.0° | " | " | " |
| 5 | Shibui-do (O. H. H. H.) | — | 10.0° | — | 0.95 | 12.0° | " | " 26, " | " |
| 6 | Tomoe-Kōsen (M. H. H. H.) | common salt | 16.9° | 4.72 | 1.30 | 19.8° | Schm. | Apr. 20, " | M. Komori |
| 7 | " No. 2 | " | 17.0° | 4.81 | 1.32 | 18.6° | " | " 25, " | " |

* The spring issues from the Granite.

† The spring is located at *Fukutsutsura*, 3 m. S. E. from *Ishioka*.

‡ Tomoe-Kōsen, located in *Fukutsutsura*, Kashima-gun, is grouped for convenience in this table.

MORIGASAKI

Location.—Omori-machi, Ebara-gun, Province Musashi, Tōkyō-fu.

The springs issue from the Alluvium.

The cold spring "Morigasaki-Kōsen"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1900.

Temperature: 17° C. (air temp. 18° C.).

Total residue: ca. 6.08.

Flow of water: ca. 3370 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|---------------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0,0866 | 2,2119 | 2,2119 |
| Sodium ion (Na ⁺) | 1,5071 | 65,3882 | 65,3882 |
| Calcium ion (Ca ⁺⁺) | 0,3245 | 8,0923 | 16,1846 |
| Magnesium ion (Mg ⁺⁺) | 0,2780 | 11,4121 | 22,8242 |
| Ferrous ion (Fe ⁺⁺) | 0,0062 | 0,1109 | 0,2218 |
| | | | 100,8307 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 3,6589 | 103,2147 | 103,2147 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0,2206 | 3,7152 | 3,6158 |
| | 6,0819 | 104,0459 | 106,8305 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0057 | | |
| | 6,1475 | | |
| Free carbon dioxide (CO ₂) | 0,3950 | 8,9909 | |
| | 6,5431 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|---|---------------|
| Potassium chloride (KCl) | 0,1650 |
| Sodium chloride (NaCl) | 3,8250 |
| Calcium chloride (CaCl ₂) | 0,8989 |
| Magnesium chloride (MgCl ₂) | 0,9250 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,2482 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0198 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0656 |
| | 6,1475 |
| Free carbon dioxide (CO ₂) | 0,3956 |
| | 6,5431 |

Besides these trace of sulphate and hydrophosphate ion

The spring thus may be classified as “**earth-muriated common salt spring**”.

Radio-activity. 0,40 Mache's units in 1 litre of water at 15,8° C.

(Kohl.-Löw. contact.; Dr R. Ishizu, Oct. 24, 1913).

ANAMORI

Location.—Haneda-machi, Ebara-gun, Province Musashi, Tōkyō-fu.

The springs issue from the Alluvium.

The cold spring “Anamori-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1896.

Temperature: 17° C.

Total residue: ca. 9,38.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|---------------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0,0897 | 2,2912 | 2,2912 |
| Sodium ion (Na ⁺) | 2,9468 | 127,8438 | 127,8438 |
| Calcium ion (Ca ⁺⁺) | 0,7183 | 4,6708 | 0,5416 |
| Magnesium ion (Mg ⁺⁺) | 0,2191 | 8,9942 | 17,9884 |
| Ferrous ion (Fe ⁺⁺) | 0,0140 | 0,2544 | 0,5088 |
| Aluminium ion (Al ⁺⁺⁺) | 0,0109 | 0,4002 | 1,2006 |
| | | | 159,1744 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 5,2424 | 147,8843 | 147,8843 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0,6905 | 11,3177 | 11,3177 |
| | 6,4177 | 169,2020 | 159,2022 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0071 | | |
| | 6,4708 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|---|---------------|
| Potassium chloride (KCl) | 0,1708 |
| Sodium chloride (NaCl) | 7,4792 |
| Calcium chloride (CaCl ₂) | 0,5188 |
| Magnesium chloride (MgCl ₂) | 0,3424 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,7914 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0445 |
| Aluminium chloride (AlCl ₃) | 0,0536 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0701 |
| | 9,4708 |

Besides these trace of hydrophosphate ion and boric acid.

The spring thus may be classified as “**iron carbonate common salt spring**”.

Radio-activity. $1,62 \times 10^{-10}$ curies in 1 litre of water at 21° C.

(Schmidt's electrom.; H. Kakehi, June 1, 1914).

0,45 Mache's units (recalculated).

Radio-activity of mineral springs and well waters in Tōkyō,
Yokohama and their neighbourhood.

Table 45.

| No. | Location | Spring or well | Classification | Temp. of spring
or well
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-------------------------------|---|-------------------------------|-------------------------------------|--|---------------------|---------------------|--------------------|--|---------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Maché's units | at
(temp. in C.) | | | |
| 1 | Tōkyō (<i>Honjō-ku</i>) | Well (<i>K. Fujii</i>) | — | air temp. | 1.10 | 0.30 | 21.0° | Schm. | July 16, 1914 | M. Komori |
| 2 | „ (<i>Ushigome-ku</i>) | „ (<i>R. Shibata</i>) | — | — | 2.00 | 0.55 | 19.0° | „ | May 8, „ | H. Kakehi |
| 3 | „ (<i>Shiba-ku</i>) | „ (<i>H. Eizawa</i>) | — | 16.0° | 1.81 | 0.50 | 15.5° | „ | Apr. 9, „ | M. Komori |
| 4 | „ (<i>Azabu-ku</i>) | „ (<i>S. Okabayashi</i>) | — | — | — | 0.78 | 18.5° | K. & L. | Dec. 19, 1913 | „ |
| 5 | „ „ | „ (<i>K. Kobayashi</i>) | — | — | — | 0.75 | 14.0° | „ | Nov. 25, „ | I. Wakita |
| 6 | Shibuya | „ (<i>R. Hata</i>) | — | — | 1.27 | 0.35 | 15.4° | Schm. | March 3, 1914 | H. Kakehi |
| 7 | Shinagawa | „ (<i>Y. Sakurai</i>) | — | — | 1.51 | 0.41 | 13.5° | „ | Feb. 28, „ | M. Komori |
| 8 | Osaki (<i>Gotanda</i>) | „ (<i>T. Suzuki</i>) | — | — | — | 0.57 | 18.5° | K. & L. | Oct. 29, 1913 | I. Wakita |
| 9 | „ (<i>Minohara</i>) | „ (<i>H. Kamada</i>) | — | — | 1.55 | 0.43 | 14.0° | Schm. | Feb. 6, 1914 | M. Komori |
| 10 | Ikegami | „ (<i>S. Shimizu</i>) | — | 16.0° | 1.15 | 0.32 | 16.2° | „ | Apr. 18, „ | H. Kakehi |
| 11 | Iriarai | „ (<i>C. Hirabayashi</i>) | — | 17.0° | 1.86 | 0.51 | 25.5° | „ | Sept. 7, „ | „ |
| 12 | Ōmori | „ (<i>Seikwa-en</i>) | — | 18.5° (15°) | — | 0.44 | 15.9° | K. & L. | Oct. 24, 1913 | Dr. R. Ishizu |
| 13 | „ | Morigasaki-Kōsen | earth-magnated
common salt | 17.0° (18°) | — | 0.40 | 15.8° | „ | „ | „ |
| 14 | Haneda | Anamori-Kōsen
(<i>Izumi-kan</i>) | iron carbonate
common salt | 17.0° | 1.62 | 0.45 | 21.0° | Schm. | June 1, 1914 | H. Kakehi |
| 15 | Kawasaki | Doku-mizu (<i>Kawa-
saki-Daishi</i>) | — | 16.0° (16.5°) | — | 0.41 | 18.5° | K. & L. | Oct. 24, 1913 | Dr. R. Ishizu |
| 16 | Daishigawara | Well (<i>T. Mizuno</i>) | — | 18.5° | 4.75 | 1.31 | 22.0° | Schm. | July 13, 1914 | M. Komori |
| 17 | Ōami | Cold spring <i>A</i>
(<i>C. Honma</i>) | alkaline | — | 1.70 | 0.47 | 21.0° | „ | „ 27, „ | „ |
| 18 | Yokohama
(<i>Koyasu</i>) | Well (<i>F. Fukamachi</i>) | — | — | 1.53 | 0.42 | 24.0° | „ | „ 20, „ | „ |
| 19 | Enoshima | „ (<i>Ebisu-ya</i>) | — | — | 1.68 | 0.46 | 24.5° | „ | Aug. 9, „ | „ |
| | Kawasaki | Gas evolving from
Doku-mizu | | | Emanation per litre
of gas at 0° C.
0.84 | | | K. & L. | Date when
the gas was
collected
Oct. 24, 1913 | Dr. R. Ishizu |

YUMOTO—TŌNOSAWA

Location.—Yumoto-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring "Moto-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881. Temperature: 42° C.

Total residue: ca. 0.57° C.

Flow of water: ca. 360 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0128 | 0.3269 | 0.3269 |
| Sodium ion (Na ⁺) | 0.1805 | 7.8308 | 7.8308 |
| Calcium ion (Ca ⁺⁺) | 0.0128 | 0.3192 | 0.6384 |
| | | | 8.7961 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.1967 | 5.5487 | 5.5487 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.1255 | 1.3065 | 2.6130 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0392 | 0.6425 | 0.6425 |
| | 0.5675 | 15.9746 | 8.8042 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0327 | | |
| | 0.6002 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0244 |
| Sodium chloride (NaCl) | 0.3055 |
| Sodium sulphate (Na ₂ SO ₄) | 0.1856 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0520 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0327 |
| | 0.6002 |

Besides these trace of magnesium, ferrous ion and organic substances.

The spring thus may be classified as "simple thermal".

The other noted hot springs

| Spring | Fukuzumi-no-yu | Tamano-o-no-yu | Fujiya-no-yu | Sekiguchi-no-yu |
|---|----------------|--------------------|--------------------|-----------------|
| Specific gravity. | — | 1.0004 (at 16° C.) | 1.0004 (at 18° C.) | — |
| Temperature | 44.0° C. | 47.3° C. | 45.5° C. | 42.8° C. |
| Total residue | ca. 0.57 | ca. 0.64 | ca. 0.62 | ca. 0.50 |
| Flow of water.
(in 24 hours) | ca. 540 hls. | — | — | — |

The springs, being in their composition nearly the same as that of "Moto-yu", may be classified as "simple thermal".

MIYANOSHITA

Location.—Onsen-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring "Taikō-no-yu"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0017 at 15° C.

Temperature: 67° C.

Total residue: ca. 2.28.

Flow of water: ca. 208 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.02166 | 1.82886 | 1.82886 |
| Sodium ion (Na ⁺) | 0.65505 | 28.41865 | 28.41865 |
| Ammonium ion (NH ₄ ⁺) | 0.00053 | 0.02931 | 0.02931 |
| Calcium ion (Ca ⁺⁺) | 0.07915 | 1.97875 | 3.9575 |
| Magnesium ion (Mg ⁺⁺) | 0.00742 | 0.30460 | 0.60920 |
| Ferrous ion (Fe ⁺⁺) | 0.00210 | 0.03750 | 0.07500 |
| | | | 34.91852 |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.13643 |
| Sodium chloride (NaCl) | 1.66256 |
| Ammonium chloride (NH ₄ Cl) | 0.00157 |
| Calcium chloride (CaCl ₂) | 0.12599 |
| Calcium sulphate (CaSO ₄) | 0.05585 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.07003 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.04459 |

Anions.

| | | | | | |
|--|---------|----------|-----------------|---|---------|
| Chlorine ion (Cl') | 1.15393 | 32.55001 | 32.55091 | Ferrous bicarbonate [$\text{Fe}(\text{HCO}_3)_2$] | 0.00668 |
| Sulphate ion (SO_4'') | 0.03943 | 0.41047 | 0.82094 | Boric acid (meta) (HBO_2) | 0.02129 |
| Hydrocarbonate ion (HCO_3') | 0.09449 | 1.54876 | 1.54876 | Silicic acid (meta) (H_2SiO_3) | 0.19701 |
| | 2.10370 | 07.10781 | 34.92061 | | 2.32200 |
| Boric acid (meta) (HBO_2) | 0.02129 | | | Free carbon dioxide (CO_2) | 0.02185 |
| Silicic acid (meta) (H_2SiO_3) | 0.19701 | | | | 2.34385 |
| | 2.32200 | | | | |
| Free carbon dioxide (CO_2) | 0.02185 | 0.49659 | | | |
| | 2.34385 | | | | |

The spring thus may be classified as "common salt spring".

The other noted hot springs

| Spring | Mikazuki-no-yu | Kumano-no-yu | Meiji-no-yu | Yoshida-no-yu |
|---------------------------------------|-------------------|--------------------|--------------------|---------------|
| Specific gravity. | | | 1.0017 (at 16°C.) | — |
| Temperature | 51° C. | 36° C. | 67° C. | 45.6° C. |
| Total residue | ca. 2.06 | ca. 1.08 | ca. 2.15 | ca. 1.01 |
| Flow of water (in 24 hours) | ca. 7 hls. | ca. 20 hls. | ca. 49 hls. | — |
| Spring | Taki-no-yu | Fuji-no-yu | Shinrei-no-yu | |
| Specific gravity. | 1.0014 (at 15°C.) | 1.00175 (at 15°C.) | 1.0011 (at 17° C.) | |
| Temperature | 64° C. | 81° C. | 48.4° C. | |
| Total residue | ca. 2.06 | ca. 2.35 | ca. 1.51 | |
| Flow of water (in 24 hours) | ca. 182 hls. | — | — | |

The springs, being in their composition nearly the same as that of "Taikō-no-yu", may be classified as "common salt spring".

DŌGASHIMA

Location.—Onsen-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The spring issues from the Augite Andesite and its Agglomerate.

The hot spring "Musō-no-yu"

Analysis (calculated from the salt table).

Analysed by the Yokohama Imperial Hygienic Laboratory. 1885.

Specific gravity: 1.001 at 15° C.

Temperature: 46° C.

Total residue: ca. 1.46.

Flow of water: ca. 450 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K') | 0.0150 | 0.3831 | 0.3831 |
| Sodium ion (Na') | 0.3820 | 16.5727 | 16.5727 |
| Calcium ion (Ca'') | 0.0690 | 1.7216 | 3.4432 |
| Magnesium ion (Mg'') | 0.0086 | 0.3530 | 0.7060 |
| | | | 21.1050 |
| Anions. | | | |
| Chlorine ion (Cl') | 0.6696 | 18.8886 | 18.8886 |
| Sulphate ion (SO_4'') | 0.0255 | 0.2655 | 0.5310 |
| Hydrocarbonate ion (HCO_3') | 0.1030 | 1.6883 | 1.6883 |
| | | 1.2727 | 39.8728 |
| Silicic acid (meta) (H_2SiO_3) | 0.1207 | | 21.1079 |
| | 1.3934 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0285 |
| Sodium chloride (NaCl) | 0.9696 |
| Calcium chloride (CaCl_2) | 0.1072 |
| Calcium sulphate (CaSO_4) | 0.0362 |
| Calcium bicarbonate [$\text{Ca}(\text{HCO}_3)_2$] | 0.0794 |
| Magnesium bicarbonate [$\text{Mg}(\text{HCO}_3)_2$] | 0.0518 |
| Silicic acid (meta) (H_2SiO_3) | 0.1207 |
| | 1.3934 |

Besides these trace of iodine, hydrophosphate ion, boric acid and organic substances.

The spring thus may be classified as "common salt spring".

SOKOKURA

Location.—Onsen-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring “Onjun-no-yu”

Analysis (calculated from the salt table).

Analysed by the Yokohama Imperial Hygienic Laboratory. 1885. Specific gravity: 1.00192.
Temperature: 72° C. Total residue: ca. 2.19.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|----------------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0513 | 1.3103 | 1.3103 |
| Sodium ion (Na ⁺) | 0.6332 | 27.4707 | 27.4707 |
| Calcium ion (Ca ⁺⁺) | 0.1144 | 2.8529 | 5.7058 |
| Magnesium ion (Mg ⁺⁺) | 0.0092 | 0.3777 | 0.7554 |
| | | | 35.2422 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 1.0204 | 28.7898 | 28.7898 |
| Sulphate ion (SO ₄ ^{''}) | 0.0419 | 0.4362 | 0.8724 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.3412 | 5.5925 | 5.5925 |
| | 2.2116 | 66.8301 | 35.2547 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1841 | | |
| | 2.3957 | | |
| Free carbon dioxide (CO ₂) | not determined | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0978 |
| Sodium chloride (NaCl) | 1.6071 |
| Calcium sulphate (CaSO ₄) | 0.0595 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.3920 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0552 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1841 |
| | 2.3957 |

Besides these trace of ferrous, aluminium, iodine and hydrophosphate ion, boric acid, and organic substances.

The spring thus may be classified as “common salt spring”.

The other noted hot springs

| Spring | Reisen-no-yu | Manju-no-yu | Sinrei-no-yu |
|------------------------------|--------------|-------------|--------------|
| Specific gravity. | 1.00176 | — | — |
| Temperature | 76.0° C. | 70.0° C. | 64.0° C. |
| Total residue | ca. 2.18 | ca. 1.92 | ca. 1.83 |
| Flow of water (in 24 hours). | ca. 6 hls. | ca. 78 hls. | — |

The springs, being in their composition nearly the same as that of “Onjun-no-yu”, may be classified as “common salt spring”.

KIGA

Location.—Miyagino-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring “Ōdaki-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.
Temperature: 46.7° C. Total residue: ca. 1.39.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Sodium ion (Na ⁺) | 0.2992 | 12,9805 | 12,9805 |
| Calcium ion (Ca ⁺⁺) | 0.0859 | 2,1421 | 4,2842 |
| Magnesium ion (Mg ⁺⁺) | 0.0573 | 2,3525 | 4,7050 |
| | | | 21,9697 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.5440 | 15,3456 | 15,3456 |
| Sulphate ion (SO ₄ ^{''}) | 0.1620 | 1,6864 | 3,3728 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.1980 | 3,2454 | 3,2454 |
| | 1,3464 | 37,7525 | 21,9638 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1532 | | |
| | 1,4996 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 0,7595 |
| Magnesium chloride (MgCl ₂) | 0,1125 |
| Calcium sulphate (CaSO ₄) | 0,2297 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,0738 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,1709 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,1532 |
| | 1,4996 |

Besides these trace of bromine ion and organic substances.

The spring thus may be classified as "common salt spring".

The other noted hot springs

| Spring | Kami-no-yu | Shōbu-no-yu | Tani-no-yu | Iwa-no-yu |
|--------------------------------------|-------------|-------------|------------|-------------|
| Temperature. | 40° C. | 45° C. | — | 42° C. |
| Total residue | ca. 1,12 | ca. 1,05 | ca. 1,06 | ca. 0,81 |
| Flow of water (in 24 hrs.) | ca. 35 hls. | — | — | ca. 54 hls. |

The springs, except the last one, being in their composition nearly the same as that of "Ōdaki-no-yu", may be classified as "common salt spring", but "Iwa-no-yu", containing the smaller proportion of solid matters, as "simple thermal".

KOWAKIDANI

Location.—Miyagino-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring "Kowakidani-Onsen"

Analysis (calculated from the salt table).

Analysed by the Yokohama Imperial Hygienic Laboratory. 1885. Specific gravity: 1,00078.
Temperature: 35,6° C. Total residue: ca. 1,13.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Hydrogen ion (H ⁺) | 0.0020 | 1,9801 | 1,9801 |
| Sodium ion (Na ⁺) | 0.0209 | 0,9067 | 0,9067 |
| Calcium ion (Ca ⁺⁺) | 0.0348 | 0,8678 | 1,7356 |
| Magnesium ion (Mg ⁺⁺) | 0.0125 | 0,5131 | 1,0262 |
| Ferrous ion (Fe ⁺⁺) | 0.1465 | 2,6208 | 5,2416 |
| Manganous ion (Mn ⁺⁺) | 0.0015 | 0,0616 | 0,1232 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0295 | 1,0886 | 3,2658 |
| | | | 14,2792 |
| Anions. | | | |
| Hydrosulphate ion (HSO ₄ [']) | 0.1900 | 1,9574 | 1,9574 |
| Sulphate ion (SO ₄ ^{''}) | 0.5877 | 6,1181 | 12,2362 |
| | 1,0254 | 16,1142 | 14,1936 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1324 | | |
| | 1,1578 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Sodium sulphate (Na ₂ SO ₄) | 0,0645 |
| Calcium sulphate (CaSO ₄) | 0,1178 |
| Magnesium sulphate (MgSO ₄) | 0,0624 |
| Ferrous sulphate (FeSO ₄) | 0,3982 |
| Manganous sulphate (MnSO ₄) | 0,0040 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0,1865 |
| Free sulphuric acid (H ₂ SO ₄) | 0,1920 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,1324 |
| | 1,1578 |

Besides these trace of potassium, chlorine, hydrophosphate and hydrosulphide ion, boric acid, and organic substances.

The spring thus may be classified as "acid vitriol spring".

YUNOHANA-ZAWA

Location.—Moto-hakone-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.
The spring issues from the Augite Andesite and its Agglomerate.

The hot spring “Hana-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory.

Specific gravity: 1.0006 at 15° C.

Temperature: 40° C.

Total residue: ca. 0.7.

Flow of water: ca. 691 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|---------------|----------------|---------------------------|
| Hydrogen ion (H ⁺) | 0.0047 | 4.6530 | 4.6530 |
| Potassium ion (K ⁺) | 0.0001 | 0.2324 | 0.2324 |
| Sodium ion (Na ⁺) | 0.0213 | 0.9241 | 0.9241 |
| Calcium ion (Ca ⁺⁺) | 0.0303 | 0.7556 | 1.5112 |
| Magnesium ion (Mg ⁺⁺) | 0.0194 | 0.7964 | 1.5928 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0226 | 0.8339 | 2.5017 |
| | | | 11.4152 |
| Anions. | | | |
| Hydrosulphate ion (HSO ₄ [']) | 0.4557 | 4.6946 | 4.6946 |
| Sulphate ion (SO ₄ ^{''}) | 0.3262 | 3.3058 | 6.7916 |
| | 0.8893 | 16.2858 | 11.4862 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0180 | | |
| | 0.9073 | | |
| Free hydrogen sulphide (H ₂ S) | 0.1205 | | |
| | 1.0278 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|---------------|
| Potassium sulphate (K ₂ SO ₄) | 0.0202 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0657 |
| Calcium sulphate (CaSO ₄) | 0.1026 |
| Magnesium sulphate (MgSO ₄) | 0.0772 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.1432 |
| Free sulphuric acid (H ₂ SO ₄) | 0.4604 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0180 |
| | 0.2073 |
| Free hydrogen sulphide (H ₂ S) | 0.1205 |
| | 1.0278 |

Besides these trace of ferrous, chlorine and hydrophosphate ion, and organic substances.

The spring thus may be classified as “acid hydrogen sulphide spring”.

SENGOKU-BARA

Location.—Sengokubara-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.
The springs issue from the Augite Andesite and its Agglomerate.

The hot spring “Sengoku-shin-yu”

Analysis (calculated from the salt table).

Analysed by the Yokohama Imperial Hygienic Laboratory. 1885.

Specific gravity: 1.00095 at 15° C.

Total residue: ca. 1.87.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K) | 0.0064 | 0.1635 | 0.1635 |
| Sodium ion (Na ⁺) | 0.0999 | 4.3341 | 4.3341 |
| Calcium ion (Ca ⁺⁺) | 0.2544 | 6.3441 | 12.6882 |
| Magnesium ion (Mg ⁺⁺) | 0.0649 | 2.6642 | 5.3284 |
| Ferrous ion (Fe ⁺⁺) | 0.0014 | 0.0250 | 0.0500 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0042 | 0.1550 | 0.4650 |
| | | | 23.0292 |
| Anions. | | | |
| Sulphate ion (SO ₄ ^{''}) | 1.1088 | 11.5428 | 23.0856 |
| | 1.5400 | 25.2287 | 23.0856 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2726 | | |
| | 1.8126 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|---------------|
| Potassium sulphate (K ₂ SO ₄) | 0.0142 |
| Sodium sulphate (Na ₂ SO ₄) | 0.3083 |
| Calcium sulphate (CaSO ₄) | 0.8625 |
| Magnesium sulphate (MgSO ₄) | 0.3246 |
| Ferrous sulphate (FeSO ₄) | 0.0038 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.0266 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2726 |
| | 1.5126 |

Besides these trace of manganous, chlorine and hydrophosphate ion, and organic substances.

The spring thus may be classified as “sulphated bitter spring”.

UBAKO

Location.—Moto-hako e-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The spring issues from the Augite Andesite and its Agglomerate.

The hot spring "Ubako-yu"

Analysis (calculated from the salt table).

Analysed by the Yokohama Imperial Hygienic Laboratory. 1885.

Specific gravity: 1,0006 at 13° C.

Temperature: 40° C

Total residue: ca. 0.65.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0,0054 | 0,1379 | 0,1379 |
| Sodium ion (Na ⁺) | 0,0284 | 1,2321 | 1,2321 |
| Calcium ion (Ca ⁺⁺) | 0,0739 | 1,8429 | 3,6858 |
| Magnesium ion (Mg ⁺⁺) | 0,0198 | 0,8128 | 1,6256 |
| Aluminium ion (Al ⁺⁺⁺) | 0,0074 | 0,2731 | 0,8193 |
| | | | 7,5007 |
| Anions. | | | |
| Sulphate ion (SO ₄ '') | 0,3609 | 3,7570 | 7,5140 |
| | 0,4958 | 8,0558 | 7,5140 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,2031 | | |
| | 0,6989 | | |

| | Grams |
|---|--------|
| Potassium sulphate (K ₂ SO ₄) | 0,0121 |
| Sodium sulphate (Na ₂ SO ₄) | 0,0875 |
| Calcium sulphate (CaSO ₄) | 0,2506 |
| Magnesium sulphate (MgSO ₄) | 0,0989 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0,0467 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,2031 |
| | 0,6989 |

Besides these trace of ferrous, chlorine ion, and organic substances.

The spring thus may be classified as "simple thermal".

YUGAWARA

Location.—Tohi-mura, Ashigara-shimo-gun, Province Sagami, Kanagawa-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring "Kawashita-no-yu"

Analysis.

Analysed by the Tōkyō Imperial Hygienic Laboratory. Temperature: 73° C.

Total residue: ca. 2,36.

Flow of water: ca. 1458 hectolitres in 24 hours.

| | Grams | | Grams |
|---|--------|--|--------|
| Potassium chloride (KCl) | 0,0510 | Silica (SiO ₂) | 0,1068 |
| Sodium chloride (NaCl) | 1,4447 | | 2,3624 |
| Calcium chloride (CaCl ₂) | 0,0412 | Carbon dioxide (free and in form of bicar- | |
| Magnesium chloride (MgCl ₂) | 0,0546 | bonates) (CO ₂) | 0,0090 |
| Calcium sulphate (CaSO ₄) | 0,5916 | | 2,3714 |
| Calcium carbonate (CaCO ₃) | 0,0725 | | |

Besides these trace of iron, aluminium, phosphoric acid and boric acid.

The spring may be classified as "common salt spring".

Table 46.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Effluatation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------------------|----------------|--------------------------|---|------------------------------------|-------------------|--------------------|----------------------------------|---------------|
| | | | | | in
Mache's units | at
temp. in C. | | | |
| 1 | Nakanishiya-no-yu | common salt | 82,5° | — | 0,22 | 22,0° | K. & L. | Feb. 19, 1913 | Dr. R. Ishizu |
| 2 | Uenoya-no-yu | .. | 88,5° | — | 0,18 | 9,5° | .. | .. | .. |
| 3 | Izuya-no-yu | .. | 80,0° | — | 0,14 | 10,0° | .. | .. 20, .. | .. |
| 4 | Yuya-no-yu | .. | 54,5° | — | 0,15 | 19,0° | .. | .. 19, .. | Y. Kinugasa |
| 5 | Okura-no-yu | .. | 43,0° | — | 0,07 | 10,0° | .. | .. 20, .. | Dr. R. Ishizu |
| 6 | Kawashita-no-yu | .. | 73,0° | 1458 | — | — | — | — | — |
| 7 | Nangan-no-yu | .. | 52,0° | 1458 | — | — | — | — | — |
| 8 | Mamane-no-yu
(<i>Kami-no-yu</i>) | .. | 56,0° | 1458 | — | — | — | — | — |
| 9 | Yakushi-no-yu | .. | 34-46° | 1458 | — | — | — | — | — |
| 10 | Shimo-no-yu (<i>Sō-yu</i>) | .. | 72,5° | 1458 | — | — | — | — | — |
| 11 | Kawanaka-no-yu | .. | 72,0° | — | — | — | — | — | — |
| 12 | Kawara-no-yu (<i>Mae-no-yu</i>) | .. | 41,0° | — | — | — | — | — | — |

IZUSAN

Location.—Atami-machi, Takata-gun, Province Izu, Shizuoka-ken.

The spring issues from the Augite Andesite and its Agglomerate.

The hot spring “Hashiri-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1901.

Temperature: 60° C.

Flow of water: abundant.

Total residue: ca. 1,42.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols. | Milligram-
equivalents |
|--|--------|-----------------|---------------------------|
| Potassium ion (K') | 0,0114 | 0,2912 | 0,2912 |
| Sodium ion (Na') | 0,2239 | 9,7137 | 9,7137 |
| Calcium ion (Ca'') | 0,2072 | 5,1671 | 10,3342 |
| Magnesium ion (Mg'') | 0,0046 | 0,1888 | 0,3776 |
| | | | 20,7167 |
| Anions. | | | |
| Chlorine ion (Cl') | 0,2620 | 7,3907 | 7,3907 |
| Sulphate ion (SO ₄ '') | 0,6403 | 6,6656 | 13,3212 |
| | 1,3494 | 29,4171 | 20,7119 |
| Silicic acid (meta) (H ₂ SiO ₃) . . | 0,0494 | | |
| | 1,3988 | | |

| | Grams |
|--|--------|
| Sodium chloride (NaCl) | 0,4323 |
| Potassium sulphate (K ₂ SO ₄) | 0,0254 |
| Sodium sulphate (Na ₂ SO ₄) | 0,1654 |
| Calcium sulphate (CaSO ₄) | 0,7036 |
| Magnesium sulphate (MgSO ₄) | 0,0227 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0494 |
| | 1,3988 |

Besides these trace of ferrous, aluminium and hydrophosphate ion.

The spring thus may be classified as “sulphated bitter spring”.

Radio-activity. 0,10 Mache's units in 1 litre of water at 18,2° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 21, 1913).

ATAMI

Location.—Atami-machi, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Augite Andesite and its Agglomerate.

The hot spring “ $\bar{O}$ -yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 108° C.

Flow of water: ca. 1296 hectolitres in 24 hours.

Total residue: ca. 9.24.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.1859 | 4.7484 | 4.7484 |
| Sodium ion (Na ⁺) | 2.1312 | 92.4599 | 92.4599 |
| Calcium ion (Ca ⁺⁺) | 1.0831 | 27.0099 | 54.0198 |
| Magnesium ion (Mg ⁺⁺) | 0.0037 | 0.1519 | 0.3038 |
| Ferrous ion (Fe ⁺⁺) | 0.0006 | 0.0107 | 0.0214 |
| | | | 151.5533 |
| Anions. | Grams | Milli-
mols | Milligram-
equivalents |
| Chlorine ion (Cl ⁻) | 5.3053 | 149.6559 | 149.6559 |
| Sulphate ion (SO ₄ ^{''}) | 0.0926 | 0.9640 | 1.9280 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0014 | 0.0229 | 0.0229 |
| | | | 8.8038 |
| | | | 275.0236 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.6813 | | 151.6068 |
| | | | 9.4851 |

| | Grams |
|--|--------|
| Potassium chloride (KCl) | 0.3540 |
| Sodium chloride (NaCl) | 5.4090 |
| Calcium chloride (CaCl ₂) | 2.8930 |
| Magnesium chloride (MgCl ₂) | 0.0145 |
| Calcium sulphate (CaSO ₄) | 0.1313 |
| Ferrous sulphate (FeSO ₄) | 0.0020 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.6813 |
| | 9.4851 |

Besides these trace of manganous, bromine ion and organic substances.

The spring thus may be classified as “earth-muriated common salt spring”.

Radio-activity. 0.04 Mache's units in 1 litre of water at 15° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 23, 1913).

The hot spring “Me-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Specific gravity: 1.007 at 13° C.

Temperature: 48° C.

Flow of water: ca. 140 hectolitres in 24 hours.

Total residue: ca. 8.10.

The spring, being in its composition nearly the same as that of “ $\bar{O}$ -yu”, may be classified as “earth-muriated common salt spring”.

Radio-activity. 0.08 Mache's units in 1 litre of water at 20° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 21, 1913).

Table 47.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|----------------------|-------------------------------|--------------------------|---|---------------------------------|-------------------|--------------------|----------------------------------|-------------|
| | | | | | in
Maché's units | at
temp. in C. | | | |
| 1 | Ō-yu | earth-muriated
common salt | 108.0° | ca.
1296 | 0.04 | 15.0° | K. & L. | March 23, 1913 | Y. Kinugasa |
| 2 | Me-no-yu | " | 48.0° | 140 | 0.08 | 20.0° | " | " 21, " | " |
| 3 | Ozawa-no-yu No. 2 | " | 97.0° | 466 | 0.14 | 18.2° | " | " | " |
| 4 | Komatsu-no-yu | " | 93.0° | 241 | 0.08 | 20.2° | " | " | " |
| 5 | Kawara-no-yu | " | 81.0° | — | 0.02 | 15.0° | " | " 22, " | " |
| 6 | " (<i>Ishi-yu</i>) | " | 77.0° | — | 0.14 | 14.2° | " | " | " |
| 7 | Furuya-no-yu | " | 80.0° | 86 | 0.18 | 14.8° | " | " | " |
| 8 | Takasagoya-no-yu | " | 91.0° | — | 0.26 | 14.2° | " | " | " |

ITŌ

Location.—Itō-machi, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Quaternary.

The hot spring "Matsubara-Onsen"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1889.

Specific gravity: 1.0008 at 15° C.

Total residue: ca. 0.98.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram

| Composition of a solution containing 11.2147 grams | | | |
|---|--------|----------------|---------------------------|
| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
| Potassium ion (K ⁺). | 0.0150 | 0.3831 | 0.3831 |
| Sodium ion (Na ⁺) | 0.2585 | 11.2147 | 11.2147 |
| Calcium ion (Ca ⁺⁺) | 0.0407 | 1.0150 | 2.0300 |
| Magnesium ion (Mg ⁺⁺). | 0.0057 | 0.2298 | 0.4596 |
| | | | 14.0874 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.2758 | 7.7799 | 7.7799 |
| Sulphate ion (SO ₄ ⁼⁼). | 0.8030 | 3.1545 | 6.3090 |
| | 0.8987 | 23.7770 | 14.0883 |
| Silicic acid (meta) (H ₂ SiO ₃). | 0.0653 | | |
| | 0.9640 | | |
| | | | 0.9640 |

Besides these trace of ferrous, aluminium and hydrophosphate ion, and boric acid.

The spring thus may be classified as "simple thermal".

Table 48.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---|----------------|--------------------------|---------------------------------|---------------------|--------------------|--|---------------|
| | | | | in
Maché's units | at
(temp. in C.) | | | |
| 1 | Matsubara-moto-yu * | simple | 47.8° | 0.29 | 17.0° | K. & L. | Feb. 22, 1913 | Dr. R. Ishizu |
| 2 | Matsubara-shin-yu * | " | 46.0° | 0.16 | 20.0° | " | " | " |
| 3 | Private bath (<i>Yoshida-gankwa-in</i>) | — | 48.5° | 0.31 | 20.5° | " | " 23, " | " |
| 4 | " (<i>H. Hirose</i>) | — | 48.5° | 0.47 | 13.5° | " | " | " |
| 5 | " (<i>M. Suzuki</i>) | — | 47.0° | 0.27 | 19.0° | " | " | Y. Kinugasa |
| 6 | Yumoto-kwan-no-yu | — | 50.5° | 0.34 | 13.8° | " | " | " |
| 7 | Private bath (<i>Kobayashi</i>) | — | 43.5° | 0.21 | 13.0° | " | " | " |
| 8 | Shishido-Onsen * | simple | 47.0° | 0.45 | 21.0° | " | " 22, " | Dr. R. Ishizu |
| 9 | Wada-Onsen * | salt | 37.5° | 0.40 | 20.0° | " | " | " |
| 10 | Kusumi-sō-yu (<i>Onnayū</i>) * | " | 50.0° | 0.46 | 19.8° | " | " 23, " | Y. Kinugasa |
| 11 | Kusumi-Onsen * | " | 49.5° | 0.36 | 13.5° | " | " | Dr. R. Ishizu |
| 12 | Me-no-yu * | " | 37.5° | 0.33 | 19.5° | " | " | " |
| 13 | Yuda-Onsen * | " | 33.5° | 0.11 | 19.5° | " | " 22, " | " |
| 14 | Private bath (<i>S. Kitazato</i>) | — | 34.0° | 0.42 | 20.0° | " | " | " |
| 15 | " (<i>S. Kumagi</i>) | — | 50.0° | 0.48 | 13.5° | " | " 23, " | " |
| | Gas evolving from
Matsubara-moto-yu | — | — | 1.19 | — | " | Date when the
gas was
collected
Feb. 23, 1913 | Y. Kinugasa |

* These springs are for public use.

SHUZENJI

Location.—Shuzenji-mura, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Andesite and its Tuff.

The hot spring "Ishi-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1907.

Temperature: 59.5° C.

Total residue: ca. 1.13.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents | | Grams |
|----------------------|--------|----------------|---------------------------|--|--------|
| Potassium ion (K') | 0.0073 | 0.1865 | 0.1865 | Sodium chloride (NaCl) | 0.5646 |
| Sodium ion (Na') | 0.3183 | 13.8091 | 13.8091 | Potassium sulphate (K ₂ SO ₄) | 0.0162 |
| Calcium ion (Ca'') | 0.0553 | 1.3791 | 2.7582 | Sodium sulphate (Na ₂ SO ₄) | 0.2261 |
| Magnesium ion (Mg'') | 0.0006 | 0.0246 | 0.0492 | Sodium bicarbonate (NaHCO ₃) | 0.0821 |
| | | | 16.8030 | Calcium sulphate (CaSO ₄) | 0.1873 |

Anions.

| | | | | | |
|--|----------------|---------|---------------|--|--------|
| Chlorine ion (Cl') | 0,3421 | 9,6502 | 9,6502 | Magnesium sulphate (MgSO_4) | 0,0027 |
| Sulphate ion (SO_4'') | 0,2058 | 3,0773 | 6,1586 | Silicic acid (meta) (H_2SiO_3) | 0,1063 |
| Hydrocarbonate ion (HCO_3') | 0,0596 | 0,9769 | 0,9769 | | 1,1853 |
| | 1,0770 | 22,1057 | 10,7857 | | |
| Silicic acid (meta) (H_2SiO_3) | 0,1063 | | | | |
| | 1,1853 | | | | |
| Free carbon dioxide (CO_2) | not determined | | | | |

Besides these trace of ferrous, manganous, aluminium and hydrophosphate ion, and boric acid.

The spring thus may be classified as “saline common salt spring”.

Radio-activity. 1,35 Mache's units in 1 litre of water at 9,5° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 25, 1913).

8,73 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 26, 1913).

The hot spring “Dokko-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881. Temperature: 69° C.

Flow of water: ca. 432 hectolitres in 24 hours.

Total residue: ca. 1,12.

The spring, being in its composition nearly the same as that of “Ishi-yu”, may be classified as “saline common salt spring”.

Radio-activity. 0,64 Mache's units in 1 litre of water at 9,7° C.

4,11 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, Feb. 25, 1913).

The hot spring “Ma-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 64° C.

Total residue: ca. 1,21.

The spring, being in its composition nearly the same as that of “Ishi-yu”, may be classified as “saline common salt spring”.

Radio-activity. 0,13 Mache's units in 1 litre of water at 11,0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 26, 1913).

Table 49.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|--------------------|-------------------------------------|-----------------------|--------------------------|---|---------------------------------|---------------------|--------------------|----------------------------------|--|
| | | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Ishi-yu (<i>Public bath</i>) | saline
common salt | 59.5° | ca. — | 1.35 | 9.5° | K. & L. | Feb. 25, 1913 | Dr. R. Ishizu |
| 2 | Dokko-no-yu („) | „ | 69.0° | 432 | 0.64 | 9.7° | „ | „ | Y. Kinugasa |
| 3 | Ma-yu („) | „ | 64.0° | — | 0.13 | 11.0° | „ | „ 26, „ | Dr. R. Ishizu |
| 4 | Hako-yu („) | „ | 64.0° | 327 | 0.64 | 9.5° | „ | „ 25, „ | „ |
| 5 | Chigo-no-yu („) | „ | 55.0° | 519 | 1.08 | 9.5° | „ | „ | „ |
| 6 | Ayame-no-yu | „ | 77.0° | 1400 | 0.36 | 11.0° | „ | „ 26, „ | „ |
| 7 | Katsura-no-yu | „ | 64.0° | — | 0.34 | 12.5° | „ | „ | „ |
| 8 | Kikuya-no-yu <i>No. 1</i> | „ | 72.0° | — | 0.31 | 11.0° | „ | „ | „ |
| 9 | „ <i>No. 2</i> | „ | 70.0° | — | 0.50 | 9.5° | „ | „ 25, „ | Y. Kinugasa |
| 10 | Sugi-no-yu | „ | 60.0° | 156 | 0.50 | 9.5° | „ | „ | Dr. R. Ishizu |
| 11 | Ume-no-yu | „ | 60.0° | 156 | 0.66 | 9.0° | „ | „ | Y. Kinugasa |
| 12 | Asabarō-no-yu | „ | 63.0° | — | 0.76 | 11.0° | „ | „ 26, „ | Dr. R. Ishizu |
| 13 | Meiji-no-yu | „ | 66.0° | 149 | 0.46 | 9.5° | „ | „ 25, „ | „ |
| 14 | Iwa-no-yu | „ | 60.0° | 520 | 0.31 | 11.0° | „ | „ 26, „ | „ |
| 15 | Hana-no-yu | „ | 67.0° | 80 | 0.15 | 11.0° | „ | „ | „ |
| 16 | Taki-no-yu | „ | 65.0° | — | 1.00 | 11.0° | „ | „ | „ |
| 17 | Kawara-no-yu (<i>Public bath</i>) | „ | 63.0° | 108 | — | — | — | — | — |
| Gas evolving from: | | | | | | | | | Emanation per litre
of gas at 0° C. |
| 1 | Ishi-yu | — | — | — | 8.73 | — | K. & L. | Feb. 26, 1913 | Dr. R. Ishizu |
| 2 | Dokko-no-yu | — | — | — | 4.11 | — | „ | „ 25, „ | Y. Kinugasa |
| 3 | Kikuya-no-yu <i>No. 2</i> | — | — | — | 7.07 | — | „ | „ 26, „ | Dr. R. Ishizu |

NAGAOKA

Location.—Kawanishi-mura, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Tertiary Tuff Breccia.

The hot spring “Nagaoka-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1907.

Specific gravity: 1.0005 at 15° C.

Temperature: 52.7° C.

Flow of water: ca. 816 hectolitres in 24 hours.

Total residue: ca. 0.70.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-mols | Milli-gram-equivalents |
|--|--------|------------|------------------------|
| Potassium ion (K ⁺) | 0,0016 | 0,0409 | 0,0409 |
| Sodium ion (Na ⁺) | 0,2077 | 9,0108 | 9,0108 |
| Calcium ion (Ca ⁺⁺) | 0,0045 | 0,1122 | 0,2244 |
| | | | 0,2761 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0,1137 | 3,2073 | 3,2073 |
| Sulphate ion (SO ₄ '') | 0,2653 | 2,7610 | 5,5238 |
| Hydrocarbonate ion (HCO ₃ ') | 0,0332 | 0,5442 | 0,5442 |
| | 0,6260 | 15,6773 | 9,2753 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0909 | | |
| | 0,7169 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | |
|--|--------|
| Sodium chloride (NaCl) | 0,1876 |
| Potassium sulphate (K ₂ SO ₄) | 0,0036 |
| Sodium sulphate (Na ₂ SO ₄) | 0,1737 |
| Sodium bicarbonate (NaHCO ₃) | 0,0458 |
| Calcium sulphate (CaSO ₄) | 0,0153 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0909 |
| | 0,7169 |

Besides these a small quantity of aluminium ion, a very small quantity of ferrous, hydrophosphate ion, and boric acid.

The spring thus may be classified as "simple thermal".

Table 50.

| No. | Spring | Classification | Temp. of spring in C. | Emanation per litre of water | | Apparatus employed | Date when the water was taken | Examined by |
|-----|-------------------------------|----------------|-----------------------|-------------------------------------|----------------|--------------------|---------------------------------|-------------|
| | | | | in Maché's units | at temp. in C. | | | |
| 1 | Nagaoka-Onsen No. 1 | simple | 45,0° | 0,10 | 12,0° | K & L. | March 13, 1913 | Y. Kinugasa |
| 2 | " No. 2 | " | 50,0° | 0,19 | 14,6° | " | " | " |
| 3 | " No. 3 | " | 49,0° | 0,29 | 17,0° | " | " | " |
| 4 | " No. 4 | " | 53,0° | 0,18 | 16,3° | " | " | " |
| 5 | Tamon-yu No. 1 (Private bath) | " | 48,0° | 0,27 | 10,4° | " | " 14, " | " |
| 6 | " No. 2 (") | " | 49,0° | 0,33 | 14,6° | " | " | " |
| 7 | " No. 3 (") | " | 48,0° | 0,27 | 15,2° | " | " | " |
| 8 | " No. 4 (") | " | 44,0° | 0,51 | 13,0° | " | " | " |
| 9 | Private bath (S. Matsumoto) | " | 41,0° | 0,38 | 11,0° | " | " | " |
| 10 | " (S. Sugiyama) | " | 49,0° | 0,26 | 13,5° | " | " 13, " | " |
| 11 | Kona-Onsen (Public bath) | " | 52,0° | 0,17 | 15,0° | " | " 14, " | " |
| | Gas evolving from: | | | Emanation per litre of gas at 0° C. | | | Date when the gas was collected | |
| 1 | Nagaoka-Onsen No. 4 | | | 1,29 | | " | March 13, 1913 | " |
| 2 | Tamon-yu No. 3 | | | 3,34 | | " | " 14, " | " |

NAKAISHIDA

Location.—Ooka-mura, Shuntō-gun, Province Suruga, Shizuoka-ken.

The cold spring “Kametsuru-Onsen”

Temperature: 15.0° C.

Total residue: ca. 0.14.

No accurate analysis has ever been made, but the spring seems to be “sulphur spring”.

Radio-activity. 0.14 Mache's units in 1 litre of water at 13.7° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 19, 1913).

HATAGE

Location.—Kwan-nami-mura, Takata-gun, Province Izu, Shizuoka-ken.

The hot spring “Hatage-Onsen”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1911.

Specific gravity: 1.000.

Temperature: 43.3° C.

Flow of water: ca. 508 hectolitres in 24 hours.

Total residue: ca. 0.11.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents | | Grams |
|--|---------|----------------|---------------------------|---|---------|
| Potassium ion (K') | 0.00097 | 0.02478 | 0.02478 | Potassium chloride (KCl) | 0.00185 |
| Sodium ion (Na') | 0.04025 | 1.74620 | 1.74620 | Sodium chloride (NaCl) | 0.02942 |
| Ammonium ion (NH ₄ ') | 0.00021 | 0.01162 | 0.01162 | Ammonium chloride (NH ₄ Cl) | 0.00062 |
| Calcium ion (Ca'') | 0.00771 | 0.19275 | 0.38550 | Sodium sulphate (Na ₂ SO ₄) | 0.06894 |
| | | | 2.16810 | Sodium bicarbonate (NaHCO ₃) | 0.02298 |
| Anions. | | | | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.03122 |
| Chlorine ion (Cl') | 0.01912 | 0.53935 | 0.53935 | Silicic acid (meta) (H ₂ SiO ₃) | 0.02701 |
| Sulphate ion (SO ₄ '') | 0.04658 | 0.48491 | 0.96982 | | 0.18204 |
| Hydrocarbonate ion (HCO ₃ ') | 0.04020 | 0.65891 | 0.65891 | Free carbon dioxide (CO ₂) | 0.04442 |
| | 0.15504 | 3.65852 | 2.16808 | | 0.22646 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.02701 | | | | |
| | 0.18205 | | | | |
| Free carbon dioxide (CO ₂) | 0.04442 | 1.00954 | | | |
| | 0.22647 | | | | |

The spring thus may be classified as “simple thermal”.

Table 51.

| No. | Spring | Classification | Temp. of spring
in C. | Evaporation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------|----------------|--------------------------|-----------------------------------|-------------------|--------------------|----------------------------------|-------------|
| | | | | in
Maché's mat. | at
temp. in C. | | | |
| 1 | Hatage-Onsen No. 1 | Simple | 40.0° | 0.37 | 10.4 | K. & L. | March 10, 1913. | Y. Kinugasa |
| 2 | " No. 2 | " | 40.0° | 0.44 | 14.2° | " | " 15, " | " |
| 3 | " No. 3 | " | 41.0° | 0.44 | 15.4° | " | " " | " |
| 4 | " No. 4, a | " | 38.0° | 0.35 | 15.5° | " | " 16, " | " |
| 5 | " No. 4, b | " | 38.0° | 0.35 | 14.8° | " | " " | " |
| 6 | " No. 5 | " | 39.0° | 0.48 | 11.4° | " | " 15, " | " |
| 7 | " No. 6 | " | 39.0° | 0.35 | 10.7 | " | " 10, " | " |
| 8 | " No. 7 | " | 40.0° | 0.41 | 16.2° | " | " " | " |
| 9 | " No. 8 | " | 39.0° | 0.39 | 15.5° | " | " " | " |

FUNABARA

Location.—Naka-kano-mura, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Andesite.

The hot spring "Kōgyoku-no-yu" No. 1

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1909.

Specific gravity: 1.0006 at 18° C.

Temperature: 47° C.

Flow of water: ca. 684 hectolitres in 24 hours.

Total residue: ca. 1.09.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
moles | Milliequiva-
lents | | Grams |
|--|--------|-----------------|-----------------------|---|--------|
| | | | | | |
| Potassium ion (K ⁺) | 0.0037 | 0.0939 | 0.0939 | Potassium chloride (KCl) | 0.0070 |
| Sodium ion (Na ⁺) | 0.2682 | 11.6357 | 11.6359 | Sodium chloride (NaCl) | 0.2840 |
| Calcium ion (Ca ⁺⁺) | 0.0742 | 1.8504 | 3.7008 | Sodium sulphate (Na ₂ SO ₄) | 0.4662 |
| Magnesium ion (Mg ⁺⁺) | 0.0004 | 0.0164 | 0.0328 | Sodium bicarbonate (NaHCO ₃) | 0.0192 |
| | | | 15.4734 | Calcium sulphate (CaSO ₄) | 0.2516 |
| Anions. | | | | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0024 |
| Chlorine ion (Cl ⁻) | 0.1754 | 4.0434 | 4.9481 | Silicic acid (meta) (H ₂ SiO ₃) | 0.0414 |
| Sulphate ion (SO ₄ ^{''}) | 0.4926 | 5.1270 | 10.2540 | | 1.0718 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0159 | 0.2613 | 0.2613 | Free carbon dioxide (CO ₂) | 0.0208 |
| | 1.0794 | 25.6339 | 15.4734 | | 1.0926 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0414 | | | | |
| | 1.0718 | | | | |
| Free carbon dioxide (CO ₂) | 0.0208 | 0.4727 | | | |
| | 1.0926 | | | | |

Besides these a small quantity of ferrous, aluminium ion and trace of boric acid.

The spring thus may be classified as “saline bitter spring”.

Radio-activity. 0.20 Mache's units in 1 litre of water at 8.0° C.

1.77 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 11, 1913).

Table 52.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--|----------------|--------------------------|---|---------------------------------|---------------------|--------------------|---|-------------|
| | | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kōgyoku-no-yu No. 1 | saline bitter | 47.0° | ca. 68.4 | 0.20 | 8.0° | K. & L. | March 11, 1913 | Y. Kinugasa |
| 2 | „ No. 2 | „ | 35.0° | — | 0.23 | 7.5° | „ | „ | „ |
| 3 | „ No. 3 | „ | 40.0° | — | 0.15 | 7.0° | „ | „ | „ |
| 4 | Kumano-no-yu | salt | 46.0° | 1530 | 0.30 | 5.0° | „ | „ | „ |
| | Gas evolving from
Kōgyoku-no-yu No. 1 | — | — | — | 1.77 | — | „ | Date when the
gas was
collected
March 11, 1913 | „ |

YOSHINA

Location.—Kami-kano-mura, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Andesite.

The hot spring “Yoshina-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1897.

Total residue: ca. 1.24.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | | Grams | Milli-
mols | Milligram-
equivalents | | Grams |
|--|--------|---------|----------------|---------------------------|--|--------|
| Potassium ion (K ⁺) | 0.0090 | 0.2299 | 0.2299 | | Sodium chloride (NaCl) | 0.1823 |
| Sodium ion (Na ⁺) | 0.2727 | 11.8308 | 11.8308 | | Potassium sulphate (K ₂ SO ₄) | 0.0200 |
| Calcium ion (Ca ⁺⁺) | 0.0874 | 2.1802 | 4.3604 | | Sodium sulphate (Na ₂ SO ₄) | 0.6200 |
| | | | 16.4211 | | Calcium sulphate (CaSO ₄) | 0.2963 |
| Anions. | | | | | Silicic acid (meta) (H ₂ SiO ₃) | 0.0867 |
| Chlorine ion (Cl ⁻) | 0.1105 | 3.1171 | 3.1171 | | | 1.2053 |
| Sulphate ion (SO ₄ ^{''}) | 0.6390 | 6.6520 | 13.3040 | | | |
| | 1.1186 | 24.0100 | 16.4211 | | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0867 | | | | | |
| | 1.2053 | | | | | |

Besides these trace of magnesium, ferrous, aluminium and hydrophosphate ion.

The spring thus may be classified as “saline bitter spring”.

Table 53.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------------|----------------|--------------------------|---|---------------------------------|-------------------|--------------------|----------------------------------|-------------|
| | | | | | in
Maché's units | at
temp. in C. | | | |
| 1 | Ō-yu (<i>Public bath</i>) | saline bitter | 50,0° | ca.
265 | 0,30 | 7,0 | K. & L. | March 11, 1913 | Y. Kinugasa |
| 2 | Tōfuya-no-yu | " | 49,0° | 126 | 0,15 | 7,2° | " | " | " |
| 3 | Tōfuya-shin-yu | " | 41,0° | — | 0,36 | 7,0° | " | " | " |

YUGASHIMA

Location.—Kami-kano-mura, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Andesite and its Tuff.

Table 54.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--|----------------|--------------------------|---|---------------------------------|-------------------|--------------------|----------------------------------|-------------|
| | | | | | in
Maché's units | at
temp. in C. | | | |
| 1 | Seko-no-O-yu (<i>Public bath</i>) | salt | 55,0° | ca.
216 | 1,93 | 10,5° | K. & L. | March 3, 1913 | Y. Kinugasa |
| 2 | Seko-no-Moto-yu | " | 49,0° | 130 | 0,81 | 10,5° | " | " | " |
| 3 | Kidachi-yu (<i>Public bath</i>) | carbonated | 41,0 | 80 | 0,15 | 9,9 | " | " | " |
| 4 | Spring No. 2 (") | — | 49,0° | — | 0,70 | 8,5 | " | " | " |
| 5 | Yumoto-kwan-no-yu
(<i>Spring No. 7</i>) | carbonated | 56,0° | 130 | 1,27 | 8,0 | " | " | " |
| 6 | Yumoto-kwan-Taki-yu
(<i>Spring No. 8</i>) | — | 55,0° | — | 1,09 | 9,0 | " | " | " |
| 7 | Ochiai-rō-no-yu | salt | 64,0° | 173 | 1,31 | 7,5 | " | " 4, " | " |
| 8 | Spring No. 3 (<i>S. Adachi</i>) | — | 49,0° | — | 1,00 | 8,3 | " | " 3, " | " |
| 9 | Spring No. 9 (<i>K. Adachi</i>) | — | 56,0° | — | 1,13 | 8,0 | " | " | " |
| 10 | Spring No. 4 & 5 (") | — | 56,0° | — | 0,77 | 9,7 | " | " 2, " | " |
| 11 | Kidachiya-no-yu | carbonated | 55,0 | 130 | 0,38 | 10,2 | " | " 3, " | " |

YUGANO

Location.—Kami-kawazu-mura, Kamo-gun, Province Izu, Shizuoka-ken.

The springs issue from the Andesite.

Table 55.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-------------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Yugano-Onsen (<i>Public bath</i>) | salt | 52,0° | ca.
270 | 0,21 | 12,0° | K. & L. | March 4, 1913 | Y. Kinugasa |
| 2 | Konabe-Onsen („) | „ | 42,0° | 126 | 0,17 | 12,2° | „ | „ | „ |

YATSU

Location.—Shimo-kawazu-mura, Kamo-gun, Province Izu, Shizuoka-ken.

The springs issue from the Tertiary.

Table 56.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---|----------------|--------------------------|---------------------------------|--------------------|--------------------|----------------------------------|-------------|
| | | | | in
Mache's units | at
temp. in C.) | | | |
| 1 | Yatsu-Kyōdō-Onsen
(<i>Public bath</i>) | salt | 56,0° | 0,75 | 15,0° | K. & L. | March 5, 1913 | Y. Kinugasa |
| 2 | Ishida-Onsen | „ | 46,0° | 0,47 | 15,8° | „ | „ | „ |
| 3 | Ishida-shin-Onsen | „ | 52,0° | 0,80 | 12,0° | „ | „ | „ |
| 4 | Mageya-no-yu | „ | 50,0° | 0,35 | 13,8° | „ | „ | „ |
| 5 | Yamada-Onsen | „ | 70,0° | 0,36 | 14,0° | „ | „ | „ |
| 6 | Misawa-no-yu | „ | 65,0° | 0,42 | 13,3° | „ | „ | „ |

RENDAIJI

Location.—Inōzawa-mura Kamo-gun Province Izu, Shizuoka-ken.

The springs issue from the Tertiary.

Table 57.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--|----------------|--------------------------|---------------------------------|-------------------|--------------------|----------------------------------|-------------|
| | | | | in
Maché's units | at
temp. in C. | | | |
| 1 | Kami-fujiwara no-yu (<i>Public bath</i>) | salt | 53.0° | 0.43 | 15.0° | K. & I. | March 7, 1913 | Y. Kinugasa |
| 2 | Shimo-fujiwara-no-yu (") | " | 50.0° | 0.44 | 14.7° | " | " | " |
| 3 | Bath of Zuyō middle school | " | 47.0° | 0.27 | 15.2° | " | " | " |
| 4 | Junji-kwan-no-yu | " | 53.0° | 1.27 | 14.6° | " | " | " |
| 5 | Kakezukaya-no-yu | " | 45.0° | 0.36 | 14.8° | " | " 6, " | " |
| 6 | Aizu-kwan-no-yu | " | 43.0° | 0.72 | 16.1° | " | " | " |
| 7 | Shin-yoshirō-no-yu | " | 42.0° | 0.44 | 17.6° | " | " | " |
| 8 | Idoya-no-yu | " | 48.0° | 1.97 | 12.4° | " | " 8, " | " |
| 9 | Private bath (<i>C. Murayama</i>) | " | 53.0° | 0.50 | 13.5° | " | " | " |
| 10 | " (<i>I. Yoshimura</i>) | " | 45.0° | 0.31 | 17.0° | " | " 6, " | " |
| 11 | " (<i>R. Ogawa</i>) | " | 46.0° | 0.34 | 14.2° | " | " 7, " | " |
| 12 | " (<i>E. Ishibashi</i>) | " | 47.0° | 0.43 | 13.0° | " | " | " |
| 13 | " (<i>K. Murayama</i>) | " | 47.0° | 0.37 | 13.0° | " | " | " |
| 14 | Kami-no-yu (<i>Public bath</i>) | " | 30.0° | 0.42 | 15.7° | " | " | " |
| 15 | Shimo-no-yu (") | " | 48.0° | 0.52 | 13.7° | " | " | " |
| 16 | Fuji-no-yu * | " | 31.0° | 0.25 | 10.8° | " | " 8, " | " |
| 17 | Akama-Onsen No. 1 * | — | 29.0° | 0.35 | 9.9° | " | " | " |
| 18 | " No. 2 * | — | 22.0° | 0.33 | 9.2° | " | " | " |

* These springs, located in *Shimoda*, about 2.5 m. from Rendaiji are grouped for convenience in this table.

SHIMOGAMO

Location.—Minami-naka-mura, Kamo-gun, Province Izu, Shizuoka-ken.

The springs issue from the Tertiary.

Table 58.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------------------------|----------------|--------------------------|---------------------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kanō-Kyōdō-yu (<i>Public bath</i>) | Salt | 71.0° | 0.53 | 13.6° | K. & L. | March 9, 1913 | Y. Kinugasa |
| 2 | Hizume-Kyōdō-yu („) | „ | 79.0° | 0.89 | 12.8° | „ | „ | „ |
| 3 | Shimogamo-Kyōdō-yu („) | „ | 77.0° | 0.29 | 11.5° | „ | „ | „ |
| 4 | Seitō-Onsen (<i>M. Takahashi</i>) | „ | 63.0° | 2.62 | 11.2° | „ | „ | „ |
| 5 | Kinokuniya-no-yu | „ | 73.0° | 0.25 | 11.5° | „ | „ | „ |

TOHI

Location.—Tohi-mura, Takata-gun, Province Izu, Shizuoka-ken.

The springs issue from the Tertiary.

The hot spring “Ō-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1904

Specific gravity: 1.001 at 17° C.

Total residue: ca. 1.37.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols. | Milligram-
equivalents |
|--|--------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0079 | 0.2018 | 0.2018 |
| Sodium ion (Na ⁺) | 0.1360 | 5.9002 | 5.9002 |
| Calcium ion (Ca ⁺⁺) | 0.2624 | 6.5436 | 13.0872 |
| | | | 19.1892 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.0685 | 1.9323 | 1.9323 |
| Sulphate ion (SO ₄ ^{''}) | 0.8290 | 8.6299 | 17.2598 |
| | | 1.3038 | 23.2078 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0506 | | |
| | | | 1.3544 |

| | Grams |
|--|--------|
| Potassium chloride (KCl) | 0.0150 |
| Sodium chloride (NaCl) | 0.1013 |
| Sodium sulphate (Na ₂ SO ₄) | 0.2967 |
| Calcium sulphate (CaSO ₄) | 0.8908 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0506 |
| | 1.3544 |

Besides these trace of magnesium, ferrous, aluminium and hydrophosphate ion.

The spring thus may be classified as “sulphated bitter spring”.

The hot spring “Ana-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1904.

Specific gravity: 1.001 at 17° C.

Temperature: 48° C.

Flow of water: ca. 194 hectolitres in 24 hours.

Total residue: ca. 1.62.

The spring, being in its composition nearly the same as that of “Ō-yu,” may be classified as “sulphated bitter spring”.

Radio-activity. 0.48 Mache's units in 1 litre of water at 19.5° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 18, 1913)

The hot spring "Tohi-kwan-no-yu" No. 1

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1910.

Temperature: 53.5° C.

Specific gravity: 1.0018 at 12.5° C.

Total residue: ca. 1.41.

The spring, being in its composition nearly the same as that of "O-yu", may be classified as "**sulphated bitter spring**".

Radio-activity. 0.40 Mache's units in 1 litre of water at 16.0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 18, 1913).

The hot spring "Tohi-kwan-no-yu" No. 2

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1910.

Temperature: 56° C.

Specific gravity: 1.0022 at 10° C.

Total residue: ca. 1.69.

The spring, being in its composition nearly the same as that of "O-yu", may be classified as "**sulphated bitter spring**".

Radio-activity. 0.85 Mache's units in 1 litre of water at 15.3° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, March 18, 1913).

Table 59.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|------------------------------------|------------------|--------------------------|---|---------------------------------|-------------------|--------------------|----------------------------------|-------------|
| | | | | | in
Mache's units | at
temp. in C. | | | |
| 1 | O-yu <i>Public bath Vent No. 1</i> | sulphated bitter | 46.0° | ca. — | 0.64 | 16.7° | K. & L. | March 17, 1913 | Y. Kinugasa |
| 2 | " <i>Vent No. 2</i> | " | 57.0° | — | 0.47 | 15.3° | " | " | " |
| 3 | Kyōdō-yu <i>Suzuki Public bath</i> | " | 55.0° | — | 0.29 | 16.5° | " | " 18. " | " |
| 4 | Hamano-Kyōdō-yu (") | " | 55.0° | — | 0.73 | 17.5° | " | " | " |
| 5 | Ana-yu (") | " | 48.0° | 104 | 0.48 | 19.5° | " | " | " |
| 6 | Tohi kwan-no-yu <i>No. 1</i> | " | 53.5° | — | 0.40 | 16.0° | " | " | " |
| 7 | " <i>No. 2</i> | " | 56.0° | — | 0.85 | 15.3° | " | " | " |
| 8 | Takasagoya-no-yu | " | 56.0° | — | 0.28 | 14.3° | " | " 17. " | " |
| 9 | Asaka-no-yu <i>No. 1</i> | " | 55.5° | — | 0.74 | 17.4° | " | " | " |
| 10 | " <i>No. 2</i> | " | 52.0° | — | 0.20 | 17.0° | " | " 18. " | " |
| 11 | Private bath <i>M. Ichiwara</i> | " | 43.0° | — | 0.25 | 18.7° | " | " | " |
| 12 | " <i>S. Asaka</i> | " | 52.0° | — | 0.47 | 17.0° | " | " | " |
| 13 | " <i>T. Yoshimura</i> | " | 55.0° | — | 0.23 | 17.6° | " | " | " |
| 14 | " <i>(T. Torizawa)</i> | " | 66.0° | — | 0.69 | 21.9° | " | " | " |
| 15 | " <i>Z. Yoda</i> | " | 57.0° | — | 0.27 | 17.2° | " | " | " |
| 16 | " <i>K. Hasegawa</i> | " | 51.5° | — | 0.40 | 18.0° | " | " | " |

YUMURA

Location.—Ōmiya-mura, Nishi-yamanashi-gun, Province Kai, Yamanashi-ken.

Table 60.

| No. | Spring | Geology | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-------------------------------|---|----------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Tani-Onsen <i>Public bath</i> | Alluvium | salt | 35.5° | ca.
— | 5.17 | 1.42 | 33.0° | Schm. | March 19, 1914 | Y. Kinugasa |
| 2 | Washi-Onsen " | Angite Andesite
and its
Agglomerate | " | 42.0° | 313 | — | 1.80 | 16.0° | K. & L. | " 18, " | " |
| 3 | Kōbōzue-Onsen | " | " | 33.8° | — | — | 1.86 | 14.8° | " | " " | " |
| 4 | Meiji-Onsen | — | " | 38.0° | — | — | 2.41 | 12.9° | " | " 19, " | " |

KAMI-SEKISUIJI

Location.—Aikawa-mura, Nishi-yamanashi-gun, Province Kai, Yamanashi-ken.

Table 61.

| No. | Spring | Geology | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-------------|---|----------------|--------------------------|---|---------------------------------|---------------------|--|--------------------|----------------------------------|-------------|
| | | | | | | in
Mache's units | at
(temp. in C.) | | | | |
| 1 | Yōsō-Onsen | Granite | — | 10.5° | ca.
— | 2.95 | 14.4° | | K. & L. | March 20, 1914 | Y. Kinugasa |
| 2 | Yōgai-Onsen | Angite Andesite
and its
Agglomerate | simple | 11.0° | 36 | 1.55 | 15.2° | | " | " 21, " | " |

SHIMOBE

Location.—Tomisato-mura, Nishi-yatsushiro-gun, Province Kai, Yamanashi-ken.

The springs issue from the Tertiary Shale.

The cold spring "Shimobe-Onsen"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 0.9998 at 19° C.

Temperature: 35° C.

Total residue: ca. 0.46.

Flow of water: ca. 6480 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milli-gram-equivalents |
|--|---------|------------|------------------------|
| Potassium ion (K ⁺) | 0.00024 | 0.00613 | 0.00613 |
| Sodium ion (Na ⁺) | 0.08059 | 3.49631 | 3.49631 |
| Ammonium ion (NH ₄ ⁺) | 0.00021 | 0.01162 | 0.01162 |
| Calcium ion (Ca ⁺⁺) | 0.05714 | 1.42850 | 2.85700 |
| Magnesium ion (Mg ⁺⁺) | 0.00031 | 0.01273 | 0.02546 |
| Ferrous ion (Fe ⁺⁺) | 0.00140 | 0.02500 | 0.05000 |
| | | | 0.44652 |

Anions.

| | | | |
|--|---------|---------|---------|
| Chlorine ion (Cl ⁻) | 0.06676 | 1.88322 | 1.88322 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.20531 | 2.13731 | 4.27462 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.01759 | 0.28831 | 0.28831 |
| | | 0.42955 | 9.28913 |
| Boric acid (meta) (HBO ₂) | 0.01419 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.03052 | | |
| | 0.47426 | | |
| Free carbon dioxide (CO ₂) | 0.00730 | 0.16591 | |
| | 0.48156 | | |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.00046 |
| Sodium chloride (NaCl) | 0.10913 |
| Ammonium chloride (NH ₄ Cl) | 0.00062 |
| Sodium sulphate (Na ₂ SO ₄) | 0.11591 |
| Calcium sulphate (CaSO ₄) | 0.17987 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.01725 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.00155 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.00445 |
| Boric acid (meta) (HBO ₂) | 0.01419 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.03052 |
| | 0.47375 |
| Free carbon dioxide (CO ₂) | 0.00730 |
| | 0.48125 |

The spring thus may be classified as "simple cold spring".

Table 62.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water * | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|---------------------|----------------|-----------------------|---|--------------------------------|------------------|------------------|--------------------|-------------------------------|-------------|
| | | | | | in 10 ⁻¹⁰ curies | in Mache's units | at (temp. in C.) | | | |
| 1 | Shimobe-Onsen No. 1 | simple | 35.0° | ca. 6480 | 0.85 | 0.23 | 21.5° | Schm. | May 25, 1914 | H. Kakehi |
| 2 | " No. 2 | " | 35.0° | 6480 | 0.48 | 0.13 | 27.0° | " | Aug. 1, " | M. Komori |
| 3 | " No. 3 | " | 36.0° | 6480 | 0.27 | 0.07 | 28.0° | " | " | " |

* Measurement was made with the sample brought to the laboratory by applicant, taking into account decay of radium emanation with the time.

MASUTOMI

Location.—Masutomi-mura, Kita-koma-gun, Province Kai, Yamanashi-ken.

The springs issue from the Granite.

The cold spring "Nibuzawa" No. 1

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0070 at 10° C.

Temperature: 23.5° C. (air temp. 4.5° C.).

Total residue: ca. 7.41.

Flow of water: ca. 77 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.13765 | 3.51596 | 3.51596 |
| Sodium ion (Na ⁺) | 2.43382 | 105.58872 | 105.58872 |
| Ammonium ion (NH ₄ ⁺) | 0.00106 | 0.05863 | 0.05863 |
| Calcium ion (Ca ⁺⁺) | 0.23677 | 5.91925 | 11.83850 |
| Magnesium ion (Mg ⁺⁺) | 0.02085 | 0.85591 | 1.71182 |
| Ferrous ion (Fe ⁺⁺) | 0.00140 | 0.02500 | 0.05000 |
| | | | 122.76363 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 3.14674 | 88.76587 | 88.76587 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.46988 | 4.89153 | 9.78306 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 1.47765 | 24.21980 | 24.21980 |
| | 7.92582 | 233.84067 | 122.76873 |
| Boric acid (meta) (HBO ₂) | 0.16910 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.16762 | | |
| | 8.26254 | | |
| Free carbon dioxide (CO ₂) | 0.26432 | 6.00727 | |
| | 8.52686 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.2623 |
| Sodium chloride (NaCl) | 4.9835 |
| Ammonium chloride (NH ₄ Cl) | 0.0031 |
| Sodium sulphate (Na ₂ SO ₄) | 0.6954 |
| Sodium bicarbonate (NaHCO ₃) | 0.8927 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.9590 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.1253 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0045 |
| Boric acid (meta) (HBO ₂) | 0.1691 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1676 |
| | 8.2625 |
| Free carbon dioxide (CO ₂) | 0.2643 |
| | 8.5268 |

The spring thus may be classified as "earthy common salt spring".

Radio-activity. 1433.17×10^{-10} curies in 1 litre of water at 15.0° C.

(Schmidt's electrom.; Dr. R. Ishizu, Dec. 6, 1913).

394.12 Mache's units (recalculated).

The cold spring "Kamigawara" No. 1

Analysis.

Analysed by the Hygienic Laboratory of Yamanashi-ken. 1914. Temperature: 21.5° C. (air temp. 23.5° C.).

Total residue: ca. 5.98.

Flow of water: ca. 22 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Nibuzawa" No. 1, may be classified as "earthy common salt spring".

Radio-activity. 3012.14×10^{-10} curies in 1 litre of water at 21.5° C.

(Schmidt's electrom.; Y. Kinugasa, Sept. 9, 1914).

828.34 Mache's units (recalculated).

The cold spring "Yunokubo-gawara-no-yu"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914. Specific gravity: 1.0066 at 23° C.

Temperature: 15.0° C. (air temp. 10.5° C.).

Total residue: ca. 9.86.

The spring, being in its composition nearly the same as that of "Nibuzawa" No. 1, may be classified as "earthy common salt spring".

Radio-activity. 883.39×10^{-10} curies in 1 litre of water at 12.0° C.

(Schmidt's electrom.; Y. Kinugasa, March 23, 1914).

243.82 Mache's units (recalculated).

1871.31×10^{-10} curies in 1 litre of gas at 0° C.

(Schmidt's electrom.; Y. Kinugasa, April 3, 1914).

514.61 Mache's units (recalculated).

The cold spring "Tsugane-yu" No. 1 (A)

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0070 at 11° C.

Temperature: 17.5° C. (air temp. 7° C.).

Total residue: ca. 7.94.

Flow of water: ca. 45 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Nibuzawa" No. 1, may be classified as "earthy common salt spring".

Radio-activity. 856.83×10^{-10} curies in 1 litre of water at 14.5° C.

(Schmidt's electrom.; Y. Kinugasa, April 1, 1914).

235.63 Mache's units (recalculated).

The cold spring "Kuridaira" No. 1

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0028 at 10° C.

Temperature: 16.0° C. (air temp. 7.0° C.).

Total residue: ca. 2.18.

Flow of water: ca. 40 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Nibuzawa" No. 1, differing only in the smaller proportion of hydrocarbonates of alkaline-earth metals from the latter, may be classified as "common salt spring".

Radio-activity. 2147.05×10^{-10} curies in 1 litre of water at 5.0° C.

(Schmidt's electrom.; Dr. R. Ishizu, Dec. 7, 1913).

590.44 Mache's units (recalculated).

5506.30×10^{-10} curies in 1 litre of gas at 0° C.

(Schmidt's electrom.; Dr. R. Ishizu, Dec. 8, 1913).

1514.23 Mache's units (recalculated).

The cold spring "Tochikubo" No. 1

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0085 at 11° C.

Temperature: 24.5° C. (air temp. 9.0° C.).

Total residue: ca. 9.20.

Flow of water: ca. 66 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of "Nibuzawa" No. 1, may be classified as "earthy common salt spring".

Radio-activity. 25.87×10^{-10} curies in 1 litre of water at 14.0° C.

(Schmidt's electrom.; Dr. R. Ishizu, Dec. 9, 1913).

7.11 Mache's units (recalculated).

86.41×10^{-10} curies in 1 litre of gas at 0° C.

(Schmidt's electrom.; Dr. R. Ishizu, Dec. 13, 1913).

23.76 Mache's units (recalculated).

Composition of the gas (analysed in 1914).

1000 c.c. of gas* evolving from the spring consist of:

| | Kuridaira No. 1 | Tochikubo No. 1 |
|---|-----------------|-----------------|
| Carbon dioxide (CO ₂) | 870 c.c. | 960 c.c. |
| Oxygen (O ₂). | 10 " | 10 " |
| The rest | not determined | not determined |

* The analysis is not accurate, the numbers giving only the approximate value.

Discharge of Radium Emanation.

| No. | Spring | Per minute
Emanation | | | Per hour
Emanation | | | Per day (24 hours)
Emanation | | |
|-----|----------------------|----------------------------|-------------------------------|---------------------|----------------------------|-------------------------------|--|---------------------------------|--|--|
| | | Flow of water
in litres | in
10 ⁻⁸ curies | in
Mache's units | Flow of water
in litres | in
10 ⁻⁸ curies | in
10 ³ Mache's
units | Flow of water
in hectolitres | in
10 ⁻⁶ curies
(microcuries) | in
10 ³ Mache's
units |
| 1 | Kamigawara No. 1 | 1,5 | 46 | 1266 | 92 | 2761 | 76 | 22 | 663 | 1822 |
| 2 | Kuridaira No. 1 | 2,8 | 60 | 1640 | 167 | 3578 | 98 | 40 | 859 | 2362 |
| 3 | " No. 1 Vent A | 7,0 | 80 | 2210 | 417 | 4822 | 133 | 100 | 1157 | 3183 |
| 4 | " No. 1 Vent B | 7,0 | — | 2027 | 417 | — | 122 | 100 | — | 2919 |
| 5 | " No. 1 Vent C | 0,9 | — | 370 | 54 | — | 22 | 13 | — | 532 |
| 6 | " No. 3 | 0,9 | 9 | 259 | 54 | 565 | 16 | 13 | 136 | 373 |
| 7 | Tsugane-yu No. 1 (A) | 3,1 | 27 | 736 | 188 | 1607 | 44 | 45 | 386 | 1060 |
| 8 | " No. 1 (B) | 0,9 | — | 78 | 54 | — | 5 | 13 | — | 112 |
| 9 | Nibuzawa No. 1 | 5,4 | 77 | 2108 | 321 | 4598 | 127 | 77 | 1104 | 3035 |

Table 63.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|------------------------------|--------------------------|----------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|---------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Ōshiba No. 1 (I. Ōshiba) | — | air temp.
19,0° (14,5°) | ca. | 1008,76 | 277,41 | 8,5° | Schm. | March 26, 1914 | Y. Kinugasa |
| 2 | " | — | " | — | — | 298,33 | 17,4° | K. & L. | " 24, " | " |
| 3 | " No. 2 (T. Fujiwara) | simple | 14,5° (14°) | — | — | 1,85 | 15,7° | " | " | " |
| 4 | Ginsen-tō | — | 29,5° (7°) | — | 372,63 | 102,47 | 23,0° | Schm. | Dec. 10, 1913 | Dr. R. Ishizu |
| 5 | " | — | 30,0° (14°) | — | — | 109,96 | 15,4° | K. & L. | March 24, 1914 | Y. Kinugasa |
| 6 | Kamigawara No. 1 (K. Fujita) | earthy
common
salt | 21,5° (23,5°) | 22 | 3012,14 | 828,34 | 21,5° | Schm. | Sept. 9, " | " |
| 7 | " No. 2 (") | " | 14,0° | — | 2390,02 | 657,26 | 8,5° | " | March 28, " | " |
| 8 | Kuridaira No. 1 (S. Haki) | common
salt | 16,0° (7°) | 40 | 2147,05 | 590,44 | 5,0° | " | Dec. 7, 1913 | Dr. R. Ishizu |
| 9 | " No. 1 Vent A (") | " | 20,0° (4°) | 100 | 1157,29 | 318,25 | 13,3° | " | March 31, 1914 | Y. Kinugasa |
| 10 | " | " | " | " | — | 388,14 | 7,8° | K. & L. | " 27, " | " |
| 11 | " No. 1 Vent C (") | " | 21,5° (4°) | 13 | — | 409,31 | 7,6° | " | " | " |
| 12 | " No. 1 Vent B (") | " | 20,0° | 100 | — | 291,93 | 8,7° | " | " 28, " | " |
| 13 | " No. 2 (A) (") | — | 26,0° (17°) | 15 | 7,42 | 2,04 | 14,5° | Schm. | " 25, " | " |
| 14 | " No. 2 (C) (") | — | 20,0° (17°) | 11 | — | 1,92 | 14,0° | K. & L. | " | " |
| 15 | " No. 2 (D) (") | — | 20,0° (9,5°) | 41 | 6,79 | 1,87 | 8,0° | Schm. | Dec. 11, 1913 | Dr. R. Ishizu |

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|--------------------|------------------------------------|--------------------|--------------------------|---|---------------------------------|---------------------|---------------------------------------|--------------------|----------------------------------|---------------|
| | | | | | in
milli-
curies | in
Maché's units | at
temp. in C. | | | |
| 16 | Kuridaira No. 3 (S. Hakii) | — | 20,0° (10,5°) | ca. 13 | 1042,17 | 286,60 | 12,0° | Schm. | March 30, 1914 | Y. Kinugasa |
| 17 | Tsugane-yu No. 1 (A) | earthy common salt | 17,5° (7°) | 45 | 856,83 | 235,63 | 14,5° | " | Apr. 1, " | " |
| 18 | " | " | " | " | — | 240,20 | 9,6° | K. & L. | March 26, " | " |
| 19 | " No. 1 (B) | " | 19,0° (7°) | 13 | — | 86,27 | 9,1° | " | " | " |
| 20 | " No. 1 (C) | " | 16,0° (7°) | — | — | 83,35 | 9,2° | " | " | " |
| 21 | Yunokubo No. 1
(M. Shirakura) | — | 17,0° | — | 900,12 | 247,53 | 7,0° | Schm. | " 27, " | " |
| 22 | " | — | 18,0° | — | — | 298,22 | 14,1° | K. & L. | " 25, " | " |
| 23 | Yunokubo-gawara-no-yu
(K. Arii) | earthy common salt | 15,0° (10,5°) | — | 883,39 | 243,82 | 12,0° | Schm. | " 23, " | " |
| 24 | " | " | " | — | — | 239,22 | 16,0° | K. & L. | " | " |
| 25 | Tochikubo No. 1 (S. Hakii) | " | 24,5° (9°) | 66 | 25,87 | 7,11 | 14,0° | Schm. | Dec. 9, 1913 | Dr. R. Ishizu |
| 26 | Hatchō-jyaya-shita No. 1 | — | 18,5° | — | — | 146,97 | 7,2° | K. & L. | March 27, 1914 | Y. Kinugasa |
| 27 | Yunomukai No. 1 | — | 10,0° | — | — | 75,62 | 11,8° | " | " 31, " | " |
| 28 | Umamichi-zawa No. 1 | — | 12,0° | — | — | 2,86 | 11,2° | " | " | " |
| 29 | Yunosawa No. 1 | — | 17,0° (6°) | — | 5,76 | 1,58 | 8,0° | Schm. | Dec. 5, 1913 | Dr. R. Ishizu |
| 30 | Kinsen-tō | — | 31,5° (5,5°) | 362 | 7,58 | 2,08 | 7,5° | " | " 12, " | " |
| 31 | " | — | 33,0° | " | — | 2,29 | 17,0° | K. & L. | Apr. 2, 1914 | Y. Kinugasa |
| 32 | Nibuzawa No. 1 (Z. Gotō) | earthy common salt | 23,5° (4,5°) | 77 | 1433,17 | 394,12 | 15,0° | Schm. | Dec. 6, 1913 | Dr. R. Ishizu |
| 33 | " | " | 22,5° (4°) | " | — | 312,54 | 9,6° | K. & L. | March 28, 1914 | Y. Kinugasa |
| 34 | Taikaizawa No. 1 | — | 8,2° | — | — | 6,54 | 7,0° | " | " 29, " | " |
| 35 | " No. 2 | — | 8,2° | — | — | 8,77 | 7,2° | " | " | " |
| 36 | " No. 3 | — | 4,5° | — | — | 8,42 | 7,4° | " | " | " |
| 37 | Matsudaira No. 1 | — | 3,9° | — | — | 2,12 | 8,2° | " | " | " |
| 38 | Toyanosawa No. 1 | — | 9,0° | — | — | 4,03 | 8,9° | " | " | " |
| 39 | Akashibu No. 1 | — | 13,0° | — | — | 2,61 | 14,9° | " | Apr. 1, " | " |
| 40 | " No. 2 | — | 12,0° | — | — | 2,97 | 15,6° | " | " | " |
| 41 | " No. 3 | — | 10,5° | — | — | 3,15 | 15,2° | " | " | " |
| 42 | Itaya No. 1 | — | 14,0° | — | — | 2,92 | 14,0° | " | " | " |
| 43 | " No. 2 | — | 12,0° | — | — | 2,13 | 15,0° | " | " | " |
| Gas evolving from: | | | | | | | | | | |
| | | | | Emanation per litre
of gas at 0° C. | | | Date when
the gas was
collected | | | |
| 1 | Kamigawara No. 2 (K. Fujita) | — | — | — | 2037,42 | 560,29 | — | Schm. | Apr. 2, 1914 | " |
| 2 | Yunokubo-gawara-no-yu
(K. Arii) | — | — | — | 1871,31 | 514,61 | — | " | " 3, " | " |
| 3 | " | — | — | — | — | 539,99 | — | K. & L. | March 31, " | " |
| 4 | Tochikubo No. 1 (S. Hakii) | — | — | — | 86,41 | 23,76 | — | Schm. | Dec. 13, 1913 | Dr. R. Ishizu |
| 5 | Kuridaira No. 1 | — | — | — | 5506,30 | 1514,23 | — | " | " 8, " | " |

SHIBU

Location.—Kitayama-mura, Suwa-gun, Province Shinano, Nagano-ken.

The springs issue from the Andesite.

The cold spring “Shibu-no-yu”

Analysis.

Analysed by Dr. R. Ishizu. 1904.
Total residue: ca. 0.86.
Flow of water: abundant.

Specific gravity: 1.0003 at 15° C.
Temperature: 27° C.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Potassium chloride (KCl) | 0.0275 | Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.0976 |
| Sodium chloride (NaCl) | 0.0923 | Free sulphuric acid (H ₂ SO ₄) | 0.6692 |
| Potassium sulphate (K ₂ SO ₄) | 0.1624 | Silica (SiO ₂) | 0.0890 |
| Calcium sulphate (CaSO ₄) | 0.0326 | | 1.1854 |
| Magnesium sulphate (MgSO ₄) | 0.0046 | Free hydrogen sulphide (H ₂ S) | 0.0167 |
| Ferrous sulphate (FeSO ₄) | 0.0102 | | 1.2021 |

The spring may be classified as “acid hydrogen sulphide spring”.

Radio-activity. 1.22×10^{-10} curies in 1 litre of water* at 23° C.

(Schmidt's electrom.; M. Komori, Oct. 6, 1914).

0.34 Mache's units (recalculated).

Composition of the mineral salt “Hana-jio” (analysed in 1904).

100 parts of the brownish crystalline crust, weak acid and saline in taste, consist of:

| | | | |
|--|--------------|---|--------------|
| Potassium chloride (KCl) | 0.4167 parts | Ferrous sulphate (FeSO ₄) | 9.5390 parts |
| Sodium chloride (NaCl) | 0.1053 „ | Aluminium sulphate [Al ₂ (SO ₄) ₃] | 16.1674 „ |
| Sodium sulphate (Na ₂ SO ₄) | 15.4596 „ | Sulphuric acid (H ₂ SO ₄) | 2.2395 „ |
| Calcium sulphate (CaSO ₄) | 1.2250 „ | Silica (SiO ₂) | 1.2504 „ |
| Magnesium sulphate (MgSO ₄) | 0.7968 „ | Water | 47.7850 „ |

Composition of the sinter-deposit “Yu-no-hana” (analysed in 1904).

100 parts of the yellowish-white powder, acid in taste, consist of:

| | | | |
|--|---------------|--|--------------|
| Sulphur (S) | 77.9959 parts | Calcium sulphate (CaSO ₄) | 2.8422 parts |
| Potassium sulphate (K ₂ SO ₄) | 2.1258 „ | Ferric oxide (Fe ₂ O ₃) | 2.7703 „ |

Besides these a small quantity of magnesia, alumina, chlorine, silica, etc.

SUWA

Location.—Suwa-gun, Province Shinano, Nagano-ken.

The springs issue from the Alluvium.

The hot spring “Suwa-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1890.
Specific gravity: 1.00195 at 15° C.

Temperature: 64.7° C.
Total residue: ca. 1.06.

* Measurement was made with the sample brought to the laboratory by applicant, taking into account decay of radium emanation with the time.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0,0163 | 0,4163 | 0,4163 |
| Sodium ion (Na ⁺) | 0,2451 | 10,6334 | 10,6334 |
| Calcium ion (Ca ⁺⁺) | 0,0738 | 1,8404 | 3,6808 |
| | | | 14,7305 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0,2443 | 6,8014 | 6,8914 |
| Sulphate ion (SO ₄ ^{''}) | 0,3765 | 3,0104 | 7,8388 |
| | 0,9560 | 23,7009 | 14,7302 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0681 | | |
| | 1,0241 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Gram |
|--|--------|
| Sodium chloride (NaCl) | 0,4031 |
| Potassium sulphate (K ₂ SO ₄) | 0,064 |
| Sodium sulphate (Na ₂ SO ₄) | 0,2664 |
| Calcium sulphate (CaSO ₄) | 0,2501 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0681 |
| | 1,0241 |

Besides these trace of hydrophosphate ion and boric acid.

The spring thus may be classified as “**weak muriated saline bitter spring**”.

Table 64.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water in Maché's units at temp. in C. | Apparatus employed | Date when the water was taken | Examined by |
|-----|---------------|----------------|-----------------------|---|--|--------------------|-------------------------------|-------------|
| 1 | Public bath : | | | ca. | | | | |
| 1 | Shōjin-yu | simple | 67,5° | 130 | 1,29 5,4° | K & L. | Apr. 6, 1914 | Y. Kinugasa |
| 2 | Hira-no-yu | .. | 59,0° | 130 | 0,94 6,4° | .. | .. | .. |
| 3 | Tajiku-no-yu | .. | 83,0° | 130 | 1,13 6,0° | .. | .. | .. |
| 4 | Wata-no-yu | .. | 47,5° | 2700 | 1,61 6,0° | .. | .. 7, .. | .. |
| 5 | Ko-yu | .. | 52,0° | 1800 | 0,92 7,0° | .. | .. | .. |
| 6 | Tankwa-no-yu | .. | 67,0° | 3600 | 1,32 6,6° | .. | .. | .. |

ASAMA

Location.—Hongō-mura, Higashi-chikuma-gun, Province Shinano, Nagano-ken.

The springs issue from the Tertiary.

The hot spring “Asama-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1893.

Specific gravity: 1,00029 at 15° C.

Total residue: ca. 0,44.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|---|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0,0094 | 0,2401 | 0,2401 |
| Sodium ion (Na ⁺) | 0,0880 | 3,8178 | 3,8178 |
| Calcium ion (Ca ⁺⁺) | 0,0371 | 0,9252 | 1,8504 |
| | | | 5,9085 |

| | Grams |
|--|--------|
| Sodium chloride (NaCl) | 0,0475 |
| Potassium sulphate (K ₂ SO ₄) | 0,0209 |
| Sodium sulphate (Na ₂ SO ₄) | 0,2139 |
| Calcium sulphate (CaSO ₄) | 0,1258 |

Anions.

| | | | | | |
|--|--------|--------|--------|--|--------|
| Chlorine ion (Cl') | 0,0288 | 0,8124 | 0,8124 | Silicic acid (meta) (H ₂ SiO ₃) | 0,0491 |
| Sulphate ion (SO ₄ '') | 0,2448 | 2,5484 | 5,0968 | | 0,4572 |
| | 0,4081 | 8,3439 | 5,9092 | | |
| Silicic acid (meta) (H ₂ SiO ₃) . . . | 0,0491 | | | | |
| | 0,4572 | | | | |

Besides these a small quantity of ferrous and aluminium ion, trace of magnesium and hydrophosphate ion.

The spring thus may be classified as "simple thermal".

Table 65.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------------------|----------------|--------------------------|---------------------------------|---------------------|--------------------|----------------------------------|-------------------------|
| | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Me-no-yu | simple | 36,5° (air temp. 29°) | 2,27 | 27,6° | K. & L. | Aug. 4, 1914 | Y. Kinugasa
T. Saitō |
| 2 | Matsu-no-yu | " | 53,0° (30°) | 0,73 | 27,0° | " | " | " |
| 3 | Taki-no-yu (<i>Public bath</i>) | " | 50,0° (28°) | 0,93 | 27,4° | " | " | " |
| 4 | Kizu-no-yu | " | 44,0° (26°) | 1,24 | 27,4° | " | " | " |

YAMABE

Location.—Sato-yamabe-mura, Higashi-chikuma-gun, Province Shinano, Nagano-ken.

The springs issue from the Tertiary.

The hot spring "Shiraito-Onsen" No. 1

Analysis.

Analyst: Unknown. Specific gravity: 1,0003 at 15° C. Total residue: ca. 0,45.
Flow of water: ca. 130 hectolitres in 24 hours. Temperature: 42° C. (air temp. 30,2° C.).

In 1 kilogram of the mineral water are contained:

| Hypothetical form of combination. | |
|---|--------|
| Grams | Grams |
| Potassium (K) | 0,0314 |
| Sodium (Na) | 0,0208 |
| Calcium (Ca) | 0,0450 |
| Magnesium (Mg) | 0,0250 |
| Chlorine (Cl) | 0,0406 |
| Sulphuric anhydride (SO ₃) | 0,1745 |
| Silica (SiO ₂) | 0,0324 |
| Carbon dioxide (CO ₂) | 0,1000 |
| Potassium sulphate (K ₂ SO ₄) | 0,0542 |
| Sodium sulphate (Na ₂ SO ₄) | 0,0556 |
| Potassium carbonate (K ₂ CO ₃) | 0,0454 |
| Sodium carbonate (Na ₂ CO ₃) | 0,0725 |
| Calcium sulphate (CaSO ₄) | 0,0474 |
| Magnesium sulphate (MgSO ₄) | 0,0642 |
| Silica (SiO ₂) | 0,0324 |
| | 0,3717 |

Besides these a small quantity of phosphoric acid and trace of iron.

The spring may be classified as "simple thermal".

Radio-activity. 1,38 Mache's units in 1 litre of water at 27,8° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 4, 1914).

Table 66.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
temp. in C.) | Apparatus employed | Date when the
water was taken | Examined by |
|-----|----------------------|----------------|----------------------------|---|--|--------------------|----------------------------------|---------------------------|
| 1 | Shiraito-Onsen No. 1 | simple | air temp.
42.0° (30.2°) | ca. 130 | 1.38 27.8° | K. & L. | Aug. 4, 1914 | Y. Kinugasa
(T. Saito) |
| 2 | „ No. 2 | „ | 42.0° (30.2°) | 350 | 1.51 27.8° | „ | „ | „ |
| 3 | Mitarashi-no-yu | — | 28.0° (30°) | — | 1.19 27.3° | „ | „ | „ |
| 4 | Oboke-no-yu | — | 29.0° (30°) | — | 0.58 27.6° | „ | „ | „ |

SHIRAHONE

Location.—Azumi-mura, Minami-azumi-gun, Province Shinano, Nagano-ken.

The hot spring “Ō-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1897.

Temperature: 52° C. (air temp. 24° C.). Flow of water: ca. 236 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0290 | 0.7407 | 0.7407 |
| Sodium ion (Na ⁺) | 0.1284 | 5.5705 | 5.5705 |
| Calcium ion (Ca ⁺⁺) | 0.1403 | 3.4962 | 6.9924 |
| Magnesium ion (Mg ⁺⁺) | 0.0467 | 1.9174 | 3.8348 |
| | | | 17.1384 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.1011 | 2.8519 | 2.8519 |
| Sulphate ion (SO ₄ ^{''}) | 0.0649 | 0.6756 | 1.3512 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.7892 | 12.9353 | 12.9353 |
| | 1.2596 | 28.1876 | 17.1384 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0571 | | |
| | 1.3567 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0552 |
| Sodium chloride (NaCl) | 0.1236 |
| Sodium bicarbonate (NaHCO ₃) | 0.2905 |
| Calcium sulphate (CaSO ₄) | 0.0920 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.4576 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.2807 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0571 |
| | 1.3567 |

Besides these trace of ferrous, aluminium and hydrophosphate ion.

The spring thus may be classified as “**earthy alkaline spring**”.

Radio-activity. 0.37 Mache's units in 1 litre of water at 24.6° C.

1.05 Mache's units in 1 litre of gas at 0° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 6, 1914).

Table 67.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------|-----------------|--------------------------|---|--|---------------------|--------------------|---|---------------------------|
| | | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | O-yu <i>Public bath</i> | earthy alkaline | 52.0°
(air temp. 24°) | ca. 236 | 0.37 | 24.6° | K. & L. | Aug. 6, 1914 | { Y. Kinugasa
T. Saitō |
| 2 | Chi-no-yu | " | 48.0° (26°) | | 0.86 | 26.3° | " | " | " |
| 3 | Senki-no-yu | " | 52.0° (28°) | 236 | 0.57 | 26.0° | " | " | " |
| 4 | Wata-no-yu | " | 51.0° (28°) | 126 | 0.97 | 25.8° | " | " | " |
| | Gas evolving from
O-yu | | — | — | Emanation per litre
of gas at 0° C.
1.05 | | " | Date when the
gas was
collected
Aug. 6, 1914 | " |

KAMIKŌCHI

Location.—Azumi-mura, Minami-azumi-gun, Province Shinano, Nagano-ken.

The spring issues from the Granite.

The hot spring “Kamikōchi-Onsen”

Temperature: 53.5° C. (air temp. 20° C.).

Flow of water: ca. 778 hectolitres in 24 hours.

No accurate analysis has ever been made, but the spring seems to be “simple thermal”.

Radio-activity. 1.28 Mache's units in 1 litre of water at 21.0° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 7, 1914).

NAKABUSA

Location.—Ariake-mura, Minami-azumi-gun, Province Shinano, Nagano-ken.

The springs issue from the Granite.

The hot spring “Shirataki-no-yu”

Analysis.

Analysed by the Imperial Geological Survey.

Specific gravity: 1.00 at 15° C.

Temperature: 59.5° C. (air temp. 29° C.).

Total residue: ca. 0.55.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|--|--------|
| Potassium (K) | 0.0362 | Sulphate ion (SO ₄ '') | 0.0930 |
| Sodium (Na) | 0.0955 | Silica (SiO ₂) | 0.1758 |
| Calcium (Ca) | 0.0110 | Organic substances | 0.0284 |
| Magnesium (Mg) | 0.0002 | Carbon dioxide (CO ₂) | 0.0790 |
| Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃) | 0.0034 | Hydrogen sulphide (H ₂ S) | 0.0090 |
| Chlorine (Cl) | 0.0798 | | |

The spring may be classified as “sulphur spring”.

Radio-activity. 1.85 Mache's units in 1 litre of water at 27.7° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 11, 1914).

The hot spring “Danjō-no-yu”

Analysis.

Analysed by the Imperial Geological Survey.

Temperature: 96° C. (air temp. 28° C.).

Specific gravity: 1.00 at 15° C.

Total residue: ca. 0.71.

The spring, being in its composition nearly the same as that of “Shirataki-no-yu”, may be classified as “alkaline sulphur spring”.

Radio-activity. 0.79 Mache's units in 1 litre of water at 28.1° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 11, 1914).

The hot spring “Taki-no-yu”

Analysis.

Analysed by the Imperial Geological Survey.

Temperature: 95.5° C. (air temp. 28° C.).

Specific gravity: 1.00 at 15° C.

Total residue: ca. 1.14.

The spring, being in its composition nearly the same as that of “Shirataki-no-yu”, may be classified as “alkaline sulphur spring”.

Radio-activity. 0.77 Mache's units in 1 litre of water at 27° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 11, 1914).

The hot spring “Oyogi-no-yu”

Analysis.

Analysed by the Imperial Geological Survey.

Temperature: 74° C. (air temp. 30° C.).

Specific gravity: 0.99 at 15° C.

Total residue: ca. 0.46.

The spring, being in its composition nearly the same as that of “Shirataki-no-yu”, may be classified as “alkaline sulphur spring”.

Radio-activity. 1.28 Mache's units in 1 litre of water at 22.6° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 11, 1914).

Table 68.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water
in
Mache's units
at
(temp. in C.) | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------|------------------|-------------------------------------|---|-------|--------------------|----------------------------------|---------------------------|
| 1 | Yakushi-no-yu No. 1 | alkaline sulphur | 92.0° ^{air temp.}
(27°) | 1.06 | 26.8° | K. & L. | Aug. 11, 1914 | { Y. Kinugasa
T. Saito |
| 2 | „ No. 2 | „ | 93.0° (26.5°) | 0.98 | 27.4 | „ | „ | „ |
| 3 | Danjō-no-yu | „ | 96.0° (28°) | 0.79 | 28.1 | „ | „ | „ |
| 4 | Myōken-no-yu | „ | 96.0° (27.5°) | 1.14 | 27.9 | „ | „ | „ |
| 5 | Taki-no-yu | „ | 95.5 (28°) | 0.77 | 27.0 | „ | „ | „ |
| 6 | Shira-taki-no-yu | sulphur | 59.5 (20°) | 1.85 | 27.7 | „ | „ | „ |
| 7 | Tamura-no-yu | alkaline sulphur | 94.0 (20°) | 1.50 | 23.2 | „ | „ | „ |
| 8 | Oyogi-no-yu | „ | 74.0 (30°) | 1.28 | 22.6 | „ | „ | „ |

KUZU

Location.—Taira-mura, Kita-azumi-gun, Province Shinano, Nagano-ken.

The springs issue from the Granite.

Table 69.

| No. | Spring | Classification | Temp of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------|----------------|-------------------------|---------------------------------|---------------------|--------------------|----------------------------------|---------------------------|
| | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kanatsubo-no-yu | salt | 62.0° (27°) | 1.98 | 26.2° | K. & L. | Aug. 9, 1914 | { Y. Kinugasa
T. Saitō |
| 2 | Gorin-no-yu | " | 63.0° (28°) | 2.01 | 27.0° | " | " | " |
| 3 | Shin-yu No. 2 | " | 57.0° (31°) | 1.45 | 25.0° | " | " | " |
| 4 | Spring A | sulphur (?) | 88.0° (29°) | 1.42 | 26.2° | " | " | " |

HIRAO

Location.—Hirao-mura, Shimo-takai-gun, Province Shinano, Nagano-ken.

YUDANAKA

The springs issue from the Misaka Series.

The hot spring "Ō-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1886. Total residue: ca. 1.53.

Temperature: 76° (air temp. 27.5° C.)

Flow of water: ca. 1703 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | | Milli-
mols | Milligram-
equivalents | | Grams | |
|--|---------|----------|----------------|---------------------------|--|---------|--|
| | | | | | | | |
| Potassium ion (K ⁺) | 0.02735 | 0.69859 | 0.69859 | | Sodium chloride (NaCl) | 0.69534 | |
| Sodium ion (Na ⁺) | 0.27396 | 11.88546 | 11.88546 | | Potassium sulphate (K ₂ SO ₄) | 0.06091 | |
| Calcium ion (Ca ⁺⁺) | 0.19197 | 4.78728 | 9.57456 | | Calcium sulphate (CaSO ₄) | 0.65185 | |
| | | | 22.15861 | | Silicic acid (meta) (H ₂ SiO ₃) | 0.15394 | |
| | | | | | | 1.56204 | |
| Anions. | | | | | | | |
| Chlorine ion (Cl ⁻) | 0.42138 | 11.88660 | 11.88660 | | | | |
| Sulphate ion (SO ₄ ^{''}) | 0.49344 | 5.13679 | 10.27358 | | | | |
| | 1.40810 | 34.39472 | 22.16018 | | | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.15394 | | | | | | |
| | 1.56204 | | | | | | |

Besides these trace of magnésium, ferrous and iodine ion.

The spring thus may be classified as "muriated sulphated bitter spring".

Radio-activity. 1.33 Mache's units in 1 litre of water (taken from Vent No. 1) at 27.4° C.

(Kohl.-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 6, 1914).

ANDAI

The springs issue from the Diluvium.

The hot spring “Ō-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1898.

Temperature: 56° C.

Flow of water: ca. 648 hectolitres in 24 hours.

Total residue: ca. 0.95.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|--------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0188 | 0.4803 | 0.4803 |
| Sodium ion (Na ⁺) | 0.2181 | 9.4577 | 9.4577 |
| Calcium ion (Ca ⁺⁺) | 0.0735 | 1.8329 | 3.6658 |
| | | | 13.6038 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.2746 | 7.7461 | 7.7461 |
| Sulphate ion (SO ₄ ^{''}) | 0.2812 | 2.9273 | 5.8546 |
| | | 0.8662 | 22.4443 |
| Silicic acid (meta) (H ₂ SiO ₃) . . . | 0.1064 | | 13.6007 |
| | 0.9726 | | |

| | |
|--|--------|
| Sodium chloride (NaCl) | 0.4531 |
| Potassium sulphate (K ₂ SO ₄) | 0.0418 |
| Sodium sulphate (Na ₂ SO ₄) | 0.1221 |
| Calcium sulphate (CaSO ₄) | 0.2492 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1064 |
| | 0.9726 |

Besides these trace of magnesium, ferrous, aluminium and hydrophosphate ion.

The spring thus may be classified as “simple thermal”.

Radio-activity. 0.97 Mache's units in 1 litre of water at 27.6° C.

(Kohl-Low. fontact.; Y. Kinugasa and T. Saitō, Sept. 6, 1914).

SHIBU

The springs issue from the Diluvium.

The hot spring “Ō-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1887.

Specific gravity: 1.00203 at 14° C.

Total residue: ca. 1.21.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|--------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0058 | 0.1481 | 0.1481 |
| Sodium ion (Na ⁺) | 0.2639 | 11.4499 | 11.4490 |
| Calcium ion (Ca ⁺⁺) | 0.1052 | 2.6190 | 5.2380 |
| Ferrous ion (Fe ⁺⁺) | 0.0035 | 0.0680 | 0.1360 |
| | | | 19.0711 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.3236 | 7.1273 | 9.1283 |
| Sulphate ion (SO ₄ ^{''}) | 0.3705 | 3.8519 | 7.7138 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0082 | 0.1310 | 0.1310 |
| | 1.0810 | 27.3753 | 19.0731 |
| Silicic acid (meta) (H ₂ SiO ₃) . . . | 0.1752 | | |
| | 1.2562 | | |

| | |
|--|--------|
| Sodium chloride (NaCl) | 0.5340 |
| Potassium sulphate (K ₂ SO ₄) | 0.0130 |
| Sodium sulphate (Na ₂ SO ₄) | 0.1650 |
| Calcium sulphate (CaSO ₄) | 0.3570 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] . . | 0.0120 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1752 |
| | 1.2562 |

Besides these trace of magnesium, aluminium, iodine and hydrophosphate ion, and boric acid.

The spring thus may be classified as “sulphated common salt spring”.

Table 70.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------------------|---------------------------|------------------------------------|---|---------------------------------|---------------------|--------------------|----------------------------------|---------------------------|
| | | | | | in
Mache's units | at
(temp. in C.) | | | |
| | Yudanaka | | | | | | | | |
| 1 | Ō-yu <i>Vent. No. 1</i> | muriated sulphated bitter | 76.0° ^{air temp.} (27.5°) | ca. 1703 | 1.33 | 27.4° | K. & L. | Sept. 6, 1914 | { Y. Kinugasa
T. Saitō |
| 2 | " <i>Vent. No. 2</i> | " | 74.0° (27°) | | 1.27 | 28.2° | " | " | |
| 3 | Washi-no-yu | — | 74.5° (27.5°) | 304 | 1.22 | 27.7° | " | " | |
| | Andai | | | | | | | | |
| 4 | Ō-yu * | simple | 56.0° | 648 | 0.97 | 27.6° | " | " | " |
| 5 | Kaikwa-no-yu | " | 55.0° | 194 | 0.84 | 27.5° | " | " | " |
| | Shibu | | | | | | | | |
| 6 | Ō-yu * <i>Vent. No. 1</i> | sulphated common salt | 74.0° (28°) | — | 1.11 | 28.0° | " | " 5, " | " |
| 7 | " <i>Vent. No. 4</i> | " | 72.0° (27.7°) | — | 1.46 | 27.0° | " | " | " |
| 8 | " <i>Vent. No. 5</i> | " | 70.5° (27.5°) | — | 0.81 | 26.2° | " | " | " |
| 9 | " <i>Vent. No. 8</i> | " | 62.0 (29°) | — | 1.51 | 27.0° | " | " | " |
| 10 | " <i>Vent. No. 12</i> | " | 76.0° (27°) | — | 1.40 | 26.7° | " | " | " |
| 11 | Hatsu-yu * | " | 73.5° (26°) | 519 | 1.46 | 27.1° | " | " | " |
| 12 | Wata-no-yu * | " | 66.0 (26°) | — | 1.20 | 27.6° | " | " 6, " | " |
| 13 | Mearai-no-yu | " | 58.0° (27°) | 519 | 0.82 | 24.5° | " | " | " |
| 14 | Mujina-no-yu | " | 45.0° (26°) | — | 0.91 | 24.7° | " | " | " |
| 15 | Nanakuri-no-yu* <i>Vent. 1</i> | sulphur | 60.0 (24.5°) | | 0.94 | 27.0° | " | " | " |
| | Kambayashi | | | | | | | | |
| 16 | Tsuru-no-yu * | salt | 55.0 | 501 | 0.84 | 26.0° | " | " | " |
| | Kakuma | | | | | | | | |
| 17 | Ō-yu *† | common salt | 65.0° | 735 | 0.82 | 26.0° | " | " | " |
| 18 | Taki-no-yu *† | " | 54.0° | 389 | — | — | — | — | — |
| 19 | Shin-yu *† | " | 52.0° | 389 | — | — | — | — | — |

* These springs are for public use.
† These springs, located at *Hanami-mura*, about 1.2 m. S. from Shibu, are grouped for convenience in this table.

TAKAYAMA

Location.—Fukuoka-mura, Ena-gun, Province Mino, Gifu-ken.

The springs issue from the Granite.

The cold spring "Ena-Kōsen" (Yunoshima)

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory, 1914. Specific gravity 1.001 at 15° C.

Temperature: 10° C. (air temp. 13° C.).

Total residue: ca. 0.12.

In 1 kilogram of the mineral water are contained

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-moles | Milli-equivalents |
|--|---------|-------------|-------------------|
| Potassium ion (K ⁺) | 0.00032 | 0.00817 | 0.00817 |
| Sodium ion (Na ⁺) | 0.02773 | 1.20303 | 1.20303 |
| Ammonium ion (NH ₄ ⁺) | 0.00016 | 0.00885 | 0.00885 |
| Calcium ion (Ca ⁺⁺) | 0.00714 | 0.017850 | 0.035700 |
| Magnesium ion (Mg ⁺⁺) | 0.00022 | 0.00903 | 0.01806 |
| Ferrous ion (Fe ⁺⁺) | 0.00140 | 0.02500 | 0.05000 |
| | | | 1.64511 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.03538 | 0.99803 | 0.99803 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.00123 | 0.01280 | 0.02560 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.03801 | 0.62301 | 0.62301 |
| | 0.11159 | 3.06642 | 1.64664 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.02597 | | |
| | 0.13756 | | |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.00061 |
| Sodium chloride (NaCl) | 0.05739 |
| Ammonium chloride (NH ₄ Cl) | 0.00047 |
| Sodium sulphate (Na ₂ SO ₄) | 0.00182 |
| Sodium bicarbonate (NaHCO ₃) | 0.0132 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.02897 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.00136 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.00445 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.02597 |
| | 0.13756 |

Besides these a small quantity of free carbon dioxide and trace of boric acid.

The spring thus may be classified as "simple cold spring".

Radio-activity. 1022.15×10^{-10} curies in 1 litre of water at 11.0° C.

(Schmidt's electrom.; Y. Kinugasa, April 13, 1914).

281.09 Mache's units (recalculated).

Table 71.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------|----------------|----------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | in
10 ¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Ena-Kōsen (Yunoshima) | simple | 10.0° ^{air temp.} | 1022.15 | 281.09 | 11.0° | Schm. | Apr. 13, 1914 | Y. Kinugasa |
| 2 | " Dairi-yakushi * | " | 12.0° | 220.64 | 60.68 | 13.5° | " | " 15, " | " |
| 3 | " Shimizu-id. | " | 13.0° | 213.21 | 58.63 | 14.7° | " | " 14, " | " |
| 4 | Ichi-no-sawa No. 1 | " | 10.0° (21°) | | 23.70 | 13.7° | K. & L. | " 11, " | " |
| 5 | " No. 2 | " | 12.5° (21°) | — | 2.95 | 13.4° | " | " " | " |
| 6 | " No. 3 | " | 9.5° | — | 17.08 | 13.1° | " | " " | " |
| 7 | Okuhora | " | 13.0° | | 31.90 | 12.7° | " | " " | " |
| 8 | Shimo-issiki (Dairi)* | " | 12.0° (30°) | | 25.77 | 14.0° | " | " 12, " | " |
| 9 | Shimoyama No. 1 | " | 12.0° | | 19.42 | 13.2° | " | " " | " |
| 10 | Yamaki-no-yu | " | 11.0° | | 25.16 | 12.9° | " | " " | " |
| 11 | Inari-no-yu | " | 13.0° | | 30.36 | 13.3° | " | " " | " |
| 12 | Kami-no-ido | " | | | 18.48 | 10.3° | " | " 13, " | " |
| 13 | Shishi-no-neya | " | | | 14.20 | 10.7° | " | " " | " |
| 14 | Bunsaku-ido | " | | | 17.13 | 10.0° | " | " " | " |

* These springs, located at *Yunoshima*, about 2.5 m. S. W. from Tōkyō, are supposed to be connected with this spring.

The springs issue from the Liparite.

The hot spring “Ogawa-Onsen”* No. 5

* The waters were taken at the source on the springs at *Yamazaki-mura*, about 7.5 m. distant from the present bath-place located in the town of *Tonari*.

FUTAMI

Location.—Aimoto-mura, Shimo-niikawa-gun, Province Etchū, Toyama-ken.

The springs issue from the Granite.

The hot spring “Dai-ichi-no-yu”

Analysis.

Analysed by the Toyama Red Cross Hospital. 1913. Specific gravity: 1,0004 at 15° C.
Temperature: 64,5° (air temp. 29° C.). Total residue: ca. 0,50.

In 1 kilogram of the mineral water are contained

| | Grams | | Grams |
|---|--------|---|--------|
| Potassium chloride (KCl) | 0,0314 | Aluminium sulphate $Al_2(SO_4)_3$ | 0,0273 |
| Sodium chloride (NaCl) | 0,1514 | Boric acid (meta) (HBO_2) | 0,0204 |
| Sodium bicarbonate $(NaHCO_3)$ | 0,2057 | Silicic acid (meta) (H_2SiO_3) | 0,1664 |
| Calcium sulphate $(CaSO_4)$ | 0,0126 | | 0,6232 |
| Calcium bicarbonate $[Ca(HCO_3)_2]$ | 0,0038 | Free carbon dioxide (CO_2) | 0,0067 |
| Magnesium bicarbonate $[Mg(HCO_3)_2]$ | 0,0037 | Free hydrogen sulphide (H_2S) | 0,0004 |
| Ferrous bicarbonate $[Fe(HCO_3)_2]$ | 0,0005 | | 0,6303 |

The spring may be classified as “sulphur spring”.

Radio-activity. 2,60 Mache's units in 1 litre of water at 25,5° C.
(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 3, 1914).

The hot spring “Dai-ni-no-yu”

Analysis.

Analysed by the Toyama Red Cross Hospital. 1913. Specific gravity: 1,0006 at 15° C.
Temperature: 85° C. (air temp. 24° C.). Total residue: ca. 0,61.

The spring, being in its composition nearly the same as that of “Dai-ichi-no-yu”, may be classified as “sulphur spring”.

Radio-activity. 1,38 Mache's units in 1 litre of water at 24,2° C.
(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 3, 1913).

Table 73.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water
in
Mache's units
at
temp. in C. | Apparatus employed | Date when the
water was taken | Examined by |
|-----|----------------|----------------|--------------------------|---|--------------------|----------------------------------|-------------------------|
| 1 | Dai-ichi-no-yu | sulphur | 64,5° (29°) | 2,60 25,5 | K. & L. | Sept. 3, 1914 | Y. Kinugasa
(1.8.14) |
| 2 | Dai-ni-no-yu | .. | 85,0° (24°) | 1,38 24,2 | .. | .. | .. |
| 3 | Dai-san-no-yu | .. | 85,0° | 1,11 24,1 | .. | .. | .. |
| 4 | Spring A | .. | 12,5° | 0,61 26,0° | .. | .. | .. |
| 5 | Otoko-yu | sulphur | 88,5° (27°) | 1,94 25,0° | .. | .. | .. |
| 6 | Onna-yu | .. | 83,0° | 2,21 24,5 | .. | .. | .. |

KANETSURI

Location.—Katakaidani-mura, Shimo-niikawa-gun, Province Etchū, Toyama-ken.

The spring issues from the Contact Zone of Granite and Limestone.

The hot spring “Kanetsuri-Onsen”

Temperature: 49° C. (air temp. 29,5° C.).

Flow of water: ca. 180 hectolitres in 24 hours.

No accurate analysis has ever been made, but the spring seems to be “simple thermal”.

Radio-activity. 2,20 Mache's units in 1 litre of water at 26,8° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Sept. 3, 1914).

ŌMAKI

Location.—Toga-mura, Higashi-tonami-gun, Province Etchū, Toyama-ken.

The spring issues from the Porphyrite Dyke.

The hot spring “Ōmaki-Onsen”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1911.

Specific gravity: 1,0127 at 23° C.

Temperature: 49° C. (air temp. 25,5° C.).

Total residue: ca. 3,29.

Flow of water: ca. 346 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K') | 0,01895 | 0,48404 | 0,48404 |
| Sodium ion (Na') | 0,73404 | 31,84555 | 31,84555 |
| Ammonium ion (NH ₄ ') | 0,00048 | 0,02655 | 0,02655 |
| Calcium ion (Ca'') | 0,39442 | 9,86050 | 19,72100 |
| | | | 52,07714 |
| Anions. | | | |
| Chlorine ion (Cl') | 1,41621 | 39,94951 | 39,94951 |
| Sulphate ion (SO ₄ '') | 0,57774 | 6,01437 | 12,02874 |
| Hydrocarbonate ion (HCO ₃ ') | 0,00603 | 0,09883 | 0,09883 |
| | | 3,14787 | 88,27935 |
| Boric acid (meta) (HBO ₂) | 0,08514 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,05539 | | |
| | 3,28840 | | |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0,03611 |
| Sodium chloride (NaCl) | 1,86306 |
| Ammonium chloride (NH ₄ Cl) | 0,00142 |
| Calcium chloride (CaCl ₂) | 0,42093 |
| Calcium sulphate (CaSO ₄) | 0,81833 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,00810 |
| Boric acid (meta) (HBO ₂) | 0,08514 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,05539 |
| | 3,28848 |

Besides these a small quantity of free carbon dioxide.

The spring thus may be classified as “sulphated common salt spring”.

Radio-activity. 2,11 Mache's units in 1 litre of water at 25,7° C.

(Kohl-Löw. fontact.; Y. Kinugasa and T. Saitō, Aug. 16, 1914).

WAKURA

Location.—Hashemura, Kaslini-gun, Province Noto, Ishikawa-ken.

The springs issue from the Tertiary Tufaceous Sandstone.

The hot spring “Wakura-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1897.

Specific gravity: 1.08 at 7.8° C.

Temperature: 82° C.

Total residue: ca. 20.93.

Flow of water: ca. 2160 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.2050 | 5.2366 | 5.2366 |
| Sodium ion (Na ⁺) | 4.3456 | 188.5303 | 188.5303 |
| Calcium ion (Ca ⁺⁺) | 3.1918 | 79.5910 | 159.1820 |
| | | | 352.9489 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 12.3680 | 348.8857 | 348.8857 |
| Sulphate ion (SO ₄ ^{''}) | 0.1943 | 2.0229 | 4.0458 |
| | 20.3047 | 624.2665 | 352.9315 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0922 | | |
| | 20.3969 | | |

| | Grams |
|--|---------|
| Potassium chloride (KCl) | 0.3905 |
| Sodium chloride (NaCl) | 11.0295 |
| Calcium chloride (CaCl ₂) | 8.6091 |
| Calcium sulphate (CaSO ₄) | 0.2756 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0922 |
| | 20.3969 |

Besides these trace of magnesium, ferrous, aluminium, bromine, iodine and hydrophosphate ion.

The spring thus may be classified as “earth-muriated common salt spring”.

AWAZU

Location.—Awazu-mura, Nomi-gun, Province Kaga, Ishikawa-ken.

The springs issue from the Brecciated Liparite.

The hot spring “Awazu-Onsen”

Analysis.

Analysed by the Hygienic Laboratory of Ishikawa-ken. 1912.

Specific gravity: 1.0032 at 15° C.

Temperature: 47° C. (air temp. 4° C.).

Total residue: ca. 2.24.

Flow of water: ca. 2700 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams |
|---|--------|
| Potassium oxide (K ₂ O) | 0.0354 |
| Sodium oxide (Na ₂ O) | 0.3840 |
| Calcium oxide (CaO) | 0.4595 |
| Magnesia (MgO) | 0.0098 |
| Chlorine (Cl) | 0.2359 |
| Sulphuric anhydride (SO ₃) | 0.4727 |
| Silica (SiO ₂) | 0.0121 |
| Carbon dioxide (total) (CO ₂) | 0.6534 |
| Free hydrogen sulphide (H ₂ S) | 0.0073 |

| Hypothetical form of combination | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0541 |
| Sodium chloride (NaCl) | 0.3471 |
| Sodium bicarbonate (NaHCO ₃) | 0.5937 |
| Calcium sulphate (CaSO ₄) | 0.8032 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.3725 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0360 |
| Silicic acid (H ₂ SiO ₃) | 0.0121 |
| | 2.2187 |
| Free carbon dioxide (CO ₂) | 0.1457 |
| Free hydrogen sulphide (H ₂ S) | 0.0173 |
| | 2.3817 |

Besides these trace of iron oxide and alumina.

The spring may be classified as “sulphur spring”.

KATAYAMAZU

Location.—Sakumi-mura, Enuma-gun, Province Kaga, Ishikawa-ken.

The springs issue from the Tertiary.

The hot spring “Katayamazu-Onsen”

Analysis.

Analysed by the Kanazawa Garrison Hospital. 1909.

Specific gravity: 1.01024 at 15° C.

Temperature: 74° C.

Total residue: ca. 15.79.

Flow of water: ca. 630 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|--|---------|
| Potassium chloride (KCl) | 5.1311 | Magnesium sulphate (MgSO ₄) | 0.6513 |
| Sodium chloride (NaCl) | 7.2840 | Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃) . . | 0.1147 |
| Sodium carbonate (Na ₂ CO ₃) | 0.0029 | Silica (SiO ₂) | 0.1606 |
| Calcium chloride (CaCl ₂) | 1.3009 | | 15.7825 |
| Calcium sulphate (CaSO ₄) | 1.1370 | | |

Besides these trace of phosphoric and boric acid.

The spring may be classified as “common salt spring”.

YAMASHIRO

Location.—Yamashiro-machi, Enuma-gun, Province Kaga, Ishikawa-ken.

The springs issue from the Tertiary Shale.

The hot spring “Yamashiro-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1905.

Specific gravity: 1.0000 at 25° C.

Temperature: 66° C.

Total residue: ca. 1.66.

Flow of water: ca. 2142 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents | | Grams |
|--|----------------|------------|-----------------------|--|--------|
| Potassium ion (K ⁺) | 0.0125 | 0.3556 | 0.3556 | Potassium chloride (KCl) | 0.0238 |
| Sodium ion (Na ⁺) | 0.3904 | 17.3275 | 17.3275 | Sodium chloride (NaCl) | 0.4178 |
| Calcium ion (Ca ⁺⁺) | 0.1175 | 2.9302 | 5.8604 | Sodium sulphate (Na ₂ SO ₄) | 0.6874 |
| Magnesium ion (Mg ⁺⁺) | 0.0016 | 0.0657 | 0.1314 | Sodium bicarbonate (NaHCO ₃) | 0.0439 |
| | | | 23.6749 | Magnesium chloride (MgCl ₂) | 0.0063 |
| Anions. | | | | Calcium sulphate (CaSO ₄) | 0.3983 |
| Chlorine ion (Cl ⁻) | 0.2691 | 7.5910 | 7.5910 | Silicic acid (meta) (H ₂ SiO ₃) | 0.0876 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.7455 | 7.7608 | 15.5216 | | 1.6651 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0.0319 | 0.3286 | 0.3286 | Free hydrogen sulphide (H ₂ S) | 0.0016 |
| | 1.5775 | 36.3594 | 23.4412 | | 1.6667 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0876 | | | | |
| | 1.6651 | | | | |
| Free carbon dioxide (CO ₂) | not determined | | | | |
| Free hydrogen sulphide (H ₂ S) | 0.0016 | | | | |
| | 1.6667 | | | | |

Besides these trace of ammonium, ferrous, aluminium and hydrophosphate ion and boric acid.

The spring thus may be classified as “saline bitter sulphur spring”.

YAMANAKA

Location.—Yamanaka-machi, Tama-gun, Province Kaga, Ishikawa-ken.

The spring issues from the Tertiary Tuff Breccia.

The hot spring ‘Yamanaka-Onsen’

Analysis (calculated from the salt table).

Analysed by the Kanazawa Garrison Hospital.

Specific gravity: 1.0016 at 15° C.

Temperature: 49° C.

Total residue: ca. 1.65.

Flow of water: ca. 1800 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|----------------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.1698 | 4.3372 | 4.3372 |
| Sodium ion (Na ⁺) | 0.1416 | 6.1432 | 6.1432 |
| Calcium ion (Ca ⁺⁺) | 0.2747 | 6.8504 | 13.7008 |
| Magnesium ion (Mg ⁺⁺) | 0.0076 | 0.3120 | 0.6240 |
| | | | 24.8052 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.0799 | 2.2539 | 2.2539 |
| Sulphate ion (SO ₄ ^{''}) | 1.0802 | 11.2451 | 22.4902 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0023 | 0.0377 | 0.0377 |
| | | 1.7561 | 31.1795 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0441 | | 24.7818 |
| | 1.8002 | | |
| Free carbon dioxide (CO ₂) | not determined | | |
| Free hydrogen sulphide (H ₂ S) | 0.0010 | | |
| | 1.8012 | | |

| | Grams |
|--|--------|
| Sodium chloride (NaCl) | 0.1318 |
| Potassium sulphate (K ₂ SO ₄) | 0.3781 |
| Sodium sulphate (Na ₂ SO ₄) | 0.2741 |
| Sodium bicarbonate (NaHCO ₃) | 0.0032 |
| Calcium sulphate (CaSO ₄) | 0.9311 |
| Magnesium sulphate (MgSO ₄) | 0.0378 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0441 |
| | 1.8002 |
| Free hydrogen sulphide (H ₂ S) | 0.0010 |
| | 1.8012 |

Besides these trace of ferrous, aluminium and hydrophosphate ion and boric acid.

The spring thus may be classified as “sulphated bitter sulphur spring”.

YUWAKU

Location.—Yuwakudani-mura, Ishikawa-gun, Province Kaga, Ishikawa-ken.

The springs issue from the Tertiary.

The hot spring “Yuwaku-Onsen”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1912.

Specific gravity: 1.0020 at 17° C.

Temperature: 41° C.

Flow of water: ca. 1800 hectolitres in 24 hours

Total residue: ca. 3.31.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.01287 | 0.32874 | 0.32874 |
| Sodium ion (Na ⁺) | 0.92424 | 40.09718 | 40.09718 |
| Ammonium ion (NH ₄ ⁺) | 0.00317 | 0.17533 | 0.17533 |
| Calcium ion (Ca ⁺⁺) | 0.23096 | 5.77400 | 11.54890 |
| Magnesium ion (Mg ⁺⁺) | 0.00175 | 0.07184 | 0.14368 |
| Ferrous ion (Fe ⁺⁺) | 0.00140 | 0.02500 | 0.05000 |
| Aluminium ion (Al ⁺⁺⁺) | 0.00317 | 0.11700 | 0.35100 |
| | | 52.69333 | |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.02452 |
| Sodium chloride (NaCl) | 2.15305 |
| Ammonium chloride (NH ₄ Cl) | 0.03411 |
| Sodium sulphate (Na ₂ SO ₄) | 0.3341 |
| Calcium sulphate (CaSO ₄) | 0.70586 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.09494 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.01052 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.00426 |

Anions.

| | | | |
|--|---------|----------|-----------------|
| Chlorine ion (Cl') | 1,32261 | 37,30917 | 37,30917 |
| Sulphate ion (SO ₄ '') | 0,67343 | 7,01051 | 14,02102 |
| Hydrocarbonate ion (HCO ₃ '') | 0,08319 | 1,36356 | 1,36356 |
| | 3,23170 | 92,27233 | 52,69375 |
| Boric acid (meta) (HBO ₂) | 0,11329 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,04923 | | |
| | 3,41931 | | |

| | |
|---|---------|
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0,02003 |
| Boric acid (meta) (HBO ₂) | 0,11329 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,04923 |
| | 3,41925 |

Besides these a small quantity of free carbon dioxide.

The spring thus may be classified as "sulphated common salt spring".

AWARA

Location.—Awara-mura, Sakai-gun, Province Echizen, Fukui-ken.

The springs issue from the Alluvium.

The hot spring "Beniya-no-yu" (Funatsu No. 10)

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1,0085 at 12° C.

Temperature: 76° C.

Flow of water: ca. 152 hectolitres in 24 hours.

Total residue: ca. 10,03.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|-----------------------------------|---------|------------|-----------------------|
| Potassium ion (K') | 0,11510 | 2,94000 | 2,94000 |
| Sodium ion (Na') | 2,32621 | 100,92225 | 100,92025 |
| Ammonium ion (NH ₄ '') | 0,00212 | 0,11726 | 0,11726 |
| Calcium ion (Ca'') | 1,28974 | 32,24350 | 64,48700 |
| Magnesium ion (Mg'') | 0,00438 | 0,17980 | 0,35960 |
| Ferrous ion (Fe'') | 0,00070 | 0,01250 | 0,02500 |
| | | | 168,84911 |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0,2193 |
| Sodium chloride (NaCl) | 5,9041 |
| Ammonium chloride (NH ₄ Cl) | 0,0063 |
| Calcium chloride (CaCl ₂) | 3,1501 |
| Calcium sulphate (CaSO ₄) | 0,5221 |
| Magnesium sulphate (MgSO ₄) | 0,0213 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,0005 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0022 |
| Boric acid (meta) (HBO ₂) | 0,1407 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0888 |
| | 10,0554 |
| Free carbon dioxide (CO ₂) | 0,0537 |
| | 10,1091 |

| Anions. | Grams | Milli-mols | Milligram-equivalents |
|--|----------|------------|-----------------------|
| Chlorine ion (Cl') | 5,70011 | 160,79295 | 160,79295 |
| Sulphate ion (SO ₄ '') | 0,38556 | 4,01374 | 8,02748 |
| Hydrocarbonate ion (HCO ₃ '') | 0,00175 | 0,02868 | 0,02868 |
| | 9,82567 | 301,24868 | 168,84911 |
| Boric acid (meta) (HBO ₂) | 0,14069 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,08884 | | |
| | 10,05520 | | |
| Free carbon dioxide (CO ₂) | 0,05374 | 1,22136 | |
| | 10,10894 | | |

Besides these trace of bromine, nitrate and hydrophosphate ion.

The spring thus may be classified as "earth-muriated common salt spring".

Radio-activity. 3,21 $\frac{1}{2}$ Mache's units in 1 litre of water at 17,1° C.

(Eng.-Siev. fontact.; Y. Kinugasa, Dec. 12, 1913).

39,43 $\frac{1}{2}$ Mache's units in 1 litre of gas at 0° C.

(Eng.-Siev. fontact.; Y. Kinugasa, Dec. 13, 1913).

† Correction for the absorption of the radiation by the wall of the ionisation chamber and also for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was made.

Table 74.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | | Apparatus employed | Date when the
water was taken | Examined by |
|--------------------|--|----------------------------|--------------------------|---|--|-------|-------|---------------|--------------------|---------------------------------------|---------------|
| | | | | | in
Maché's units | | | | | | |
| | | | | | I* | II | III | (Temp. in C.) | | | |
| 1 | Beniya-no-yu
<i>Fuutsu</i> —No. 10 | Carbonated
concentrated | 76.0° | ca.
152 | 2.72 | 3.15 | 3.21 | 17.1 | E. & S. | Dec. 12, 1913 | V. K. G. & A. |
| 2 | " " No. 9 | " | 70.0° | " | 1.83 | 2.12 | 2.16 | 10.5 | " | " | " |
| 3 | Kakawatei-no-yu " No. 6 | " | 74.0° | " | 1.08 | 1.25 | 1.28 | 8.2 | " | " 13 " | " |
| 4 | Fukuirō-no-yu " No. 21 | " | 70.5° | " | 2.36 | 2.73 | 2.79 | 13.0 | " | " 13 " | " |
| 5 | Oda-no-yu (<i>Nippon</i> —No. 21) | " | 53.0° | " | 1.51 | 1.75 | 1.79 | 12.7 | " | " | " |
| 6 | Haiya-no-yu " No. 28 | " | 75.0° | " | 2.26 | 2.61 | 2.66 | 8.0 | " | " 16 " | " |
| 7 | Toichirō-no-yu
<i>Tsuika-naka</i> —No. 22 | " | 63.0° | " | 1.54 | 1.77 | 1.81 | 12.4 | " | " 13 " | " |
| 8 | Ishizukaya-no-yu " No. 6 | " | 68.2° | " | 1.85 | 2.14 | 2.18 | 7.8 | " | " 16 " | " |
| Gas evolving from: | | | | | | | | | | | |
| 1 | Beniya-no-yu
<i>Fuutsu</i> —No. 10 | | | | Emanation per
litre of gas at 0° C. | | | | | Date when the
gas was
collected | |
| 2 | " " No. 9 | — | — | — | 33.55 | 28.69 | 29.43 | — | — | Dec. 13, 1913 | " |
| | | | | | 22.45 | 25.88 | 26.37 | — | — | " 15 " | " |

* Determined according to the authors' original direction.
† Correction for the absorption of the radiation by the wall of the ionisation chamber was made.
‡ Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made.

KOMONO

Location.—Komono-mura, Mie-gun, Province Ise, Mie-ken.

The springs issue from the Granite.

The cold spring "Shika-no-yu"

Analysis (calculated from the salt table).

Analysed by the Osaka Imperial Hygienic Laboratory, 1917.

Specific gravity: 1.0006.

Temperature: 29° C. (air temp. 13° C.)

Total residue: ca. 0.14.

Flow of water: abundant.‡

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its

Cations.

| | Grams | Milli-mols | Milli-grams-equivalents | | Grams |
|---------------------------------|--------|------------|-------------------------|--|--------|
| Potassium ion (K ⁺) | 0.0026 | 0.0064 | 0.0074 | Potassium chloride (KCl) | 0.0039 |
| Sodium ion (Na ⁺) | 0.0392 | 1.7031 | 1.7031 | Sodium chloride (NaCl) | 0.0392 |
| Calcium ion (Ca ⁺⁺) | 0.0025 | 0.0023 | 0.1147 | Sodium bicarbonate (NaHCO ₃) | 0.0191 |
| | | | 1.8241 | Calcium sulphate (CaSO ₄) | 0.0025 |

‡ The waters flow down constantly from iron pipe, 5 metres in diameter, at the bottom of the spring about 200 m. from the surface.

Anions.

| | | | |
|--|--------|--------|--------|
| Chlorine ion (Cl') | 0.0133 | 0,3752 | 0,3752 |
| Sulphate ion (SO ₄ '') | 0.0021 | 0,0219 | 0,0438 |
| Hydrocarbonate ion (HCO ₃ '') | 0.0901 | 1,4768 | 1,4768 |
| | 0,1498 | 3,7057 | 1,8958 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0221 | | |
| | 0,1719 | | |

| | |
|---|--------|
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0065 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0221 |
| | 0,1719 |

Besides these a small quantity of ferrous and aluminium ion and trace of magnesium ion, boric acid and organic substances.

The spring thus may be classified as “simple cold spring”.

Radio-activity. 8.18 Mache's units in 1 litre of water at 13,4° C.

(Kohl.-Löw. fontact.; Y. Kinugasa, April 17, 1914).

Table 75.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-------------|----------------|--------------------------|---------------------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Shika-no yu | simple | air temp.
29,0° (100) | 8,18 | 13,4° | K. & L. | Apr. 17, 1914 | Y. Kinugasa |
| 2 | Spring A | , | 10,0° (17°) | 13,32 | 11,6° | " | " | " |

ROKUYŌ

Location.—Tōichi-mura, Soekami-gun, Province Yamato, Nara-ken.

The spring issues from the Alluvium.

The cold spring “Nanto-rokuyō-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1907.

Specific gravity: 1,005.

Total residue: ca. 3,79.

Temperature: 20,5° C.

Flow of water: ca. 45 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

Cations.

| | Grams | Milli-
mols | Milligram-
equivalents |
|-----------------------|--------|----------------|---------------------------|
| Potassium ion (K') | 0,0492 | 1,2567 | 1,2567 |
| Sodium ion (Na') | 0,0623 | 2,7028 | 2,7028 |
| Calcium ion (Ca'') | 0,0156 | 0,3890 | 0,7780 |
| Ferrous ion (Fe'') | 1,0751 | 19,2325 | 38,4650 |
| Aluminium ion (Al''') | 0,0190 | 0,7011 | 2,1033 |
| | | | 45,3058 |

Anions.

| | | | |
|--|--------|---------|---------|
| Chlorine ion (Cl') | 0,1308 | 3,6897 | 3,6897 |
| Sulphate ion (SO ₄ '') | 1,9990 | 20,8099 | 41,6198 |
| | 3,3510 | 48,7817 | 45,3095 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,2739 | | |
| | 3,6249 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0,0740 |
| Sodium chloride (NaCl) | 0,1580 |
| Potassium sulphate (K ₂ SO ₄) | 0,0230 |
| Calcium sulphate (CaSO ₄) | 0,0530 |
| Ferrous sulphate (FeSO ₄) | 2,9230 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0,1200 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,2739 |
| | 3,6249 |

Besides these a small quantity of hydrogen and hydrosulphate ion, and trace of hydrophosphate ion.

The spring thus may be classified as “**vitriol spring**”.

Radio-activity. 1.86×10^{-10} curies in 1 litre of water at 17° C.

(Schmidt's electrom.; H. Kibezaki, Sept. 12, 1913).

0.51 Mache's units (recalculated).

SHIRAYA

Location.—Kawakami-mura, Yoshino-gun, Province Yamato, Nara-ken.

The springs issue from the Chichibu Series.

The cold spring “Shiraya-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1904.

Specific gravity: 1.0026 at 15° C.

Total residue: ca. 1.74.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | GRAMS | Milli-
mols | Milligram-
equivalents | | GRAMS |
|--|--------|----------------|---------------------------|---|--------|
| Potassium ion (K') | 0.0105 | 0.2682 | 0.2682 | Potassium chloride (KCl) | 0.0200 |
| Sodium ion (Na') | 0.2577 | 11.1801 | 11.1801 | Sodium chloride (NaCl) | 0.0540 |
| Calcium ion (Ca'') | 0.3553 | 8.8604 | 17.7208 | Sodium bicarbonate (NaHCO ₃) | 0.8622 |
| Magnesium ion (Mg') | 0.0114 | 0.7964 | 1.5928 | Calcium bicarbonate [Ca(HCO ₃) ₂] | 1.4369 |
| Ferrous ion (Fe'') | 0.0015 | 0.0268 | 0.0536 | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.1163 |
| Manganous ion (Mn'') | 0.0010 | 0.0182 | 0.0364 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0046 |
| | | | 30.8519 | Manganous bicarbonate [Mn(HCO ₃) ₂] | 0.0031 |
| Anions. | | | | Silicic acid (meta) (H ₂ SiO ₃) | 0.0082 |
| Chlorine ion (Cl') | 0.0422 | 1.1904 | 1.1904 | | 2.5053 |
| Hydrocarbonate ion (HCO ₃ ') | 1.8095 | 29.6597 | 29.6597 | Free carbon dioxide (CO ₂) | 1.4180 |
| | 2.4971 | 52.0002 | 30.8501 | | 3.9233 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0082 | | | | |
| | 2.5053 | | | | |
| Free carbon dioxide (CO ₂) | 1.4180 | 32.2273 | | | |
| | 3.9233 | | | | |

Besides these trace of lithium, aluminium, iodine, nitrate, sulphate and hydrophosphate ion, boric acid and organic substances.

The spring thus may be classified as “**alkaline earthy carbondioxated spring**”.

SEDO-NO-KANAYAMA

Location.—Sedo-no-kanayama-mura, Nishi-muro-gun, Province Kii, Wakayama-ken.

The springs issue from the Younger Tertiary.

The hot spring “Hama-no-yu”

Analysis.

Analysed by S. Hirohashi. 1901.

Specific gravity: 1.00465 at 22° C.

Temperature: 52° C.

Total residue: ca. 4.82.

Flow of water: ca. 156 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|---|---------|--|---------|
| Sodium chloride (NaCl) | 2,03794 | Silica (SiO ₂) | 0,13605 |
| Potassium sulphate (K ₂ SO ₄) | 0,00943 | | 6,56536 |
| Sodium bicarbonate (NaHCO ₃) | 4,16764 | Total carbon dioxide (CO ₂) | 3,86100 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,12259 | Carbon dioxide (free and in form of bicarbonates) (CO ₂) | 2,58900 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,08471 | | |
| Alumina and trace of iron oxide (Al ₂ O ₃ +Fe ₂ O ₃) | 0,00700 | | |

Besides these a small quantity of potassium chloride.

The spring may be classified as “**muriated alkaline carbondioxated spring**”.

Radio-activity. $0,43 \times 10^{-10}$ curies in 1 litre of water at 52° C.

(Schmidt's electrom.; H. Kibezaki, March 28, 1913).

0,12 Mache's units (recalculated).

The hot spring “Moto-no-yu”

Analysis.

Analysed by S. Hirohashi. 1901.

Specific gravity: 1,00425 at 22° C.

Temperature: 49° C.

Flow of water: ca. 87 hectolitres in 24 hours.

Total residue: ca. 4,26.

The spring, being in its composition nearly the same as that of “Hama-no-yu”, may be classified as “**muriated alkaline carbondioxated spring**”.

Radio-activity. $0,53 \times 10^{-10}$ curies in 1 litre of water at 49° C.

(Schmidt's electrom.; H. Kibezaki, March 29, 1913).

0,14 Mache's units (recalculated).

The hot spring “Saki-no-yu”

Analysis.

Analysed by S. Hirohashi. 1901.

Specific gravity: 1,00421 at 22° C.

Temperature: 60° C.

Flow of water: ca. 260 hectolitres in 24 hours.

Total residue: ca. 4,31.

The spring, being in its composition nearly the same as that of “Hama-no-yu”, may be classified as “**muriated alkaline carbondioxated spring**”.

Radio-activity. $0,36 \times 10^{-10}$ curies in 1 litre of water at 60° C.

(Schmidt's electrom.; H. Kibezaki, March 30, 1913).

0,10 Mache's units (recalculated).

The hot spring “Yagata-yu”

Analysis.

Analysed by S. Hirohashi. 1901.

Specific gravity: 1,00413 at 22° C.

Temperature: 42° C.

Flow of water: ca. 260 hectolitres in 24 hours.

Total residue: ca. 4,09.

The spring, being in its composition nearly the same as that of “Hama-no-yu”, may be classified as “**muriated alkaline carbondioxated spring**”.

Radio-activity. $0,49 \times 10^{-10}$ curies in 1 litre of water at 42° C.

(Schmidt's electrom.; H. Kibezaki, March 30, 1913).

0,14 Mache's units (recalculated).

Table 76.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|--------------------------|---------------------------------------|-----------------------|---|------------------------------|------------------|------------------|--------------------|-------------------------------|-------------|
| | | | | | in 10-10 curies | in Mache's units | at (temp. in C.) | | | |
| 1 | Moto-no-yu (Public bath) | mineral alk. (sulf. ca.) bicarbonated | 49,0° | ca. 87 | 0,53 | 0,14 | 49,0° | Schm. | March 20, 1913 | H. Kibezaki |
| 2 | Saki-no-yu (") | " | 60,0° | 260 | 0,36 | 0,10 | 60,0° | " | " 30. " | " |
| 3 | Yagata-yu (") | " | 42,0° | 260 | 0,49 | 0,14 | 42,0° | " | " " | " |
| 4 | Hamano-no-yu (") | " | 52,0° | 156 | 0,43 | 0,12 | 52,0° | " | " 28, " | " |
| 5 | Senki-yu | " | 48,0° | 69 | — | — | — | — | — | — |
| 6 | Mabu-yu | " | 45,0° | 225 | — | — | — | — | — | — |
| 7 | Awa-yu | " | 48,0° | 78 | — | — | — | — | — | — |
| 8 | Vorozuya-no-yu | " | 43,0° | 130 | — | — | — | — | — | — |

YUNOMINE

Location.—Shi-mura, Higashi-muro-gun, Province Kii, Wakayama-ken.

The springs issue from the Tertiary.

The hot spring "Yunomine-Onsen"

Analysis (calculated from the original numbers).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1914.

Temperature: 60° C.

Specific gravity: 1,0012 at 15° C.

Total residue: ca. 1,38.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|----------------|------------|-----------------------|
| Potassium ion (K ⁺) | 0,0396 | 1,0102 | 1,0102 |
| Sodium ion (Na ⁺) | 0,3236 | 14,0695 | 14,0695 |
| Calcium ion (Ca ⁺⁺) | 0,0451 | 1,1255 | 2,2510 |
| Ferrous ion (Fe ⁺⁺) | 0,0030 | 0,0537 | 0,1074 |
| | | | 17,4381 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0,1845 | 5,2030 | 5,2030 |
| Sulphate ion (SO ₄ ⁻²) | 0,5870 | 6,1101 | 12,2202 |
| | 1,1828 | 27,5720 | 17,4232 |
| Silicic acid (meta) (H ₂ SiO ₃) . . | 0,1805 | | |
| | 1,3633 | | |
| Free carbon dioxide (CO ₂) . . | not determined | | |
| Free hydrogen sulphide (H ₂ S) | 0,01014 | | |
| | 1,37344 | | |

| | Grams |
|--|---------|
| Potassium chloride (KCl) | 0,07530 |
| Sodium chloride (NaCl) | 0,24520 |
| Sodium sulphate (Na ₂ SO ₄) | 0,70070 |
| Calcium sulphate (CaSO ₄) | 0,15350 |
| Ferrous sulphate (FeSO ₄) | 0,00810 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,18050 |
| | 1,66330 |
| Free hydrogen sulphide (H ₂ S) | 0,01014 |
| | 1,67344 |

The spring thus may be classified as "saline bitter sulphur spring".

KATSU-URA

Location.—Katsu-ura-machi, Higashi-muro-gun, Province Kii, Wakayama-ken.

The springs issue from the Tertiary.

The hot spring “Soto-no-yu”

Analysis.

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1900.

Specific gravity: 1.0008 at 15° C.

Temperature: 38.5° C.

Flow of water: ca. 900 hectolitres in 24 hours.

Total residue: ca. 0.93.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|---------|
| Potassium chloride (KCl) | 0.0078 | Silica (SiO ₂) | 0.0051 |
| Sodium chloride (NaCl) | 0.6567 | | 0.0057 |
| Calcium chloride (CaCl ₂) | 0.1244 | Carbon dioxide (free and in form of | |
| Magnesium chloride (MgCl ₂) | 0.0037 | bicarbonates) (CO ₂) | 0.0045 |
| Calcium sulphate (CaSO ₄) | 0.0524 | Free hydrogen sulphide (H ₂ S) | 0.00798 |
| Calcium carbonate (CaCO ₃) | 0.0352 | | 0.01818 |
| Ferric oxide (Fe ₂ O ₃) | 0.0002 | | |
| Alumina (Al ₂ O ₃) | 0.0002 | | |

Besides these trace of lithium, manganese, iodine and nitric, phosphoric and boric acid.

The spring thus may be classified as “sulphur spring”.

Radio-activity. 3.69×10^{-10} curies in 1 litre of water at 38.5° C.

(Schmidt's electrom.; H. Kibezaki, April 8, 1914)

0.85 Mache's units (recalculated).

Table 77.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------------------|------------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Soto-no-yu (<i>Public bath</i>) | sulphur | 38.5° | ca.
900 | 3.69 | 0.85 | 38.5° | Schm | Apr. 8, 1914 | H. Kibezaki |
| 2 | Nakaneshima-Onsen | — | 27.0° | — | 10.36 | 2.85 | 27.0° | .. | .. 9, .. | .. |
| 3 | Akashima-Onsen | alkaline sulphur | 45.0° | 900 | 6.55 | 1.80 | 45.0° | .. | .. 10, .. | .. |
| 4 | Penten-yu | — | 17.0° | — | 2.26 | 0.62 | 17.0° | .. | .. 11, .. | .. |

YUKAWA

Location.—Nachi-mura, Higashi-muro-gun, Province Kii, Wakayama-ken.

The springs issue from the Misaka Series.

Table 78.

| No. | Spring | Classification | Temperature
in C. | Flow of water
in 24 hrs.
in hectolitres | Concentration per
litre of water | | | Apparent
specific gravity | Date of analysis | Analysed by |
|-----|----------------------------|----------------|----------------------|---|-------------------------------------|----------------------------|--------------------------|------------------------------|------------------|-------------|
| | | | | | Total
solids | Total
mineral
matter | Temp. at
analysis, C. | | | |
| 1 | Onatono-yu <i>Pakikōji</i> | — | 22.0° | — | 5.31 | 1.46 | 22.0 | Schm. | Apr. 12, 1914 | H. Kasagi |
| 2 | Koshi-Onsen | alkaline | 28.5° | 108 | — | — | — | — | — | — |
| 3 | Kosse-Onsen | sulphur | 24.5° | 108 | — | — | — | — | — | — |
| 4 | Naka-no-yu | simple | 30.3° | 300 | — | — | — | — | — | — |
| 5 | Hamano-miya-yu | .. | 40.0° | 300 | — | — | — | — | — | — |

KASAGI

Location.—Kasagi-mura, Sōraku-gun, Province Yamashiro, Kyōto-fu.

The springs issue from the Granite.

The cold spring “Kasagi-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Osaka Imperial Hygienic Laboratory. 1900.

Specific gravity: 1.0044 at 15° C.

Temperature: 13.6° C. (air temp. 13° C.).

Flow of water: ca. 27–36 hectolitres in 24 hours.

Total residue: ca. 5.08.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

Cations.

| | Grams | Milli-grams | Milli-moles |
|---|--------|-------------|-------------|
| Sodium ion (Na ⁺) | 1.8077 | 78.4252 | 78.4252 |
| Calcium ion (Ca ⁺⁺) | 0.0752 | 1.8753 | 0.0009 |
| Magnesium ion (Mg ⁺⁺) | 0.0263 | 1.0777 | 0.0002 |
| Ferrous ion (Fe ⁺⁺) | 0.0029 | 0.0517 | 0.0009 |
| Manganous ion (Mn ⁺⁺) | 0.0009 | 0.0164 | 0.0002 |
| | | | 84.4714 |

| | |
|---|--------|
| Sodium chloride (NaCl) | 1.6130 |
| Sodium bicarbonate (NaHCO ₃) | 4.2781 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.3046 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.1584 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0092 |
| Manganous bicarbonate [Mn(HCO ₃) ₂] | 0.0030 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1272 |
| | 6.4935 |

Anions.

| | | | |
|--|----------------|----------|---------|
| Chlorine ion (Cl ⁻) | 0.9775 | 27.5710 | 27.5710 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 3.4758 | 56.9710 | 56.9710 |
| | 6.3663 | 165.9933 | 84.5450 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1272 | | |
| | 6.4935 | | |
| Free carbon dioxide (CO ₂) | not determined | | |

Besides these trace of lithium, aluminium, bromine and hydrophosphate ion and organic substances.

The spring thus may be classified as “**muriated alkaline spring**”.

Table 79.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------|----------------------|-------------------------------------|---------------------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kasagi-Kōsen No. 1 | muriated
alkaline | 13.6° ^{air temp.}
(13°) | 1.01 | 15.0° | K. & L. | Apr. 18, 1914 | Y. Kinugasa |
| 2 | " No. 2 | " | 13.6° (13°) | 0.84 | 14.0° | " | " | " |

DAITŌ

Location.—Mikkaichi-mura, Minami-kawachi-gun, Province Kawachi, Osaka-fu.

The spring issues from the Tertiary.

The cold spring "Daitō-Kōsen"

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1890.

Specific gravity: 1.012.

Total residue: ca. 15.64.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|----------------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0.1619 | 4.1354 | 4.1354 |
| Sodium ion (Na ⁺) | 4.2157 | 182.8937 | 182.8937 |
| Ammonium ion (NH ₄ ⁺) | 0.0060 | 0.3326 | 0.3326 |
| Calcium ion (Ca ⁺⁺) | 0.8282 | 20.6534 | 41.3068 |
| Magnesium ion (Mg ⁺⁺) | 0.3844 | 15.7800 | 31.5600 |
| Ferrous ion (Fe ⁺⁺) | 0.0154 | 0.2755 | 0.5510 |
| | | | 260.7795 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 7.1471 | 201.6107 | 201.6107 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 3.6100 | 59.1705 | 59.1705 |
| | 16.3687 | 484.8518 | 260.7812 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0909 | | |
| | 16.4596 | | |
| Free carbon dioxide (CO ₂) | not determined | | |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.3083 |
| Sodium chloride (NaCl) | 10.6998 |
| Ammonium chloride (NH ₄ Cl) | 0.0178 |
| Calcium chloride (CaCl ₂) | 0.7911 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 2.1927 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 2.3100 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0490 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0909 |
| | 16.4596 |

Besides these trace of aluminium, bromine, sulphate and hydrophosphate ion, and boric acid.

The spring thus may be classified as "earthy iron carbonate common salt spring".

Radio-activity. 1.60×10^{-10} curies in 1 litre of water* at 28° C.

(Schmidt's electrom.; H. Kibezaki, July, 6, 1914).

0.44 Mache's units (recalculated).

* Measurement was made with the sample brought to the laboratory by applicant, taking into account decay of radium emanation with the time.

| No. | Spring | Classification | Temp. of spring
in °C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-------------------------|----------------|---------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | in
10 ⁻¹⁰ curies | in
Maché's units | at
(temp. in C.) | | | |
| 5 | Kurakuen-Kōsen No. 5 | simple | — | 14.73 | 4.05 | — | Schm. | Oct. 25, 1913 | H. Kibezaki |
| 6 | " No. 6 | " | — | 31.02 | 8.53 | — | " | " 27, " | " |
| 7 | " No. 7 | " | — | 18.80 | 5.17 | — | " | Nov. 7, " | " |
| 8 | " No. 8 | " | 17.0° | 19.02 | 5.23 | 17.0° | " | " 9, " | " |
| 9 | " No. 9 | " | 15.0° | 17.93 | 4.93 | 15.0° | " | Jan. 26, 1914 | " |
| 10 | " No. 10 | " | 15.0° | 60.15 | 16.54 | 15.0° | " | " 25, " | " |
| 11 | " No. 11 | " | 11.0° | 33.67 | 9.26 | 11.0° | " | " " | " |
| 12 | " No. 12 | " | 12.0° | 23.83 | 5.72 | 12.0° | " | " 28, " | " |
| 13 | " No. 13 | " | 14.5° | 16.15 | 4.44 | 14.5° | " | June 5, " | " |
| 14 | " No. 14 | " | 18.0° | 13.93 | 3.83 | 18.0° | " | " 8, " | " |
| 15 | Saunaru-ishi-yama No. 1 | " | 14.5° | 56.80 | 15.62 | 14.5° | " | Dec. 20, 1913 | " |
| 16 | " No. 2 | " | 12.0° | 51.46 | 14.15 | 12.0° | " | Jan. 22, 1914 | " |
| 17 | " No. 3 | " | 14.0° | 58.33 | 16.04 | 14.0° | " | " 23, " | " |
| 18 | " No. 4 | " | 14.0° | 22.22 | 6.11 | 14.0° | " | " 24, " | " |
| 19 | " No. 5 | " | 14.5° | 44.29 | 12.18 | 14.5° | " | May 23, " | " |
| 20 | " No. 6 | " | 14.0° | 35.71 | 9.82 | 14.0° | " | " 22, " | " |
| 21 | Myōbandani-ue No. 1 | " | 13.0° | 1.53 | 0.42 | 13.0° | " | Jan. 31, " | " |
| 22 | " No. 2 | " | 12.0° | 9.42 | 2.59 | 12.0° | " | Feb. 1, " | " |
| 23 | Myōbandani No. 1 | vitriol | — | 7.35 | 2.02 | — | " | Dec. 19, 1913 | " |
| 24 | " No. 2 | " | — | 4.36 | 1.20 | — | " | " 21, " | " |
| 25 | Fukae-Onsen * | — | 14.5° | 4.22 | 1.16 | 14.5° | " | " 11, " | " |

* The spring, located at *Honjō-mura*, is grouped for convenience in this table.

TAKARAZUKA

Location.—Ryōgen-mura, Muko-gun, Province Settsu, Hyōgo-ken.

The springs issue from the Granite.

The cold spring "Takarazuka-Kōsen"

Analysis (calculated from the salt table).

Analyzed by the Osaka Imperial Hygienic Laboratory. 1891.

Specific gravity: 1.011.

Temperature: 18.5° C.

Total residue: ca. 13.76.

Flow of water: ca. 360 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.416 | 10.7178 | 10.7178 |
| Sodium ion (Na ⁺) | 4.6092 | 199.9740 | 199.9749 |
| Ammonium ion (NH ₄ ⁺) | 0.0013 | 0.0719 | 0.0719 |
| Calcium ion (Ca ⁺⁺) | 0.0004 | 0.0010 | 0.0020 |
| Magnesium ion (Mg ⁺⁺) | 0.1238 | 5.0821 | 10.1642 |
| | | | 220.9299 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 7.1210 | 20.8714 | 200.8914 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 1.2260 | 20.0951 | 20.0951 |
| | 13.5019 | 40.9665 | 220.9865 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0774 | | |
| | | | 13.5993 |
| Free carbon dioxide (CO ₂) | 0.9338 | 21.2227 | |
| | | | 14.5331 |

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.7777 |
| Sodium chloride (NaCl) | 11.1252 |
| Sodium bicarbonate (NaHCO ₃) | 0.8245 |
| Ammonium bicarbonate (NH ₄ HCO ₃) | 0.0054 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0015 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.7460 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0974 |
| | 13.5993 |
| Free carbon dioxide (CO ₂) | 0.9338 |
| | 14.5331 |

Besides these trace of ferrous, aluminium, bromine, iodine, sulphate and hydrophosphate ion and boric acid.

The spring thus may be classified as **"carbondioxated * common salt spring"**.

Radio-activity. 2.66×10^{-10} curies in 1 litre of water at 18.5° C.

(Schmidt's electrom.; H. Kibezaki, Sept. 13, 1913).

0.73 Mache's units (recalculated).

7.16×10^{-10} curies in 1 litre of gas.

(Schmidt's electrom.; H. Kibezaki, Oct. 7, 1913).

1.97 Mache's units (recalculated).

The cold spring "Takarazuka-Tansan-sui"

Analysis (calculated from the salt table).

Analysed by the Osaka Imperial Hygienic Laboratory. 1901.

Temperature: 18.5° C.

Specific gravity: 1.0015.

Total residue: ca. 0.58.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0184 | 0.5235 | 0.5235 |
| Sodium ion (Na ⁺) | 0.1689 | 7.3193 | 7.3193 |
| Calcium ion (Ca ⁺⁺) | 0.0213 | 0.5312 | 1.0624 |
| Magnesium ion (Mg ⁺⁺) | 0.0021 | 0.0021 | 0.1642 |
| Ferrous ion (Fe ⁺⁺) | 0.0015 | 0.0268 | 0.0536 |
| | | | 0.1230 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.2440 | 6.8829 | 6.8829 |
| Sulphate ion (SO ₄ ⁼⁼) | 0.0036 | 0.0375 | 0.0750 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0.1310 | 2.1472 | 2.1472 |
| | 0.5908 | 17.5505 | 9.1051 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0360 | | |
| | 0.6268 | | |
| Free carbon dioxide (CO ₂) | 1.5360 | 34.9091 | |
| | | | 2.1628 |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0351 |
| Sodium chloride (NaCl) | 0.3747 |
| Sodium bicarbonate (NaHCO ₃) | 0.0783 |
| Calcium sulphate (CaSO ₄) | 0.0051 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0800 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0128 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0048 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0360 |
| | 0.6268 |
| Free carbon dioxide (CO ₂) | 1.5360 |
| | 2.1628 |

Besides these trace of manganous, aluminium and hydrophosphate ion.

The spring thus may be classified as **"simple carbondioxiated spring"**.

Radio-activity. 12.03×10^{-10} curies in 1 litre of water at 18.5° C.

(Schmidt's electrom.; H. Kibezaki, June 17, 1913).

3.31 Mache's units (recalculated).

* As it is probable that the actual amount of free carbon dioxide will be 1.5360 and not 1.5360, the water may be classified more properly as **"carbondioxated common salt spring."**

HIRANO

Location.—Tada-mura, Kawabe-gun, Province Settsu, Hyōgo-ken.

The springs issue from the Chichibu Series.

The cold spring “Hirano-Kōsen” (A)

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1893.

Temperature: 27° C.

Specific gravity: 1.0048.

Total residue: ca. 4.59.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0333 | 0.8506 | 0.8506 |
| Sodium ion (Na ⁺) | 1.4227 | 61.7223 | 61.7223 |
| Calcium ion (Ca ⁺⁺) | 0.2201 | 5.4888 | 10.9776 |
| Magnesium ion (Mg ⁺⁺) | 0.0804 | 3.3005 | 6.6010 |
| Ferrous ion (Fe ⁺⁺) | 0.0107 | 0.1914 | 0.3828 |
| | | | 80.5343 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 1.5867 | 44.7588 | 44.7588 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 2.1865 | 35.8384 | 35.8384 |
| | 5.5404 | 152.1508 | 80.5972 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1704 | | |
| | 5.7108 | | |
| Free carbon dioxide (CO ₂) | 1.1924 | 27.1000 | |
| | 6.9032 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0635 |
| Sodium chloride (NaCl) | 2.5685 |
| Sodium bicarbonate (NaHCO ₃) | 1.4990 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.8910 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.4842 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0342 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1704 |
| | 5.7108 |
| Free carbon dioxide (CO ₂) | 1.1924 |
| | 6.9032 |

Besides these trace of aluminium ion and boric acid.

The spring thus may be classified as “earthy alkaline iron carbonate carbondioxated common salt spring”.

TAKEDAO

Location.—Shioze-mura, Arima-gun, Province Settsu, Hyōgo-ken.

The springs issue from the Quartz Porphyry.

The cold spring “Takedao-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1902.

Specific gravity: 1.0005 at 15° C.

Total residue: ca. 1.13.

Flow of water: ca. 90 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|---|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0040 | 0.1033 | 0.1033 |
| Sodium ion (Na ⁺) | 0.2885 | 12.5163 | 12.5163 |
| Calcium ion (Ca ⁺⁺) | 0.0952 | 2.3738 | 4.7476 |
| Magnesium ion (Mg ⁺⁺) | 0.0037 | 0.1519 | 0.3038 |
| | | | 17.6710 |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0077 |
| Sodium chloride (NaCl) | 0.7323 |
| Calcium chloride (CaCl ₂) | 0.0907 |
| Calcium sulphate (CaSO ₄) | 0.0064 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.2446 |

Anions.

| | | | | | |
|--|----------------|---------|----------------|---|--------|
| Chlorine ion (Cl') | 5.54 | 14.2567 | 14.2567 | Magnesium bicarbonate [$\text{Mg}(\text{HCO}_3)_2$] . . . | 0.0222 |
| Sulphate ion (SO_4'') | 0.0045 | 0.0468 | 0.0936 | Silicic acid (meta) (H_2SiO_3) | 0.0374 |
| Hydrocarbonate ion (HCO_3') . | 0.2026 | 3.3207 | 3.3207 | | 1.1413 |
| | 1.1437 | 32.7005 | 17.7713 | Free hydrogen sulphide (H_2S). | 0.0014 |
| Silicic acid (meta) (H_2SiO_3) . | 0.0374 | | | | 1.1427 |
| | 1.1413 | | | | |
| Free carbon dioxide (CO_2) . . | not determined | | | | |
| Free hydrogen sulphide (H_2S) | 0.0014 | | | | |
| | 1.1427 | | | | |

Besides these trace of lithium, ferrous, manganous, aluminium, bromine, iodine and hydrophosphate ion and boric acid.

The spring thus may be classified as “**muriated sulphur spring**”.

Table 81.

| No. | Spring | Classification | Temp. of spring
in $^{\circ}\text{C}$. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------------------|-------------------------------|--|---|---------------------------------|---------------------|-------------------------------------|--------------------|---------------------------------------|-------------|
| | | | | | in
10 ¹⁰ curies | in
Maché's units | at
temp. in $^{\circ}\text{C}$. | | | |
| 1 | Takarazuka-Kōsen | carbondioxated
common salt | 18.7° | ca.
360 | 2.09 | 0.58 | 18.7° | Schm. | June 19, 1913 | H. Kibezaki |
| 2 | “ | “ | 18.5° | “ | 2.66 | 0.73 | 18.5° | “ | Sep. 13, “ | “ |
| 3 | Takarazuka-Tansan-sui | simple carbon-
dioxated | 18.5° | — | 12.03 | 3.31 | 18.5° | “ | June 17, “ | “ |
| 4 | Yu-moto <i>Public bath</i> | muriated
sulphur | 19.5° | — | 19.91 | 5.47 | 19.5° | “ | May 29, “ | “ |
| 5 | Moto-no-yu | “ | 23.5 | — | 11.34 | 3.12 | 23.5° | “ | “ 30, “ | “ |
| | Gas evolving from
Takarazuka-Kōsen | — | — | — | Emanation per
litre of gas | | | “ | Date when the
gas was
collected | “ |
| | | | | | 7.16 | 1.97 | — | | Oct. 7, 1913 | |

ARIMA

Location.—Arima-machi, Arima-gun, Province Settsu, Hyōgo-ken.

The springs issue from the Quartz Porphyry.

The hot spring “Ichi-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881. Specific gravity: 1.012 at 23° C.
Temperature: 53.4 C. Total residue: ca. 19.66.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|----------------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.6725 | 17,1775 | 17,1775 |
| Sodium ion (Na ⁺) | 5.7675 | 251,5017 | 251,5617 |
| Calcium ion (Ca ⁺⁺) | 1,0455 | 26,0723 | 52,1446 |
| Ferrous ion (Fe ⁺⁺) | 0.1912 | 3,4707 | 6,9414 |
| Manganous ion (Mn ⁺⁺) | 0,0426 | 0,7745 | 1,5490 |
| Aluminium ion (Al ⁺⁺⁺) | 0,0059 | 0,2177 | 0,5331 |
| | | | 330,0273 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 11,4009 | 324,5766 | 321,5966 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0,5121 | 8,3937 | 8,3937 |
| | | 19,6689 | 629,2647 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0753 | | |
| | | | 19,7442 |
| Free carbon dioxide (CO ₂) | not determined | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 1,2810 |
| Sodium chloride (NaCl) | 14,7170 |
| Calcium chloride (CaCl ₂) | 2,8960 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,6088 |
| Manganous bicarbonate [Mn(HCO ₃) ₂] | 0,1371 |
| Aluminium chloride (AlCl ₃) | 0,0290 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0753 |
| | 19,7442 |

Besides these trace of lithium, magnesium ion and organic substances.

The spring thus may be classified as “earth-muriated iron carbonate common salt spring”.

The hot spring “Ikenobō-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1901.

Temperature : 47° C.

Total residue : ca. 64,01.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------------|------------|-----------------------|
| Potassium ion (K ⁺) | 3,9485 | 100,8341 | 100,8341 |
| Sodium ion (Na ⁺) | 17,4712 | 757,9609 | 757,9609 |
| Lithium ion (Li ⁺) | 0,0256 | 3,6714 | 3,6714 |
| Ammonium ion (NH ₄ ⁺) | 0,0098 | 0,5432 | 0,5432 |
| Calcium ion (Ca ⁺⁺) | 3,7688 | 93,9850 | 187,9200 |
| Magnesium ion (Mg ⁺⁺) | 0,0435 | 1,7460 | 3,4938 |
| Ferrous ion (Fe ⁺⁺) | 0,2569 | 4,5941 | 9,1882 |
| Manganous ion (Mn ⁺⁺) | 0,0632 | 1,1527 | 2,3054 |
| Aluminium ion (Al ⁺⁺⁺) | 0,0053 | 0,1956 | 0,5868 |
| | | | 1066,5038 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 36,7229 | 1035,9069 | 1035,9069 |
| Bromine ion (Br ⁻) | 0,0854 | 1,0768 | 1,0768 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 1,8221 | 29,8656 | 29,8656 |
| | 64,2232 | 2031,5332 | 1066,8493 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0714 | | |
| | 64,2946 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 7,5210 |
| Sodium chloride (NaCl) | 43,2070 |
| Ammonium chloride (NH ₄ Cl) | 0,0290 |
| Lithium chloride (LiCl) | 0,1550 |
| Sodium bromide (NaBr) | 0,1100 |
| Sodium bicarbonate (NaHCO ₃) | 1,5438 |
| Calcium chloride (CaCl ₂) | 10,4400 |
| Magnesium chloride (MgCl ₂) | 0,1700 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,8182 |
| Manganous bicarbonate [Mn(HCO ₃) ₂] | 0,2032 |
| Aluminium chloride (AlCl ₃) | 0,0260 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0714 |
| | 64,2946 |

Besides these trace of nitrate and hydrophosphate ion, boric acid and organic substances.

The spring thus may be classified as “earth-muriated iron carbonate concentrated common salt spring”.

Table 82.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | At this spring | Date when the
water was taken | Examined by |
|-----|--------------------------------------|---|--------------------------|---------------------------------|---------------------|-------------------|----------------|----------------------------------|-------------|
| | | | | in
10-30 cubes | in
Maché's units | at
temp. in C. | | | |
| 1 | Arima-Onsen * | common salt | 51.5 | 0.92 | 0.25 | 51.5 | Schim. | May 20, 1913 | H. Kikuchi |
| 2 | Maruyama-Kōsen | — | 19.0° | 30.07 | 8.27 | 19.0° | " | " 27, " | " |
| 3 | Arima-tansan-sen (<i>Tepōsui</i>)* | simple carbon-
dioxated | 17.0° | 6.34 | 1.75 | 17.0° | " | " 28, " | " |
| 4 | Zuihōji-Onsen | — | 28.3° | 107.67 | 29.61 | 28.3° | " | " | " |
| 5 | Ichi-no-yu * | earth-muriated
iron-carbonate
common salt | 53.4° | — | — | — | — | — | — |
| 6 | Ni-no-yu * | " (?) | 53.4° | — | — | — | — | — | — |
| 7 | Mearai-yu * | " (?) | 34.5° | — | — | — | — | — | — |
| 8 | Hana-no-yu | " (?) | 43.6° | — | — | — | — | — | — |
| 9 | Ikenobō-Kōsen | earth-muriated
iron-carbonate
concentrated
common salt | 47.0° | — | — | — | — | — | — |

* These springs are for public use.

KINOSAKI

Location.—Kinosaki-machi, Kinosaki-gun, Province Tajima, Hyōgo-ken.

The springs issue from the Tertiary.

The hot spring "Ichi-no-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881. Total residue: ca. 5.32.

Flow of water: ca. 259 hectolitres in 24 hours.

Specific gravity: 1.00404 at 8.5° C.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milli-ferrous
equivalents |
|--|----------------|----------------|------------------------------|
| Potassium ion (K ⁺) | 0.0387 | 0.9885 | 0.5885 |
| Sodium ion (Na ⁺) | 1.3108 | 56.8677 | 56.8677 |
| Calcium ion (Ca ⁺⁺) | 0.5994 | 14.9476 | 29.8952 |
| Magnesium ion (Mg ⁺⁺) | 0.0018 | 0.0739 | 0.1478 |
| | | | 87.8992 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 2.8446 | 80.2426 | 80.2426 |
| Bromine ion (Br ⁻) | 0.0005 | 0.0063 | 0.0063 |
| Sulphate ion (SO ₄ ^{''}) | 0.3026 | 3.1504 | 6.3002 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0850 | 1.3932 | 1.3932 |
| | 5.1834 | 157.6699 | 87.9423 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0898 | | |
| | 5.2732 | | |
| Free carbon dioxide (CO ₂) | not determined | | |

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 3.0165 |
| Potassium sulphate (K ₂ SO ₄) | 0.0862 |
| Sodium sulphate (Na ₂ SO ₄) | 0.3774 |
| Calcium chloride (CaCl ₂) | 1.5910 |
| Magnesium bromide (MgBr ₂) | 0.0006 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.1013 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0111 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0898 |
| | 5.2732 |

Besides these trace of ferrous, aluminium and iodine ion and boric acid.

The spring thus may be classified as "earth-muriated common salt spring".

Table 83.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------------|-------------------------------|--------------------------|---|---------------------------------|--------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mach's units | at
(temp. in C.) | | | |
| 1 | Ichi-no-yu <i>Vent No. 1</i> | earth-muriated
common salt | 50.5° | } ca.
259 | 9.91 | 2.73 | 50.5° | Schm. | May 23, 1913 | H. Kibezaki |
| 2 | " <i>Vent No. 2</i> | " | 59.5° | | 10.08 | 2.77 | 59.5° | " | " 24, " | " |
| 3 | Kō-no-yu | " | 57.0° | 204 | 9.37 | 2.58 | 57.0° | " | " 21, " | " |
| 4 | Mandara-no-yu <i>Vent No. 1</i> | " | 58.0° | } 207 | 5.17 | 1.42 | 58.0° | " | " 19, " | " |
| 5 | " <i>Vent No. 2</i> | " | 47.0° | | 11.08 | 3.05 | 47.0° | " | " 20, " | " |
| 6 | Gosho-no-yu <i>Vent No. 1</i> | " | 56.5° | } 220 | 12.77 | 3.51 | 56.5° | " | " 22, " | " |
| 7 | " <i>Vent No. 2</i> | " | 60.3° | | 30.57 | 8.41 | 60.3° | " | " | " |
| 8 | Yanagi-no-yu | " | 59.5° | 216 | 13.58 | 3.74 | 59.5° | " | " 18, " | " |
| 9 | Jizō-no-yu | " | 48.5° | 207 | 5.21 | 1.43 | 48.5° | " | " 20, " | " |

YUMURA

Location.—Onsen-mura, Mikata-gun, Province Tajima, Hyōgo-ken.

The springs issue from the Granite overlaid by Alluvium.

The hot spring "Kabu-yu"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Specific gravity: 1.00092 at 7° C.

Temperature: 90.5° C.

Total residue: ca. 0.99.

Flow of water: ca. 900 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0358 | 0.9144 | 0.9144 |
| Sodium ion (Na ⁺) | 0.2685 | 11.6486 | 11.6486 |
| Calcium ion (Ca ⁺⁺) | 0.0179 | 0.4464 | 0.8928 |
| Magnesium ion (Mg ⁺⁺) | 0.0008 | 0.0328 | 0.0656 |
| Ferrous ion (Fe ⁺⁺) | 0.0034 | 0.0608 | 0.1216 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0023 | 0.0849 | 0.2547 |
| | | | 13.8977 |
| Anions. | Grams | Milli-
mols | Milligram-
equivalents |
| Chlorine ion (Cl ⁻) | 0.1408 | 3.9718 | 3.9718 |
| Sulphate ion (SO ₄ ^{''}) | 0.2134 | 2.2215 | 4.4430 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.3346 | 5.4843 | 5.4843 |
| | 1.0175 | 24.8655 | 13.8991 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1869 | | |
| | 1.2044 | | |
| Free carbon dioxide (CO ₂) | 1.2354 | 20.2491 | |
| | 2.4398 | | |

| | |
|---|--------|
| Sodium chloride (NaCl) | 0.2323 |
| Potassium sulphate (K ₂ SO ₄) | 0.0798 |
| Sodium sulphate (Na ₂ SO ₄) | 0.2329 |
| Sodium bicarbonate (NaHCO ₃) | 0.3702 |
| Calcium bicarbonate [Ca(HCO ₃) ₂]. | 0.0723 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0046 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0111 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.0143 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.1869 |
| | 1.2044 |
| Free carbon dioxide (CO ₂). | 1.2354 |
| | 2.4398 |

Besides these a very small quantity of boric acid, and trace of hydrogen sulphide and organic substances.

The spring thus may be classified as "alkaline carbondioxated spring".

Radio-activity. 3.09×10^{-12} curies in 1 litre of water at 12.8° C.

(Schmidt's electrom.; Dr. R. Ishizu, March 14, 1914).

0.85 Mache's units (recalculated).

Table 84.

| No. | Spring | Classification | Temp of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------------|------------------------------|-------------------------|---|--|---------------------|---------------------|--------------------|---|---------------|
| | | | | | in
10 ⁻¹² curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kabu-yu <i>Public bath</i> | alkaline carbon-
dioxated | 90.5° | 900 | 3.09 | 0.85 | 12.8° | Schm. | March 14, 1914 | Dr. R. Ishizu |
| 2 | " | " | " | " | — | 0.95 | 11.8° | K. & L. | " | " |
| 3 | Ara-yu | " | 95.0° | abundant | — | 1.76 | 11.5° | " | " | " |
| 4 | Drinking water at Chimizu | — | — | — | — | 3.09 | 11.0° | " | " | " |
| 5 | " at Shin-machi | — | — | — | — | 2.02 | 12.2° | " | " | " |
| | Gas evolving from
Ara-yu | — | — | — | Emanation per litre
of gas at 0° C. | | | " | Date when the
gas was
collected
March 14, 1914 | " |
| | | | | | — | 25.98 | — | | | |

ŌDAKE

Location.—Yamada-mura, Muko-gun, Province Settsu, Hyōgo-ken.

The cold spring "Ōdake-Kōsen"

Analysis (calculated from the original numbers).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1913.

Specific gravity: 1.0022 at 15° C.

Temperature: 17.5° C.

Total residue: ca. 3.73.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Mili-
moles | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0860 | 2.1994 | 2.1994 |
| Sodium ion (Na ⁺) | 0.9442 | 41.0521 | 41.0521 |
| Calcium ion (Ca ⁺⁺) | 0.3015 | 7.5268 | 15.0536 |
| Magnesium ion (Mg ⁺⁺) | 0.0218 | 0.8964 | 1.7928 |
| Ferrous ion (Fe ⁺⁺) | 0.0126 | 0.2256 | 0.4512 |
| | | | 60.5491 |
| Anions. | Grams | Mili-
moles | Milligram-
equivalents |
| Chlorine ion (Cl ⁻) | 1.4468 | 40.0000 | 40.0000 |
| Sulphate ion (SO ₄ ^{''}) | 0.0644 | 0.6703 | 1.3406 |
| Hydrocarbonate ion (HCO ₃ [']) | 1.1662 | 19.1155 | 19.1155 |
| | 4.0435 | 111.6861 | 60.5491 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0780 | | |
| | 4.1215 | | |
| Free carbon dioxide (CO ₂) | 2.4000 | 54.5454 | |
| | 6.5215 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.1640 |
| Sodium chloride (NaCl) | 2.2569 |
| Sodium bicarbonate (NaHCO ₃) | 0.2490 |
| Calcium sulphate (CaSO ₄) | 0.0913 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 1.1111 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.1311 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0401 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0780 |
| | 4.1215 |
| Free carbon dioxide (CO ₂) | 2.4000 |
| | 6.5215 |

Besides these a very small quantity of aluminium ion, and trace of hydrophosphate ion.

The spring thus may be classified as "earthy iron carbonate carbondioxiated common salt spring".

Radio-activity. 13.56×10^{-10} curies in 1 litre of water at 17.5° C.

(Schmidt's electrom.; H. Kibezaki, June 27, 1913).

3.73 Mache's units (recalculated).

IWAI

Location.—Iwai-mura, Iwami-gun, Province Inaba, Tottori-ken.

The springs issue from the Older Tertiary.

The hot spring “Moto-yu”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1888.

Specific gravity: 1.002 at 15° C.

Temperature: 58.5° C.

Total residue: ca. 1.91.

Flow of water: ca. 4066 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents | | Grams |
|--|--------|------------|-----------------------|--|--------|
| Potassium ion (K ⁺) | 0.0303 | 0.7739 | 0.7739 | Sodium chloride (NaCl) | 0.3416 |
| Sodium ion (Na ⁺) | 0.2573 | 11.1626 | 11.1626 | Potassium sulphate (K ₂ SO ₄) | 0.0675 |
| Calcium ion (Ca ⁺⁺) | 0.2894 | 7.2170 | 14.4340 | Sodium sulphate (Na ₂ SO ₄) | 0.3782 |
| Magnesium ion (Mg ⁺⁺) | 0.0078 | 0.3202 | 0.6404 | Calcium sulphate (CaSO ₄) | 0.9828 |
| | | | 27.0109 | Magnesium sulphate (MgSO ₄) | 0.0387 |
| | | | | Silicic acid (meta) (H ₂ SiO ₃) | 0.0623 |
| | | | | | 1.8711 |
| Anions. | Grams | Milli-mols | Milligram-equivalents | | |
| Chlorine ion (Cl ⁻) | 0.2070 | 5.8392 | 5.8392 | | |
| Sulphate ion (SO ₄ ⁼⁼) | 1.0170 | 10.5871 | 21.1742 | | |
| | 1.8088 | 35.9000 | 27.0134 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0623 | | | | |
| | 1.8711 | | | | |

Besides these trace of ferrous, aluminium and hydrophosphate ion, boric acid and organic substances.

The spring thus may be classified as “saline sulphated bitter spring”.

Radio-activity. 2.29 Mache's units in 1 litre of water at 17.2° C.

(Kohl.-Löw. fontact. ; Dr. R. Ishizu, March 13, 1914).

Table 85.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | Apparatus employed | Date when the water was taken | Examined by |
|-----|----------------------------|-------------------------|-----------------------|---|------------------------------|------------------|--------------------|-------------------------------|---------------|
| | | | | | in Mache's units | at (temp. in C.) | | | |
| 1 | Moto-yu <i>Public bath</i> | saline sulphated bitter | 58.5° | ca. 4066 | 2.29 | 17.2° | K. & L. | March 13, 1914 | Dr. R. Ishizu |
| 2 | Bizenya-no-yu | „ | 50.0° | 448 | 1.46 | 18.8° | „ | „ | „ |
| 3 | Hanaya-no-yu | „ | 50.0° | 538 | 1.77 | 18.2° | „ | „ | „ |
| 4 | Iwaiya-no-yu | „ | 60.0° | 749 | 1.15 | 19.0° | „ | „ | „ |
| 5 | Nakajimaya-no-yu | „ | 46.0° | — | 2.87 | 16.7° | „ | „ | „ |
| 6 | Iwami-kwan-no-yu | „ | 46.0° | — | 4.17 | 17.8° | „ | „ | „ |
| 7 | Shimo-nakagawara-no-yu * | — | 37.0° | — | 2.93 | 16.7° | „ | „ | „ |

Measurement was made with the sample brought to the examiner by applicant, taking into account decay of radium emanation with the time.

TOTTORI

Location.—Tottori-shi, Province Inaba, Tottori-ken.

The springs issue from the Alluvium.

YOSHIKATA

The hot spring “Tottori-Onsen”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1906.

Specific gravity: 1.0045 at 15.5° C.

Flow of water: ca. 36 hectolitres in 24 hours.

Temperature: 47.5° C. (air temp. 15° C.).

Total residue: ca. 4.62.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligramm-equivalents |
|--|--------|------------|------------------------|
| Potassium ion (K ⁺) | 0.1657 | 4.2324 | 4.2324 |
| Sodium ion (Na ⁺) | 1.4670 | 63.6432 | 63.6432 |
| Calcium ion (Ca ⁺⁺) | 0.0971 | 2.4214 | 4.8428 |
| Magnesium ion (Mg ⁺⁺) | 0.0158 | 0.6486 | 1.2972 |
| | | | 74.0156 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 1.0217 | 28.8209 | 28.8209 |
| Sulphate ion (SO ₄ ^{''}) | 1.6354 | 17.0247 | 34.0494 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.6801 | 11.1472 | 11.1472 |
| | 5.0828 | 127.9384 | 74.0175 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.4413 | | |
| | 5.5241 | | |

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 1.6850 |
| Potassium sulphate (K ₂ SO ₄) | 0.3690 |
| Sodium sulphate (Na ₂ SO ₄) | 1.7760 |
| Sodium bicarbonate (NaHCO ₃) | 0.8274 |
| Calcium sulphate (CaSO ₄) | 0.3290 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0954 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.4413 |
| | 5.5241 |

Besides these trace of ferrous, manganous, aluminium, bromine, iodine and hydrophosphate ion and boric acid.

The spring thus may be classified as “muriated saline bitter spring”.

Radio-activity. 11.91×10^{-10} curies in 1 litre of water at 18.5° C.

(Schmidt's electrom.; Dr. R. Ishizu, Feb. 17, 1914).

3.28 Mache's units (recalculated).

The hot spring “Takasago-Onsen”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1905.

Specific gravity: 1.0044 at 15° C.

Flow of water: ca. 25 hectolitres in 24 hours.

Temperature: 43° C. (air temp. 15° C.).

Total residue: ca. 4.03.

The spring, being in its composition nearly the same as that of “Tottori-Onsen”, may be classified as “muriated saline bitter spring”.

Radio-activity. 3.45 Mache's units in 1 litre of water at 16.3° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 17, 1914).

TERAMACHI

The cold spring “Kasuga-Onsen”

Analysis (calculated from the original numbers).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1912.

Specific gravity: 1.0023 at 15° C.

Total residue: ca. 3.09.

Temperature: 28.5° C. (air temp. 10° C.).

The spring, being in its composition nearly the same as that of "Tottori-Onsen", may be classified as "muriated saline bitter spring".

Radio-activity. 2.21×10^{-10} curies in 1 litre of water at 18.9° C.

(Schmidt's electrom.; Dr. R. Ishizu, Feb. 16, 1914).

0.61 Mache's units (recalculated).

8.76×10^{-10} curies in 1 litre of gas at 0° C.

(Schmidt's electrom.; Dr. R. Ishizu, Feb. 17, 1914).

2.41 Mache's units (recalculated).

Table 86.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|------------------------------------|---------------------------|----------------------------|---|--|---------------------|---------------------|--------------------|--|---------------|
| | | | | | in
10^{-10} curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kasuga-Onsen | muriated
saline bitter | 28.5°
(air temp. 10.0°) | ca. — | 2.21 | 0.61 | 18.9° | Schm. | Feb. 16, 1914 | Dr. R. Ishizu |
| 2 | Ebisu-Onsen | — | 26.0° (13°) | — | — | 0.98 | 14.2° | K. & L. | " 17. " | " |
| 3 | Yorō-no-yu | — | — | — | — | 0.25 | 13.7° | " | " | " |
| 4 | Tottori-Onsen (<i>Uchi-yu</i>) | muriated
saline bitter | 47.5° (15°) | 36 | 11.91 | 3.28 | 18.5° | Schm. | " | " |
| 5 | " | " | " | " | — | 3.12 | 15.1° | K. & L. | " | " |
| 6 | Takasago-Onsen | " | 43.0° (15°) | 25 | — | 3.45 | 16.3° | " | " | " |
| 7 | Matsukawa Onsen (<i>Soto-yu</i>) | — | 47.0° (15°) | — | — | 4.86 | 12.2° | " | " | " |
| 8 | Shiratama-Onsen | — | 35.6° (14°) | — | — | 3.87 | 13.5° | " | " | " |
| 9 | Kinka-Onsen | — | 24.4° (15°) | 25 | — | 5.83 | 16.5° | " | " | " |
| 10 | Kinoe-Onsen No. 1 | — | — | 15 | — | 3.33 | 13.9° | " | " | " |
| 11 | " No. 2 | — | — | — | — | 1.35 | 13.3° | " | " 16. " | " |
| 12 | Iroha-no-yu | — | — | 15 | — | — | — | — | — | — |
| 13 | Sangai-ryokwan-no-yu | — | — | — | — | — | — | — | — | — |
| 14 | Marufuku-no-yu | — | — | 15 | — | — | — | — | — | — |
| | Gas evolving from
Kasuga-Onsen | — | — | — | Emanation per litre
of gas at 0° C. | | | Schm. | Date when the
gas was
collected
Feb. 17, 1914 | Dr. R. Ishizu |
| | | | | | 8.76 | 2.41 | — | | | |

YUDANI

Location.—Meiji-mura, Yazu-gun, Province Inaba, Tottori-ken.

The spring issues from the Tertiary.

The cold spring "Kabu-yu"

Analysis (calculated from the salt table).

Analysed by the Osaka Imperial Hygienic Laboratory. 1888.

Specific gravity: 1.0029 at 15° C.

Temperature: 32° C. (air temp. 11° C.).

Flow of water: ca. 90 hectolitres in 24 hours.

Total residue: ca. 2.75.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-moles | Milligram-equivalents |
|--|--------|-------------|-----------------------|
| Potassium ion (K ⁺) | 0.0557 | 1.4227 | 1.4227 |
| Sodium ion (Na ⁺) | 0.4974 | 41.1887 | 41.1887 |
| Calcium ion (Ca ⁺⁺) | 0.0589 | 1.4688 | 2.9376 |
| Ferrous ion (Fe ⁺⁺) | 0.0025 | 0.0417 | 0.0894 |
| | | | 45.7554 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.9506 | 26.8152 | 26.8152 |
| Sulphate ion (SO ₄ ^{''}) | 0.1070 | 1.1139 | 2.2278 |
| Hydrocarbonate ion (HCO ₃ [']) | 1.0123 | 19.524 | 16.5924 |
| | | | 45.7554 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0831 | | |
| | 3.2195 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | |
|---|--------|
| Potassium chloride (KCl) | 0.1060 |
| Sodium chloride (NaCl) | 1.4857 |
| Sodium bicarbonate (NaHCO ₃) | 1.3273 |
| Calcium sulphate | |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0578 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0080 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0831 |
| | 3.2195 |

Besides these trace of ammonium, magnesium, bromine and iodine ion, boric acid and organic substances.

The spring thus may be classified as "alkaline common salt spring".

Radio-activity. 15.42 × 10⁻¹⁰ curies in 1 litre of water at 13.6° C.

4.24 Mache's units (recalculated).

86.54 × 10⁻¹⁰ curies in 1 litre of gas at 0° C.

23.80 Mache's units (recalculated).

(Schmidt's electrom.; Dr. R. Ishizu, Feb. 20, 1914).

YOSHIOKA

Location.—Yoshioka-mura, Kedaka-gun, Province Inaba, Tottori-ken.

The springs issue from the Tertiary.

The hot spring "Kamei-dono-no-yu"

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1889. Temperature: 50° C.

Specific gravity: 1.003 at 16.5° C.

Total residue: ca. 0.56.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-moles | Milligram-equivalents |
|--|--------|-------------|-----------------------|
| Potassium ion (K ⁺) | 0.0136 | 0.3474 | 0.3474 |
| Sodium ion (Na ⁺) | 0.1653 | 7.1714 | 7.1714 |
| Calcium ion (Ca ⁺⁺) | 0.0170 | 0.4239 | 0.8478 |
| Magnesium ion (Mg ⁺⁺) | 0.0011 | 0.0457 | 0.0914 |
| | | | 8.4580 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.1119 | 3.1566 | 3.1566 |
| Sulphate ion (SO ₄ ^{''}) | 0.1462 | 1.5220 | 3.0440 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.1377 | 2.2574 | 2.2574 |
| | 0.5928 | 14.9244 | 8.4580 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0727 | | |
| | 0.6655 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | |
|---|--------|
| Sodium chloride (NaCl) | 0.1847 |
| Potassium sulphate (K ₂ SO ₄) | 0.0302 |
| Sodium sulphate (Na ₂ SO ₄) | 0.1917 |
| Sodium bicarbonate (NaHCO ₃) | 0.1108 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0688 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0067 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0727 |
| | 0.6656 |

Besides these trace of ferrous, hydrophosphate ion and boric acid.

The spring thus may be classified as "simple thermal".

Radio-activity. 4.36 Mache's units in 1 litre of water at 10.2° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 18, 1914).

The hot spring "Naka-no-yu"

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1888. Specific gravity: 1,0010 at 15° C.

Temperature: 52,5° C. (air temp. 15,5° C.). Total residue: ca. 0,58.

The spring, being in its composition nearly the same as that of "Kamei-dono-no-yu", may be classified as "simple thermal".

Radio-activity. 3,19 Mache's units in 1 litre of water at 9,6° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 18, 1914).

The hot spring "Shimo-no-yu"

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1888. Specific gravity: 1,0005 at 15° C.

Temperature: 42,5° C. (air temp. 8° C.). Total residue: ca. 0,68.

The spring, being in its composition nearly the same as that of "Kamei-dono-no-yu", differing only in containing hydrogen sulphide from the latter, may be classified as "sulphur spring".

Radio-activity. 3,07 Mache's units in 1 litre of water at 10,2° C.

9,63 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 18, 1914).

Table 87.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|--|--------------------|----------------|----------------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|---------------|
| | | | | in
10 ¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kabu-yu * | simple | 56,5 ^{an temp.}
(18) | 10,33 | 2,84 | 9,5° | Schm. | Feb. 18, 1914 | Dr. R. Ishizu |
| 2 | Kamei-dono-no-yu * | " | 50,0° | — | 4,36 | 10,2° | K. & L. | " | " |
| 3 | Naka-no-yu * | " | 52,5° (15,5°) | — | 3,19 | 9,6° | " | " | " |
| 4 | Nakajimaya-no-yu | " | 53,5° | — | 3,17 | 10,2° | " | " | " |
| 5 | Shimo-no-yu * | sulphur | 42,5° (8°) | — | 3,07 | 10,2° | " | " | " |
| 6 | Tonosama-no-yu * | simple | — | — | — | — | — | — | — |
| 7 | Kadoya-no-yu | " | — | — | — | — | — | — | — |
| 8 | Kagi-no-yu | " | — | — | — | — | — | — | — |
| 9 | Hiiragiya-no-yu | " | — | — | — | — | — | — | — |
| Emanation per litre
of gas at 0° C. | | | | | | | | | |
| 1 | Kabu-yu * | — | — | 99,13 | 27,26 | — | Schm. | Feb. 18, 1914 | Dr. R. Ishizu |
| 2 | Shimo-no-yu * | — | — | — | 9,63 | — | K. & L. | " | " |

* These springs are for public use.

HAMAMURA

Location.—Seijō-mura, Kedaka-gun, Province Inaba, Tottori-ken.

The springs issue from the Alluvium with Granite Base.

KACHIMI

The hot spring “Nakataya-no-yu”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1888.

Temperature: 56° C.

Specific gravity: 1.0013 at 15° C.

Total residue: ca. 1.36.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|--------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0304 | 0.7765 | 0.7765 |
| Sodium ion (Na ⁺) | 0.3146 | 13.6486 | 13.6486 |
| Calcium ion (Ca ⁺⁺) | 0.0608 | 1.5162 | 3.0324 |
| Magnesium ion (Mg ⁺⁺) | 0.0037 | 0.1519 | 0.3038 |
| | | | 17.7613 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.2894 | 8.1636 | 8.1636 |
| Sulphate ion (SO ₄ ^{''}) | 0.4612 | 4.8012 | 9.6024 |
| | 1.1601 | 29.0580 | 17.7660 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.3300 | | |
| | 1.4901 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|--|--------|
| Sodium chloride (NaCl) | 0.4775 |
| Potassium sulphate (K ₂ SO ₄) | 0.0678 |
| Sodium sulphate (Na ₂ SO ₄) | 0.3903 |
| Calcium sulphate (CaSO ₄) | 0.2062 |
| Magnesium sulphate (MgSO ₄) | 0.0183 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.3300 |
| | 1.4901 |

Besides these a small quantity of free carbon dioxide and organic substances and trace of aluminium, bromine and hydrophosphate ion.

The spring thus may be classified as “muriated saline bitter spring”.

Radio-activity. 31.18 × 10⁻¹⁰ curies in 1 litre of water at 16.3° C.

8.57 Mache's units (recalculated).

197.62 × 10⁻¹⁰ curies in 1 litre of gas at 0° C.

54.35 Mache's units (recalculated).

(Schmidt's electrom.; Dr. R. Ishizu, Feb. 19, 1914).

HAMAMURA

The hot spring “Suzukiya-no-yu”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1888.

Temperature: 49° C.

Specific gravity: 1.0008.

Total residue: ca. 1.07.

The spring, being in its composition nearly the same as that of “Nakataya-no-yu”, may be classified as “muriated saline bitter spring”.

Radio-activity. 5.35 Mache's units in 1 litre of water at 13.0° C.

(Kohl-Löw. contact.; Dr. R. Ishizu, Feb. 19, 1914).

Table 88.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---|---------------------------|--------------------------|---|--|---------------------|---------------------|--------------------|---------------------------------------|---------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Sagi no-yu (<i>Public bath</i>) | — | 51.5° | ca.
126 | — | 5.89 | 14.1° | K. & L. | Feb. 19, 1914 | Dr. R. Ishizu |
| 2 | Nakataya-no-yu | muriated
saline bitter | 50.0° | — | 31.18 | 8.57 | 16.3° | Schm. | " | " |
| 3 | Private bath (<i>Y. Kinoshita</i>) | — | 56.0° | — | — | 7.01 | 13.0° | K. & L. | " | " |
| 4 | " (<i>Kurezumi Kōsha</i>
<i>Vent No. 1</i>) | — | 53.0° | — | — | 7.15 | 13.0° | " | " | " |
| 5 | Kyado-yu (<i>Vent No. 1</i>
<i>(Public bath)</i>) | — | 48.0° | — | — | 1.90 | 13.4° | " | " | " |
| 6 | Suzukiya no-yu | muriated
saline bitter | 49.0° | — | — | 5.35 | 13.0° | " | " | " |
| 7 | Kadoya-no-yu | — | 45.0° | — | — | 3.70 | 12.9° | " | " | " |
| 8 | Tabakoya-no-yu | — | 45.0° | — | — | 3.89 | 13.6° | " | " | " |
| | | | | | Emanation per litre
of gas at 0° C. | | | | Date when
the gas was
collected | |
| 1 | Gas evolving from:
Nakataya-no-yu | — | — | — | 197.62 | 54.35 | — | Schm. | Feb. 19, 1914 | " |
| 2 | Private bath (<i>Kurezumi-
Kōsha</i>) <i>Vent No. 2</i> | — | 51.0° | — | — | 13.37 | — | K & L. | " | " |

TÔGÔ

Location.—Tôgô-mura, Tôhaku-gun, Province Hôki, Tottori-ken.

The springs issue from the Andesite overlaid by Alluvium.

The hot spring "Yôjyô-kwan-no-yu"

Analysis (calculated from the salt table).

Analysed by the Osaka Imperial Hygienic Laboratory. 1888.

Specific gravity: 1.0007.

Total residue: ca. 0.93.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|--------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0204 | 0.5201 | 0.5201 |
| Sodium ion (Na ⁺) | 0.2420 | 10.4945 | 10.4945 |
| Calcium ion (Ca ⁺⁺) | 0.0504 | 1.2569 | 2.5138 |
| Magnesium ion (Mg ⁺⁺) | 0.0112 | 0.4598 | 0.9196 |
| Ferrous ion (Fe ⁺⁺) | 0.0013 | 0.0233 | 0.0466 |
| | | | 14.4946 |
| Anions. | Grams | Milli-
moles | Milligram-
equivalents |
| Chlorine ion (Cl ⁻) | 0.3634 | 10.2511 | 10.2511 |
| Sulphate ion (SO ₄ ⁻²) | 0.1387 | 1.4439 | 2.8878 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0.0824 | 1.3503 | 1.3503 |
| | | | 14.4892 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0750 | | |
| | 0.9848 | | |

| | Grams |
|---|--------------|
| Potassium chloride (KCl) | 0.0389 |
| Sodium chloride (NaCl) | 0.5691 |
| Sodium bicarbonate (NaHCO ₃) | 0.0650 |
| Calcium sulphate (CaSO ₄) | 0.1707 |
| Magnesium sulphate (MgSO ₄) | 0.0231 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0390 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0040 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0750 |
| | <hr/> 0.9848 |

Besides these trace of ammonium, aluminium, bromine and hydrophosphate ion and organic substances.

The spring thus may be classified as "simple thermal".

Table 89.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|------------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|-------------------|--------------------|----------------------------------|---------------|
| | | | | | in
10 ¹⁰ curies | in
Mache's units | at
temp. in C. | | | |
| 1 | Yōjyō-kwan-no-yu <i>Jet No. 1</i> | simple | 50.0° | ca.
1730 | 11.17 | 3.07 | 10.6° | Schm | Feb. 23, 1914 | Dr. R. Ishizu |
| 2 | " | " | " | — | — | 4.06 | 10.2° | K. & L. | " | " |
| 3 | " <i>Jet No. 3</i> | " | 45.0° | — | — | 5.02 | 10.2° | " | " | " |
| 4 | " <i>Jet No. 5</i> | " | 31.0° | — | — | 3.04 | 10.0° | " | " | " |
| 5 | " <i>Jet No. 6</i> | " | 46.5° | — | — | 1.64 | 9.6° | " | " | " |
| 6 | Matsuzaki-kwan-no-yu | salt | 36.0° | — | — | 5.87 | 11.2° | " | " | " |
| 7 | Spring <i>in Matsuzaki station</i> | " | 32.0° | — | — | 4.55 | 11.7° | " | " | " |

ASŌZU

Location.—Asōzu-mura, Tōhaku-gun, Province Hōki, Tottori-ken.

The springs issue from the Andesite overlaid by Alluvium.

The hot spring “Moto-yu”

Analysis.

Analyst: Unknown. Specific gravity: 1.00017 at 15° C. Temperature: 56° C.

Total residue: ca. 1.64. Flow of water: ca. 360 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Sodium chloride (NaCl) | 1.2056 | Calcium sulphate (CaSO ₄) | 0.0648 |
| Potassium sulphate (K ₂ SO ₄) | 0.0945 | Silica (SiO ₂) | 0.0572 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0106 | | 1.4327 |

Besides these trace of alumina, phosphoric and boric acid and organic substances.

The spring may be classified as “common salt spring”.

Radio-activity. 4.33 Mache's units in 1 litre of water at 11.0° C.

(Kohl-Löw. fontact.; Dr. R. Ishizu, Feb. 21, 1914).

The hot spring “Asahi-kwan-no-yu”

Analysis.

Analyst: Unknown. Specific gravity: 1.00013 at 15° C. Temperature: 56° C.

Total residue: ca. 1.91. Flow of water: ca. 360 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of “Moto-yu”, differing only in containing hydrogen sulphide from the latter, may be classified as “muriated sulphur spring”.

Radio-activity. 3.56 Mache's units in 1 litre of water at 11.3° C.

(Kohl-Löw. fontact.; Dr. R. Ishizu, Feb. 21, 1914).

Table 90.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water
in
Mache's units
at
(temp. in C.) | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------|---------------------|--------------------------|---|---|-------|--------------------|----------------------------------|---------------|
| 1 | Moto-yu | common salt | 56,0° | ca.
360 | 4,33 | 11,0° | K. & L. | Feb. 21, 1914 | Dr. R. Ishizu |
| 2 | Asahi-kwan no-yu | muriated
sulphur | 56,0° | 360 | 3,56 | 11,3° | " | " | " |
| 3 | Sugimoto-no-yu | " | 46,0° | — | 2,73 | 12,2° | " | " | " |
| 4 | Nisshin-kwan-no-yu | — | 48,0° | — | 1,89 | 11,2° | " | " | " |

SEKIGANE

Location.—Yaokuri-mura, Tōhaku-gun, Province Hōki, Tottori-ken.

The springs issue from the Granite.

The hot spring "Kabu-yu"

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1888. Temperature: 42° C.

Specific gravity: 1,000 at 15° C.

Total residue: ca. 0,52.

In 1 kilogram of the mineral water are contained:

| | Grams | Milli-
moles | Milligram-
equivalents |
|--|----------------|-----------------|---------------------------|
| Cations. | | | |
| Potassium ion (K'). | 0,0192 | 0,4904 | 0,4904 |
| Sodium ion (Na'). | 0,1774 | 7,6963 | 7,6963 |
| Calcium ion (Ca''). | 0,0144 | 0,3591 | 0,7182 |
| Ferrous ion (Fe''). | 0,0016 | 0,0286 | 0,0572 |
| | | | 8,9621 |
| Anions. | | | |
| Chlorine ion (Cl') | 0,1276 | 3,5994 | 3,5994 |
| Sulphate ion (SO ₄ ''). | 0,1086 | 1,1306 | 2,2612 |
| Hydrocarbonate ion (HCO ₃ '). | 0,1891 | 3,0995 | 3,0995 |
| | 0,6379 | 16,4039 | 8,9601 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0867 | | |
| | 0,7246 | | |
| Free carbon dioxide (CO ₂) | not determined | | |
| Free hydrogen sulphide (H ₂ S) | 0,0002 | | |
| | 0,7248 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Sodium chloride (NaCl). | 0,2106 |
| Potassium sulphate (K ₂ SO ₄) | 0,0427 |
| Sodium sulphate (Na ₂ SO ₄) | 0,0751 |
| Sodium bicarbonate (NaHCO ₃) | 0,2558 |
| Calcium sulphate (CaSO ₄) | 0,0487 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0050 |
| Silicic acid (meta) (H ₂ SiO ₃). | 0,0867 |
| | 0,7246 |
| Free hydrogen sulphide (H ₂ S). | 0,0002 |
| | 0,7248 |

Besides these trace of magnesium, aluminium and hydrophosphate ion, boric acid and a small quantity of organic substances.

The spring thus may be classified as "sulphur spring".

Radio-activity. 33,47 Mache's units in 1 litre of water at 9,4° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 23, 1914).

The hot spring “Kami-jyaya-no-yu”

Analysis (calculated from the salt table).

Analysed by the Osaka Imperial Hygienic Laboratory. 1888.

Specific gravity: 1,0003.

Temperature: 44,5° C.

Flow of water: ca. 270 hectolitres in 24 hours.

Total residue: ca. 0,54.

The spring, being in its composition nearly the same as that of “Kabu-yu”, may be classified as “sulphur spring”.

Radio-activity. 32,41 Mache's units in 1 litre of water at 10,4° C.

(Kohl-Low, contact, Dr. R. Ishizu, Feb. 23, 1914).

Table 91.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|----------------------------|----------------|-----------------------|---|------------------------------|------------------|----------------|--------------------|-------------------------------|---------------|
| | | | | | in 10 ⁻¹⁰ curies | in Mache's units | at temp. in C. | | | |
| 1 | Kabu-yu <i>Public bath</i> | sulphur | 42,0° | ca. — | — | 33,47 | 9,4° | K. & L. | Feb. 23, 1914 | Dr. R. Ishizu |
| 2 | Shimo-jyaya-no-yu | .. | 44,0° | 126 | 109,51 | 30,12 | 10,2° | Schm. | .. | .. |
| 3 | Kami-jyaya-no-yu | .. | 44,5° | 270 | — | 32,41 | 10,4° | K. & L. | .. | .. |
| 4 | Inkyo-jyaya-no-yu | .. | 42,0° | 90 | — | 22,07 | 10,9° | .. | .. | .. |
| 5 | Kon-ya-no-yu | .. | 46,0° | 108 | — | 10,65 | 9,9° | .. | .. | .. |
| 6 | Shin-yu | .. | — | 171 | — | — | — | — | — | — |

MISASA

Location.—Misasa-mura, Tōhaku-gun, Province Hōki, Tottori-ken.

The springs issue from the Granite.

The hot spring “Naka-no-yu”

Analysis (calculated from the salt table).

Analysed by the Osaka Imperial Hygienic Laboratory. 1888.

Temperature: 56,5° C.

Specific gravity: 1,0005.

Total residue: ca. 1,24.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols. | Milli-equivs. |
|--|----------------|-------------|---------------|
| Potassium ion (K ⁺) | 0,0375 | 0,9553 | 0,9553 |
| Sodium ion (Na ⁺) | 0,4118 | 17,8655 | 17,8655 |
| Calcium ion (Ca ⁺⁺) | 0,0317 | 0,7905 | 1,5810 |
| | | | 20,4018 |
| Anions. | Grams | Milli-mols. | Milli-equivs. |
| Chlorine ion (Cl ⁻) | 0,4500 | 12,6939 | 12,6939 |
| Sulphate ion (SO ₄ ⁼⁼) | 0,0957 | 0,9963 | 1,9926 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0,3487 | 5,7153 | 5,7153 |
| | 1,3754 | 3,6968 | 20,4018 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,1019 | | |
| | 1,4773 | | |
| Free carbon dioxide (CO ₂) | not determined | | |
| Free hydrogen sulphide (H ₂ S) | 0,0023 | | |
| | 1,4796 | | |

| | Grams |
|--|--------|
| Potassium chloride (KCl) | 0,0401 |
| Sodium chloride (NaCl) | 0,7112 |
| Potassium sulphate (K ₂ SO ₄) | 0,0365 |
| Sodium bicarbonate (NaHCO ₃) | 0,4803 |
| Calcium sulphate (CaSO ₄) | 0,1073 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,1019 |
| | 1,4773 |
| Free hydrogen sulphide (H ₂ S) | 0,0023 |
| | 1,4796 |

Besides these trace of ferrous, aluminium, bromine, iodine and hydrophosphate ion, boric acid and organic substances.

The spring thus may be classified as “**muriated sulphur spring**”.

Radio-activity. 11.04 Mache's units in 1 litre of water at 12.0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 22, 1914).

The hot spring “Sakaya-no-yu” No. 2

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1889.
Temperature: 67° C.

Specific gravity: 1.0003 at 16.5° C.

Total residue: ca. 0.87.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents |
|--|--------|----------------|---------------------------|
| Potassium ion (K ⁺) | 0.0178 | 0.4547 | 0.4547 |
| Sodium ion (Na ⁺) | 0.2649 | 11.4924 | 11.4924 |
| Calcium ion (Ca ⁺⁺) | 0.0194 | 0.4838 | 0.9676 |
| Magnesium ion (Mg ⁺⁺) | 0.0003 | 0.0123 | 0.0246 |
| | | | 12.9393 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.2991 | 8.4372 | 8.4372 |
| Sulphate ion (SO ₄ ^{''}) | 0.0812 | 0.8453 | 1.6906 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.1715 | 2.8115 | 2.8115 |
| | 0.8542 | 24.5372 | 12.9393 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0922 | | |
| | 0.9464 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 0.4936 |
| Potassium sulphate (K ₂ SO ₄) | 0.0396 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0879 |
| Sodium bicarbonate (NaHCO ₃) | 0.1529 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0787 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0015 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0922 |
| | 0.9464 |

Besides these trace of ferrous, aluminium and hydrophosphate ion and boric acid.

The spring thus may be classified as “**simple thermal**”.

Radio-activity. 60.35 Mache's units in 1 litre of water at 15.5° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, March 12, 1914).

The hot spring “Kyōyūchi-no-yu”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1889.
Temperature: 71.5° C.

Specific gravity: 1.0005 at 16.5° C.

Total residue: ca. 0.86.

The spring, being in its composition nearly the same as that of “Sakaya-no-yu” No. 2, may be classified as “**simple thermal**”.

Radio-activity. 13.81 Mache's units in 1 litre of water at 12.5° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Feb. 22, 1914).

The hot spring “Tōkyōya-no-yu”

Analysis.

Analyst: Unknown.

Temperature: 72° C.

Specific gravity: 1.0009 at 15° C.

Total residue: ca. 1.16.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Sodium chloride (NaCl) | 0.7112 | Calcium sulphate (CaSO ₄) | 0.0505 |
| Potassium sulphate (K ₂ SO ₄) | 0.0785 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0060 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0325 | Silica (SiO ₂) | 0.0780 |
| Sodium carbonate (Na ₂ CO ₃) | 0.2075 | | 1.1642 |

Besides these trace of magnesia, alumina, bromine, iodine, phosphoric and boric acid and organic substances.

The spring may be classified as “**common salt spring**”.

Radio-activity. 20.23 Mache's units in 1 litre of water at 13.1° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, March 11, 1914).

Table 92.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---|------------------|------------------------------|--|---------------------|-------------------|--------------------|----------------------------------|---------------|
| | | | | in
10 cc
in
Maché's units | in
Maché's units | at
temp. in C. | | | |
| 1 | Kabu-yu (<i>Public bath</i>) | — | 45.0° ^(air temp.) | 37.19 | 10.23 | 12.0° | Schm. | Feb. 22, 1914 | Dr. R. Ishizu |
| 2 | Drinking water (<i>N. Okazaki</i>) | — | — | — | 4.19 | 15.1° | K. & L. | March 12, " | " |
| 3 | Ochaya-no-yu | — | 63.5° | — | 27.88 | 12.1° | " | Feb. 22, " | " |
| 4 | Shin-ya-no-yu | — | 44.5° | — | 27.87 | 15.9° | " | March 12, " | " |
| 5 | Mae-iyaya-no-yu | — | 61.0° | — | 6.69 | 16.0° | " | " | " |
| 6 | Naka-no-yu (<i>Public bath</i>) | muriated sulphur | 56.5° | — | 11.04 | 12.0° | " | Feb. 22, " | " |
| 7 | Hashizuya-no-yu | — | 63.0° | — | 15.52 | 15.9° | " | March 12, " | " |
| 8 | Private bath (<i>S. Mifune</i>) | — | 63.0° | — | 18.55 | 15.3° | " | " | " |
| 9 | Bun-aburaya-no-yu | — | 67.0° | — | 24.46 | 13.3° | " | " 11, " | " |
| 10 | Tōfuya-no-yu | — | 57.0° | — | 24.46 | 11.5° | " | " | " |
| 11 | Hanaya-no-yu | — | 60.0° | — | 72.65 | 12.6° | " | " | " |
| 12 | Sakaya-no-yu <i>No. 1</i> | — | 64.0° | — | 63.47 | 15.7° | " | " 12, " | " |
| 13 | " <i>No. 2</i> | simple | 67.0° | — | 60.35 | 15.5° | " | " | " |
| 14 | Private bath * (<i>K. Tsugawa</i>) | carbonated (?) | 52.0° | — | 12.85 | 12.5° | " | Feb. 22, " | " |
| 15 | Spring A (") | simple | 11.5° ^(14°) | — | 3.02 | 15.3° | " | March 12, " | " |
| 16 | Aburaya-no-yu | — | 48.5° | — | 19.20 | 12.6° | " | " 11, " | " |
| 17 | Uenoya-no-yu | — | 46.0° | — | 24.75 | 12.4° | " | " | " |
| 18 | Private bath (<i>K. Kishida</i>) | — | 49.0° | — | 58.48 | 12.6° | " | " | " |
| 19 | Iwa-yu (<i>Otoko-yu</i>) | — | 54.5° | — | 36.02 | 13.3° | " | " | " |
| 20 | " (<i>Onna-yu</i>) | — | 56.0° | — | 39.70 | 13.3° | " | " | " |
| 21 | Private bath (<i>T. Matsuiara</i>) | — | 71.0° | 516.87 | 142.14 | 16.3° | Schm. | " 12, " | " |
| 22 | " | — | " | — | 121.99 | 14.0° | K. & L. | " | " |
| 23 | Akazakiya-no-yu | — | 75.0° | — | 58.18 | 13.5° | " | " 11, " | " |
| 24 | Kyōyūchi-no-yu | simple | 71.5° | — | 13.81 | 12.5° | " | Feb. 22, " | " |
| 25 | Tokyōya-no-yu | common salt | 72.0° | — | 20.23 | 13.1° | " | March 11, " | " |
| 26 | Kiya-no-yu | — | 58.5° | — | 28.37 | 12.3° | " | " | " |
| 27 | Seitō-kwan-no-yu | — | 51.0° ^(8°) | 371.83 | 102.25 | 11.7° | Schm. | " | " |
| 28 | " | — | " | — | 107.82 | 12.2° | K. & L. | " | " |
| 29 | Spring B (<i>River bed</i>) | — | 69.5° | — | 28.57 | 15.7° | " | " 12, " | " |
| 30 | Spring C | — | 33.5° ^(10°) | — | 15.18 | 12.0° | " | Feb. 22, " | " |
| | Gas evolving from
Kabu-yu (<i>Public bath</i>) | — | | Emanation per litre
of gas at 0° C. | | | Schm. | Feb. 22, 1914 | " |
| | | | | 228.19 | 62.75 | — | | | |

* The water originates in an artesian well, about 52 ft. in depth.

TAMATSUKURI

Location.—Tamayu-mura, Yatsuka-gun, Province Izumo, Shimane-ken.

The springs issue from the Tertiary.

The hot spring “Kami-no-yu”

Analysis.

Analysed by the Matsue Garrison Hospital. 1909.

Specific gravity: 1.0023 at 15° C.

Temperature: 64° C.

Flow of water: ca. 1953 hectolitres in 24 hours.

Total residue: ca. 1.58.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Potassium chloride (KCl) | 0.0415 | Calcium sulphate (CaSO ₄) | 0.1740 |
| Sodium chloride (NaCl) | 0.9125 | Magnesium sulphate (MgSO ₄) | 0.0254 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0375 | | 1.2350 |
| Sodium bicarbonate (NaHCO ₃) | 0.0481 | | |

Besides these a small quantity of iron and trace of bromine, phosphoric and boric acid, silica and organic substances.

The spring may be classified as “common salt spring”.

Radio-activity. 10.79×10⁻¹⁰ curies in 1 litre of water at 15.1° C.

2.97 Mache's units (recalculated).

101.79×10⁻¹⁰ curies in 1 litre of gas at 0° C.

27.99 Mache's units (recalculated).

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, Feb. 25, 1914).

Table 93.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|--------------------|-------------------------------------|----------------|--------------------------|---|--|---------------------|-------------------|---------------------------------------|----------------------------------|-------------------------------|
| | | | | | in
10 ¹⁰ curies | in
Mache's units | at
temp. in C. | | | |
| 1 | Kami-no-yu (<i>Public bath</i>) | common
salt | 64.0° | ca.
1953 | 10.79 | 2.97 | 15.1° | Schm. | Feb. 25, 1914 | (Dr. R. Ishizu
& T. Saitō) |
| 2 | Tonosama-yu („) | — | 61.5° | — | — | 2.37 | 13.6° | K. & L. | „ | „ |
| 3 | Private bath (<i>S. Hasegawa</i>) | — | 54.0° | — | — | 5.14 | 14.5° | „ | „ | „ |
| Gas evolving from: | | | | | Emanation per litre
of gas at 0° C. | | | Date when the
gas was
collected | | |
| 1 | Kami-no-yu | — | — | — | 101.79 | 27.99 | — | Schm. | Feb. 25, 1914 | „ |
| 2 | River-bed <i>Vent A</i> | — | — | — | 128.04 | 35.21 | — | „ | „ | „ |
| 3 | „ <i>Vent B</i> | — | — | — | — | 24.44 | — | K. & L. | „ | „ |
| 4 | „ <i>Vent C</i> | — | — | — | — | 28.44 | — | „ | „ | „ |

HIROSE

Location.—Iinashi-mura, Nogi-gun, Province Izumo, Shimane-ken.

The springs issue from the Tertiary.

Table 94.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------------------|----------------|--------------------------|---------------------------------|---------------------|-------------------|--------------------|----------------------------------|---------------------------|
| | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
temp. in C. | | | |
| 1 | Sagi-no-yu | .. | 43.0° | 9.91 | 2.73 | 13.2° | Schm. | Feb. 27, 1914 | Dr. R. Ishizu
T. Saitō |
| 2 | Private bath (<i>Asylum</i>) | — | 37.0° | — | 2.06 | 10.2° | K. & L. | .. | " |

GAKUTŌ

Location.—Shōbara-mura, Hinokawa-gun, Province Izumo, Shimane-ken.

The springs issue from the Tertiary.

Table 95.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|----------------|----------------|--------------------------|---|---------------------------------|-------------------|--------------------|----------------------------------|---------------------------|
| | | | | | in
Mache's units | at
temp. in C. | | | |
| 1 | Yunokawa-Kōsen | salt | 27.0° | ca. 864 | 0.42 | 10.1° | K. & L. | Feb. 26, 1914 | Dr. R. Ishizu
T. Saitō |
| 2 | Sudani-yakutō | " | — | — | 0.72 | 10.1° | .. | .. | " |

USHIO

Location.—Ushio-mura, Ōhara-gun, Province Izumo, Shimane-ken.

The springs issue from the Andesite.

The hot spring "Ushio-no-yu"

Temperature: 41.5° C. (air temp. 14.8° C.).

Total residue: ca. 1.22.

Flow of water: ca. 441 hectolitres in 24 hours.

From the result of an old analysis made in 1881, the spring seems to be "bitter spring".

Radio-activity. 1.19 × 10⁻¹⁰ curies in 1 litre of water at 18.3° C.

0.33 Mache's units (recalculated).

9.03 × 10⁻¹⁰ curies in 1 litre of gas at 0° C.

2.48 Mache's units (recalculated).

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, March 1, 1914).

YUMURA

Location.—Yu-mura, Nita-gun, Province Izumo, Shimane-ken.

The springs issue from the Granite.

The hot spring “Shitsuni-no-yu”

Temperature: 43° C. (air temp. 8° C.).

Total residue: ca. 0.32.

Flow of water: ca. 3888 hectolitres in 24 hours.

From the result of an old analysis made in 1881, the spring seems to be “simple thermal”.

Radio-activity. 13.60 × 10⁻¹⁰ curies in 1 litre of water at 16.1° C.

3.74 Mache's units (recalculated).

84.30 × 10⁻¹⁰ curies in 1 litre of gas at 0° C.

23.18 Mache's units (recalculated).

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, Feb. 28, 1914).

KAWAI

Location.—Kawai-mura, Anno-gun, Province Iwami, Shimane-ken.

The springs issue from the Granite.

The cold spring “Tōkōji-no-yu”

Analysis.

Analysed by the Hygienic Laboratory of Shimane-ken. 1911.

Specific gravity: 1.0079 at 16° C.

Temperature: 14° C. (air temp. 12° C.).

Total residue: ca. 9.77.

Flow of water: ca. 29 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|---|---------|
| Potassium chloride (KCl) | 0.2196 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0440 |
| Sodium chloride (NaCl) | 5.6670 | Alumina (Al ₂ O ₃) | 0.0188 |
| Calcium chloride (CaCl ₂) | 2.3210 | Silica (SiO ₂) | 0.0580 |
| Calcium sulphate (CaSO ₄) | 0.9313 | | 10.5755 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.6607 | Free carbon dioxide (CO ₂) | 2.2815 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.6551 | | 12.8570 |

Besides these a small quantity of boric acid.

The spring may be classified as “earth-muriated iron carbonate carbondioxated common salt spring”.

Radio-activity. 1.01 Mache's units in 1 litre of water at 14.9° C.

2.03 Mache's units in 1 litre of gas at 0° C.

(Kohl-Löw. fontact.; Dr. R. Ishizu and T. Saitō, March 3, 1914).

The cold spring “Urisaka-no-yu”

Analysis.

Analysed by the Hygienic Laboratory of Shimane-ken. 1911.

Specific gravity: 1.0103 at 18° C.

Temperature: 13° C. (air temp. 12° C.).

Total residue: ca. 10.20.

Flow of water: ca. 29 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|---|---------|
| Potassium chloride (KCl) | 0.3029 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0114 |
| Sodium chloride (NaCl) | 7.2197 | Alumina (Al ₂ O ₃) | 0.0044 |
| Potassium sulphate (K ₂ SO ₄) | 0.2174 | Silica (SiO ₂) | 0.0700 |
| Magnesium chloride (MgCl ₂) | 0.4062 | | 10.3574 |
| Calcium sulphate (CaSO ₄) | 1.0545 | Free carbon dioxide (CO ₂) | 2.0736 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 1.0709 | | 12.4310 |

Beside these a very small quantity of boric acid.

The spring may be classified as “carbondioxated common salt spring”.

Radio-activity. 1.18 Mache's units in 1 litre of water at 13.2° C.

2.21 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu and T. Saitō, March 3, 1914).

SAHIME

Location.—Sahime-mura, Anno-gun, Province Iwami, Shimane-ken.

The springs issue from the Enstatite Andesite.

KOYABARA

The hot spring “Koyabara-Onsen”

Analysis.

Analysed by the Hygienic Laboratory of Shimane-ken. 1911.

Specific gravity: 1.0032 at 17° C.

Temperature: 38.2° C. (air temp. 8° C.).

Total residue: ca. 5.28.

Flow of water: ca. 540 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|---|--------|
| Sodium chloride (NaCl) | 3.9120 | Alumina (Al ₂ O ₃) | 0.0180 |
| Calcium chloride (CaCl ₂) | 0.2837 | Silica (SiO ₂) | 0.1560 |
| Calcium sulphate (CaSO ₄) | 0.3313 | | 5.8395 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.6889 | Free carbon dioxide (CO ₂) | 0.9053 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.4172 | | 6.7448 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0324 | | |

Beside these a small quantity of boric acid.

The spring may be classified as “iron carbonate common salt spring”.

Radio-activity. 13.65×10⁻¹⁰ curies in 1 litre of water at 18.7° C.

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, March 4, 1914).

3.75 Mache's units (recalculated).

29.53×10⁻¹⁰ curies in 1 litre of gas at 0° C.

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, March 3, 1914).

8.12 Mache's units (recalculated).

SHIGAKU

The springs issue from the Enstatite Andesite.

The hot spring “Shigaku-Onsen”

Analysis.

Analysed by the Hygienic Laboratory of Shimane-ken. 1911.

Specific gravity: 1.0028 at 17° C.

Temperature: 46.5° C. (air temp. 18° C.).

Total residue: ca. 2.39.

Flow of water: ca. 31100 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|---|--------|
| Sodium chloride (NaCl) | 1.5212 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0211 |
| Sodium bicarbonate (NaHCO ₃) | 0.0463 | Silica (SiO ₂) | 0.1590 |
| Calcium chloride (CaCl ₂) | 0.4151 | | 2.5469 |
| Calcium sulphate (CaSO ₄) | 0.0302 | Free carbon dioxide (CO ₂) | 0.5932 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.3540 | | 2.4444 |

Besides these trace of alumina and boric acid.

The spring may be classified as “common salt spring”.

Radio-activity. 1.06 Mache's units in 1 litre of water at 15.6° C.

1.56 Mache's units in 1 litre of gas at 0° C.

(Kohl-Löw. fontact.; Dr. R. Ishizu and T. Saitō, March 5, 1914).

Table 96.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|--------------------|-----------------------------------|------------------------------|--------------------------|---|--|---------------------|-------------------|---------------------------------------|----------------------------------|-----------------------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
temp. in C. | | | |
| 1 | Koyabara-Onsen | non carbonate
common salt | air temp.
38.2° (87°) | ca.
540 | 13.65 | 3.75 | 18.7° | Schm. | March 4, 1914 | { Dr. R. Ishizu
T. Saitō |
| 2 | Nobata-no-yu * | carbonated(?) | 24.0 (75°) | — | — | 3.14 | 16.1° | K. & L. | .. | |
| 3 | Nobata-shin-yu * | .. | 17.0° (63°) | — | 682.67 | 187.74 | 16.3° | Schm. | .. | |
| 4 | Shigaku-Onsen | common salt | 46.5° (115°) | 31100 | — | 1.06 | 15.6° | K. & L. | .. 5. " | |
| 5 | Private bath (<i>S. Nagira</i>) | — | 46.5° (115°) | — | — | 0.29 | 16.3° | .. | .. | |
| 6 | Spring <i>A</i> | — | 22.5° (73°) | — | — | 1.86 | 16.4° | .. | .. | |
| Gas evolving from: | | | | | Emanation per litre
of gas at 0° C. | | | Date when the
gas was
collected | | |
| 1 | Koyabara-Onsen | — | — | — | 29.53 | 8.12 | — | Schm. | March 3, 1914 | .. |
| 2 | Nobata-shin-yu * | .. | .. | .. | 1307.21 | 359.48 | .. | .. | .. 4. " | .. |
| 3 | Nobata-no-yu * | — | — | — | — | 18.16 | — | K. & L. | .. | .. |
| 4 | Shigaku-Onsen | — | — | — | — | 1.56 | — | .. | .. 5. " | .. |

* These springs issue from the Granite.

* These springs issue from the Granite.

YUNOTSU

Location.—Yunotsu-machi, Nima-gun, Province Iwami, Shimane-ken.

The springs issue from the Tertiary.

The hot spring “Yunotsu-Onsen”

Analysis.

Analysed by the Hygienic Laboratory of Shimane-ken. 1911.

Specific gravity: 1.0056 at 15° C.

Total residue: ca. 7.06.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|---|--------|
| Potassium chloride (KCl) | 0.1528 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0174 |
| Sodium chloride (NaCl) | 4.7941 | Alumina (Al ₂ O ₃) | 0.0900 |
| Sodium bicarbonate (NaHCO ₃) | 0.4049 | Silica (SiO ₂) | 0.0925 |
| Calcium sulphate (CaSO ₄) | 1.0986 | | 7.0611 |
| Magnesium sulphate (MgSO ₄) | 0.3185 | Free carbon dioxide (CO ₂) | 0.8813 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0923 | | 7.9424 |

Besides these trace of phosphoric and boric acid.

The spring may be classified as “sulphated common salt spring”.

FUKUMITSU

Location.—Fukumitsu-mura, Nima-gun, Province Iwama, Shimane-ken.

The spring issues from the Tertiary.

The cold spring “Yunohara-Onsen”

Analysis.

Analysed by the Hygienic Laboratory of Shimane-ken. 1911.

Specific gravity: 1.0038 at 15° C.

Temperature: 34.5° C. (air temp. 16° C.)

Total residue: ca. 4.48.

Flow of water: ca. 77 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|--|--------|
| Potassium chloride (KCl) | 0.2395 | Ferrous bicarbonate [FeHCO_3] | 0.0174 |
| Sodium chloride (NaCl) | 2.6202 | Alumina (Al_2O_3) | 0.0375 |
| Sodium bicarbonate (NaHCO_3) | 0.6258 | Silica (SiO_2) | 0.1025 |
| Calcium sulphate (CaSO_4) | 0.4415 | | 4.5521 |
| Calcium bicarbonate [$\text{Ca}(\text{HCO}_3)_2$] | 0.0616 | Free carbon dioxide (CO_2) | 1.0617 |
| Magnesium bicarbonate [$\text{Mg}(\text{HCO}_3)_2$] | 0.4061 | | 5.6138 |

Besides these trace of phosphoric and boric acid.

The spring may be classified as “carbondioxated common salt spring”.

Radio-activity. 0.12 Mache's units in 1 litre of water at 21.6° C.

0.03 Mache's units in 1 litre of gas at 0° C.

(Kohl-Löw. contact.; Dr. R. Ishizu and T. Saitō, March 7, 1914).

Table 97.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------|-------------------------------|---------------------------|---|--|---------------------|---------------------|--------------------|---------------------------------------|-----------------------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kyū-yu | sulphated
common salt | 50.0 ^{air temp.} | 700 | — | 1.61 | 20.1° | K. & L. | March 6, 1914 | Dr. R. Ishizu
(T. Saitō) |
| 2 | Shin-yu | “ | 46.0° | 544 | 18.00 | 4.95 | 24.8° | Schm. | “ | “ |
| 3 | Yunohara-Onsen | carbondioxated
common salt | 34.5° (16°) | 77 | — | 0.12 | 21.6° | K. & L. | “ 7. “ | “ |
| | Gas evolving from: | | | | Emanation per litre
of gas at 0° C. | | | | Date when the
gas was
collected | |
| 1 | Shin-yu | — | — | — | 4.92 | 1.35 | — | Schm. | March 6, 1914 | “ |
| 2 | Kyū-yu | — | — | — | — | 3.31 | — | K. & L. | “ | “ |
| 3 | Yunohara-Onsen | — | — | — | — | 0.03 | — | “ | “ 7. “ | “ |

ARIFUKU

Location.—Arifuku-mura, Naka-gun, Province Iwami, Shimane-ken.

The springs issue from the Diorite.

The hot spring “ Arifuku-Onsen ”

Analysis.

Analysed by the Hygienic Laboratory of Shimane-ken. 1912.

Specific gravity: 1,0013 at 14° C.

Total residue: ca. 0,29.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Sodium chloride (NaCl) | 0,1401 | Calcium sulphate (CaSO ₄) | 0,0243 |
| Potassium sulphate (K ₂ SO ₄) | 0,0141 | Silica (SiO ₂) | 0,0560 |
| Magnesium chloride (MgCl ₂) | 0,0098 | | 0,2443 |

Besides these a small quantity of boric acid and organic substances.

The spring may be classified as “simple thermal”.

Table 98.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------|----------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-----------------------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Arifuku-Onsen Vent. No. 1 | simple | 48,0° | ca. 3230 | 0,80 | 0,22 | 18,6° | Schm. | March 7, 1914 | Dr. R. Ishizu
(T. Saitō) |
| 2 | „ Vent. No. 2 | „ | 49,0° | — | — | 0,64 | 19,2° | K. & L. | „ | „ |
| 3 | Arifuku-shin-yu | „ | 44,5° | — | — | 1,51 | 18,9° | „ | „ | „ |

TOMBARA

Location.—Tombara-mura, Iishi-gun, Province Izumo, Shimane-ken.

The springs issue from the Granite.

The cold spring “ Kamiyasumi-Kōsen ”

Temperature: 15,7° C.

Flow of water: ca. 108 hectolitres in 24 hours.

No accurate analysis has ever been made, but the spring seems to be “carbonated spring”.

Radio-activity. 9,69 Mache's units in 1 litre of water* at 15,2° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu and T. Saitō, March 3, 1914).

KASUBUCHI

Location.—Kasubuchi-mura, Oochi-gun, Province Iwami, Shimane-ken.

The spring issues from the Liparite.

The cold spring “ Kasubuchi-no-yu ”

No accurate analysis has ever been made, but the spring seems to be “carbonated spring”.

Radio-activity. 15,77 Mache's units in 1 litre of water* at 16,0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu and T. Saitō, March 4, 1914).

* Measurement was made with the sample brought to the examiners by applicant, taking into account decay of radium emanation with the time.

YUNOGŌ

Location.—Yunogō-mura, Katsuda-gun, Province Mimasaka, Okayama-ken.

The springs issue from the Liparite.

The hot spring “Sagi-no-yu”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1914.

Specific gravity: 1.0022 at 13° C.

Total residue: ca. 2.27.

Temperature: 38° C. (air temp. 3° C.).

Flow of water: ca. 1296 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|---|---------|------------|-----------------------|
| Potassium ion (K'). | 0.02091 | 0.53410 | 0.53410 |
| Sodium ion (Na'). | 0.46393 | 20.12711 | 20.12711 |
| Calcium ion (Ca'). | 0.35208 | 8.80200 | 17.60400 |
| Magnesium ion (Mg''). | 0.00044 | 0.01800 | 0.03600 |
| Ferrous ion (Fe''). | 0.00084 | 0.01500 | 0.03000 |
| Aluminium ion (Al'''). | 0.00042 | 0.01550 | 0.04650 |
| | | | 38.37783 |
| Anions. | | | |
| Chlorine ion (Cl'). | 1.34209 | 37.85867 | 37.85867 |
| Sulphate ion (SO ₄ ''). | 0.01560 | 0.16240 | 0.32480 |
| Hydrocarbonate ion (HCO ₃ '). | 0.01184 | 0.19407 | 0.19407 |
| | 2.20815 | 67.72691 | 38.37754 |
| Boric acid (meta) (HBO ₃). | 0.01419 | | |
| Silicic acid (meta) (H ₂ SiO ₃). | 0.04598 | | |
| | 2.26832 | | |

| | Grams |
|--|---------|
| Potassium chloride (KCl). | 0.03984 |
| Sodium chloride (NaCl). | 1.17748 |
| Calcium chloride (CaCl ₂). | 0.95356 |
| Calcium sulphate (CaSO ₄). | 0.01894 |
| Calcium bicarbonate [Ca(HCO ₃) ₂]. | 0.01037 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂]. | 0.00264 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂]. | 0.00267 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃]. | 0.00265 |
| Boric acid (meta) (HBO ₃). | 0.01419 |
| Silicic acid (meta) (H ₂ SiO ₃). | 0.04598 |
| | 2.26832 |

Besides these a small quantity of free carbon dioxide.

The spring thus may be classified as “**earth-muriated common salt spring**”.

Radio-activity. 3.06×10^{-10} curies in 1 litre of water at 11.5° C.

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, March 17, 1914).

0.84 Mache's units (recalculated).

Gas evolving from the spring (collected at the vent, from which gas evolution is most abundant) consists almost of Nitrogen,

19.50×10^{-10} curies in 1 litre of gas at 0° C.

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, March 17, 1914).

5.36 Mache's units (recalculated).

The hot spring “Shin-sagi-no-yu”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1912.

Specific gravity: 1.0018 at 17° C.

Flow of water: ca. 540 hectolitres in 24 hours.

Temperature: 37.7° C. (air temp. 27.2° C.).

Total residue: ca. 2.25.

The spring, being in its composition nearly the same as that of “Sagi-no-yu”, may be classified as “**earth-muriated common salt spring**”.

Radio-activity. 0.85 Mache's units in 1 litre of water at 27.7° C.

4.56 Mache's units in 1 litre of gas at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, July 15, 1913).

Table 99.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-------------------------------------|-----------------------------------|----------------------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-----------------------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Sagi-no-yu (<i>Public bath</i>) | earth-mixed
common salt | air temp
38.0° (30°) | ca.
1296 | 3.06 | 0.84 | 11.5° | Schm. | March 17, 1914 | { Dr. R. Ishizu
T. Saitō |
| 2 | " | " | " | " | — | 1.00 | 10.0° | K. & L. | " | |
| 3 | " (<i>Onna-yu</i>) | " | 37.0° (30°) | — | — | 0.90 | 7.4° | " | " | " |
| 4 | Shin-sagi-no-yu | " | 37.7° (27.2°) | 540 | — | 0.85 | 27.7° | " | July 15, 1913 | " |
| 5 | Me-no-yu | sulphur (?) | 25.0° (10°) | — | — | 1.00 | 13.4° | " | March 17, 1914 | " |
| Emanation per litre of gas at 0° C. | | | | | | | | | | |
| 1 | Gas evolving from:
Sagi no-yu | — | — | — | 19.50 | 5.36 | — | Schm. | March 17, 1914 | " |
| 2 | " (<i>Onna-yu</i>) | — | — | — | — | 4.11 | — | K. & L. | " | " |
| 3 | Shin-sagi-no-yu | — | — | — | — | 4.56 | — | " | July 15, 1913 | " |
| 4 | Me-no-yu | — | — | — | — | 4.74 | — | " | March 17, 1914 | " |

KAIDANI

Location.—Nodani-mura, Mitsu-gun, Province Bitchū, Okayama-ken.

The spring issues from the Granite.

The cold spring "Tomada-no-yu"

Temperature: 14.5° C. (air temp. 9.2° C.)

No accurate analysis has ever been made, but the spring seems to be "simple cold spring".

Radio-activity. 48.66 × 10⁻¹⁰ curies in 1 litre of water at 11.5° C.

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, Feb. 13, 1914).

13.38 Mache's units (recalculated).

TAKEBE

Location.—Kami-takebe-mura, Mitsu-gun, Province Bitchū, Okayama-ken.

The spring issues from the Granite.

The cold spring "Yawata-Onsen"

Analysis.

Analysed by the Hygienic Laboratory of Okayama-ken. 1900.

Specific gravity: 1.0011 at 12° C.

Temperature: 28° C. (air temp. 16° C.).

Total residue: ca. 0.26.

Flow of water: ca. 390 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained .

| | Grams | | Grams |
|---|---------|---|---------|
| Potassium chloride (KCl) | 0,12410 | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,00708 |
| Sodium chloride (NaCl) | 0,13842 | Silica (SiO ₂) | 0,02692 |
| Sodium sulphate (Na ₂ SO ₄) | 0,01677 | | 0,39845 |
| Sodium bicarbonate (NaHCO ₃) | 0,04250 | Free carbon dioxide (CO ₂) | 0,00900 |
| Calcium sulphate (CaSO ₄) | 0,02499 | | 0,40745 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,01767 | | |

Besides these trace of iron, alumina, phosphoric and boric acid.

The spring may be classified as "simple cold spring".

Radio-activity. $17,66 \times 10^{-10}$ curies in 1 litre of water at 13,0° C.

(Schmidt's electrom.; Dr. R. Ishizu and T. Saitō, Feb. 14, 1914)

4,86 Mache's units (recalculated).

YUDA

Location.—Shimo-unoryō-mura, Yoshiki-gun, Province Suō, Yamaguchi-ken.

The springs issue from the Alluvium.

Table 100.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------|----------------|--------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-----------------------------|
| | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Nagamune-no-yu | salt | 45,0° | 9,72 | 2,67 | 45,0° | Schm. | Aug. 4, 1914 | H. Kibezaki
S. Nishimura |
| 2 | Matsudaya-no-yu | " | 41,0° | 9,60 | 2,64 | 41,0° | " | " 5, " | " |

KAWATANA

Location.—Kawatana-mura, Toyora-gun, Province Nagato, Yamaguchi-ken.

The springs issue from the Granite.

Table 101.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------|----------------|--------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-----------------------------|
| | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Public bath :
Juei-sen | salt | 41,0° | 36,19 | 9,95 | 41,0° | Schm. | Aug. 9, 1914 | H. Kibezaki
S. Nishimura |
| 2 | Seiryū-sen | " | 40,0° | 43,21 | 11,88 | 40,0° | " | " | " |

TAWARAYAMA

Location.—Tawarayama-mura, Ōtsu-gun, Province Nagato, Yamaguchi-ken.

The springs issue from the Mesozoic Conglomerate.

Table 102.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------------|----------------|--------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------------------------|
| | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Public bath :
Kagi-yu | alkaline | 40,0° | 3,59 | 0,99 | 40,0° | Schm. | Aug. 7, 1914 | { H. Kibezaki
S. Nishimura |
| 2 | Kawa-yu | „ | 42,0° | 6,21 | 1,71 | 42,0° | „ | „ 8 „ | „ |

FUKAWA

Location.—Fukawa-mura, Ōtsu-gun, Province Nagato, Yamaguchi-ken.

The springs issue from the Liparite.

Table 103.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--|----------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------------------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Public bath :
Yumoto-Onsen (<i>On-tō</i>) | alkaline | 40,0° | ca. | 0,45 | 0,12 | 40,0° | Schm. | Aug. 6, 1914 | { H. Kibezaki
S. Nishimura |
| 2 | „ (<i>Kai-tō</i>) | „ | 42,0° | 2520 | 0,40 | 0,11 | 42,0° | „ | „ 7 „ | „ |

DŌGO

Location.—Dōgo-yuno-machi, Onsen-gun, Province Iyo, Ehime-ken.

The springs issue from the Granite.

The hot spring “Kami-no-yu”

Analysis.

Analysed by the Hygienic Laboratory of Hiroshima-ken. 1896.

Specific gravity: 1,00106 at 15° C.

Temperature: 47° C.

Total residue: ca. 0,78.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|--|---------|---|---------|
| Sodium chloride (NaCl) | 0,15597 | Calcium sulphate (CaSO ₄) | 0,11855 |
| Potassium sulphate (K ₂ SO ₄) | 0,03775 | Magnesium sulphate (MgSO ₄) | 0,08758 |
| Sodium sulphate (Na ₂ SO ₄) | 0,19669 | | 0,78828 |
| Sodium bicarbonate (NaHCO ₃) | 0,03964 | Carbon dioxide (free and in form of bicarbonates) + CO ₂ | 0,05887 |
| Sodium phosphate (Na ₂ HPO ₄) | 0,01597 | | 0,04715 |
| Sodium silicate (Na ₄ SiO ₄) | 0,13613 | | |

Besides these a very small quantity of ferrous oxide, boric acid and trace of nitric acid.

The spring may be classified as "simple thermal".

Radio-activity. $14,47 \times 10^{-10}$ curies in 1 litre of water at 47,0° C.

(Schmidt's electrom.; H. Kibezaki, July 10, 1913).

3,98 Mache's units (recalculated).

$85,02 \times 10^{-10}$ curies in 1 litre of gas.

(Schmidt's electrom.; H. Kibezaki, July 15, 1913).

23,38 Mache's units (recalculated).

The hot spring "Yōjyō-yu"

Analysis.

Analysed by the Hygienic Laboratory of Hiroshima-ken. 1896.

Temperature: 46° C.

Specific gravity: 1,01418 at 15° C.

Total residue: ca. 0,83.

The spring, being in its composition nearly the same as that of "Kami-no-yu", may be classified as "simple thermal".

Radio-activity. $16,07 \times 10^{-10}$ curies in 1 litre of water at 46,0° C.

(Schmidt's electrom.; H. Kibezaki, July 16, 1913).

4,42 Mache's units (recalculated).

Table 104.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|---------------------------------|----------------|----------------------------|---------------------------------|---------------------|---------------------|--------------------|---------------------------------------|-------------|
| | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Kami-no-yu (Public bath) | simple | 46,5° ^{air temp.} | 16,33 | 4,49 | 46,5° | Schm. | March 24, 1913 | H. Kibezaki |
| 2 | " | " | 47,0° | 14,47 | 3,98 | 47,0° | " | July 10, " | " |
| 3 | Yōjyō-yu (") | " | 46,5° | 13,70 | 3,77 | 46,5° | " | March 25, " | " |
| 4 | " | " | 46,0° | 16,07 | 4,42 | 46,0° | " | July 16, " | " |
| 5 | Shin-Onsen (I. Iwasaki) | — | 29,0° | 23,05 | 6,34 | 29,0° | " | March 26, " | " |
| 6 | " | — | 29,0° | 24,04 | 6,61 | 29,0° | " | July 13, " | " |
| 7 | Tama-no-i | — | — | 3,90 | 1,07 | — | " | March 23, " | " |
| 8 | Well (Funaya) | — | 23,0° | 8,85 | 2,43 | 23,0° | " | " | " |
| 9 | Tama-no-yu (Public bath) | simple | 45,8° (19,5°) | — | — | — | — | — | — |
| 10 | Matsu-no-yu (") | " | 42,0° (19,5°) | — | — | — | — | — | — |
| | Gas evolving from
Kami-no-yu | — | — | Emanation per
litre of gas | | | Schm. | Date when the
gas was
collected | |
| | | | | 85,02 | 23,38 | — | | July 15, 1913 | H. Kibezaki |

MUSASHI

Location.—Futsukaichi-machi, Tsukushi-gun, Province Chikuzen, Fukuoka-ken.

The springs issue from the Granite.

The hot spring “Yakushi-no-yu”

Analysis.

Analysed by the Hygienic Laboratory of Fukuoka-ken. 1897. Specific gravity: 1.002 at 12° C.
Flow of water: ca. 377 hectolitres in 24 hours. Total residue: ca. 0.72.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|---------|--|---------|
| Potassium chloride (KCl) | 0.02160 | Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃) . | 0.00785 |
| Sodium chloride (NaCl) | 0.55285 | Silica (SiO ₂) | 0.03366 |
| Potassium sulphate (K ₂ SO ₄) | 0.00395 | | 0.72386 |
| Sodium carbonate (Na ₂ CO ₃) | 0.08244 | Free hydrogen sulphide (H ₂ S) | 0.00226 |
| Calcium sulphate (CaSO ₄) | 0.02151 | | 0.72612 |

Besides these a small quantity of boric acid and trace of magnesia and phosphoric acid.

The spring may be classified as “sulphur spring”.

Table 105.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------------------------|----------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|--|-------------------------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Yakushi-no-yu No. 1
(Public bath) | sulphur | 41.0° | 377 | 7.29 | 2.00 | 41.0° | Schm. | July 30, 1914 | { H. Kibezaki
S. Nishimura |
| 2 | “ No. 2 (“) | “ | 44.0° | | 14.78 | 4.07 | 44.0° | “ | “ 29. “ | |
| 3 | Enju-no-yu (“) | “ | 46.0° | | 13.46 | 3.70 | 46.0° | “ | “ 31. “ | |
| 4 | Osakaya-no-yu | “ | 44.5° | 207 | 11.67 | 3.21 | 44.5° | “ | “ 30. “ | “ |
| 5 | Gozen-yu (Public bath) | “ | 42.2° | 213 | — | — | — | — | — | — |
| 6 | Kawa-yu (“) | “ | 46.7° | 890 | — | — | — | — | — | — |
| | Gas evolving from
Osakaya-no-yu | — | — | — | Emanation per litre
of gas | | | Schm. | Date when the
gas was
collected
July 31, 1914 | { H. Kibezaki
S. Nishimura |
| | | | | | 118.33 | 32.54 | — | | | |

FUNAGOYA

Location.—Mizuta-mura, Yame-gun, Province Chikugo, Fukuoka-ken.

The springs issue from the Diluvium.

The cold spring “Funagoya-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1900. Temperature: 21.0° C.
Flow of water: ca. 63 hectolitres in 24 hours. Total residue: ca. 0.17.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Sodium ion (Na ⁺) | 0,0185 | 0,8026 | 0,8026 |
| Calcium ion (Ca ⁺⁺) | 0,0168 | 0,4189 | 0,8378 |
| Magnesium ion (Mg ⁺⁺) | 0,0058 | 0,2385 | 0,4770 |
| Ferrous ion (Fe ⁺⁺) | 0,0021 | 0,0376 | 0,0752 |
| | | | 2,1926 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0,0285 | 0,8142 | 0,8142 |
| Sulphate ion (SO ₄ ^{''}) | 0,0183 | 0,1905 | 0,3810 |
| Hydrocarbonate ion (HCO ₃ [']) | 0,0609 | 0,9982 | 0,9982 |
| | 0,1509 | 3,5005 | 2,1934 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0430 | | |
| | 0,1939 | | |
| Free carbon dioxide (CO ₂) | 1,9391 | 44,0705 | |
| | 2,1330 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 0,0470 |
| Calcium sulphate (CaSO ₄) | 0,0260 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,0364 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,0249 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0066 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0430 |
| | 0,1939 |
| Free carbon dioxide (CO ₂) | 0,8924 |
| | 1,0863 |

Besides these trace of hydrophosphate ion.

The spring thus may be classified as "simple carbondioxated spring".

Table 106.

| No. | Spring | Classification | Temp. of spring in C. | Flow of water in 24 hrs. in hectolitres | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|---------------------------------|------------------------|-----------------------|---|------------------------------|------------------|-----------------|--------------------|-------------------------------|-----------------------------|
| | | | | | in 10 ⁻¹⁰ curies | in Mache's units | at temp. in (°) | | | |
| 1 | Funagoya-Onsen
(Public bath) | simple carbon-dioxated | 17,5° | ca. 63 | 3,43 | 0,94 | 17,5° | Schm. | Aug. 2, 1914 | H. Kibezaki
S. Nishimura |
| 2 | Nomi-yu | " | 21,0° | " | 1,45 | 0,40 | 21,0° | " | " | " |

BEPPU

Location.—Beppu-machi, Hayami-gun, Province Bungo, Oita-ken.

The springs issue from the Andesite overlaid by Alluvium.

The hot spring "Furō-sen"

Analysis (calculated from the salt table).

Analysed by the Hygienic Laboratory of Fukuoka-ken.

Temperature: 57,5° C.

Flow of water: ca. 648 hectolitres in 24 hours.

Total residue: ca. 0,85.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|---|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0,0023 | 0,0587 | 0,0587 |
| Sodium ion (Na ⁺) | 0,1059 | 4,5944 | 4,5944 |
| Calcium ion (Ca ⁺⁺) | 0,0168 | 0,4190 | 0,8380 |
| Magnesium ion (Mg ⁺⁺) | 0,0354 | 1,4532 | 2,9064 |
| Ferrous ion (Fe ⁺⁺) | 0,0120 | 0,2147 | 0,4294 |
| | | | 8,8269 |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 0,2263 |
| Potassium sulphate (K ₂ SO ₄) | 0,0051 |
| Sodium sulphate (Na ₂ SO ₄) | 0,0516 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0,0680 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0,2210 |
| Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0,0383 |

Anions.

| | | | |
|--|----------------|---------|--------|
| Chlorine ion (Cl') | 0,1371 | 3,8646 | 3,8646 |
| Sulphate ion (SO ₄ '') | 0,0377 | 0,3925 | 0,7850 |
| Hydrogencarbonate ion (HCO ₃ ') | 0,2631 | 4,3124 | 4,3124 |
| | 0,6103 | 15,3095 | 8,9620 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,2548 | | |
| | 0,8651 | | |
| Free carbon dioxide (CO ₂) | not determined | | |

Silicic acid (meta) (H₂SiO₃) $\frac{0,2548}{0,8651}$

Besides these trace of aluminium and hydrophosphate ion.

The spring thus may be classified as “iron carbonate spring”.

Radio-activity. $0,66 \times 10^{-10}$ curies in 1 litre of water at 57,5° C.

(Schmidt's electrom.; H. Kibezaki, March 7, 1913)

0,18 Mache's units (recalculated).

KWANKAIJI

Location.—Ishikaki-mura, Hayami-gun, Province Bungo, Ōita-ken.

The spring issues from the Pyroxene Andesite and its Agglomerate.

The hot spring “Kwankaiji-Onsen”

Analysis.

Analysed by the Hygienic Laboratory of Ōita-ken. 1906.

Specific gravity: 1,0004 at 15° C.

Temperature: 57° C.

Total residue: ca. 0,77.

Flow of water: ca. 216 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|--|--------|---|--------|
| Potassium oxide (K ₂ O). | 0,0062 | Chlorine (Cl) | 0,0732 |
| Sodium oxide (Na ₂ O) | 0,0536 | Sulphuric anhydride (SO ₃). | 0,0346 |
| Calcium oxide (CaO). | 0,0818 | Silica (SiO ₂) | 0,0936 |
| Magnesia (MgO) | 0,0357 | Carbon dioxide (total) (CO ₂) | 0,5933 |
| Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃) | 0,0590 | | |

Besides these trace of phosphoric and boric acid.

The spring may be classified as “iron carbonate spring”.

Radio-activity. $0,07 \times 10^{-10}$ curies in 1 litre of water at 57,0° C.

(Schmidt's electrom.; H. Kibezaki, March 8, 1913).

0,02 Mache's units (recalculated).

KANNAWA

Location.—Asahi-mura, Hayami-gun, Province Bungo, Ōita-ken.

The springs issue from the Pyroxene Andesite and its Agglomerate.

The hot spring “Shibu-no-yu”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Specific gravity: 1,00109 at 4° C.

Temperature: 89° C.

Flow of water: ca. 414 hectolitres in 24 hours.

Total residue: ca. 2,67.

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Hydrogen ion (H ⁺) | 0.0221 | 21,8812 | 21,8812 |
| Potassium ion (K ⁺) | 0.1128 | 2,8812 | 2,8812 |
| Sodium ion (Na ⁺) | 0.2957 | 12,8286 | 12,8286 |
| Calcium ion (Ca ⁺⁺) | 0.0657 | 1,6384 | 3,2768 |
| Magnesium ion (Mg ⁺⁺) | 0.0162 | 0,6650 | 1,3300 |
| Ferrous ion (Fe ⁺⁺) | 0.0157 | 0,2809 | 0,8427 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0310 | 1,1439 | 3,4317 |
| | | | 47,4722 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 1,1402 | 32,1636 | 32,1636 |
| Sulphate ion (SO ₄ ^{''}) | 0.6801 | 7,0800 | 14,1600 |
| | 2,3795 | 80,5628 | 46,3236 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.3751 | | |
| | 2,7540 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Grams |
|--|--------|
| Sodium chloride (NaCl) | 0.0923 |
| Potassium sulphate (K ₂ SO ₄) | 0.2514 |
| Sodium sulphate (Na ₂ SO ₄) | 0.8009 |
| Calcium chloride (CaCl ₂) | 0.1820 |
| Magnesium chloride (MgCl ₂) | 0.0635 |
| Ferric chloride (FeCl ₃) | 0.0456 |
| Aluminium chloride (AlCl ₃) | 0.1528 |
| Free hydrochloric acid (HCl) | 0.7910 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.3751 |
| | 2,7540 |

Besides these a small quantity of organic substances and a very small quantity of boric acid.

The spring thus may be classified as "acid vitriol spring".

Composition of the mineral salt "Yu-no-hana"

(analysed by the Hygienic Laboratory of Oita-ken in 1900).

100 parts of the yellowish crystalline mass, almost totally soluble in water with an acid reaction and in an astringent taste, consist of :

| | Parts | | Parts |
|--|-------|---|-------|
| Potassium sulphate (K ₂ SO ₄) | 0,11 | Aluminium sulphate [Al ₂ (SO ₄) ₃] | 35,51 |
| Sodium sulphate (Na ₂ SO ₄) | 0,93 | Calcium phosphate [Ca ₂ (HPO ₄) ₂] | 0,15 |
| Calcium sulphate (CaSO ₄) | 0,29 | Boric acid (H ₃ BO ₃) | trace |
| Magnesium sulphate (MgSO ₄) | 2,20 | Insoluble matters | 1,77 |
| Ferric sulphate [Fe ₂ (SO ₄) ₃] | 16,77 | Water | 42,23 |

The salt is used for bathing purpose.

The hot spring "Umi-Jigoku"

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1912. Total residue: ca. 3.51.

Specific gravity: 1.0029 at 15° C.

Temperature: 90,5° C. (air temp. 26° C.).

In 1 kilogram of the mineral water are contained :

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------|------------|-----------------------|
| Hydrogen ion (H ⁺) | 0.00613 | 6,06930 | 6,06930 |
| Potassium ion (K ⁺) | 0.17560 | 4,48531 | 4,48531 |
| Sodium ion (Na ⁺) | 0.82521 | 35,80086 | 35,80086 |
| Ammonium ion (NH ₄ ⁺) | 0.00154 | 0,08518 | 0,08518 |
| Calcium ion (Ca ⁺⁺) | 0.08456 | 2,11400 | 4,22800 |
| Magnesium ion (Mg ⁺⁺) | 0.03183 | 1,30660 | 2,61320 |
| Ferrous ion (Fe ⁺⁺) | 0.01293 | 0,23100 | 0,46200 |
| Aluminium ion (Al ⁺⁺⁺) | 0.00788 | 0,29077 | 0,87210 |
| | | | 54,61595 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 1,26300 | 35,62764 | 35,62764 |
| Sulphate ion (SO ₄ ^{''}) | 0.91216 | 9,49573 | 18,99146 |
| | 3,32084 | 95,50630 | 54,61910 |
| Boric acid (meta) (HBO ₂) | 0.01414 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.48049 | | |
| | 3,81547 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram :

| | Gram |
|---|---------|
| Sodium chloride (NaCl) | 1,32380 |
| Ammonium chloride (NH ₄ Cl) | 0,00455 |
| Potassium sulphate (K ₂ SO ₄) | 0,39102 |
| Sodium sulphate (Na ₂ SO ₄) | 0,93622 |
| Calcium chloride (CaCl ₂) | 0,23444 |
| Magnesium chloride (MgCl ₂) | 0,12447 |
| Ferrous sulphate (FeSO ₄) | 0,03513 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0,04977 |
| Free hydrochloric acid (HCl) | 0,22140 |
| Boric acid (meta) (HBO ₂) | 0,01414 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,48049 |
| | 3,81547 |

The spring thus may be classified as "acid vitriol spring".

MYŌBAN

Location.—Asahi-mura, Hayami-gun, Province Bungo, Ōita-ken.

The springs issue from the Andesite.

The hot spring “Jizō-no-yu”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1898.

Specific gravity: 1.003 at 20° C.

Temperature: 91° C. (air temp. 22° C.).

Total residue: ca. 2.52.

Flow of water: ca. 389 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|---------------|------------|-----------------------|
| Hydrogen ion (H ⁺) | 0.0205 | 20.2970 | 20,2970 |
| Potassium ion (K ⁺) | 0.0373 | 0.9528 | 0.9528 |
| Sodium ion (Na ⁺) | 0.0480 | 2.0824 | 2.0824 |
| Calcium ion (Ca ⁺⁺) | 0.0430 | 1.0723 | 2.1446 |
| Ferrous ion (Fe ⁺⁺) | 0.0585 | 1.0465 | 2.0930 |
| Ferric ion (Fe ⁺⁺⁺) | 0.0880 | 1.5742 | 4.7226 |
| Aluminium ion (Al ⁺⁺⁺) | 0.1366 | 5.0406 | 15,1218 |
| | | | 47.4142 |

| | Grams |
|---|--------|
| Potassium sulphate (K ₂ SO ₄) | 0.0830 |
| Sodium sulphate (Na ₂ SO ₄) | 0.1480 |
| Calcium sulphate (CaSO ₄) | 0.1460 |
| Ferrous sulphate (FeSO ₄) | 0.1590 |
| Ferric sulphate [Fe ₂ (SO ₄) ₃] | 0.3150 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.8630 |
| Free sulphuric acid (H ₂ SO ₄) | 1.9870 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.4608 |
| | 4.1618 |

Anions.

| | | | |
|--|--------|---------|----------------|
| Hydrosulphate ion (HSO ₄ [']) | 1.9665 | 20.2586 | 20.2586 |
| Sulphate ion (SO ₄ ^{''}) | 1.3026 | 13.5603 | 27,1206 |
| | 3.7010 | 65.8847 | 47.3792 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.4608 | | |
| | 4.1618 | | |

Besides these trace of magnesium, nitrate and chlorine ion and boric acid.

The spring thus may be classified as “acid alum vitriol spring”.

Radio-activity. 0.13 Mache's units in 1 litre of water at 25.0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Oct. 11, 1913).

KAMEGAWA

Location.—Okoshi-machi, Hayami-gun, Province Bungo, Ōita-ken.

The springs issue from the Pyroxene Andesite and its Agglomerate.

The hot spring “Shi-no-yu”

Analysis (calculated from the salt table).

Analysed by the Ōsaka Imperial Hygienic Laboratory. 1895.

Temperature: 57.0° C. (air temp. 24° C.).

Flow of water: ca. 198 hectolitres in 24 hours.

Total residue: ca. 1.22.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|---|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0377 | 0.9630 | 0.9630 |
| Sodium ion (Na ⁺) | 0.2939 | 12.7505 | 12,7505 |
| Calcium ion (Ca ⁺⁺) | 0.0363 | 0.9052 | 1.8104 |
| Magnesium ion (Mg ⁺⁺) | 0.0085 | 0.3489 | 0.6978 |
| Ferrous ion (Fe ⁺⁺) | 0.0022 | 0.0394 | 0.0788 |
| | | | 16.3005 |

| | Grams |
|---|--------|
| Sodium chloride (NaCl) | 0.6780 |
| Potassium sulphate (K ₂ SO ₄) | 0.0840 |
| Sodium sulphate (Na ₂ SO ₄) | 0.0040 |
| Sodium bicarbonate (NaHCO ₃) | 0.0930 |
| Calcium sulphate (CaSO ₄) | 0.1230 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0510 |

nions.

| | | | |
|--|--------|---------|---------|
| Chlorine ion (Cl') | 0,4109 | 11,5910 | 11,5910 |
| Sulphate ion (SO_4'') | 0,1357 | 1,4127 | 2,8254 |
| Hydrocarbonate ion (HCO_3') | 0,1148 | 1,8817 | 1,8817 |
| | 1,0400 | 20,8024 | 10,2081 |
| Silicic acid (meta) (H_2SiO_3) | 0,2557 | | |
| | 1,2957 | | |
| Free carbon dioxide (CO_2) | 0,0310 | 0,7045 | |
| | 1,3207 | | |

| | |
|---|--------|
| Ferrous bicarbonate [$\text{Fe}(\text{HCO}_3)_2$] | 0,0070 |
| Silicic acid (meta) (H_2SiO_3) | 0,2557 |
| | 1,2957 |
| Free carbon dioxide (CO_2) | 0,0310 |
| | 1,3207 |

Besides these trace of aluminium ion.

The spring thus may be classified as “common salt spring”.

Radio-activity. 0,22 Mache's units in 1 litre of water at 24,0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Oct. 14, 1913).

The hot spring “Kiyō-sen”

Analysis.

Analysed by the Hygienic Laboratory of Ōita-ken. 1913.

Specific gravity: 1,0027 at 15° C. Temperature: 54° C. (air temp. 24,5° C.).

Total residue: ca. 2,65. Flow of water: ca. 270 hectolitres in 24 hours.

The spring, being in its composition nearly the same as that of “Shi-no-yu”, may be classified as “common salt spring”.

Radio-activity. 0,32 Mache's units in 1 litre of water at 29,0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Oct. 14, 1913).

The hot spring “Chinoike-Jigoku”

Temperature: 76° C. (air temp. 26° C.).

Flow of water: abundant.*

No accurate analysis has ever been made, but the spring seems to be “acid vitriol spring”.

Composition of the gas (analysed by the Tōkyō Imperial Hygienic Laboratory in 1913).

1000 c.c. of gas evolving from the spring consist of:

| | | | | | |
|--|-----|------|---|-------|------|
| Carbon dioxide (CO_2) | 450 | c.c. | Nitrogen (N_2) | 477,5 | c.c. |
| Oxygen (O_2) | 70 | „ | Ethane (C_2H_6) | 2,5 | „ |

The analysis is not so accurate, the number being only the approximate value.

Radio-activity. 4,82 Mache's units in 1 litre of gas, evolving from Vent No. 1, at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Oct. 10, 1913).

4,99 Mache's units in 1 litre of gas, evolving from Vent No. 2, at 0° C.

(Kohl.-Löw. fontact.; Dr. R. Ishizu, Oct. 13, 1913).

Composition of the sinter-deposit † (analysed by the Tōkyō Imperial Hygienic Laboratory in 1913).

100 parts of the ochreous mass, almost insoluble in water consist of:

| | Parts | | Parts |
|--|-------|--|-------|
| Ferric oxide (Fe_2O_3) | 42,38 | Calcium oxide (CaO) | 0,40 |
| Alumina (Al_2O_3) | 22,20 | Sulphuric acid (H_2SO_4) | trace |
| Silica (SiO_2) | 16,84 | Chlorine (Cl) | „ |
| Magnesia (MgO) | 0,58 | Water | 17,60 |

The ochreous clay is used for dyeing in the place near “Chi-no-ike”.

* The outflow of the boiling pool forms a steaming brook rushing down to the sea.

† The ochreous sinter-deposit is found on the shore of the pool.

SHIBASEKI

Location.—Okoshi-machi, Hayami-gun, Province Bungo, Ōita-ken.

The springs issue from the Pyroxine Andesite and its Agglomerate.

The hot spring “Hijiri-yu”

Analysis.

Analysed by the Hygienic Laboratory of Ōita-ken. Temperature: 69° C. (air temp. 23° C.).
Flow of water: ca. 216 hectolitres in 24 hours. Total residue: ca. 1.59.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|---------|---|---------|
| Sodium chloride (NaCl) | 0.36317 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.22072 |
| Potassium sulphate (K ₂ SO ₄) | 0.00600 | Alumina (Al ₂ O ₃) | 0.00142 |
| Sodium sulphate (Na ₂ SO ₄) | 0.43012 | Silica (SiO ₂) | 0.21802 |
| Sodium bicarbonate (NaHCO ₃) | 0.23108 | | 1.74945 |
| Calcium chloride (CaCl ₂) | 0.20002 | Free carbon dioxide (CO ₂) | 0.01531 |
| Magnesium chloride (MgCl ₂) | 0.06806 | | 1.76476 |
| Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.01084 | | |

The spring may be classified as “iron carbonate spring”.

Radio-activity. 0.31 Mache's units in 1 litre of water at 20.0° C.

1.91 Mache's units in 1 litre of gas at 0° C.

(Kohl-Löw. fontact.; Dr. R. Ishizu, Oct. 13, 1913).

Table 107.

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|--------------------------|------------------------------|--------------------------|---|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------|
| | | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Furō-sen | iron carbonate | 57.5°
air temp. | ca. 648 | 0.66 | 0.18 | 57.5° | Schm. | March 7, 1913 | H. Kibezaki |
| 2 | Kojimaya-no-yu No. 1 | carbonated
common salt | 55.0° | — | 0.75 | 0.21 | 55.0° | „ | „ 5, „ | „ |
| 3 | „ No. 2 | „ | 54.0° | — | 0.52 | 0.14 | 54.0° | „ | „ 9, „ | „ |
| 4 | Shiokyū-no-yu | „ | 57.0° | — | 0.39 | 0.11 | 57.0° | „ | „ 8, „ | „ |
| 5 | Seisen-kaku-no-yu | carbonated | 54.0° | — | 0.20 | 0.05 | 54.0° | „ | „ 9, „ | „ |
| 6 | Reichō-sen (Public bath) | simple carbon-
dioxated | 67.0° | 520 | — | — | — | — | — | — |
| 7 | Higashi-Onsen („) | common salt | 60.0° | 1167 | — | — | — | — | — | — |
| 8 | Nishi-Onsen („) | iron carbonate
carbonated | 57.0° | 783 | — | — | — | — | — | — |
| 9 | Kusunoki-Onsen („) | iron carbonate | 51.0° | 1321 | — | — | — | — | — | — |
| 10 | Takegawara-Onsen („) | carbonated | 56.0° | — | — | — | — | — | — | — |
| 11 | Azenashi-Onsen („) | simple | 51.0° | 162 | — | — | — | — | — | — |
| 12 | Asami-Onsen („) | „ (?) | 41.0° | — | — | — | — | — | — | — |
| 13 | Kwankaiji-Onsen („) | iron carbonate | 57.0° | 216 | 0.07 | 0.02 | 57.0° | Schm. | March 8, 1913 | H. Kibezaki |
| 14 | Hotta-no-yu „ | sulphur | 36.0° | 225 | — | — | — | — | — | — |

| No. | Spring | Classification | Temp. of spring
in C. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|---------------------|--|----------------------|-----------------------------------|---|--|--------------------|-------------------|---------------------------------------|----------------------------------|---------------|
| | | | | | in
10 ¹⁰ curies | in
Mach's units | at
temp. in C. | | | |
| 15 | Netsu-no-yu (<i>Public bath</i>) | carbonated | 80,0°
<small>air temp.</small> | 865 | — | — | — | — | — | — |
| 16 | Shin-yu („) | sulphur | 96,0° | 414 | — | — | — | — | — | — |
| 17 | Shibu-yu („) | acid vitriol | 89,0° | 414 | — | — | — | — | — | — |
| 18 | Mushi-buro (<i>Vapour bath</i>) | — | 69,0° | — | — | — | — | — | — | — |
| 19 | Andō-mushi („) | — | 45,0° | — | — | — | — | — | — | — |
| 20 | Umi-Jigoku | acid vitriol | 90,5° (26°) | — | — | — | — | — | — | — |
| 21 | Tsuru-no-yu (<i>Public bath</i>) | sulphur | 98,0° (18°) | 373 | — | — | — | — | — | — |
| 22 | Jizō-no-yu („) | acid alum
vitriol | 91,0° (22°) | 389 | — | 0,13 | 25,0° | K. & L. | Oct. 11, 1913 | Dr. R. Ishizu |
| 23 | Yakushi-no-yu („) | „ | 97,0° | 360 | — | — | — | — | — | — |
| 24 | Kamii-no-yu („) | „ | 63,0° (30°) | 90 | — | — | — | — | — | — |
| 25 | Kiyō-sen („) | common salt | 54,0° (24,5°) | 270 | — | 0,32 | 29,0° | K. & L. | „ 14, „ | Dr. R. Ishizu |
| 26 | Shi-no-yu („) | „ | 57,0° (24°) | 198 | — | 0,22 | 24,0° | „ | „ „ | „ |
| 27 | Kamado-no-yu | — | 99,0° (26°) | — | — | 0,22 | 22,0° | „ | „ 10, „ | „ |
| 28 | Noda-no-yu | — | 80,0° (21,5°) | — | — | 0,48 | 24,0° | „ | „ 12, „ | „ |
| 29 | Gomusō-no-yu | — | 48,0° (24°) | — | — | 0,45 | 25,0° | „ | „ „ | „ |
| 30 | Hamada-Onsen
(<i>Public bath</i>) | carbonated | 59,0° (25°) | 180 | — | 0,11 | 25,0° | „ | „ 13, „ | „ |
| 31 | Hijiri-yu | iron carbonate | 69,0° (23°) | 216 | — | 0,31 | 20,0° | „ | „ „ | „ |
| Gas evolving from : | | | | | Emanation per litre
of gas at 0° C. | | | Date when the
gas was
collected | | |
| 1 | Chinoike-Jigoku <i>Vent No. 1</i> | — | — | — | — | 4,82 | — | „ | Oct. 10, 1913 | „ |
| 2 | „ <i>Vent No. 2</i> | — | — | — | — | 4,99 | — | „ | „ 13, „ | „ |
| 3 | Hijiri-yu | — | — | — | — | 1,91 | — | „ | „ „ | „ |

OBAMA

Location.—Obama-mura, Minami-takagi-gun, Province Hizen, Nagasaki-ken.

The springs issue from the Augite Andesite.

The hot spring “Fukiage-yu” *

Analysis.

Analysed by the Hygienic Laboratory of Nagasaki-ken. 1909. Specific gravity: 1,0065 at 15° C.
Temperature: 65° C. Total residue: ca. 8,84.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|--|--------|
| Potassium chloride (KCl) | 0,3985 | Calcium sulphate (CaSO ₄) | 0,4000 |
| Sodium chloride (NaCl) | 5,4326 | Ferric oxide and alumina (Fe ₂ O ₃ +Al ₂ O ₃) . . | 0,0055 |
| Calcium chloride (CaCl ₂) | 1,2578 | Silica (SiO ₂) | 0,2200 |
| Magnesium chloride (MgCl ₂) | 0,0007 | | 8,5040 |

Besides these trace of ammonia and phosphoric acid.

* The spring is of intermittent nature and sinter cone is being formed around its jet (vide Plate IV, fig. 1).

The spring may be classified as “**earth-muriated common salt spring**”.

Radio-activity. 0.36×10^{-10} curies in 1 litre of water at 65,0° C.

(Schmidt's electrom.; H. Kibezaki and S. Nishimura, July 22, 1914).

0,10 Mache's units (recalculated).

Table 108.

| No. | Spring | Classification | Temp. of spring
in C. | Emanation per litre
of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------------------|-------------------------------|--------------------------|---------------------------------|---------------------|---------------------|--------------------|----------------------------------|-------------------------------|
| | | | | in
10 ⁻¹⁰ curies | in
Mache's units | at
(temp. in C.) | | | |
| 1 | Fukiage-yu (<i>Public bath</i>) | earth muriated
common salt | 65,0° | 0,36 | 0,10 | 65,0° | Schm. | July 22, 1914 | { H. Kibezaki
S. Nishimura |
| 2 | Gorin-yu („) | — | 94,0° | 2,31 | 0,64 | 94,0° | „ | „ 21, „ | „ |
| 3 | Meiji-yu | — | 54,0° | 0,83 | 0,23 | 54,0° | „ | „ | „ |
| 4 | Karimizu-Kōsen | carbonated | 24,0° | 3,29 | 0,90 | 24,0° | „ | „ 22, „ | „ |

UNZEN

Location.—Obama-mura, Minami-takagi-gun, Province Hizen, Nagasaki-ken.

The springs issue from the Hornblende Andesite.

The hot spring “Enryaku-yu”

Analysis.

Analysed by the Hygienic Laboratory of Nagasaki-ken. 1909.

Specific gravity: 1,0017 at 15° C.

Temperature: 65° C.

Total residue: ca. 1,20.

Flow of water: ca. 180 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|--------|---|----------------|
| Potassium sulphate (K_2SO_4) | 0,0172 | Free sulphuric acid (H_2SO_4) | 1,2887 |
| Sodium sulphate (Na_2SO_4) | 0,0613 | Silica (SiO_2) | 0,2450 |
| Calcium sulphate ($CaSO_4$) | 0,0656 | | 2,3819 |
| Magnesium sulphate ($MgSO_4$) | 0,0516 | Free hydrogen sulphide (H_2S) | 0,0035 |
| Ferrous sulphate ($FeSO_4$) | 0,2539 | | 2,3854 |
| Aluminium sulphate [$Al_2(SO_4)_3$] | 0,3986 | Free carbon dioxide (CO_2) | not determined |

Besides these a very small quantity of free hydrochloric acid and phosphoric acid and trace of ammonia.

The spring may be classified as “**acid hydrogen sulphide vitriol spring**”.

Radio-activity. 4.34×10^{-10} curies in 1 litre of water at 65,0° C.

(Schmidt's electrom.; H. Kibezaki and S. Nishimura, July 24, 1914).

1,19 Mache's units (recalculated).

The hot spring “Ko-Jigoku”

Analysis.

Analysed by the Hygienic Laboratory of Nagasaki-ken. 1910.

Specific gravity: 1,0007 at 15° C.

Temperature: 74° C.

Flow of water: ca. 260 hectolitres in 24 hours.

Total residue: 0,37.

In 1 kilogram of the mineral water are contained :

| | Grams | | Grams |
|--|--------|---|--------|
| Potassium sulphate (K_2SO_4) | 0.0483 | Free sulphuric acid (H_2SO_4) | 0.0735 |
| Sodium sulphate (Na_2SO_4) | 0.0017 | Silica (SiO_2) | 0.0020 |
| Calcium sulphate ($CaSO_4$) | 0.0729 | | 0.3634 |
| Ferric oxide and alumina ($Fe_2O_3+Al_2O_3$) | 0.0150 | | |

Besides these a very small quantity of free hydrochloric acid and phosphoric acid, trace of ammonia and magnesium sulphate.

The spring may be classified as “acid spring”.

Radio-activity. 1.46×10^{-10} curies in 1 litre of water at $74.0^\circ C$.

(Schmidt's electrom.; H. Kibezaki and S. Nishimura, July 24, 1914).

0.40 Mache's units (recalculated).

Table 109.

| No. | Spring | Classification | Temp. of spring
in $^\circ C$. | Flow of water
in 24 hrs.
in hectolitres | Emanation per
litre of water | | | Apparatus employed | Date when the
water was taken | Examined by |
|-----|-----------------------------------|-----------------------------------|------------------------------------|---|---------------------------------|---------------------|------------------------------|--------------------|----------------------------------|-------------------------------|
| | | | | | in
10^{-10} units | in
Mache's units | at
(temp. in $^\circ C$) | | | |
| 1 | Enryaku-yu (<i>Public bath</i>) | acid hydrogen
sulphide vitriol | 65.0° | ca.
180 | 4.34 | 1.19 | 65.0° | Schm. | July 24, 1914 | { H. Kibezaki
S. Nishimura |
| 2 | Shin-yu | acid hydrogen
sulphate | 38.0° | — | 8.51 | 2.34 | 38.0° | „ | „ 23. „ | „ |
| 3 | Ko-Jigoku | acid | 74.0° | 260 | 1.46 | 0.40 | 74.0° | „ | „ 24. „ | „ |
| 4 | Hiki-yu (<i>Yumei-Hotel</i>) | — | 63.0° | — | 2.69 | 0.74 | 63.0° | „ | „ 25. „ | „ |
| 5 | Shin-yu (<i>Jami-Jigoku</i>) | acid | 56.0° | 180 | — | — | — | — | — | — |
| 6 | „ (<i>Nomi-yu</i>) | acid hydrogen
sulphide | 61.0° | 180 | — | — | — | — | — | — |

MICHINO-O

Location.—Nagayo-mura, Nishi-sonogi-gun, Province Hizen, Nagasaki-ken.

The springs issue from the Tertiary Tuff.

The cold spring “Michino-o-Kōsen”

Temperature: 24.0°

Flow of water: ca. 108 hectolitres in 24 hours.

No accurate analysis has ever been made, but the spring seems to be “vitriol spring.”

Radio-activity. 5.50×10^{-10} curies in 1 litre of water at $24.0^\circ C$.

(Schmidt's electrom.; H. Kibezaki and S. Nishimura, July 27, 1914).

1.51 Mache's units (recalculated).

TAKEO

Location.—Takeo-machi, Kinoshima-gun, Province Hizen, Saga-ken.

The spring issues from the Mica Andesite.

The hot spring "Takeo-Onsen"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Temperature: 49° C.

Flow of water: ca. 1512 hectolitres in 24 hours.

Total residue: ca. 0.77.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents |
|--|--------|------------|-----------------------|
| Potassium ion (K ⁺) | 0.0046 | 0.1252 | 0.1252 |
| Sodium ion (Na ⁺) | 0.2154 | 9.3466 | 9.3466 |
| Calcium ion (Ca ⁺⁺) | 0.0036 | 0.0898 | 0.1796 |
| | | | 9.6514 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.1040 | 2.9337 | 2.9337 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.4094 | 6.7104 | 6.7104 |
| | | 0.7370 | 19.2057 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0480 | | 9.6441 |
| | 0.7850 | | |

| | Grams |
|---|--------|
| Potassium chloride (KCl) | 0.0088 |
| Sodium chloride (NaCl) | 0.1648 |
| Sodium bicarbonate (NaHCO ₃) | 0.5489 |
| Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0145 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0480 |
| | 0.7850 |

Besides these trace of magnesium and hydrosulphide ion.

The spring thus may be classified as "simple thermal".

TOCHINOKI

Location.—Nagakita-mura, Aso-gun, Province Higo, Kumamoto-ken.

The springs issue from the Augite Andesite.

Table 110.

| No. | Spring | Classification | Temp. of spring in C. | Emanation per litre of water | | | Apparatus employed | Date when the water was taken | Examined by |
|-----|-------------------------------------|------------------|-----------------------|------------------------------|-----------------|----------------|--------------------|-------------------------------|-------------|
| | | | | in 10 ⁻¹⁰ curies | in Mach's units | at temp. in C. | | | |
| 1 | Tochinoki-Onsen (Public bath) No. 1 | sulphated bitter | 45.0° | 0.66 | 0.18 | 45.0° | Schm. | March 12, 1913 | H. Kibezaki |
| 2 | " " No. 2 | " | 39.0° | 0.39 | 0.11 | 39.0° | " | " | " |
| 3 | Yunotani-Onsen (Public bath) | carbonated | 75.0° | 2.39 | 0.66 | 75.0° | " | " 13, " | " |
| 4 | Yunotani-Jigoku (Suzume-Jigoku) | — | 76.0° | 1.08 | 0.30 | 76.0° | " | " | " |
| 5 | Kami-no-yu (Public bath) | sulphur | 57.0° | 3.80 | 1.05 | 57.0° | " | " 14, " | " |
| 6 | Shimo-no-yu (") | " | 64.0° | 0.59 | 0.16 | 64.0° | " | " | " |
| 7 | Tarutama-Jigoku | — | 75.0° | 0.36 | 0.10 | 75.0° | " | " | " |

IWODANI

Location.—Makizono-mura, Aira-gun, Province Ōsumi, Kagoshima-ken.

The springs issue from the Volcanic Ash and Mud Lava.

The hot spring "Iwōdani-Onsen"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881. Temperature: 60.6° C.

Specific gravity: 1.0003 at 27° C.

Total residue: ca. 0.47.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents | | Grams |
|--|---------------|------------|-----------------------|---|--------|
| Potassium ion (K ⁺) | 0.0213 | 0.5441 | 0.5441 | Potassium chloride (KCl) | 0.0328 |
| Sodium ion (Na ⁺) | 0.0483 | 2.0955 | 2.0955 | Sodium chloride (NaCl) | 0.0110 |
| Calcium ion (Ca ⁺⁺) | 0.0188 | 0.4688 | 0.4688 | Potassium sulphate (K ₂ SO ₄) | 0.0092 |
| Magnesium ion (Mg ⁺⁺) | 0.0050 | 0.2053 | 0.4106 | Sodium bicarbonate (NaHCO ₃) | 0.1183 |
| Ferrous ion (Fe ⁺⁺) | 0.0029 | 0.0519 | 0.1038 | Sodium hydrosulphide (NaHS) | 0.0283 |
| | | | 4.0916 | Calcium sulphate (CaSO ₄) | 0.0636 |
| Anions. | | | | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0298 |
| Chlorine ion (Cl ⁻) | 0.0223 | 0.6291 | 0.6291 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0094 |
| Sulphate ion (SO ₄ ^{''}) | 0.0499 | 0.5195 | 1.0390 | Silicic acid (meta) (H ₂ SiO ₃) | 0.2151 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.1171 | 1.9195 | 1.9195 | | 0.5175 |
| Hydrosulphide ion (HS [']) | 0.0167 | 0.5040 | 0.5040 | Free carbon dioxide (CO ₂) | 0.0766 |
| | 0.3023 | 6.9377 | 4.0916 | Free hydrogen sulphide (H ₂ S) | 0.0522 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2151 | | | | 0.6463 |
| | 0.5174 | | | | |
| Free carbon dioxide (CO ₂) | 0.0766 | | 1.7418 | | |
| Free hydrogen sulphide (H ₂ S) | 0.0522 | | | | |
| | 0.6462 | | | | |

The spring thus may be classified as "hydrogen sulphide spring".

EINO-O

Location.—Makizono-mura, Aira-gun, Province Ōsumi, Kagoshima-ken.

The springs issue from the Volcanic Ash and Mud Lava.

The hot spring "Eino-o-Onsen"

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1881.

Specific gravity: 1.00048.

Temperature: 76.7° C.

Total residue: ca. 1.02.

Flow of water: ca. 5400 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equivalents | | Grams |
|--|---------------|------------|-----------------------|---|--------|
| Potassium ion (K ⁺) | 0.0128 | 0.3270 | 0.3270 | Sodium chloride (NaCl) | 0.0478 |
| Sodium ion (Na ⁺) | 0.0871 | 3.7787 | 3.7787 | Potassium sulphate (K ₂ SO ₄) | 0.0285 |
| Calcium ion (Ca ⁺⁺) | 0.0196 | 0.4888 | 0.9776 | Sodium sulphate (Na ₂ SO ₄) | 0.0816 |
| Magnesium ion (Mg ⁺⁺) | 0.0070 | 0.2874 | 0.5748 | Sodium bicarbonate (NaHCO ₃) | 0.0573 |
| Ferrous ion (Fe ⁺⁺) | 0.0019 | 0.0340 | 0.0680 | Sodium hydrosulphide (NaHS) | 0.0640 |
| | | | 5.7261 | Calcium sulphate (CaSO ₄) | 0.0665 |
| Anions. | | | | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0417 |
| Chlorine ion (Cl ⁻) | 0.0290 | 0.8181 | 0.8181 | Ferrous bicarbonate [Fe(HCO ₃) ₂] | 0.0060 |
| Sulphate ion (SO ₄ ^{''}) | 0.1178 | 1.2263 | 2.4526 | Silicic acid (meta) (H ₂ SiO ₃) | 0.2129 |
| Hydrocarbonate ion (HCO ₃ [']) | 0.0803 | 1.3164 | 1.3164 | | 0.6063 |
| Hydrosulphide ion (HS [']) | 0.0377 | 1.1390 | 1.1390 | Free carbon dioxide (CO ₂) | 0.1063 |
| | 0.3023 | 6.9377 | 5.7261 | Free hydrogen sulphide (H ₂ S) | 0.2374 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.2129 | | | | 0.9500 |
| | 0.6061 | | | | |
| Free carbon dioxide (CO ₂) | 0.1063 | | 2.4152 | | |
| Free hydrogen sulphide (H ₂ S) | 0.2374 | | | | |
| | 0.9498 | | | | |

The spring thus may be classified as "hydrogen sulphide spring".

ON-YŌ

Location.—Jinshū-men, Gazan-gun, Chūsei-nandō, Chōsen.

The springs issue from the Granite.

The hot spring “On-yō-Onsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1906.

Specific gravity: 1,0004 at 15° C.

Temperature: 38–42° C.

Total residue: ca. 0,21.

Flow of water: ca. 233 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-mols | Milligram-equiv. lents | | Grams |
|--|--------|------------|------------------------|--|--------|
| Potassium ion (K'). | 0,0051 | 0,1303 | 0,1303 | Sodium chloride (NaCl) | 0,0300 |
| Sodium ion (Na') | 0,0570 | 2,4729 | 2,4729 | Potassium bicarbonate (KHCO ₃) | 0,0130 |
| | | | 2,6032 | Sodium bicarbonate (NaHCO ₃) | 0,1648 |
| Anions. | | | | Silicic acid (meta) (H ₂ SiO ₃) | 0,0779 |
| Chlorine ion (Cl') | 0,0182 | 0,5134 | 0,5134 | | 0,2857 |
| Hydrocarbonate ion (HCO ₃ ') | 0,1275 | 2,0898 | 2,0898 | | |
| | | 0,2078 | 5,2064 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0,0779 | | | | |
| | 0,2857 | | | | |

Besides these trace of ferrous and aluminium ion.

The spring thus may be classified as “simple thermal”.

TŌRAI

Location.—Tōrai-men, Tōrai-gun, Keishō-nandō, Chōsen.

The springs issue from the Granite.

The hot spring “Tōrai-Onsen”

Analysis.

Analysed by the Government Hygienic Laboratory of Chōsen. 1908.

Specific gravity: 1,0012 at 15° C.

Temperature: 52° C.

Total residue: ca. 5,50.

Flow of water: ca. 907 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

| | Grams | | Grams |
|---|-------|---|-------|
| Sodium chloride (NaCl) | 2,961 | Ferrous oxyde and Alumina (FeO+Al ₂ O ₃) . . | 0,028 |
| Potassium carbonate (K ₂ CO ₃) | 0,068 | Silica (SiO ₂) | 0,540 |
| Sodium carbonate (Na ₂ CO ₃) | 0,186 | | 5,453 |
| Calcium sulphate (CaSO ₄) | 0,700 | Free carbon dioxide (CO ₂) | 0,066 |
| Calcium carbonate (CaCO ₃) | 0,520 | | 5,519 |
| Magnesium carbonate (MgCO ₃) | 0,450 | | |

The spring may be classified as “common salt spring”.

HOKUTŌ

Location.—Hokutō-shō, Shirai-nihō, Dohoku-shō, Taiwan (Formosa).

The springs issue from the Volcanic Rock.

The hot spring “Kaikōsha-Onsen”.

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1911.

Specific gravity: 1.0094 at 26° C.

Temperature: 51° C.

Flow of water: ca. 1,080 hectolitres in 24 hours.

Total residue: ca. 0.30.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|---------|-----------------|---------------------------|
| Potassium ion (K ⁺) | 0.00355 | 0.09068 | 0.09068 |
| Sodium ion (Na ⁺) | 0.01034 | 0.44859 | 0.44859 |
| Ammonium ion (NH ₄ ⁺) | 0.01572 | 0.86891 | 0.86891 |
| Calcium ion (Ca ⁺⁺) | 0.00679 | 0.16975 | 0.33950 |
| Magnesium ion (Mg ⁺⁺) | 0.00306 | 0.12562 | 0.25124 |
| Aluminium ion (Al ⁺⁺⁺) | 0.01458 | 0.53801 | 1.61403 |
| | | | 3.61295 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 0.00964 | 0.27193 | 0.27193 |
| Sulphate ion (SO ₄ ^{''}) | 0.16049 | 1.67073 | 3.34146 |
| | 0.22417 | 4.18422 | 3.61339 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.07582 | | |
| | 0.29999 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|---------|
| Ammonium chloride (NH ₄ Cl) | 0.01456 |
| Potassium sulphate (K ₂ SO ₄) | 0.00790 |
| Sodium sulphate (Na ₂ SO ₄) | 0.03188 |
| Ammonium sulphate [(NH ₄) ₂ SO ₄] | 0.03945 |
| Calcium sulphate (CaSO ₄) | 0.02310 |
| Magnesium sulphate (MgSO ₄) | 0.01513 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.09210 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.07582 |
| | 0.29999 |

Besides these trace of ferrous and ferric ion and boric acid.

The spring thus may be classified as “simple thermal”.

The hot spring “Taki-no-yu”

Analysis (calculated from the original numbers).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1911.

Specific gravity: 1.0152 at 27° C.

Flow of water: ca. 59530 hectolitres in 24 hours.

Temperature: 48.5° C.

In 1 kilogram of the mineral water are contained:

| Cations. | Grams | Milli-
moles | Milligram-
equivalents |
|--|---------|-----------------|---------------------------|
| Hydrogen ion (H ⁺) | 0.06575 | 65.06575 | 65.09901 |
| Potassium ion (K ⁺) | 0.41636 | 10.63499 | 10.63499 |
| Sodium ion (Na ⁺) | 1.12728 | 48.90586 | 48.90586 |
| Ammonium ion (NH ₄ ⁺) | 0.20867 | 11.54148 | 11.54148 |
| Calcium ion (Ca ⁺⁺) | 0.12665 | 3.166375 | 6.32750 |
| Magnesium ion (Mg ⁺⁺) | 0.02686 | 1.10263 | 2.20526 |
| Ferrous ion (Fe ⁺⁺) | 0.13791 | 2.46268 | 4.92536 |
| Aluminium ion (Al ⁺⁺⁺) | 0.15670 | 5.82222 | 17.34687 |
| | | | 166.98633 |
| Anions. | | | |
| Chlorine ion (Cl ⁻) | 3.76525 | 10.63499 | 106.21298 |
| Sulphate ion (SO ₄ ^{''}) | 2.91844 | 30.37143 | 60.76286 |
| Hydrophosphate ion (HPO ₄ ^{''}) | 0.00102 | 0.01062 | 0.01114 |
| | 8.95089 | 285.29772 | 166.99708 |
| Boric acid (meta) (HBO ₂) | 0.07805 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.17904 | | |
| | 9.20798 | | |

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| | Grams |
|---|---------|
| Potassium chloride (KCl) | 0.79337 |
| Sodium chloride (NaCl) | 1.10772 |
| Ammonium chloride (NH ₄ Cl) | 0.61773 |
| Sodium sulphate (Na ₂ SO ₄) | 2.13025 |
| Calcium sulphate (CaSO ₄) | 0.43078 |
| Magnesium sulphate (MgSO ₄) | 0.13277 |
| Ferrous sulphate (FeSO ₄) | 0.37445 |
| Aluminium sulphate [Al ₂ (SO ₄) ₃] | 0.98870 |
| Aluminium phosphate [Al ₂ (HPO ₄) ₃] | 0.00121 |
| Free hydrochloric acid (HCl) | 2.37365 |
| Boric acid (meta) (HBO ₂) | 0.07805 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.17904 |
| | 9.20772 |

Besides these trace of barium, lead and ferric ion.

The spring thus may be classified as “acid alum vitriol spring”.

SU-Ō

Location.— Su ō shō, Ritakukan-hō, Giran-chō, Taiwan (Formosa).

The spring issues from the Clay Slate.

The cold spring “Su-ō-Kōsen”

Analysis (calculated from the salt table).

Analysed by the Tōkyō Imperial Hygienic Laboratory. 1901.

Temperature: 23° C.

Total residue: ca. 0.14.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents | | Grams |
|--|--------|----------------|---------------------------|---|--------|
| Potassium ion (K ⁺) | 0.0006 | 0.0154 | 0.0154 | Potassium chloride (KCl) | 0.0012 |
| Sodium ion (Na ⁺) | 0.0222 | 0.9631 | 0.9631 | Sodium chloride (NaCl) | 0.0039 |
| Calcium ion (Ca ⁺⁺) | 0.0148 | 0.3691 | 0.7382 | Sodium bicarbonate (NaHCO ₃) | 0.0756 |
| Magnesium ion (Mg ⁺⁺) | 0.0048 | 0.1970 | 0.3940 | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0599 |
| Aluminium ion (Al ⁺⁺⁺) | 0.0021 | 0.0775 | 0.2325 | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0290 |
| | | | 2.3432 | Aluminium chloride (AlCl ₃) | 0.0102 |
| Anions. | | | | Silicic acid (meta) (H ₂ SiO ₃) | 0.0356 |
| Chlorine ion (Cl ⁻) | 0.0111 | 0.3131 | 0.3131 | | 0.2154 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 0.1242 | 2.0357 | 2.0357 | Free carbon dioxide (CO ₂) | 1.5682 |
| | 0.1798 | 3.9709 | 2.3488 | | 1.7836 |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0356 | | | | |
| | 0.2154 | | | | |
| Free carbon dioxide (CO ₂) | 1.5682 | 35.6409 | | | |
| | 1.7836 | | | | |

Besides these trace of ferrous and hydrophosphate ion.

The spring thus may be classified as “simple carbondioxated spring”.

KWANSHIREI

Location.—Kwanshirei-shō, Tarakaku-tōkahō, Kagi-chō, Taiwan (Formosa).

The springs issue from the Tertiary.

The hot spring “Kwanshirei-Onsen” No. 1

Analysis (calculated from the salt table).

Analysed by the Central Government Laboratory of Taiwan. 1912.

Specific gravity: 1.0090 at 22 C.

Temperature: 80° C.

Flow of water: ca. 260 hectolitres in 24 hours.

In 1 kilogram of the mineral water are contained:

The mineral water corresponds approximately in its composition to a solution containing in 1 kilogram:

| Cations. | Grams | Milli-
mols | Milligram-
equivalents | | Grams |
|--|--------|----------------|---------------------------|---|--------|
| Potassium ion (K ⁺) | 0.1865 | 4.7637 | 4.7637 | Potassium chloride (KCl) | 0.3553 |
| Sodium ion (Na ⁺) | 2.6982 | 117.0586 | 117.0586 | Sodium chloride (NaCl) | 4.1769 |
| Calcium ion (Ca ⁺⁺) | 0.0080 | 0.1995 | 0.3990 | Sodium bromide (NaBr) | 0.0220 |
| Magnesium ion (Mg ⁺⁺) | 0.0071 | 0.2915 | 0.5830 | Sodium iodide (NaI) | 0.0145 |
| | | | 122.8043 | Sodium sulphate (Na ₂ SO ₄) | 0.0311 |
| Anions. | | | | Sodium bicarbonate (NaHCO ₃) | 3.7783 |
| Chlorine ion (Cl ⁻) | 2.7000 | 76.1636 | 76.1636 | Calcium bicarbonate [Ca(HCO ₃) ₂] | 0.0324 |
| Bromine ion (Br ⁻) | 0.0171 | 0.2139 | 0.2139 | Magnesium bicarbonate [Mg(HCO ₃) ₂] | 0.0425 |
| Iodine ion (I ⁻) | 0.0123 | 0.0969 | 0.0969 | Silicic acid (meta) (H ₂ SiO ₃) | 0.0130 |
| Sulphate ion (SO ₄ ⁻) | 0.0210 | 0.2186 | 0.4372 | Organic substances | 0.0506 |
| Hydrocarbonate ion (HCO ₃ ⁻) | 2.8028 | 45.9400 | 45.9400 | | 8.5166 |
| | 8.4530 | 244.9463 | 122.8516 | | |
| Silicic acid (meta) (H ₂ SiO ₃) | 0.0130 | | | | |
| Organic substances | 0.0506 | | | | |
| | 8.5166 | | | | |

Besides these trace of ferrous and hydrophosphate ion.

The spring thus may be classified as “alkaline common salt spring”.

PART II (B)

(II) The Tables of Mineral Springs arranged according to :

- I. Radio-activity.
- II. Temperature.
- III. Flow of Water.
- IV. The Quantity of Solid Ingredients.

(III) A. The Tables of Hot Spring Spas arranged according to :

- I. Altitude.
- II. The Number of Visitors.

B. The Tables of Summer Resorts and Watering Places arranged geographically.

I. Radio-activity.

The sample of water for examination was generally taken directly from the source of each spring.

Radio-activity of the water marked with Asterisk (*) was determined with the sample brought to the laboratory by applicant, taking into account decay of radium emanation with the time.

Table 1.

The quantity of radium emanation was determined from the diminution in the charge after the **maximum current** had been reached, by employing Schmidt's electrometer standardized with a normal solution of radium.

For recalculation of Mache's unit from the international unit following factor was used:

$$1 \times 10^{-10} \text{ curie} = 0.275 \text{ Mache's units.}$$

(A) Hot Springs.

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in °C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|------------------------------|---------------------------------|---|------------------------|---|---------------------------|------------|----------|----------------------------|
| | in curies | in Mache's units (recalculated) | | | | | | | |
| 1 | 516.87 | 142.14 | ca. — | 71.0° | — | Misasa | 三 朝鳥取 | Granite | Private bath T. Matsubara |
| 2 | 371.83 | 102.25 | — | 51.0° | — | " | " | " | Seitō-kwan-no-yu |
| 3 | 109.51 | 30.12 | 126 | 44.0° | sulphur | Sekigane | 關 金 (") | " | Shimo-jyaya-no-yu |
| 4 | 94.03 | 25.85 | 1800 | 39.0° | simple | Tochiomata | 栃 尾 又 新湯 | " | Tochiomata-no-yu (No. 1) |
| 5 | 43.21 | 11.88 | — | 40.0° | salt | Kawatana | 川 棚 山口 | " | Seiryū-sen |
| 6 | 37.19 | 10.23 | — | 45.0° | — | Misasa | 三 朝鳥取 | " | Kabu-yu |
| 7 | 36.19 | 9.95 | — | 41.0° | salt | Kawatana | 川 棚 山口 | " | Juei-sen |
| 8 | 31.18 | 8.57 | — | 56.0° | mineralized saline bitter | Kachim | 珍 見 鳥取 | " | Nakataya-no-yu |
| 9 | 30.57 | 8.41 | — | 60.3° | earth mineralized common salt | Kinosaki | 城 崎 兵庫 | Tertiary | Gosho-no-yu (Part No. 2) |
| 10 | 25.21 | 6.93 | — | 93.0° | " | Wakura | 和 倉 石川 | " | Wakazaki-no-yu (No. 1) |
| 11 | 18.00 | 4.95 | 544 | 46.0° | sulphated common salt | Yunotsu | 溫 泉 津 島根 | " | Shin-yu |
| 12 | 16.07 | 4.42 | — | 46.0° | simple | Dogo | 道 後 愛媛 | Granite | Yōjō-yu |
| 13 | 15.13 | 4.16 | 504 | 72.0° | salt | Takase | 高 瀬 新湯 | Alluvium | Takase-no-yu |
| 14 | 14.78 | 4.07 | — | 44.0° | sulphur | Musashi | 武 藏 福岡 | Granite | Yakushi-no-yu (No. 2) |
| 15 | 14.47 | 3.98 | — | 47.0° | simple | Dogo | 道 後 愛媛 | " | Kami-no-yu |
| 16 | 13.65 | 3.75 | 540 | 38.2° | non-carbonated common salt | Koyabara | 小 屋 原 (島根) | Andesite | Koyabara-Onsen |
| 17 | 13.60 | 3.74 | 3888 | 43.0° | simple | Yumura (Shimane-ken) | 湯 村 " " | Granite | Shitsuni-no-yu |
| 18 | 13.58 | 3.74 | 216 | 59.5° | earth-mineralized common salt | Kinosaki | 城 崎 兵庫 | Tertiary | Yanagi-no-yu |
| 19 | 13.46 | 3.70 | 315 | 46.0° | sulphur | Musashi | 武 藏 福岡 | Granite | Enju-no-yu |
| 20 | 12.77 | 3.51 | — | 50.5 | earth-mineralized common salt | Kinosaki | 城 崎 兵庫 | Tertiary | Gosho-no-yu (Part No. 1) |
| 21 | 12.09 | 3.33 | 830 | 39.5° | carbonated | Deyu | 出 湯 新湯 | Granite | Chōsenkutsu-no-yu |
| 22 | 11.91 | 3.28 | 39 | 47.5° | mineralized saline bitter | Yoshikata | 吉 方 鳥取 | Alluvium | Tottori-Onsen |
| 23 | 11.67 | 3.21 | 207 | 44.5° | sulphur | Musashi | 武 藏 福岡 | Granite | Osakaya-no-yu |
| 24 | 11.27 | 3.10 | 65 | 50.0° | salt | Yuzawa [Onnagawa] | 湯 澤 (大) 新潟 | Alluvium | Moto-yu |
| 25 | 11.25 | 3.09 | — | 57.0° | simple | O-yu (Niigata-ken) | 大 湯 " " | Granite | O-yu (Onnagawa) |
| 26 | 11.17 | 3.07 | 1730 | 50.0° | " | Togo | 東 郷 鳥取 | Andesite | Yōjō-kwan-no-yu (No. 1) |
| 27 | 11.08 | 3.05 | — | 47.0 | earth-mineralized common salt | Kinosaki | 城 崎 兵庫 | Tertiary | Mandara-no-yu (Part No. 2) |
| 28 | 10.79 | 2.97 | 1953 | 64.0° | common salt | Tamatsukuri (Shimane-ken) | 玉 造 島根 | " | Kami-no-yu |
| 29 | 10.33 | 2.81 | 2592 | 56.5° | simple | Yoshioka | 吉 岡 鳥取 | " | Kabu-yu |
| 30 | 10.08 | 2.77 | — | 59.5° | earth-mineralized common salt | Kinosaki | 城 崎 兵庫 | " | Ichi-no-yu (Part No. 2) |
| 31 | 9.91 | 2.73 | — | 43.0 | — | Hirase (Shimane-ken) | 廣 瀬 島根 | " | Sagi-no-yu |
| 32 | 9.91 | 2.73 | — | 50.5° | earth-mineralized common salt | Kinosaki | 城 崎 兵庫 | " | Ichi-no-yu (Part No. 1) |
| 33 | 9.72 | 2.67 | — | 45.0 | salt | Yuda | 湯 田 山口 | Alluvium | Nagamune-no-yu |
| 34 | 9.69 | 2.64 | — | 41.0° | " | " | " | " | Matsudaya-no-yu |
| 35 | 9.37 | 2.58 | 204 | 57.0° | earth-mineralized common salt and hydrogen sulphate | Kinosaki | 城 崎 兵庫 | Tertiary | Kō-no-yu |
| 36 | 8.51 | 2.34 | — | 38.0° | " | Unzen | 溫 泉 (長崎) | Andesite | Shin-yu |

| No. | Flow of water per litre of water | | Flow of water in 24 hrs. in bottles | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|----------------------------------|--------------------------------|-------------------------------------|-----------------------|--------------------------------|--------------------|-----------|-----------------|---------------------------|
| | in bottles | in Machine's tank recalculated | | | | | | | |
| 37 | 7.64 | 2.10 | - | 73.5° | earth-muriated common salt | Wakura | 和倉石川 | Tertiary | Sō-yu |
| 38 | 7.48 | 2.06 | - | - | simple | Kashi | 甲子(福島) | Granite | Kashi-Onsen |
| 39 | 7.29 | 2.00 | - | 41.0° | sulphur | Musashi | 武藏(福岡) | " | Yakushi-no-yu No. 1 |
| 40 | 6.55 | 1.87 | 900 | 45.0° | alkaline sulphur | Katsu-ura | 勝浦和歌山 | Tertiary | Akashima-Onsen |
| 41 | 6.21 | 1.71 | - | 42.0° | alkaline | Tawarayama | 俵山山口 | Mesoz. Conglom. | Kawa-no-yu |
| 42 | 6.15 | 1.69 | 270 | 40.0° | simple | Kizu | 木津(京都) | Tertiary | Iwatsubo-no-yu |
| 43 | 6.15 | 1.69 | 1800 | 45.2° | sulphated bitter sulphur | Yamanaka | 山中石川 | " | Yamanaka-Onsen (Sō-yu) |
| 44 | 5.93 | 1.63 | - | 59.5° | sulphur | Nakabusa | 中房長野 | Granite | Shirataki-no-yu |
| 45 | 5.21 | 1.43 | 207 | 48.5° | earth-muriated common salt | Kinosaki | 城崎(兵庫) | Tertiary | Jizō-no-yu |
| 46 | 5.17 | 1.42 | - | 58.0° | " | " | " | " | Mandara-no-yu Vent. No. 1 |
| 47 | 4.82 | 1.33 | 340 | 49.0° | sulphated common salt | Omaki | 大牧(富山) | Porphyr. Dyke | Omaki-Onsen |
| 48 | 4.65 | 1.28 | - | 79.0° | common salt | Katayamazu | 片山津(石川) | Tertiary | Shiotsu-no-Tsubo |
| 49 | 4.34 | 1.99 | 180 | 65.0° | acid hydrogen sulphide vitriol | Unzen | 温泉(長崎) | Andesite | Enryaku-yu |
| 50 | 4.04 | 1.11 | - | - | sulphated common salt | Atsumi | 温海(山形) | Tertiary | Yuatsumi-no-yu |
| 51 | 3.89 | 1.05 | - | 57.0° | sulphur | Tarutama | 垂玉(熊本) | Andesite | Kami-no-yu |
| 52 | 3.59 | 0.99 | - | 40.0° | alkaline | Tawarayama | 俵山山口 | Mesoz. Conglom. | Kagi-yu |
| 53 | 3.54 | 0.97 | 2700 | 54.3° | sulphur | Awazu | 栗津石川 | Liparite | Awazu Onsen (Sō-yu) |
| 54 | 3.11 | 0.86 | 300 | 50.0° | saline bitter | Yujiku | 湯宿(群馬) | Tertiary | Ō-daki |
| 55 | 3.09 | 0.85 | 900 | 38.5° | sulphur | Katsu-ura | 勝浦和歌山 | " | Soto-no-yu |
| 56 | 3.09 | 0.85 | 900 | 90.5° | alkaline carbonated | Yumura (Hyōgo-ken) | 湯村(兵庫) | Granite | Kabu-yu |
| 57 | 3.06 | 0.84 | 1290 | 38.0° | earth-muriated common salt | Yunogo | 湯郷岡山 | Liparite | Sagi-no-yu |
| 58 | 3.03 | 0.83 | - | 45.0° | salt | Yuzawa [Uo-numa] | 湯澤(新潟) | Misaka Series | Moto-yu |
| 59 | 2.90 | 0.89 | 90 | 47.0° | saline bitter | Yujiku | 湯宿(群馬) | Tertiary | Ko-daki |
| 60 | 2.69 | 0.74 | - | 63.0° | - | Unzen | 温泉(長崎) | Andesite | Hiki-yu (Yumei-Hotel) |
| 61 | 2.56 | 0.70 | 648 | 79.0° | saline bitter | Yujiku | 湯宿(群馬) | Tertiary | Kubo-yu |
| 62 | 2.51 | 0.69 | - | 71.0° | saline bitter sulphur | Yamashiro | 山代石川 | " | Araya-no-yu |
| 63 | 2.39 | 0.66 | - | 75.0° | carbonated | Yunotani | 湯ノ谷熊本 | Andesite | Yunotani-Onsen |
| 64 | 2.31 | 0.64 | - | 94.0° | - | Obama | 小濱長崎 | " | Gorin-yu |
| 65 | 2.30 | 0.63 | - | 67.0° | saline bitter | Yujiku | 湯宿(群馬) | Tertiary | Sen-tō |
| 66 | 1.59 | 0.41 | - | 37.2° | " | " | " | " | Kawaburi-yu |
| 67 | 1.48 | 0.41 | 2142 | 69.0° | saline bitter sulphur | Yamashiro | 山代(石川) | " | Yamashiro-Onsen (Sō-yu) |
| 68 | 1.46 | 0.40 | 200 | 74.0° | acid | Unzen | 温泉(長崎) | Andesite | Ko-Jigoku |
| 69 | 1.19 | 0.33 | 441 | 41.5° | bitter | Ushio | 海潮(島根) | " | Ushio-no-yu |
| 70 | 1.08 | 0.30 | - | 76.0° | - | Yunotani | 湯ノ谷熊本 | " | Yunotani-Jigoku |
| 71 | 0.92 | 0.25 | - | 51.5° | common salt | Arima | 有馬(兵庫) | Quartz Porph. | Arima-Onsen |
| 72 | 0.83 | 0.23 | - | 54.0° | - | Obama | 小濱長崎 | Andesite | Meiji-yu |
| 73 | 0.80 | 0.22 | - | 48.0° | simple | Arifuku | 有福島根 | Diorite | Arifuku-Onsen Vent. No. 1 |
| 74 | 0.75 | 0.21 | - | 55.0° | carbonated common salt | Beppu | 別府大分 | Andesite | Kojimaya-no-yu No. 1 |
| 75 | 0.66 | 0.18 | - | 45.0° | sulphated bitter | Tochinoki | 栃木熊本 | " | Tochinoki-Onsen No. 1 |
| 76 | 0.66 | 0.18 | 648 | 57.5° | iron carbonate | Beppu | 別府大分 | " | Furō-sen |
| 77 | 0.59 | 0.16 | - | 64.0° | sulphur | Tarutama | 垂玉熊本 | " | Shimo-no-yu |
| 78 | 0.53 | 0.14 | 87 | 49.0° | muriated alkaline carbonated | Sedo-no-kanayama | 瀬戸鉛山(和歌山) | Tertiary | Moto-no-yu |
| 79 | 0.52 | 0.14 | - | 54.0° | common salt | Beppu | 別府大分 | Andesite | Kojimaya-no-yu No. 2 |
| 80 | 0.49 | 0.14 | 200 | 42.0° | muriated alkaline carbonated | Sedo-no-kanayama | 瀬戸鉛山(和歌山) | Tertiary | Yagata-yu |
| 81 | 0.49 | 0.14 | - | - | - | Yugano | 湯ヶ野静岡 | Andesite | Spring B |
| 82 | 0.48 | 0.13 | - | - | - | " | " | " | " A |
| 83 | 0.45 | 0.12 | - | 40.0° | alkaline | Fukawa | 深川山口 | Liparite | Yumoto-Onsen (On-tō) |
| 84 | 0.43 | 0.12 | 150 | 52.0° | muriated alkaline carbonated | Sedo-no-kanayama | 瀬戸鉛山(和歌山) | Tertiary | Hama-no-yu |
| 85 | 0.40 | 0.11 | - | 42.0° | alkaline | Fukawa | 深川山口 | Liparite | Yumoto-Onsen (Rei-tō) |
| 86 | 0.40 | 0.11 | 239 | 52.0° | earthy alkaline | Shirahone | 白根(長野) | Chichibu Series | Ō-yu |
| 87 | 0.39 | 0.11 | - | 57.0° | carbonated common salt | Beppu | 別府大分 | Andesite | Shiokyū-no-yu |
| 88 | 0.39 | 0.11 | - | 39.0° | sulphated bitter | Tochinoki | 栃木熊本 | " | Tochinoki-Onsen No. 2 |
| 89 | 0.36 | 0.10 | - | 65.0° | earth-muriated common salt | Obama | 小濱長崎 | " | Fukiage-yu |
| 90 | 0.36 | 0.10 | - | 75.0° | - | Tarutama | 垂玉熊本 | " | Tarutama-Jigoku |

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|--------------------|------------------------------|------------------------------|---|-----------------------|----------------------------|-----------------|-------|----------|----------------------|
| | in c.c. to 10 c.c. | in Maché's units, calculated | | | | | | | |
| 91 | 0.36 | 0.10 | 260 | 60.0° | carbonated, common salt | Sedo-no-kanyama | 瀬戸鉛山 | Tertiary | Saki-no-yu |
| 92 | 0.22 | 0.06 | 300 | 102.0° | common salt | Senami | 瀬波新湯 | — | Senami-Funtō |
| 93 | 0.20 | 0.05 | — | 54.0° | carbonated | Boppu | 別府大分 | Andesite | Seisenkaku-no-yu |
| 94 | 0.13 | 0.04 | — | — | — | Yugano | 湯ヶ野静岡 | „ | Spring C |
| 95 | 0.07 | 0.02 | 210 | 57.0° | iron carbonate | Kwankaji | 観海寺大分 | „ | Kwankaji-Onsen |
| Gas evolving from: | | | | | | | | | |
| 1 | 339.32 | 93.31 | — | 93.0° | carbonated common salt | Wakura | 和倉石川 | Tertiary | Wakazaki-no-yu No. 1 |
| 2 | 228.19 | 62.75 | — | 45.0° | — | Misasa | 三朝鳥取 | Granite | Kabu-yu |
| 3 | 197.62 | 54.35 | — | 50.0° | mineral saline bitter | Kachimi | 勝見 | „ | Nakataya-no-yu |
| 4 | 128.04 | 35.21 | — | — | — | Tamatsukuri | 玉造島根 | Tertiary | River bed A |
| 5 | 118.33 | 32.54 | — | 44.5° | sulphur | Musashi | 武藏福岡 | Granite | Osakaya no-yu |
| 6 | 101.79 | 27.99 | — | 64.0° | common salt | Tamatsukuri | 玉造島根 | Tertiary | Kami-no-yu |
| 7 | 99.13 | 27.26 | — | 56.5° | simple | Yoshioka | 吉岡鳥取 | „ | Kabu-yu |
| 8 | 87.91 | 24.18 | — | 79.0° | common salt | Katayamazu | 片山津石川 | „ | Shiotsu-no-Tsubo |
| 9 | 85.02 | 23.38 | — | 47.0° | simple | Dogo | 道後愛媛 | Granite | Kami-no-yu |
| 10 | 84.30 | 23.18 | — | 43.0° | „ | Yumura | 湯村島根 | „ | Shitsuni-no-yu |
| 11 | 33.36 | 9.18 | — | 67.0° | saline bitter | Yujiku | 湯宿群馬 | Tertiary | Sen-tō |
| 12 | 30.69 | 8.44 | — | 71.0° | saline bitter-sulphur | Yamashiro | 山代石川 | „ | Araya-no-yu |
| 13 | 29.53 | 8.12 | — | 38.2° | iron carbonate common salt | Koyabara | 小屋原島根 | Andesite | Koyabara-Onsen |
| 14 | 19.50 | 5.35 | — | 38.0° | carbonated common salt | Yunogō | 湯郷岡山 | Liparite | Sagi-no-yu |
| 15 | 9.03 | 2.48 | — | 41.5° | bitter | Ushio | 海潮島根 | Andesite | Ushio-no-yu |
| 16 | 4.92 | 1.35 | — | 46.0° | sulphated common salt | Yunotsu | 温泉津 | Tertiary | Shin-yu |

† Radio-activity of the gas was determined at a temperature lower than 50° C, but the correction of the volume due to temperature was not made.

(B) Cold Springs.

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|------------------------------|------------------------------|---|-----------------------|--------------------|----------|-------|---------|-----------------------------|
| | in c.c. to 10 c.c. | in Maché's units, calculated | | | | | | | |
| 1 | 3012.14 | 828.34 | 22 | 21.5° (23.5°) | earthy common salt | Masutomi | 増富山梨 | Granite | Kamigawara No. 1 |
| 2 | 2390.02 | 657.26 | — | 14.0° | „ | „ | „ | „ | „ No. 2 |
| 3 | 2147.05 | 590.44 | 40 | 10.0° (7°) | common salt | „ | „ | „ | Kuridaira No. 1 |
| 4 | 1433.17 | 394.12 | 77 | 23.5° (4.5°) | earthy common salt | „ | „ | „ | Nibuzawa No. 1 |
| 5 | 1157.29 | 318.25 | 100 | 20.0° (4°) | common salt | „ | „ | „ | Kuridaira No. 1 (cont.) |
| 6 | 1042.17 | 286.60 | 13 | 20.0° (10.5°) | — | „ | „ | „ | „ No. 3 |
| 7 | 1022.15 | 281.09 | — | 10.0° (13°) | simple | Takayama | 高田山岐阜 | „ | Ena Kosen (Yamanashi) |
| 8 | 1008.76 | 277.41 | — | 19.0° (11.5°) | — | Masutomi | 増富山梨 | „ | Oshika No. 1 |
| 9 | 900.12 | 247.53 | — | 17.0° | — | „ | „ | „ | Yunokubo No. 1 |
| 10 | 883.39 | 243.82 | — | 15.0° (10.5°) | earthy common salt | „ | „ | „ | Yunokubo-gawara no-yu |
| 11 | 856.83 | 235.63 | 45 | 17.5° (7°) | „ | „ | „ | „ | Tsugino-yu No. 1 (I) |
| 12 | 682.67 | 187.74 | — | 17.0° (12°) | carbonated (?) | Ikeda | 池田島根 | „ | Nolata-shin-yu |
| 13 | 372.63 | 102.47 | — | 29.5° (7°) | — | Masutomi | 増富山梨 | „ | Ginsen-tō |
| 14 | 220.64 | 60.68 | — | 12.0° (20°) | simple | Hirukawa | 蜷川岐阜 | „ | Ena-Kosen (Dai-2 Yamanashi) |
| 15 | 213.21 | 58.63 | — | 13.0° (1°) | „ | Takayama | 高田山 | „ | „ (Shinano) |
| 16 | 180.41 | 49.61 | 530 | 25.0° (10°) | „ | Murasugi | 村杉新潟 | „ | Murasugi-Kosen No. 1 |

| No. | Water temp. at
100 fathoms | Water temp. at
200 fathoms | Water temp. at
300 fathoms | Temp. of spring
in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|-------------------------------|-------------------------------|-------------------------------|--------------------------|--|---------------------|---------|---------------|-------------------------|
| 17 | 176.61 | 48.57 | — | 26.0° | simple | Murasugi | 村 杉新湯 | Granite | Murasugi-Kō-en No. 2 |
| 18 | 161.74 | 44.48 | 230 | 36.0° (27.5°) | — | Tochimoto | 所 尾 又 | " | Jizai kwan-uchiyu No. 1 |
| 19 | 107.67 | 29.61 | — | 28.5° | — | Arima | 有 馬兵庫 | Quartz Porph. | Zuihōji-Onsen |
| 20 | 60.15 | 16.54 | — | 15.0° | simple | Koshiki-iwa-shinden | 越水岩新田 | Granite | Kuraku-en-Kōsen No. 10 |
| 21 | 58.33 | 16.04 | — | 14.0° | " | " | " | " | Sarumaru-ishiyama No. 3 |
| 22 | 56.80 | 15.62 | — | 14.5° | " | " | " | " | " No. 1 |
| 23 | 51.46 | 14.15 | — | 12.0° | " | " | " | " | " No. 2 |
| 24 | 48.66 | 13.38 | — | 14.5° | " | Kaidani | 柏 谷岡山 | " | Tomada-no-yu |
| 25 | 44.29 | 12.18 | — | 14.5° | " | Koshiki-iwa-shinden | 越水岩新田 | " | Sarumaru-ishiyama No. 5 |
| 26 | 43.48 | 11.96 | 130 | 14.0° | sulphur | Bobata | 母 畑福島 | Gneiss | Kami-no-moto-yu |
| 27 | 35.71 | 9.82 | — | 14.0° | simple | Koshiki-iwa-shinden | 越水岩新田 | Granite | Sarumaru-ishiyama No. 6 |
| 28 | 34.51 | 9.49 | — | 19.0° | " | " | " | " | Kuraku-en-Kōsen No. 1 |
| 29 | 33.67 | 9.26 | — | 11.0° | " | " | " | " | " No. 11 |
| 30 | 31.02 | 8.53 | — | 16.8° (13.5°) | " | " | " | " | " No. 6 |
| 31 | 30.07 | 8.27 | — | 19.0° | — | Arima | 有 馬 | Quartz Porph. | Maruyama-Kōsen |
| 32 | 26.22 | 7.21 | — | 18.0° | simple | Koshiki-iwa-shinden | 越水岩新田 | Granite | Kuraku-en-Kōsen No. 3 |
| 33 | 25.87 | 7.11 | 66 | 24.5° (9°) | earthy common salt | Masutomi | 増 富山梨 | " | Tochikubo No. 1 |
| 34 | 24.04 | 6.61 | — | 29.0° | — | Dogo | 道 後愛媛 | " | Shin-Onsen |
| 35 | 22.22 | 6.11 | — | 14.0° | simple | Koshiki-iwa-shinden | 越水岩新田 | " | Sarumaru-ishiyama No. 4 |
| 36 | 22.04 | 6.06 | — | 18.0° | " | " | " | " | Kuraku-en-Kōsen No. 2 |
| 37 | 21.47 | 5.91 | 22 | — | " | Nekonaki | 猫 崎福島 | Gneiss | Nekonaki-no-yu |
| 38 | 21.31 | 5.86 | — | 24.0° | " | Okoto | 雄 琴滋賀 | Alluvium | Spring B S. Inai |
| 39 | * 20.84 | 5.73 | — | — | " | Ono | 大 野 | Diluvium | Spring A (I. Kishimoto) |
| 40 | 20.80 | 5.72 | — | 12.0° | " | Koshiki-iwa-shinden | 越水岩新田 | Granite | Kuraku-en-Kōsen No. 12 |
| 41 | 19.91 | 5.47 | — | 19.5° | simple | Takedao | 武 田尾 | Quartz Porph. | Yu-moto |
| 42 | 19.02 | 5.23 | — | 17.0° | simple | Koshiki-iwa-shinden | 越水岩新田 | Granite | Kuraku-en-Kōsen No. 8 |
| 43 | * 18.85 | 5.18 | — | — | carbonated | Okawa | 大 桑長野 | " | Shika-no-yu |
| 44 | 18.80 | 5.17 | — | 18.0° | simple | Koshiki-iwa-shinden | 越水岩新田 | " | Kuraku-en-Kōsen No. 7 |
| 45 | * 18.73 | 5.15 | — | — | " | Nishino | 西 野 | Tertiary | Spring B K. Miyazaki |
| 46 | 17.93 | 4.93 | — | 15.0° | " | Koshiki-iwa-shinden | 越水岩新田 | Granite | Kuraku-en-Kōsen No. 9 |
| 47 | 17.66 | 4.85 | 300 | 28.0° | " | Takebe | 建 部岡山 | " | Yawata-Onsen |
| 48 | 16.15 | 4.44 | — | 14.5° | " | Koshiki-iwa-shinden | 越水岩新田 | " | Kuraku-en-Kōsen No. 13 |
| 49 | 15.42 | 4.24 | 90 | 32.0° | alkaline common salt | Yudani | 湯 谷鳥取 | Tertiary | Kabu-yu |
| 50 | 14.73 | 4.05 | — | 15.0° (13.5°) | simple | Koshiki-iwa-shinden | 越水岩新田 | Granite | Kuraku-en-Kōsen No. 5 |
| 51 | 13.93 | 3.83 | — | 18.0° | " | " | " | " | " No. 14 |
| 52 | 13.56 | 3.73 | — | 17.5° | earthy iron con-
taining com-
mon salt | Odake | 大 岳 | Quartz Porph. | Odake-Kōsen |
| 53 | 12.84 | 3.53 | — | 18.5° | simple | Koshiki-iwa-shinden | 越水岩新田 | Granite | Kuraku-en Kōsen No. 4 |
| 54 | 12.05 | 3.31 | 285 | 27.0° | " | Bobata | 母 畑福島 | Gneiss | Shimo-no-yu No. 1 |
| 55 | 12.03 | 3.31 | — | 18.5° | simple carbon-
ated No. 1 | Takarazuka | 寶 塚兵庫 | Granite | Takarazuka-Tansan-sui |
| 56 | 11.45 | 3.14 | — | — | simple | Koto-ura | 琴 浦岡山 | " | Spring C (K. Miki) |
| 57 | 11.41 | 3.14 | — | — | " | Bobata | 母 畑福島 | Gneiss | Kami-no-shin-yu |
| 58 | * 11.38 | 3.13 | — | — | simple | Nishinomiya | 西 ノ 宮兵庫 | Granite | Kōro-en-Kōsen |
| 59 | 11.34 | 3.12 | — | 23.5° | simple | Takedao | 武 田尾 | Quartz Porph. | Moto-no-yu |
| 60 | * 10.75 | 2.96 | — | 18.5° | — | Hokaizawa | 法 界 澤青森 | Tertiary | Natsumazawa Kōsen |
| 61 | 10.36 | 2.85 | — | 27.0° | " | Katsu-ura | 勝 浦和歌山 | " | Nakanoshima-Onsen |
| 62 | 9.64 | 2.65 | — | — | simple | Nishino | 西 野兵庫 | " | Spring A K. Miyazaki |
| 63 | 9.42 | 2.59 | — | 14.0° | " | Koshiki-iwa-shinden | 越水岩新田 | Granite | Myōbandani-ue No. 2 |
| 64 | 8.85 | 2.43 | — | 23.5° | " | Dogo | 道 後愛媛 | " | Well Furayama |
| 65 | 8.58 | 2.36 | — | — | simple | Bobata | 母 畑福島 | Gneiss | Naka-no-yu No. 1 |
| 66 | 8.39 | 2.31 | — | — | carbonated | Sawaguchi | 澤 口秋田 | Diluvium | Ojigasawa-Kōsen No. 3 |
| 67 | * 8.26 | 2.27 | — | — | " | Koto-ura | 琴 浦岡山 | Granite | Spring D (K. Miki) |
| 68 | 7.58 | 2.08 | 300 | 31.5° | " | Masutomi | 増 富山梨 | " | Kinsen-tō |
| 69 | 7.42 | 2.04 | 15 | 26.0° (17°) | — | " | " | " | Kuridaira No. 2 (A) |

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名 府縣名 | Geology | Spring |
|-----|------------------------------|-----------------|---|-----------------------|---|-----------------------------------|------------|------------------|--------------------------|
| | in curies | in Mach's units | | | | | | | |
| 70 | 7.35 | 2.02 | ca. | — | vitriol | Koshakawa (Iwate) | 越木岩新田 | Granite | Myōbandani No. 1 |
| 71 | 6.79 | 1.87 | 41 | 20.0° (9.5°) | — | Masutomi | 増 富山梨 | " | Kuradaira No. 2 |
| 72 | 6.78 | 1.87 | — | — | — | Koto-ura | 琴 浦 岡山 | " | Spring B (M. Miki) |
| 73 | 6.73 | 1.85 | — | — | earthy common salt | Hoshio | 星 尾 群馬 | Chichibu Series | Shiomizu-Kōsen |
| 74 | 6.34 | 1.75 | — | 17.0° | simple carbon-dioxated | Arima | 有 馬 兵庫 | Quartz Porph. | Arakawa (M. Miki) |
| 75 | 5.81 | 1.60 | — | 13.9° | simple | Toyohira | 豊 平 長野 | Andesite | Karasawa-Kōsen A |
| 76 | 5.76 | 1.58 | — | 17.0° (6) | — | Masutomi | 増 富山梨 | Granite | Yamawata No. 1 |
| 77 | 5.50 | 1.51 | 108 | 24.0° | vitriol | Michi-no-o | 道 之 尾 長崎 | Tertiary | Michi-no-o-Kōsen |
| 78 | 5.31 | 1.45 | — | 22.0° | — | Yukawa | 湯 川 和歌山 | Misaka Series | Ōhata-no-yu |
| 79 | 5.17 | 1.42 | — | 35.5° | salt | Yumura | 湯 村 山梨 | Alluvium | Tani-Onsen |
| 80 | 5.10 | 1.40 | — | — | — | Yamanashi-ken
Tokyo-Shitaya-ku | 下 谷 東京 | Diluvium | Well (C. Sasaki) |
| 81 | 4.81 | 1.32 | — | 17.0° | common salt | Tomoe | 巴 茨城 | " | Tomoe-Kōsen No. 2 |
| 82 | 4.75 | 1.31 | — | 18.5° | — | Daishi-gawara | 大師河原 (神奈川) | Alluvium | Well (T. Miki) |
| 83 | 4.72 | 1.30 | 270 | 16.9° | common salt | Tomoe | 巴 茨城 | Diluvium | Tomoe-Kōsen No. 1 |
| 84 | 4.36 | 1.20 | — | — | vitriol | Koshakawa-land | 越木岩新田 | Granite | Myōbandani No. 2 |
| 85 | 4.22 | 1.16 | — | 14.5° | — | Honjō | 本 庄 茨城 | Alluvium | Fukae-Kōsen |
| 86 | 4.12 | 1.13 | — | — | earthy common salt | Atsushio | 熱 鹽 (福島) | " | Atsushio-Kōsen No. 1 |
| 87 | 3.90 | 1.07 | — | — | — | Dōgo | 道 後 愛媛 | Granite | Tama-no-i |
| 88 | 3.78 | 1.04 | — | 9.0° | — | Isobe (Gunma-ken) | 磯 部 群馬 | Tertiary | Ji-no-yu |
| 89 | 3.73 | 1.03 | — | — | — | Shirakawa | 白 河 福島 | Alluvium | Emmyōji-Kōsen |
| 90 | 3.43 | 0.94 | 63 | 17.5° | simple carbon-dioxated | Funagoya | 船 小 屋 福岡 | Diluvium | Funagoya-Onsen |
| 91 | 3.29 | 0.90 | — | 24.0° | carbonated | Obama | 小 濱 長崎 | Andesite | Karimizu-Kōsen |
| 92 | 3.23 | 0.89 | — | — | — | Ōi | 大 井 (東京) | Diluvium | Well (G. Aochi) |
| 93 | 3.21 | 0.88 | — | 13.0° | simple | Nekonaki | 猫 啼 (福島) | Gneiss | Tomaki-no-yu |
| 94 | 3.20 | 0.88 | — | — | earthy sulphated common salt | Natsu-abura | 夏 油 岩手 | Liparite | Natsu-abura-Kōsen A |
| 95 | 3.14 | 0.86 | — | — | alkaline carbon-dioxated concentrated common salt | Onishi | 鬼 石 群馬 | Chrystal. Schist | Yashio-Kōsen A (T. Miki) |
| 96 | 2.91 | 0.89 | — | — | — | Koto-ura | 琴 浦 岡山 | Granite | Spring A (M. Sasaki) |
| 97 | 2.82 | 0.78 | — | — | earthy common salt | Atsushio | 熱 鹽 (福島) | Alluvium | Atsushio-Kōsen No. 2 |
| 98 | 2.70 | 0.74 | — | — | — | Omi | 青 海 新潟 | " | Taumi-Kōsen |
| 99 | 2.66 | 0.73 | 300 | 18.5° | carbonated common salt | Takarazuka | 寶 塚 兵庫 | Granite | Takarazuka-Kōsen |
| 100 | 2.64 | 0.73 | 270 | 17.5° (11) | common salt | Shingu | 新 宮 石川 | " | Shingu-Kōsen No. 1 |
| 101 | 2.51 | 0.69 | — | — | — | Tomoe | 巴 茨城 | Diluvium | Momiji-Kōsen |
| 102 | 2.40 | 0.66 | 58 | 18.0° | carbonated | Notsubarū | 野 津 原 大分 | Volc. Ejectam. | Tsukano-no-yu |
| 103 | 2.26 | 0.62 | — | 17.0° | — | Katsu-ura | 勝 浦 (和歌山) | Tertiary | Benten-yu |
| 104 | 2.21 | 0.61 | — | 28.5° (10°) | mineral saline bitter | Tera machi (Tottori) | 寺 町 鳥取 | Alluvium | Kasuga-Onsen |
| 105 | 2.10 | 0.58 | — | — | — | Shinagawa | 品 川 (東京) | Diluvium | Well (in Ebura Jinja) |
| 106 | 2.00 | 0.55 | — | — | — | Tōkyō-Utsunomiya-ken | 牛 込 (東京) | " | " (K. Sasaki) |
| 107 | 1.98 | 0.54 | — | — | earthy common salt | Atsushio | 熱 鹽 (福島) | Alluvium | Atsushio-Kōsen No. 2 |
| 108 | 1.94 | 0.53 | — | — | — | Hirotsu | 廣 津 長野 | Tufec. shale | Hirotsu-Kōsen |
| 109 | 1.86 | 0.51 | — | 17.0° | — | Iriarai | 入 新 井 東京 | Alluvium | Well (C. Hirabayashi) |
| 110 | 1.86 | 0.51 | 45 | 20.5° | vitriol | Rokuyō | 鹿 野 園 奈良 | " | Nanto-rokuyō-Kōsen |
| 111 | 1.81 | 0.50 | — | 16.0° | — | Tōkyō-Shiba-ku | 芝 (東京) | Diluvium | Well (A. Kikuchi) |
| 112 | 1.76 | 0.48 | — | — | — | Haneda | 羽 田 神奈川 | Alluvium | Well (S. Motohashi) |
| 113 | 1.70 | 0.47 | — | — | alkaline | Omi-Kanagawa-ken | 大 網 神奈川 | " | Spring A (C. Hirotsu) |
| 114 | 1.68 | 0.46 | — | — | — | Enoshima | 江 ノ 島 (東京) | Tertiary | Well (Ebisuya) |
| 115 | 1.62 | 0.45 | — | 17.0° | iron carbonate solution salt | Haneda | 羽 田 東京 | Alluvium | Well (in Ebura Jinja) |
| 116 | 1.60 | 0.44 | — | — | earthy iron carbonate common salt | Daito | 大 塚 大阪 | Tertiary | Daito-Kōsen |
| 117 | 1.56 | 0.43 | — | 24.0° | simple | Okoto | 雄 琴 滋賀 | Alluvium | Spring A (S. Iwano) |
| 118 | 1.55 | 0.43 | — | — | — | Osaki | 大 崎 東京 | Diluvium | Well (H. Kamada) |
| 119 | 1.55 | 0.43 | — | — | earthy sulphated common salt | Natsu-abura | 夏 油 岩手 | Liparite | Natsu-abura-Kōsen B |
| 120 | 1.53 | 0.42 | — | — | — | Yokohama-Koyasu | 横 濱 神奈川 | Alluvium | Well (T. Miki) |
| 121 | 1.53 | 0.42 | — | 13.0 | simple | Koshakawa-land | 越木岩新田 | Granite | Myōbandani No. 2 |

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|---------------------|------------------------------|-------------|---|-----------------------|---|----------------------|---------|-------------|-----------------------|
| | in 10 c.c. | in 100 c.c. | | | | | | | |
| 122 | 1.52 | 0.42 | ca. | air temp. | — | Chōshi | 銚子千葉 | — | Well K. Watanabe |
| 123 | 1.51 | 0.42 | — | — | — | Shinagawa | 品川東京 | Alluvium | " (Y. Sakurai) |
| 124 | 1.45 | 0.40 | — | — | — | Haneda | 羽田 " | " | " A. Furutani |
| 125 | 1.45 | 0.40 | 0.3 | 21.0° | simple carbonated | Funagoya | 船小屋(福岡) | Diluvium | Nomi-yu |
| 126 | 1.40 | 0.39 | — | — | — | Futagawa | 二川愛知 | " | Well K. Matsushima |
| 127 | 1.40 | 0.39 | — | — | earthy carbonated | Sawaguchi | 澤口秋田 | " | Yunotai-Kōsen |
| 128 | 1.39 | 0.38 | — | — | — | Shinagawa | 品川東京 | " | Well (S. Ōkubo) |
| 129 | 1.35 | 0.37 | — | — | — | Tōshimo | 東下茨城 | Alluvium | " U. Nomaka |
| 130 | 1.33 | 0.37 | — | — | carbonated | Sawaguchi | 澤口秋田 | Diluvium | Ojigasawa-Kōsen No. 2 |
| 131 | 1.29 | 0.36 | — | — | — | Naguri | 名栗埼玉 | — | Spring A |
| 132 | 1.27 | 0.35 | — | — | — | Shibuya | 澁谷東京 | Diluvium | Well R. Hara |
| 133 | 1.22 | 0.34 | — | 27.0° | cool hydrogen sulphide | Shibu [Suwa] | 澁(諏訪)長野 | Andesite | Shibu-no-yu |
| 134 | 1.15 | 0.32 | — | 10.0° | — | Ikegami | 池上(東京) | Diluvium | Well S. Shiraishi |
| 135 | 1.10 | 0.30 | — | — | — | Tōkyō—Honjō-ku | 本所(" | Alluvium | " K. Fujii |
| 136 | 1.08 | 0.30 | — | — | — | Nakano | 中野茨城 | " | " K. Yabu |
| 137 | 1.04 | 0.29 | — | — | — | Unakami | 海上千葉 | — | Kakine-Kōsen |
| 138 | 1.04 | 0.29 | — | 14.5° | — | Goka | 伍賀長野 | Volc. Detr. | Toai-Kōsen A |
| 139 | 0.98 | 0.27 | — | 12.5° | — | " | " | " | " B |
| 140 | 0.95 | 0.26 | — | 10.0° | common salt ? | Narutō | 成東(千葉) | — | Narutō-Kōsen |
| 141 | 0.94 | 0.26 | — | — | — | Sawaguchi | 澤口秋田 | Diluvium | Ojigasawa-Kōsen No. 4 |
| 142 | 0.93 | 0.26 | — | 14.0° | — | Uruga | 浦賀神奈川 | — | Aramaki-Kōsen |
| 143 | 0.89 | 0.25 | — | 11.7° | — | Matsu-ida | 松井田群馬 | Tertiary | Spring A (T. Ezaka) |
| 144 | 0.85 | 0.23 | 6480 | 35.0° | simple | Shimobe | 下都山梨 | Tert. Shale | Shimobe-Onsen No. 1 |
| 145 | 0.81 | 0.22 | — | — | — | Tōkyō—Azabu-ku | 麻布東京 | Diluvium | Well (T. Yagi) |
| 146 | 0.79 | 0.22 | — | — | — | Shinagawa | 品川 " | " | " S. Ishii |
| 147 | 0.78 | 0.22 | — | 13.0° | — | Hitomi | 人見群馬 | Volc. Detr. | Spring A S. Hidaka |
| 148 | 0.77 | 0.21 | 864 | 27.0° | salt | Gakutō | 學頭鳥根 | Tertiary | Yunokawa-Kōsen |
| 149 | 0.48 | 0.13 | 6480 | 34.0° | simple | Shimobe | 下都山梨 | Tert. Shale | Shimobe-Onsen No. 2 |
| 150 | 0.47 | 0.13 | 720 | 15.5° | alkaline carbon-dioxated concentrated common salt | Isobe (Gumma-ken) | 磯部群馬 | Tertiary | Isobe-Kōsen |
| 151 | 0.47 | 0.13 | — | 20.0° | — | Taura | 田浦神奈川 | " | Ura-no-gō-Kōsen |
| 152 | 0.41 | 0.11 | — | — | alkaline common salt | Hitose (Myagi-ken) | 廣瀬宮城 | — | Nanatsu-ishi-Kōsen |
| 153 | 0.30 | 0.08 | 6480 | 36.0° | simple | Shimobe | 下都山梨 | Tert. Shale | Shimobe-Onsen No. 3 |
| 154 | 0.28 | 0.08 | — | — | — | Mobara | 茂原千葉 | Diluvium | Well K. Inoue |
| 155 | 0.27 | 0.07 | — | — | — | Ta-ura | 田浦(神奈川) | Tertiary | Yaochi-Kōsen |
| 156 | 0.25 | 0.07 | — | — | — | Tōkyō—Asakusa-ku | 浅草東京 | Alluvium | Well K. Kamigawa |
| 157 | 0.22 | 0.06 | — | 32.0° | alkaline | Higashi-nagakura | 東長倉長野 | Volc. Detr. | Aka-iwa-Kōsen |
| Gas evolving from : | | | | | | | | | |
| 1 | 5506.30 | 1514.23 | — | 16.0° | common salt | Masutomi | 増富山梨 | Granite | Kuridaira No. 1 |
| 2 | 2037.42 | 560.29 | — | 14.0° | — | " | " | " | Kamigawara No. 2 |
| 3 | 1871.31 | 514.61 | — | 15.0° | " | " | " | " | Yunokubo-gawara-no-yu |
| 4 | 1307.21 | 359.48 | — | 17.0° (12°) | carbonated (?) | Ikeda | 池田鳥根 | " | Nobata-shin-yu |
| 5 | 86.54 | 23.89 | — | 32.0° | — | Yudani | 湯谷鳥取 | Tertiary | Kabu-yu |
| 6 | 86.41 | 23.76 | — | 24.5° | — | Masutomi | 増富山梨 | Granite | Tochikubo No. 1 |
| 7 | 35.02 | 9.63 | — | — | carbonated (?) | Kawakami | 川上大阪 | " | Spring A (K. Fujiki) |
| 8 | 9.38 | 2.58 | — | — | — | Yokoyama | 横山 " | Tertiary | Yokoyama-Kōsen |
| 9 | 8.76 | 2.41 | — | 28.5° | — | Tera-machi [Tottori] | 寺町鳥取 | Alluvium | Kasuga-Onsen |
| 10 | 7.42 | 2.04 | — | 9.0° | — | Isobe (Gumma-ken) | 磯部群馬 | Tertiary | Ji-no-yu |
| 11 | 7.16 | 1.97 | — | — | — | Takarazuka | 寶塚兵庫 | Granite | Takarazuka-Kōsen |
| 12 | 1.41 | 0.39 | — | 15.5° (30°) | alkaline carbon-dioxated concentrated common salt | Isobe (Gumma-ken) | 磯部群馬 | Tertiary | Isobe-Kōsen |

* Reductivity of the gas was determined at a temperature lower than 20°C. but the correction of the volume due to temperature was not made.

Table 2.

The determination of radium emanation was carried on with the fontactoscope of C. Engler and H. Sieveking modified by Kohlrausch and Löwenthal exactly according to the authors' original instructions.

A Hot Springs.

| No. | Emanation per litre of water in Nache's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|---|---|-----------------------|--------------------|------------|----------|----------|-----------------------------|
| 1 | 121.99 | ca. | 71.0° | — | Misasa | 三朝鳥取 | Granite | Private bath (T. Matsuda) |
| 2 | 107.82 | — | 51.0° | — | " | " | " | Seitō-kwan-no-yu |
| 3 | 72.65 | — | 60.0° | — | " | " | " | Hanaya-no-yu |
| 4 | 63.47 | — | 64.0° | — | " | " | " | Sakaya-no-yu No. 1 |
| 5 | 60.35 | — | 67.0° | simple | " | " | " | " No. 2 |
| 6 | 58.48 | — | 40.0° | — | " | " | " | Private bath (K. Kishida) |
| 7 | 58.18 | — | 75.0° | — | " | " | " | Akazakiya-no-yu |
| 8 | 39.70 | — | 56.0° | — | " | " | " | Iwa-no-yu (Ono-yu) |
| 9 | 36.02 | — | 54.5° | — | " | " | " | " (Ono-yu) |
| 10 | 33.47 | — | 42.0° | sulphur | Sekigane | 關金 | " | Kabu-yu |
| 11 | 32.41 | 270 | 44.5° | " | " | " | " | Kamijaya-no-yu |
| 12 | 28.57 | — | 60.5° | — | Misasa | 三朝 (") | " | Spring B. River bed |
| 13 | 28.37 | — | 58.5° | — | " | " | " | Kiya-no-yu |
| 14 | 27.88 | — | 63.5° | — | " | " | " | Ochaya-no-yu |
| 15 | 27.87 | — | 44.5° | — | " | " | " | Shin-ya-no-yu |
| 16 | 25.15 | 1800 | 39.0° | simple | Tochiomata | 栃尾又 (新湯) | " | Tochiomata-no-yu No. 1 |
| 17 | 24.75 | — | 46.0° | — | Misasa | 三朝鳥取 | " | Uenoya-no-yu |
| 18 | 24.46 | — | 57.0° | — | " | " | " | Tōfuya-no-yu |
| 19 | 24.46 | — | 67.0° | — | " | " | " | Bun-aburaya-no-yu |
| 20 | 24.05 | 236 | 38.0° | simple | Tochiomata | 栃尾又新湯 | " | Jizai-kwan-uchi-yu No. 2 |
| 21 | 23.33 | — | 39.0° | " | " | " | " | Tochiomata-no-yu No. 2 |
| 22 | 22.07 | 90 | 42.0° | sulphur | Sekigane | 關金鳥取 | " | Inkyōjaya-no-yu |
| 23 | 20.23 | — | 72.0° | common salt | Misasa | 三朝 (") | " | Tōkyōya-no-yu |
| 24 | 19.20 | — | 48.5° | — | " | " | " | Aburaya-no-yu |
| 25 | 18.55 | — | 63.0° | — | " | " | " | Private bath (S. Mifune) |
| 26 | 15.52 | — | 63.0° | — | " | " | " | Hashizuya-no-yu |
| 27 | 13.81 | — | 71.5° | simple | " | " | " | Kyōyūchi-no-yu |
| 28 | 13.21 | — | 48.0° | earthy common salt | Yunokawa | 湯川北海道 | Liparite | Senshin-kwan-no-yu No. 2 |
| 29 | 12.85 | — | 52.0° | carbonated ? | Misasa | 三朝鳥取 | Granite | Private bath (K. Tsugawa) |
| 30 | 11.32 | 216 | 46.0° | earthy common salt | Yunokawa | 湯川北海道 | Liparite | Hōmei-kwan-no-yu |
| 31 | 11.24 | — | 40.5° | " | " | " | " | Senshin-kwan-no-yu No. 1 |
| 32 | 11.04 | — | 56.5° | neutral sulphur | Misasa | 三朝鳥取 | Granite | Naka-no-yu |
| 33 | 10.65 | 108 | 46.0° | sulphur | Sekigane | 關金 (") | " | Kon-ya-no-yu |
| 34 | 10.28 | 126 | 62.0° | common salt | Owani | 大鰐青森 | Liparite | Umeke-no-yu |
| 35 | 9.69 | 144 | 63.0° | salt | " | " | " | Hie-no-yu |
| 36 | 9.27 | 126 | 69.0° | " | " | " | " | Kagasuke-no-yu |
| 37 | 8.77 | 144 | 62.5° | " | " | " | " | Netsu-no-yu |
| 38 | 8.48 | — | 48.0° | earthy common salt | Yunokawa | 湯川北海道 | " | Shin-yu No. 1 (Rinchō-kwan) |
| 39 | 8.21 | — | 49.0° | " | " | " | " | Chōju-yu (") |
| 40 | 7.96 | 216 | 42.5° | " | " | " | " | Toyokawa-no-yu |
| 41 | 7.50 | 108 | 67.0° | common salt | Ōwani | 大鰐青森 | " | Yamabuki-no-yu |
| 42 | 7.47 | 110 | 50.0° | earthy common salt | Yunokawa | 湯川北海道 | " | Mura-yu |
| 43 | 7.35 | 216 | 50.0° | " | " | " | " | Taki-no-yu (Rinchō-kwan) |
| 44 | 7.15 | — | 53.0° | — | Kuchino | 善見鳥取 | Granite | Private bath (") |
| 45 | 7.01 | — | 56.0° | — | " | " | " | " (") |
| 46 | 6.69 | — | 61.0° | — | Misasa | 三朝 (") | " | Macjaya-no-yu |

| No. | Evaporation per litre of water in Maclure's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|---|-----------------------|----------------------------|--------------------------------|-----------|----------------------|--------------------------------------|
| 47 | 6.51 | ca. | 50.0° | earthy common salt | Yunokawa | 湯川(北海道) | Liparite | Shin-yu No. 2 (<i>Rinchō-kwan</i>) |
| 48 | 5.80 | 126 | 51.5° | " | Kachimi | 勝見(鳥取) | Granite | Sagi-no-yu |
| 49 | 5.78 | " | 70.0° | earthy common salt | Onogawa | 小野川(山形) | Tertiary | Taki-no-yu |
| 50 | 5.76 | " | 50.0° | common salt | Shiobara—Fukuwata | 鹽原—福渡(栃木) | " | Hadaka-no-yu |
| 51 | 5.54 | " | 66.0° | earthy common salt | Nezaki | 根崎(北海道) | Liparite | Rikuzō-no-yu |
| 52 | 5.35 | " | 49.0° | earthy saline bitter | Hamamura | 濱村(鳥取) | Granite | Suzukiya-no-yu |
| 53 | 5.14 | " | 54.0° | " | Tamatsukuri | 玉造(鳥根) | Tertiary | Private bath (<i>S. Hasegawa</i>) |
| 54 | 5.02 | " | 45.0° | simple | Tōgō | 東郷(鳥取) | Andesite | Yōjō kwan-no-yu No. 3 |
| 55 | 4.86 | " | 47.0° | " | Yoshikata | 吉方(") | Alluvium | Matsukawa-Onsen |
| 56 | 4.77 | " | 42.0° | simple | Yumura (Shimane-ken) | 湯村(鳥根) | Granite | Spring A |
| 57 | 4.77 | 504 | 72.0° | salt | Takase | 高瀬(新潟) | Alluvium | Takase-no-yu |
| 58 | 4.59 | " | 51.0° | simple | Kashi | 甲子(福島) | Granite | Moto-yu |
| 59 | 4.59 | " | 42.0° | saline bitter | Higashiyama | 東山(") | Andesite or Tertiary | Fudō-no-yu |
| 60 | 4.38 | " | 69.0° | common salt | Kuradate | 蔵館(青森) | Alluvium | Kami-no-yu |
| 61 | 4.37 | 544 | 46.0° | sulphated common salt | Yunotsu | 温泉津(鳥根) | Tertiary | Shin-yu |
| 62 | 4.36 | " | 50.0° | simple | Yoshioka | 吉岡(鳥取) | " | Kamei-dono-no-yu |
| 63 | 4.33 | 360 | 56.0° | common salt | Asōzu | 浅津(") | Andesite | Moto-yu |
| 64 | 4.17 | " | 46.0° | saline sulphated bitter | Iwai | 岩井(") | Tertiary | Iwami-kwan-no-yu |
| 65 | * 4.17 | " | 37.0° | " | Mimata | 美又(鳥根) | Granite | Mimata-Onsen |
| 66 | 4.15 | " | 57.5° | saline bitter | Shiobara—Ōami | 鹽原—大網(栃木) | Tertiary | Ishi-ai-no-yu |
| 67 | 4.14 | " | 56.0° | simple | Ōyu (Niigata-ken) | 大湯(新潟) | Granite | Kawagishi-no-yu No. 1 |
| 68 | 4.11 | " | 55.0° | saline bitter | Shiobara—Ōami | 鹽原—大網(栃木) | Tertiary | Kawara-no-yu |
| 69 | 4.10 | 65 | 50.0° | salt | Yuzawa [Omazawa] (Niigata-ken) | 湯澤[女川]—新潟 | Alluvium | Moto-yu |
| 70 | 4.06 | 1730 | 50.0° | simple | Tōgō | 東郷(鳥取) | Andesite | Yōjō kwan-no-yu No. 1 |
| 71 | 4.00 | 979 | 63.0° | common salt | Takanosu | 鷹之巢(新潟) | Tertiary | Takanosu-Onsen |
| 72 | 3.89 | " | 45.0° | " | Hamamura | 濱村(鳥取) | Granite | Tabakoya-no-yu |
| 73 | 3.83 | " | 56.0° | simple | Ōyu (Niigata-ken) | 大湯(新潟) | " | Kawagishi-no-yu No. 2 |
| 74 | 3.70 | " | 45.0° | " | Hamamura | 濱村(鳥取) | " | Kadoya-no-yu |
| 75 | 3.62 | " | 56.0° | simple | Ōyu (Niigata-ken) | 大湯(新潟) | " | Sakashita-no-yu |
| 76 | 3.60 | " | 48.0° | salt | Yuzawa [Omazawa] (Niigata-ken) | 湯澤[女川]— " | Alluvium | Sugai-no-yu |
| 77 | 3.59 | " | 57.0° | simple | Ōyu (Niigata-ken) | 大湯(") | Granite | Ō-yu (<i>Otoko-yu</i>) |
| 78 | 3.56 | 360 | 56.0° | muriated sulphur | Asōzu | 浅津(鳥取) | Andesite | Asahi-kwan-no-yu |
| 79 | 3.55 | " | 43.5° | earthy common salt | Nezaki | 根崎(北海道) | Liparite | Taishō-kwan-no-yu |
| 80 | 3.52 | " | 56.0° | simple | Ōyu (Niigata-ken) | 大湯(新潟) | Granite | Ō-yu (<i>Onna-yu</i>) |
| 81 | 3.45 | " | 68.0° | earthy common salt | Onogawa | 小野川(山形) | Tertiary | Suzuki-no-yu |
| 82 | 3.45 | 25 | 43.0° | muriated saline bitter | Yoshikata | 吉方(鳥取) | Alluvium | Takasago-Onsen |
| 83 | 3.43 | " | 53.0° | simple | Ōyu (Niigata-ken) | 大湯(新潟) | Granite | Yakushi-no-yu |
| 84 | 3.43 | 830 | 39.5° | carbonated | Deyu | 出湯(") | " | Chōsenkutsu-no-yu |
| 85 | 3.39 | " | 43.0° | simple | Yumura (Shimane-ken) | 湯村(鳥根) | " | Spring B |
| 86 | 3.38 | " | 50.0° | " | Kashi | 甲子(福島) | " | Yujin-no-yu |
| 87 | 3.24 | " | 70.5° | earthy common salt | Wakura | 和倉(石川) | Tertiary | Wakazaki no-yu No. 3 |
| 88 | 3.23 | 180 | 56.5° | saline bitter | Higashiyama | 東山(福島) | Andesite or Tertiary | Sō-yu |
| 89 | 3.20 | " | 43.0° | " | Hirose (Shimane-ken) | 瀬(鳥根) | Tertiary | Sagi-no-yu |
| 90 | 3.20 | 180 | 43.0° | saline bitter | Higashiyama | 東山(福島) | Andesite or Tertiary | Saru-no-yu |
| 91 | 3.20 | 3888 | 43.0° | simple | Yumura (Shimane-ken) | 湯村(鳥根) | Granite | Shitsuni-no-yu |
| 92 | 3.20 | " | 52.0° | salt | Yuzawa [Omazawa] (Niigata-ken) | 湯澤[女川]—新潟 | Alluvium | Takahashi-kwan-no-yu |
| 93 | 3.19 | " | 52.5° | simple | Yoshioka | 吉岡(鳥取) | Tertiary | Naka-no-yu |
| 94 | 3.17 | " | 53.5° | " | " | " | " | Nakajimaya-no-yu |
| 95 | 3.14 | " | 50.0° | saline bitter | Higashiyama | 東山(福島) | Andesite or Tertiary | Sugi-no-yu |
| 96 | 3.12 | 36 | 47.5° | earthy common salt | Yoshikata | 吉方(鳥取) | Alluvium | Tottori-Onsen |
| 97 | 3.07 | " | 42.5° | sulphur | Yoshioka | 吉岡(") | Tertiary | Shimo-no-yu |
| 98 | 3.02 | 210 | 45.0° | earth-muriated common salt | Yunokawa | 湯川(北海道) | Liparite | Tōyō-kwan-no-yu |
| 99 | 3.02 | 180 | 58.0° | saline bitter | Higashiyama | 東山(福島) | Andesite or Tertiary | Saza-nami-no-yu |

| No. | Emission per litre of water in Maube's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|--|---|-----------------------|----------------------------|----------------------|-----------|--|-------------------------|
| 100 | *2.93 | — | 37.0 | — | Iwai | 岩井島取 | Tertiary | Shimo-nakagawara-no-yu |
| 101 | 2.87 | — | 40.0 | saline sulphated bitter | " | " | " | Nakajimaya-no-yu |
| 102 | 2.86 | 450 | 72.5 | common salt | Kuradate | 蔵館青森 | Alluvium | Shimo-no-yu |
| 103 | 2.86 | — | 48.5 | simple | Kashii | 甲子福島 | Granite | Tengu-no-yu |
| 104 | 2.77 | — | 48.0 | common salt | Shiodara-Fukuwata | 篠原一福渡戸栃木 | Tertiary | Iwa-no-yu |
| 105 | 2.73 | — | 37.0° | salt | Yuzawa (Yamanaka) | 湯澤魚沼新潟 | Misaka Series | Kiridōshi-no-yu |
| 106 | 2.73 | — | 40.0 | — | Asōzu | 淺津島取 | Andesite | Sugimoto-no-yu |
| 107 | 2.71 | — | 60.0° | alkaline common salt | Ogawa | 小川富山 | Liparite | Ogawa-Onsen No. 5 |
| 108 | 2.62 | — | 63.0 | salt | Shimogamo | 下賀茂静岡 | Tertiary | Seitō-Onsen |
| 109 | 2.61 | — | 57.0° | alkaline common salt | Ogawa | 小川富山 | Liparite | Ogawa-Onsen No. 6 |
| 110 | 2.60 | 540 | 38.2° | non-carbonated common salt | Koyabara | 小屋原(島根) | Andesite | Koyabara-Onsen |
| 111 | 2.60 | 540 | 72.0° | common salt | Kuradate | 蔵館青森 | Alluvium | Ō-yu |
| 112 | 2.60 | — | 64.5° | sulphur | Futami | 二見富山 | Granite | Dai-ichi-no-yu |
| 113 | 2.46 | — | 49.0° | alkaline common salt | Ogawa | 小川 | Liparite | Ogawa-Onsen No. 2 |
| 114 | 2.41 | — | 38.0 | salt | Yumura | 湯村山梨 | | Meiji-Onsen |
| 115 | 2.37 | — | 61.5 | — | Tamat-sukuri | 玉造島根 | Tertiary | Tonosama-yu |
| 116 | 2.36 | — | 44.5° | common salt | Shiobara-Fukuwata | 篠原一福渡戸栃木 | " | Awa-no-yu |
| 117 | 2.29 | 4066 | 58.5° | saline sulphated bitter | Iwai | 岩井島取 | " | Moto-yu |
| 118 | 2.28 | — | 45.2° | sulphated bitter salt | Yamanaka | 山中(石川) | " | Kiku-no-yu (Onna-yu) |
| 119 | 2.23 | — | 66.5° | earth-nitrated common salt | Onogawa | 小野川(山形) | " | Ogiya-no-yu No. 2 |
| 120 | 2.21 | — | 83.0° | sulphur | Kuronagi | 黒薙富山 | Granite | Onna-yu |
| 121 | 2.20 | 180 | 49.0° | simple | Kanetsuri | 鐘釣(山形) | Contact Zone of Granite & Trondhjemite | Kanetsuri-Onsen |
| 122 | *2.19 | 4500 | 48.0° | " | Yubara (Okayama-ken) | 湯原岡山 | Granite | Yakuō-tō |
| 123 | *2.18 | 2700 | 37.5° | " | Maga | 真賀(山形) | " | Kin-yu |
| 124 | 2.17 | 152 | 76.0° | earth-nitrated common salt | Awara | 蘆原福井 | Alluvium | Funatsu No. 10 |
| 125 | 2.17 | 488 | 61.5° | bitter | Kaminoyama | 上山(山形) | Liparite | Kawashimaya-no-yu No. 2 |
| 126 | 2.11 | 346 | 49.0° | sulphated common salt | Omaki | 大牧富山 | Porphyry Dyke | Omaki-Onsen |
| 127 | 2.09 | 1953 | 64.0° | common salt | Tamatsukuri | 玉造(島根) | Tertiary | Kami-no-yu |
| 128 | 2.08 | — | 62.0° | hyrogen sulphate | Nikkō-Yumoto | 日光湯本(栃木) | Quartz Porph. | Tsuru-no-yu |
| 129 | 2.07 | — | 54.0° | simple | Nasu-Benten | 那須-辨天(山形) | Andesite | Kawa-no-yu |
| 130 | 2.06 | — | 37.0° | — | Hirose (Shimane-ken) | 廣瀬(島根) | Tertiary | Private bath (Asakura) |
| 131 | 2.01 | — | 63.0° | salt | Kuzu | 葛(長野) | Granite | Gorin-no-yu |
| 132 | 1.98 | — | 62.0° | " | " | " | " | Kanatsubo-no-yu |
| 133 | 1.97 | — | 48.0° | " | Rendaiji | 蓮臺寺静岡 | Tertiary | Idoya-no-yu |
| 134 | 1.97 | — | 88.0° | simple | Yubiso | 湯檜曾群馬 | Granite | Yubiso-no-yu No. 2 |
| 135 | *1.95 | — | — | — | Ginzan-daira | 银山平新潟 | | Koyu-no-mata-Onsen |
| 136 | 1.95 | — | 73.5° | earth-nitrated common salt | Onogawa | 小野川(山形) | Tertiary | Ama-no-yu |
| 137 | 1.95 | — | 38.5° | alkaline | Goshiki | 五色(山形) | " | Goshiki-Onsen No. 2 |
| 138 | 1.94 | — | 88.5° | sulphur | Kuronagi | 黒薙富山 | Granite | Otoko-yu |
| 139 | 1.93 | 216 | 55.0° | salt | Yugashima | 湯ヶ島静岡 | Andes. and Tuff | Seko-no-Ō-yu |
| 140 | 1.93 | — | 42.0° | earthy common salt | Aoyama | 青嶺山北海道 | Andesite | Furōkaku-no-yu No. 2 |
| 141 | 1.90 | — | 48.0° | — | Hamamura | 青嶺村島取 | Granite | Kyōdo-yu (No. 1) |
| 142 | 1.89 | 152 | 70.5° | earth-nitrated common salt | Awara | 蘆原福井 | Alluvium | Funatsu No. 21 |
| 143 | 1.89 | — | 48.0° | — | Asōzu | 淺津島取 | Andesite | Nisshin-kwan-no-yu |
| 144 | 1.87 | — | 78.0° | common salt | Kuradate | 蔵館青森 | Alluvium | Hoyōen-no-yu |
| 145 | *1.86 | — | — | — | Ginzan-daira | 银山平新潟 | | Rōhai-no-yu |
| 146 | 1.85 | — | 59.5° | sulphur | Nakabusa | 中房長野 | Granite | Shirataki-no-yu |
| 147 | 1.83 | — | 38.5° | saline common salt | Kamasaki | 鎌先宮城 | Andesite | Mogamiya-no-yu |
| 148 | 1.83 | 3944 | 62.0° | bitter | Kaminoyama | 上山(山形) | Liparite | Ilie-no-yu |
| 149 | 1.81 | — | 45.8° | sulphated bitter salt | Yamanaka | 山中(石川) | Tertiary | Kiku-no-yu (Otoko-yu) |
| 150 | 1.80 | 313 | 42.0° | salt | Yumura | 湯村山梨 | Andesite | Washi-Onsen |
| 151 | 1.78 | — | 56.0° | sulphur | Awazu | 粟津石川 | Liparite | Zetsumoto-no-yu |

| No. | Emanation per litre of water in Maeter's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|--|---|-----------------------|----------------------------|-------------------------------|------------|-----------------|-----------------------|
| 152 | 1.77 | 5.38 | 50.0° | saline sulphated bitter | Iwai | 岩井(鳥取) | Tertiary | Hanaya-no-yu |
| 153 | 1.76 | 5.41 | 65.0° | alkaline carbon-dioxated | Yumura [Hōgo-ken] | 湯村(兵庫) | Granite | Ara-yu |
| 154 | 1.75 | 15.2 | 75.0° | earth-muriated common salt | Awara | 蘆原(福井) | Alluvium | Nimen No. 28 |
| 155 | 1.73 | 270 | 66.0° | bitter | Asamushi | 浅虫(青森) | Andesite | Ō-yu |
| 156 | 1.73 | — | 64.0° | hydrogen sulphide | Nikko-Yumoto | 日光湯本(栃木) | Quartz Porph. | Kawara-no-yu |
| 157 | 1.70 | 3944 | 61.0° | bitter | Kaminoyama | 上ノ山(山形) | Liparite | Atatamari-no-yu |
| 158 | 1.69 | 15.2 | 68.2° | earth-muriated common salt | Awara | 蘆原(福井) | Alluvium | Tanaka-naka No. 6 |
| 159 | 1.64 | — | 46.5° | simple | Togo | 東郷(鳥取) | Andesite | Yojo-kwan-no-yu No. 6 |
| 160 | 1.62 | — | 45.0° | salt | Yuzawa [Uonuma] (Niigata-ken) | 湯澤[魚沼](新潟) | Misaka Series | Moto-yu |
| 161 | 1.62 | — | 75.5° | earth-muriated common salt | Wakura | 和倉(石川) | Tertiary | Sō-yu |
| 162 | 1.61 | 700 | 50.0° | sulphated common salt | Yunotsu | 温泉津(島根) | Tertiary | Kyū-yu |
| 163 | 1.61 | 2700 | 47.5° | simple | Shimo-Suwa | 下諏訪(長野) | Alluvium | Wata-no-yu |
| 164 | 1.59 | 15.2 | 70.0° | earth-muriated common salt | Awara | 蘆原(福井) | " | Funatsu No. 9 |
| 165 | 1.58 | — | 66.0° | hydrogen sulphide | Nikko-Yumoto | 日光湯本(栃木) | Quartz Porph. | Gosho-no-yu |
| 166 | 1.52 | — | 57.0° | muriated alkaline | Shiobara - Hataori | 鹽原一畑下(") | Tertiary | Hato-no-yu |
| 167 | 1.51 | — | 62.0° | sulphated common salt | Shibu [Hirao] | 澁[平穩](長野) | Diluvium | Ō-yu Tent. No. 3 |
| 168 | 1.51 | 356 | 42.0° | simple | Yamabe | 山邊(") | Tertiary | Shiraito-Onsen No. 2 |
| 169 | 1.51 | — | 44.5° | " | Arifuku | 有福(島根) | Diorite | Shin-yu |
| 170 | 1.50 | — | 63.5° | muriated alkaline | Shiobara - Hataori | 鹽原一畑下(栃木) | Tertiary | Mujina-no-yu |
| 171 | 1.50 | — | 94.0° | alkaline sulphur | Nakabusa | 中房(長野) | Granite | Tamura-no-yu |
| 172 | 1.46 | 448 | 50.0° | saline sulphated bitter | Iwai | 岩井(鳥取) | Tertiary | Bizenya-no-yu |
| 173 | 1.46 | — | 72.0° | sulphated common salt | Shibu [Hirao] | 澁[平穩](長野) | Diluvium | Ō-yu Tent. No. 4 |
| 174 | 1.46 | 519 | 73.5° | " | " | " | " | Hatsu-yu |
| 175 | 1.45 | 908 | 58.0° | earth-muriated common salt | Akayu | 赤湯(山形) | Tertiary | Ō-yu |
| 176 | 1.45 | — | 45.0° | salt | Yuzawa [Uonuma] (Niigata-ken) | 湯澤[魚沼](新潟) | Misaka Series | Kami-no-yu |
| 177 | 1.45 | — | 57.0° | " | Kuzu | 葛(長野) | Granite | Shin-yu No. 2 |
| 178 | 1.42 | — | 50.0° | common salt | Shiobara - Fukuwata | 鹽原一福渡戸(栃木) | Tertiary | Hiie-no-yu |
| 179 | 1.42 | — | 88.0° | sulphur (?) | Kuzu | 葛(長野) | Granite | Spring A |
| 180 | 1.40 | — | 76.0° | sulphated common salt | Shibu [Hirao] | 澁[平穩](") | Diluvium | Ō-yu Tent. No. 12 |
| 181 | 1.39 | 277.2 | 44.0° | earth common salt | Aoyama | 青山(北海道) | Andesite | Furokaku-no-yu No. 1 |
| 182 | 1.39 | — | 62.0° | common salt | Ikariga-seki | 碓ヶ関(青森) | Volc. Ash | Bun-kwan-no-yu |
| 183 | 1.38 | 130 | 42.0° | simple | Yamabe | 山邊(長野) | Tertiary | Shiraito-Onsen No. 1 |
| 184 | 1.38 | — | 85.8° | sulphur | Futami | 二見(富山) | Granite | Dai-ni-no-yu |
| 185 | 1.38 | — | 43.0° | sulphated bitter sulphur | Yamanaka | 山中(石川) | Tertiary | Mitaniya-no-yu |
| 186 | 1.38 | 179 | 49.0° | earth-muriated common salt | Akayu | 赤湯(山形) | " | Ama-yu |
| 187 | 1.35 | — | 59.5° | saline common salt | Shuzenji | 修善寺(静岡) | Andes. and Tuff | Ishi-yu |
| 188 | 1.33 | 207 | 50.0° | earth-muriated common salt | Akayu | 赤湯(山形) | Tertiary | Tamba-yu |
| 189 | 1.33 | — | 76.0° | muriated sulphated bitter | Yudanaka | 湯田中(長野) | Misaka Series | Ō-yu Tent. No. 1 |
| 190 | 1.32 | 3009 | 67.0° | simple | Shimo-Suwa | 下諏訪(") | Alluvium | Tankwa-no-yu |
| 191 | 1.32 | 97.2 | 81.0° | " | Yubiso | 湯檜曾(群馬) | Granite | Yubiso-no-yu No. 1 |
| 192 | 1.32 | — | 57.0° | " | Yubara | 湯原(") | " | Fujiya-no-yu |
| 193 | 1.31 | 173 | 64.0° | salt | Yagashima | 湯ヶ島(静岡) | Andes. and Tuff | Ochiai-rō-no-yu |
| 194 | 1.30 | — | 42.0° | " | Yuzawa [Uonuma] (Niigata-ken) | 湯澤[魚沼](新潟) | Misaka Series | Chubu-no-yu |
| 195 | 1.29 | 130 | 67.5° | simple | Kami-Suwa | 上諏訪(長野) | Alluvium | Shōjin-yu |
| 196 | 1.28 | — | 74.0° | alkaline sulphur | Nakabusa | 中房(") | Granite | Oyogi-no-yu |
| 197 | 1.28 | — | 57.0° | sulphur | Awazu | 粟津(石川) | Liparite | Kametani-no-yu |
| 198 | 1.28 | 778 | 53.5° | simple | Kamikochi | 上高地(長野) | Granite | Kamikochi-Onsen |
| 199 | 1.27 | 130 | 56.0° | carbonated | Yagashima | 湯ヶ島(静岡) | Andes. and Tuff | Yumoto-kwan-no-yu |
| 200 | 1.27 | 520 | 83.5° | common salt | Shima | 四万(群馬) | Misaka Series | Iwane-no-yu |
| 201 | 1.27 | 630 | 61.0° | " | Ikariga-seki | 碓ヶ関(青森) | Volc. Ash | Taki-no-yu |
| 202 | 1.27 | — | 74.0° | muriated sulphated bitter | Yudanaka | 湯田中(長野) | Misaka Series | Ō-yu Tent. No. 2 |
| 203 | 1.27 | — | 53.0° | salt | Rendaiji | 蓮臺寺(静岡) | Tertiary | Junji-kwan-no-yu |

| No. | Emission per litre of water in Mach's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|---|---|-----------------------|----------------------------|-------------------------------|--------------|-------------------|--------------------------|
| 204 | 1.26 | ca. | 79.0° | common salt | Katayamazu | 片山津石川 | Tertiary | Niiho-no-Tsubo |
| 205 | 1.25 | 778 | 44.5° | alkaline | Goshiki | 五色山形 | " | Goshiki-Onsen No. 1 |
| 206 | 1.24 | — | 44.0° | simple | Asama | 淺間長野 | " | Kizu-no-yu |
| 207 | 1.23 | — | 41.5° | salt | Yuzawa [Yuzawa] (Niigata-ken) | 湯澤魚沼新潟 | Miyako Series | Tamago-no-yu |
| 208 | 1.23 | — | 43.0° | " | " | " | " | Shin-yu |
| 209 | 1.22 | 304 | 74.5° | — | Yudanaka | 湯田中長野 | " | Washi-no-yu |
| 210 | 1.21 | — | 41.0° | simple | Innai-Yunosawa | 院内湯澤秋田 | Tertiary | Taki-no-yu |
| 211 | 1.20 | — | 60.0° | sulphate common salt | Shibu [Hirao] | 澁 [平穩] 長野 | Diluvium | Wata-no-yu |
| 212 | 1.20 | 558 | 61.0° | common salt | Ikariga-seki | 碓ヶ関青森 | Volc. Ash | Netsu-no-yu |
| 213 | 1.15 | 749 | 60.0° | same sulphated salt | Iwai | 岩井鳥取 | Tertiary | Iwaiya-no-yu |
| 214 | 1.15 | 108 | 67.5° | simple | Dai | 臺岩手 | Diluvium | Tsuru-no-yu |
| 215 | 1.15 | — | 54.0° | " | Nasu-Kita | 那須北栃木 | Andesite | Ai-no-yu |
| 216 | 1.15 | — | 63.0° | sulphated bitter | Asamushi | 浅虫青森 | " | Nambuya-shin-yu |
| 217 | 1.14 | — | 96.0° | alkaline sulphur | Nakabusa | 中房長野 | Granite | Myōken-no-yu |
| 218 | 1.13 | — | 56.0° | — | Yugashima | 湯ヶ島静岡 | And. s. and Tuff | Spring No. 2 (N. Adachi) |
| 219 | 1.13 | — | 79.0° | sulphated bitter | Asamushi | 浅虫青森 | Andesite | Tsukaki-no-yu No. 1 |
| 220 | 1.13 | 130 | 83.0° | simple | Kami Suwa | 上諏訪長野 | Alluvium | Tajiku-no-yu |
| 221 | 1.11 | — | 74.0° | sulphated common salt | Shibu [Hirao] | 澁 [平穩] (") | Diluvium | O-yu No. 1 |
| 222 | 1.11 | — | 50.5° | alkaline common salt | Ogawa | 小川富山 | Liparite | Ogawa-Onsen No. 4 |
| 223 | 1.11 | — | 69.0° | common salt | Shiobara-Fukuwata | 鹽原一福渡戸栃木 | Tertiary | Iwa-no-yu |
| 224 | 1.11 | — | 79.0° | " | Katayamazu | 片山津石川 | " | Shibayama-no-Tsubo |
| 225 | 1.11 | — | 95.0° | sulphur | Futami | 二見富山 | Granite | Dai-san-no-yu |
| 226 | 1.09 | — | 55.0° | — | Yugashima | 湯ヶ島静岡 | Andes. and Tuff | Yumoto-kwan-Taki-no-yu |
| 227 | 1.08 | 519 | 55.0° | saline common salt | Shuzenji | 修善寺 (") | " | Chigo-no-yu |
| 228 | 1.07 | — | 48.3° | — | Shiobara-Furumachi | 鹽原一古町栃木 | Tertiary | Asahi-no-yu |
| 229 | 1.06 | 31100 | 46.5° | common salt | Shigaku | 志學島根 | Andesite | Shigaku-Onsen |
| 230 | 1.06 | — | 92.0° | alkaline sulphur | Nakabusa | 中房長野 | Granite | Yakushi-no-yu No. 1 |
| 231 | 1.06 | 24024 | 40.5° | simple | Innai-Yunosawa | 院内湯澤秋田 | Tertiary | Yunosawa-Onsen |
| 232 | 1.06 | — | 58.0° | sulphur | Awazu | 粟津石川 | Liparite | Kamiya-no-yu |
| 233 | *1.04 | 864 | 62.0° | salt | Kaigake | 貝掛新潟 | — | Kaigake-Onsen |
| 234 | 1.04 | — | 62.0° | earthy alkaline sulphur | Akakura | 赤倉 (") | Andesite | Akakura-Onsen J. 1 No. 2 |
| 235 | 1.03 | — | 50.0° | common salt | Shiobara-Monzen | 鹽原一門前(栃木) | Tertiary | Jirakubō-no-yu |
| 236 | 1.01 | — | 40.5° | sulphur | Naruko | 鳴和子宮城 | Andesite | Taki-no-yu No. 1 |
| 237 | 1.01 | — | 80.0° | earth-muriated common salt | Wakura | 和倉石川 | Tertiary | Wakura-aki-no-yu No. 2 |
| 238 | 1.00 | — | 72.0° | sulphated bitter | Asamushi | 浅虫青森 | Andesite | Udo-no-yu |
| 239 | 1.00 | 1296 | 38.0° | earth-muriated common salt | Yunogō | 湯郷岡山 | Liparite | Sagi-no-yu |
| 240 | 1.00 | — | 65.0° | saline common salt | Shuzenji | 修善寺静岡 | Andes. and Tuff | Taki-no-yu |
| 241 | 1.00 | — | 49.0° | — | Yugashima | 湯ヶ島 (") | " | Spring No. 3 (N. Adachi) |
| 242 | 0.98 | 630 | 58.0° | common salt | Oyu Akita-ken | 大湯秋田 | Alluv. Pum. Layer | Kami-no-yu |
| 243 | 0.98 | — | 93.0° | alkaline sulphur | Nakabusa | 中房(長野) | Granite | Yakushi-no-yu No. 2 |
| 244 | 0.97 | 126 | 51.0° | earthy alkaline | Shirahone | 白骨 (") | Chichibu Series | Wata-no-yu |
| 245 | 0.97 | 648 | 56.0° | simple | Andai | 安代 (") | Diluvium | Ō-yu |
| 246 | 0.96 | — | 79.0° | common salt | Katayamazu | 片山津石川 | Tertiary | Shiotsu-no-Tsubo |
| 247 | 0.95 | 102 | 78.0° | acid | Dai | 臺岩手 | Diluvium | Matsu-no-yu |
| 248 | 0.95 | 900 | 90.5° | alkaline carbon- | Yumura (Hyōgo-ken) | 湯村兵庫 | Granite | Kō-yu |
| 249 | 0.94 | 152 | 71.0° | earth-muriated common salt | Awara | 魚原福井 | Alluvium | Funatsu No. 6 |
| 250 | 0.94 | — | 60.0° | sulphur | Shibu [Hirao] | 澁 [平穩] 長野 | Diluvium | Nakakuri-no-yu No. 1 |
| 251 | 0.94 | 130 | 59.0° | simple | Kami Suwa | 上諏訪 (") | Alluvium | Hira-no-yu |
| 252 | 0.93 | 6012 | 61.0° | sulphur | Oshiki | 大澁秋田 | Alluv. Tal. Dep. | Tsuru-no-yu |
| 253 | 0.93 | — | 50.0° | simple | Asama | 淺間長野 | Tertiary | Taki-no-yu |
| 254 | 0.92 | 1800 | 52.0° | " | Shimo-Suwa | 下諏訪 (") | Alluv. | Ko-yu |
| 255 | 0.92 | — | 95.5° | common salt | Ogawa | 小野川山形 | Tertiary | Ōgiya-no-yu No. 1 |
| 256 | 0.92 | 432 | 58.5° | simple | Yuno | 湯野福島 | " | Anabara-no-yu |

| No. | Quantity per litre of water in March, 1911 | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|--|---|-----------------------|-------------------------------|---------------------|------------|-------------------|---------------------------------------|
| 257 | 0.92 | 378 | 91.0° | common salt | Jōzankei | 定山溪(北海道) | Liparite | Moto-no-yu |
| 258 | 0.92 | — | 55.0 | " | Ikariga-seki | 碓ヶ関(青森) | Volc. Ash | Fuji-no-yu |
| 259 | 0.91 | — | 45.0 | sulphated common salt | Shibu [Hirao] | 澁[平穩](長野) | Diluvium | Mujina-no-yu |
| 260 | 0.91 | — | 62.5° | simple | Shiobara—Samaki | 鹽原—街巻(栃木) | Tertiary | Taki-no-yu |
| 261 | 0.90 | — | 37.0 | saline mineral common salt | Yunogō | 湯郷(岡山) | Liparite | Sagi-no-yu (<i>Onna-yu</i>) |
| 262 | 0.89 | — | 51.5 | simple | Yubara (Gumma-ken) | 湯原(群馬) | Granite | Ubukata-no-yu |
| 263 | 0.89 | 414 | 69.5° | common salt | Oyu Akita-ken | 大湯秋田 | Alluv. Pum. Layer | Shimo-no-yu |
| 264 | 0.89 | — | 79.0° | salt | Shimogamo | 下賀茂(静岡) | Tertiary | Hizume-Kyōdō-Onsen |
| 265 | 0.88 | — | 69.0° | alkaline sulphate | Nikkō-Yumoto | 日光湯本(栃木) | Quartz Proph. | Ara-yu |
| 266 | 0.87 | 311 | 55.0 | salt | Shima | 四萬(群馬) | Misaka Series | Shinkoku-tō |
| 267 | 0.87 | 3150 | 48.0° | saline common salt | Kamasaki | 鎌先(宮城) | Andesite | Ichijō-no-yu |
| 268 | 0.87 | — | 71.5 | common salt | Shiobara—Shio-no-yu | 鹽原—鹽之湯(栃木) | Tertiary | Naka-no-yu |
| 269 | 0.86 | — | 48.0° | earthy alkaline | Shirahone | 白骨(長野) | Chichibu Series | Chi-no-yu |
| 270 | 0.86 | 270 | 45.5° | hydrogen sulphide common salt | Ōyuzawa | 大湯澤(秋田) | Diluvium | <i>No. 35 Ō-yu.</i> |
| 271 | 0.85 | 540 | 37.7° | saline mineral common salt | Yunogō | 湯郷(岡山) | Liparite | Shin-Sagi-no-yu |
| 272 | 0.85 | — | 56.0° | sulphated bitter | Tohi | 土肥(静岡) | Tertiary | Tohi-kwan-no-yu <i>No. 2</i> |
| 273 | 0.84 | 194 | 55.0° | simple | Andai | 安代(長野) | Diluvium | Kaikwa-no-yu |
| 274 | 0.84 | 501 | 55.0 | salt | Kanabayashi | 上林(„) | Misaka Series | Tsuru-no-yu |
| 275 | 0.82 | 340 | 40.5° | simple | Shin-Goshiki | 新五色(山形) | Tertiary | Shin-goshiki-Onsen |
| 276 | 0.82 | 735 | 65.0° | common salt | Kakuma | 角間(長野) | Diluvium | Ō-yu |
| 277 | 0.82 | 524 | 74.0° | salt | Shima | 四萬(群馬) | Misaka Series | Shio-no-yu (<i>Seizen-kwan</i>) |
| 278 | 0.82 | 510 | 58.0° | sulphated common salt | Shibu [Hirao] | 澁[平穩](長野) | Diluvium | Mearai-no-yu |
| 279 | 0.82 | 635 | 62.0° | " | Shima | 四萬(群馬) | Misaka Series | Tokiwa-no-yu |
| 280 | 0.81 | 619 | 46.0° | common salt | Oyu Akita-ken | 大湯秋田 | Alluv. Pum. Layer | Arase-no-yu |
| 281 | 0.81 | 130 | 49.0° | salt | Yugashima | 湯ヶ島(静岡) | Andes. and Tuff | Seko-no-Moto-yu |
| 282 | 0.81 | — | 70.5° | sulphated common salt | Shibu [Hirao] | 澁[平穩](長野) | Diluvium | <i>O-yu Vent. No. 5</i> |
| 283 | 0.80 | — | 43.0° | saline common salt | Kamasaki | 鎌先(宮城) | Andesite | Kimuraya-no-yu |
| 284 | 0.80 | — | 52.0° | salt | Yatsu | 谷津(静岡) | Tertiary | Ishida-Shin-Onsen |
| 285 | 0.79 | — | 54.0° | common salt | Shiobara—Monzen | 鹽原—門前(栃木) | " | Kawara-no-yu |
| 286 | 0.79 | — | 96.0° | alkaline sulphur | Nakabusa | 中房(長野) | Granite | Danjō-no-yu |
| 287 | 0.78 | 195 | 42.0° | saline mineral common salt | Akayu | 赤湯(山形) | Tertiary | Mori-no-yu |
| 288 | 0.77 | 810 | 80.0° | simple | Jōzankei | 定山溪(北海道) | Liparite | Naka-no-yu |
| 289 | 0.77 | 300 | 61.5° | sulphated bitter | Asamushi | 浅虫(青森) | Andesite | Hadaka-no-yu |
| 290 | 0.77 | — | 56.0° | — | Yugashima | 湯ヶ島(静岡) | Andes. and Tuff | Spring <i>No. 1 & 5 K. Adachi</i> |
| 291 | 0.77 | — | 95.5° | alkaline sulphur | Nakabusa | 中房(長野) | Granite | Taki-no-yu <i>No. 1</i> |
| 292 | 0.77 | — | 42.0° | " | Shiobara—Furumachi | 鹽原—古町(栃木) | Tertiary | Takara-no-yu |
| 293 | 0.76 | — | 63.0° | saline mineral salt | Shuzenji | 修善寺(静岡) | Andes. and Tuff | Asabarō-no-yu |
| 294 | 0.75 | — | 67.0° | saline bitter | Yujiku | 湯宿(群馬) | Tertiary | Sen-tō |
| 295 | 0.75 | — | 50.0° | salt | Yatsu | 谷津(静岡) | " | Yatsu-Kyōdō-Onsen |
| 296 | 0.74 | — | 76.0° | " | Ōwari | 大鰐(青森) | Liparite | Kawara-no-yu |
| 297 | 0.74 | — | 55.5° | sulphated bitter | Tohi | 土肥(静岡) | Tertiary | Asaka-no-yu |
| 298 | 0.74 | — | 42.0° | sulphur | Tsubame | 燕土肥(新潟) | Andesite | Tsubame-Onsen <i>J. 1 No. 3</i> |
| 299 | 0.73 | — | 55.0° | sulphated bitter | Tohi | 土肥(静岡) | Tertiary | Hama-no-Kyōdō-yu |
| 300 | 0.73 | — | 53.0° | simple | Asama | 浅間(長野) | " | Matsu-no-yu |
| 301 | 0.73 | — | 74.0° | hydrogen sulphide | Nasu Yumoto | 那須湯本(栃木) | Andesite | Taki-no-yu |
| 302 | 0.72 | — | 51.0 | simple | Osawa | 大澤(岩手) | Tertiary | Ō-yu <i>Vent. No. 1</i> |
| 303 | 0.72 | — | 42.5° | hydrogen sulphide | Nikkō-Yumoto | 日光湯本(栃木) | Quartz Porph. | Donsu-no-yu |
| 304 | 0.72 | — | 51.5° | alkaline | Shiobara—Furumachi | 鹽原—古町(„) | Tertiary | Gosho-no-yu |
| 305 | 0.72 | — | 43.0° | salt | Kendōji | 蓮臺寺(静岡) | " | Aizu-kwan-no-yu |
| 306 | 0.71 | — | 62.0° | bitter | Ōtaki | 大滝(秋田) | Alluv. Tal. Dep. | Susuki-no-yu |
| 307 | 0.71 | 144 | 50.0° | simple | Iizaka | 飯坂(福島) | Tertiary | Tennōji-yu |
| 308 | 0.70 | — | 49.0° | " | Yugashima | 湯ヶ島(静岡) | Andes. and Tuff | Spring <i>No. 2</i> |
| 309 | 0.69 | — | 66.0° | sulphated bitter | Tohi | 土肥(„) | Tertiary | Private bath (<i>T. Terizawa</i>) |

| No. | Emanation per
litre of water
in Mahe's units | Flow of water
in 24 hrs.
in hectolitres | Temp. of spring
in C. | Classification | Location | 地名府縣名 | Geology | Spring |
|-----|--|---|--------------------------|-----------------------------|-----------------|-----------|-------------------|-------------------------------------|
| 310 | 0.68 | 180 | 94.0° | common salt | Noboribetsu | 登別 北海道 | Tertiary | Kami-no-yu |
| 311 | 0.68 | 180 | 59.0° | simple | Tizaka | 飯坂 福島 | " | Taki-no-yu |
| 312 | 0.67 | — | 46.0° | sulphated bitter | Ikao | 伊香保 群馬 | Andesite | Nomi-yu |
| 313 | 0.67 | — | 56.5° | simple | Iizaka | 飯坂 福島 | Tertiary | Spring <i>Senat. F. D. Army</i> |
| 314 | 0.66 | 150 | 60.0° | common salt | Shuzenji | 修善寺 靜岡 | Andes. and Tuff | Ume-no-yu |
| 315 | 0.66 | 634 | 70.5° | common salt | Oyu (Akita-ken) | 大湯 秋田 | Alluv. Pum. Layer | Kawara-no-yu |
| 316 | 0.65 | — | 71.0° | saline bitter
sulphur | Yamashiro | 山代 石川 | Tertiary | Araya-no-yu |
| 317 | 0.64 | — | 44.0° | simple | Arifuku | 有戸 福島 | Diorite | Arifuku-Onsen <i>Unit No. 2</i> |
| 318 | 0.64 | 380 | 76.0° | bitter | Shido-daira | 志戸平 岩手 | Tertiary | Warabi-no-yu |
| 319 | 0.64 | 635 | 84.0° | salt | Shima | 志茂 群馬 | Misaka Series | Ryūgū-no-yu |
| 320 | 0.64 | 432 | 69.0° | common salt | Shuzenji | 修善寺 靜岡 | Andes. and Tuff | Dokko-no-yu |
| 321 | 0.64 | 327 | 64.0° | " | " | " | " | Hako-yu |
| 322 | 0.64 | — | 46.0° | sulphated bitter | Tohi | 上肥 靜岡 | Tertiary | O-yu <i>Vent No. 1</i> |
| 323 | 0.63 | — | 61.0° | common salt | Katayamazu | 片山津 石川 | " | Higashi no Tsubo |
| 324 | 0.63 | 300 | 54.0° | " | Ikariga saki | 碓ヶ関 青森 | Volc. Ash | Hie-no-yu (<i>Onna-yu</i>) |
| 325 | 0.57 | 360 | 50.0° | saline bitter | Yujiku | 湯宿 群馬 | Tertiary | Ō-daki |
| 326 | 0.57 | 210 | 58.0° | simple | Karurusu | カラス 北海道 | " | Tsuru-no-yu <i>Unit No. 2</i> |
| 327 | 0.57 | 236 | 52.0° | earthy alkaline | Shirahone | 白根 長野 | Chichibu Series | Senki-no-yu |
| 328 | 0.56 | — | 67.5° | salt | Shin-kuruma | 新車 宮城 | Andesite | Moto-yu |
| 329 | 0.56 | — | 43.0° | acid hydrogen
sulphate | Nasu-Yumoto | 那須一湯本 栃木 | " | Komatsu-no-yu |
| 330 | 0.55 | 648 | 79.0° | saline bitter | Yujiku | 湯宿 群馬 | Tertiary | Kubo-yu |
| 331 | 0.54 | — | 62.0° | earthy alkaline
sulphate | Akakura | 赤倉 新潟 | Andesite | Akakura-Onsen <i>Jet No. 3</i> |
| 332 | 0.54 | 1530 | 42.0° | earthy common salt | Miyagawa | 宮川 北海道 | " | Miyagawa-no-yu |
| 333 | 0.54 | — | 54.0° | salt | Shin-kuruma | 新車 宮城 | " | Takashige-no-yu |
| 334 | 0.53 | — | 71.0° | " | Shimogamo | 下賀茂 靜岡 | Tertiary | Kano-Kyōdo-yu |
| 335 | 0.53 | 72 | 54.0° | simple | Dai | 臺 岩手 | Diluvium | Suzume-no-yu |
| 336 | 0.53 | — | 42.5° | carbonated | Kawatabi | 川渡 宮城 | Tertiary | Suzukino-yu |
| 337 | 0.52 | — | 48.0° | salt | Kōchi | 河内 靜岡 | Andesite | Shimo-no-yu |
| 338 | 0.52 | — | 70.0° | simple | Iizaka | 飯坂 福島 | Tertiary | Shin-Sabako-no-yu |
| 339 | 0.51 | — | 44.0° | " | Nagaoaka | 長岡 靜岡 | " | Tamon-yu <i>No. 4</i> |
| 340 | 0.51 | — | 45.0° | sulphated bitter | Ikao | 伊香保 群馬 | Andesite | Fukiage-no-yu |
| 341 | 0.50 | 108 | 56.5° | simple | Iizaka | 飯坂 福島 | Tertiary | Hako-yu |
| 342 | 0.50 | — | 53.0° | salt | Rendaiji | 蓮臺寺 靜岡 | " | Private bath (<i>C. Maruyama</i>) |
| 343 | 0.50 | 150 | 60.0° | common salt | Shuzenji | 修善寺 (") | Andes. and Tuff | Sugi-no-yu |
| 344 | 0.50 | — | 70.0° | " | " | " | " | Kikuya-no-yu <i>No. 2</i> |
| 345 | 0.49 | 270 | 53.0° | simple | Onokobe | 鬼首 宮城 | Tertiary | Todoroki-no-yu |
| 346 | 0.48 | 194 | 48.0° | sulphated bitter | Tohi | 上肥 靜岡 | " | Ana-yu |
| 347 | 0.48 | — | 89.0° | " | Kanegawa | 龜川 大分 | Andesite | Noda-no-yu |
| 348 | 0.48 | — | 39.0° | simple | Hatare | 畑毛 靜岡 | Volc. Detr. | Hatige-Onsen <i>No. 3</i> |
| 349 | 0.48 | — | 50.0° | " | Ito | 伊東 靜岡 | Quaternary | Private bath (<i>S. Kumagi</i>) |
| 350 | 0.47 | 360 | 63.0° | sulphated bitter | Asamushi | 淺虫 青森 | Andesite | Yanagi-no-yu |
| 351 | 0.47 | — | 48.5° | " | Ito | 伊東 靜岡 | Quaternary | Private bath (<i>H. Hirose</i>) |
| 352 | 0.47 | — | 46.0° | salt | Yatsu | 谷津 靜岡 | Tertiary | Ishida-Onsen |
| 353 | 0.47 | — | 59.0° | simple | Karurusu | カラス 北海道 | " | Matsu-no-yu |
| 354 | 0.47 | — | 52.0° | sulphated bitter | Tohi | 上肥 靜岡 | " | Private bath (<i>S. Asaka</i>) |
| 355 | 0.47 | — | 57.0° | " | " | " | " | O-yu <i>Vent No. 2</i> |
| 356 | 0.47 | — | 47.0° | " | Ikao | 伊香保 群馬 | Andesite | Ohaguro-no-yu |
| 357 | 0.46 | — | 50.0° | salt | Itō | 伊豆 靜岡 | Quaternary | Kusumi-sō-yu (<i>Onna-yu</i>) |
| 358 | 0.46 | 140 | 66.0° | saline common salt | Shuzenji | 修善寺 (") | Andes. and Tuff | Meiji-no-yu |
| 359 | 0.46 | 630 | 76.0° | common salt | Noboribetsu | 登別 北海道 | Tertiary | Shimo-no-yu |
| 360 | 0.45 | — | 47.0° | simple | Itō | 伊東 靜岡 | Quaternary | Shishido-Onsen |
| 361 | 0.45 | — | 48.0° | " | Kanegawa | 龜川 大分 | Andesite | Gomusō-no-yu |
| 362 | 0.44 | — | 50.0° | salt | Rendaiji | 蓮臺寺 靜岡 | Tertiary | Shimo-fujiwara-no-yu |
| 363 | 0.44 | — | 40.0° | simple | Hatare | 畑毛 靜岡 | Volc. Detr. | Hatare-Onsen <i>No. 2</i> |

| No. | Emanation per litre of water in Maché's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|---|-----------------------|------------------------------------|--------------|-------------|-----------------|--------------------------------------|
| 364 | 0.44 | — | 41.0° | simple | Hatage | 畑毛 (静岡) | Volc. Detr. | Hatage-Onsen No. 3 |
| 365 | 0.44 | — | 42.0° | salt | Rendaiji | 蓮臺寺 („) | Tertiary | Shin-yoshirō-no-yu |
| 366 | 0.43 | — | 51.0° | simple | Ōsawa | 大澤 (岩手) | „ | Ō-yu Vent No. 2 |
| 367 | 0.43 | 247.3 | 53.0° | salt | Shima | 四萬 (群馬) | Misaka Series | Hinatami-no-yu |
| 368 | 0.43 | — | 53.0° | „ | Rendaiji | 蓮臺寺 (静岡) | Tertiary | Kami-fujiwara-no-yu |
| 369 | 0.43 | — | 47.0° | „ | „ | „ | „ | Private bath (<i>E. Ishibashi</i>) |
| 370 | 0.43 | — | 50.0° | acid hydrogen sulphide common salt | Ogawa | 小川 (富山) | Liparite | Ogawa-Onsen No. 1 |
| 371 | 0.43 | 126 | 58.0° | Acid | Dai | 臺 (岩手) | Diluvium | Ō-yu |
| 372 | 0.42 | — | 39.0° | salt | Kōchi | 河内 (静岡) | Andesite | Kami-no-yu |
| 373 | 0.42 | — | 65.0° | „ | Yatsu | 谷津 („) | Tertiary | Misawa-no-yu |
| 374 | 0.42 | — | 71.0° | simple | Na-u—Daimaru | 那須一大丸 (栃木) | Andesite | Sakura-no-yu |
| 375 | 0.41 | — | 74.5° | acid hydrogen sulphide vitriol | „ —Yumoto | „ —湯本 („) | „ | Shika-no-yu |
| 376 | 0.41 | — | 40.0° | simple | Hatage | 畑毛 (静岡) | Volc. Detr. | Hatage-Onsen No. 7 |
| 377 | 0.40 | — | 51.5° | sulphated bitter | Tohi | 土肥 („) | Tertiary | Private bath (<i>K. Hasegawa</i>) |
| 378 | 0.40 | — | 37.5° | salt | Itō | 伊東 („) | Quaternary | Wada-Onsen |
| 379 | 0.40 | — | 53.5° | sulphated bitter | Tohi | 土肥 („) | Tertiary | Tohi-kwan-no-yu No. 1 |
| 380 | 0.39 | — | 39.0° | simple | Hatage | 畑毛 („) | Volc. Detr. | Hatage-Onsen No. 8 |
| 381 | 0.39 | 432 | 57.0° | „ | Yuno | 湯野 (福島) | Tertiary | Kiri-no-yu |
| 382 | 0.39 | 540 | 50.0° | „ | Noboribetsu | 登別 (北海道) | „ | Manju-no-yu No. 1 |
| 383 | 0.38 | 130 | 55.0° | carbonated | Yugashina | 湯ヶ島 (静岡) | Andes. and Tuff | Kidachiya-no-yu |
| 384 | 0.38 | 2160 | 56.0° | simple | Yuno | 湯野 (福島) | Tertiary | Hashimoto-no-yu |
| 385 | 0.38 | 216 | 60.0° | „ | Karurusu | カルルス (北海道) | „ | Tsuru-no-yu Vent No. 1 |
| 386 | 0.38 | — | 41.0° | „ | Nagaoka | 長岡 (静岡) | „ | Private bath (<i>S. Matsumoto</i>) |
| 387 | 0.37 | 108 | 52.0° | „ | Karurusu | カルルス (北海道) | „ | Kame-no-yu |
| 388 | 0.37 | — | 47.0° | salt | Rendaiji | 蓮臺寺 (静岡) | „ | Private bath (<i>K. Murayama</i>) |
| 389 | 0.37 | — | 40.0° | simple | Hatage | 畑毛 („) | Volc. Detr. | Hatage-Onsen No. 1 |
| 390 | 0.37 | 236 | 52.0° | earthy alkaline | Shirahone | 白骨 (長野) | Chichibu Series | Ō-yu |
| 391 | 0.37 | 25400 | 44.5° | sulphated bitter | Ikao | 伊香保 (群馬) | Andesite | Ōseki-no-yu |
| 392 | 0.36 | — | 41.0° | saline bitter | Yoshina | 吉奈 (静岡) | „ | Tōfuya-shin-yu |
| 393 | 0.36 | — | 45.0° | salt | Rendaiji | 蓮臺寺 („) | Tertiary | Kakezukaya-no-yu |
| 394 | 0.36 | — | 70.0° | „ | Yatsu | 谷津 („) | „ | Yamada-Onsen |
| 395 | 0.36 | 540 | 49.0° | simple | Yuno | 湯野 (福島) | „ | Kitsune-yu |
| 396 | 0.36 | 1400 | 77.0° | due common salt | Shuzenji | 修善寺 (静岡) | Andes. and Tuff | Ayame-no-yu |
| 397 | 0.36 | — | 49.5° | salt | Itō | 伊東 („) | Quaternary | Kusumi-Onsen |
| 398 | 0.35 | — | 39.0° | simple | Hatage | 畑毛 („) | Volc. Detr. | Hatage-Onsen No. 6 |
| 399 | 0.35 | — | 38.0° | „ | „ | „ | „ | „ No. 4 A |
| 400 | 0.35 | — | 38.0° | „ | „ | „ | „ | „ No. 4 B |
| 401 | 0.35 | — | 50.0° | salt | Yatsu | 谷津 („) | Tertiary | Mageya-no-yu |
| 402 | 0.34 | — | 50.5° | — | Itō | 伊東 („) | Quaternary | Yumoto-kwan-no-yu |
| 403 | 0.34 | — | 86.5° | salt | Moto-kuruma | 元車 (宮城) | Andesite | Ubuchi-no-yu |
| 404 | 0.34 | — | 64.0° | due common salt | Shuzenji | 修善寺 (静岡) | Andes. and Tuff | Katsura-no-yu |
| 405 | 0.34 | — | 59.5° | alkaline common salt | Ogawa | 小川 (富山) | Liparite | Ogawa-Onsen No. 3 |
| 406 | 0.34 | — | 46.0° | salt | Rendaiji | 蓮臺寺 (静岡) | Tertiary | Private bath (<i>R. Ogawa</i>) |
| 407 | 0.33 | — | 49.0° | simple | Nagaoka | 長岡 („) | „ | Tamon-yu No. 2 |
| 408 | 0.33 | — | 55.5° | earthy alkaline sulphur | Akakura | 赤倉 (新潟) | Andesite | Akakura-Onsen Jet No. 1 |
| 409 | 0.33 | — | 37.5° | salt | Itō | 伊東 (静岡) | Quaternary | Me-no-yu |
| 410 | 0.32 | 270 | 51.0° | sulphur | Naruko | 鳴丁 (宮城) | Andesite | Genzō-no-yu |
| 411 | 0.32 | 270 | 54.0° | common salt | Kamegawa | 龜川 (大分) | „ | Kiyō-sen |
| 412 | 0.31 | — | 72.0° | alkaline common salt | Shuzenji | 修善寺 (静岡) | Andes. and Tuff | Kikuya-no-yu No. 1 |
| 413 | 0.31 | 441 | 41.5° | bitter | Ushio | 海潮 (島根) | Andesite | Ushio-no-yu |
| 414 | 0.31 | 216 | 69.0° | iron carbonate | Shibasaki | 芝石 (大分) | „ | Hijiri-yu |
| 415 | 0.31 | — | 45.0° | salt | Rendaiji | 蓮臺寺 (静岡) | Tertiary | Private bath (<i>I. Yoshimura</i>) |
| 416 | 0.31 | — | 48.5° | — | Itō | 伊東 („) | Quaternary | Private bath (<i>Yoshida</i>) |
| 417 | 0.31 | 520 | 60.0° | due common salt | Shuzenji | 修善寺 („) | Andes. and Tuff | Iwa-no-yu |

| No. | Fumigation per litre of water in Maché's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|--|---|-----------------------|-------------------------|--------------------|----------|-----------------|-----------------------------------|
| 418 | 0.30 | ca. 450 | 61.0° | simple | Noboribetsu | 登別(北海道) | Tertiary | Okawa-no-yu |
| 419 | 0.30 | 54 | 48.0° | salt | Naruko | 鳴子(宮城) | Andesite | Kawara-no-yu |
| 420 | 0.30 | — | 49.5° | acid alum vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Sa-no-kawara-no-yu (Kumano-no-yu) |
| 421 | 0.30 | 1530 | 46.0° | salt | Funabara | 船原(静岡) | Andesite | Kumano-no-yu |
| 422 | 0.30 | 205 | 50.0° | saline bitter | Yoshina | 吉奈(静岡) | " | O-yu |
| 423 | 0.29 | — | 46.5° | — | Shigaku | 志學(島根) | " | Private bath (S. Nagira) |
| 424 | 0.29 | — | 47.8° | simple | Ito | 伊東(静岡) | Quaternary | Matsubara-Moto-yu |
| 425 | 0.29 | — | 77.0° | salt | Shimogamo | 下賀茂(静岡) | Tertiary | Shimogamo-Kyôdô-yu |
| 426 | 0.29 | — | 55.0° | sulphated bitter | Tohi | 土肥(静岡) | " | Kyôdô-yu (Suzuki) |
| 427 | 0.29 | — | 49.0° | simple | Nagaoka | 長岡(静岡) | " | Nagaoka-Onsen (No. 1) |
| 428 | 0.28 | — | 58.0° | carbonated | Akayu (Miyagi-ken) | 赤湯(宮城) | Andesite | Goten-no-yu |
| 429 | 0.27 | — | 100.0° | common salt | Shimogamo | 下賀茂(静岡) | Tertiary | Spring near Jiunji temple |
| 430 | 0.27 | — | 47.0° | — | Itô | 伊東(静岡) | Quaternary | Private bath (M. Suzuki) |
| 431 | 0.27 | — | 57.0° | sulphated bitter | Tohi | 土肥(静岡) | Tertiary | " Z. Yudo |
| 432 | 0.27 | — | 47.0° | salt | Rendaiji | 蓮臺寺(静岡) | " | Bath of Zuyô middle school |
| 433 | 0.27 | — | 48.0° | simple | Nagaoka | 長岡(静岡) | " | Tamon-yu (No. 1) |
| 434 | 0.27 | — | 48.0° | " | " | " | " | " (No. 3) |
| 435 | 0.26 | — | 49.0° | " | " | " | " | Private bath (S. Sugiyama) |
| 436 | 0.26 | — | 91.0° | earth-mixed common salt | Atami | 熱海(静岡) | Andesite | Takasagoya-no-yu |
| 437 | 0.25 | — | 73.0° | salt | Shimogamo | 下賀茂(静岡) | Tertiary | Kinokuniya-no-yu |
| 438 | 0.25 | — | 43.0° | sulphated bitter | Tohi | 土肥(静岡) | " | Private bath (M. Takasago) |
| 439 | 0.24 | abundant | 92.0° | acid vitriol | Naruko | 鳴子(宮城) | Andesite | Taki-no-yu No. 2 |
| 440 | 0.23 | — | 55.0° | sulphated bitter | Tohi | 土肥(静岡) | Tertiary | Private bath (T. Yoshimura) |
| 441 | 0.23 | — | 71.5° | saline bitter sulphur | Yamashiro | 山代(石川) | " | Kuraya-no-yu |
| 442 | 0.23 | — | 44.0° | sulphur | Tsubame | 燕(新潟) | Andesite | Tsubame-Onsen (No. 2) |
| 443 | 0.22 | — | 82.5° | common salt | Yugawara | 湯河原(神奈川) | " | Nakanishiya-no-yu |
| 444 | 0.22 | — | 58.0° | — | Kusatsu | 草津(群馬) | Volc. Detr. | Shirahata-no-yu |
| 445 | 0.22 | 198 | 57.0° | common salt | Kamegawa | 龜川(大分) | Andesite | Shi-no-yu |
| 446 | 0.22 | — | 99.0° | — | " | " | " | Kamado-no-yu |
| 447 | 0.21 | — | 48.0° | sulphur | Noboribetsu | 登別(北海道) | Tertiary | Manju-no-yu No. 2 |
| 448 | 0.21 | — | 43.5° | — | Itô | 伊東(静岡) | Quaternary | Private bath (Kobayashi) |
| 449 | 0.21 | 270 | 52.0° | salt | Yugano | 湯ヶ野(静岡) | Andesite | Yugano-Onsen |
| 450 | 0.20 | 684 | 47.0° | saline bitter | Funabara | 船原(静岡) | " | Kogyoku-no-yu (No. 1) |
| 451 | 0.20 | — | 52.0° | sulphated bitter | Tohi | 土肥(静岡) | Tertiary | Asaka-no-yu No. 2 |
| 452 | 0.19 | — | 50.0° | simple | Nagaoka | 長岡(静岡) | " | Nagaoka-Onsen No. 2 |
| 453 | 0.18 | — | 43.0° | acid alum vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Sa-no-kawara-no-yu (Kumano-no-yu) |
| 454 | 0.18 | 108 | 53.0° | acid | Dai | 大井(岩手) | Diluvium | Yakushi-no-yu |
| 455 | 0.18 | — | 88.5° | common salt | Yugawara | 湯河原(神奈川) | Andesite | Uenoya-no-yu |
| 456 | 0.18 | — | 53.0° | simple | Nagaoka | 長岡(静岡) | Tertiary | Nagaoka-Onsen (No. 1) |
| 457 | 0.18 | 54000 | 71.5° | acid vitriol | Noboribetsu | 登別(北海道) | " | Taki-no-yu |
| 458 | 0.18 | 80 | 80.0° | earth-mixed common salt | Atami | 熱海(静岡) | Andesite | Furuya-no-yu |
| 459 | 0.17 | — | 52.0° | simple | Kona | 古奈(静岡) | Tertiary | Kona-Onsen |
| 460 | 0.17 | — | 56.0° | — | Kusatsu | 草津(群馬) | Volc. Detr. | Chiyo-no-yu |
| 461 | 0.17 | — | 93.0° | alkaline | Naruko | 鳴子(宮城) | Andesite | Shin-unagi-no-yu |
| 462 | 0.17 | 126 | 42.0° | salt | Konabe | 鍋(静岡) | " | Konabe-Onsen |
| 463 | 0.16 | — | 56.0° | acid alum vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Netsu-no-yu |
| 464 | 0.16 | — | 46.0° | simple | Itô | 伊東(静岡) | Quaternary | Matsubara-Shin-yu |
| 465 | 0.15 | — | 54.5° | common salt | Yugawara | 湯河原(神奈川) | Andesite | Yuya-no-yu |
| 466 | 0.15 | 126 | 49.0° | saline bitter | Yoshina | 吉奈(静岡) | " | Tôfuya-no-yu |
| 467 | 0.15 | 80 | 41.0° | carbonated | Yugashima | 湯ヶ島(静岡) | Andes. and Tuff | Kidachi-yu |
| 468 | 0.15 | — | 40.0° | saline bitter | Funabara | 船原(静岡) | Andesite | Kogyoku-no-yu (No. 2) |
| 469 | 0.15 | 80 | 67.0° | saline common salt | Shuzenji | 修善寺(静岡) | Andes. and Tuff | Hana-no-yu |
| 470 | 0.14 | — | 80.0° | common salt | Yugawara | 湯河原(神奈川) | Andesite | Izuya-no-yu |
| 471 | 0.14 | 466 | 97.0° | earth-mixed common salt | Atami | 熱海(静岡) | " | Okawa-no-yu (No. 2) |

| No. | Emanation per litre of water in cubic units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-------------------------------------|---|---|-----------------------|-------------------------------------|---------------------------|----------|-----------------|--|
| 472 | 0.14 | — | 56.5° | acid alum vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Jizō-no-yu |
| 473 | 0.14 | — | 77.0° | earth-muriated common salt | Atami | 熱海(静岡) | Andesite | Ishi-yu |
| 474 | 0.13 | — | 58.0° | acid hydrogen sulphide alum vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Yubatake-no-yu |
| 475 | 0.13 | 380 | 91.0° | acid alum vitriol | Myōban | 明礬(大分) | Andesite | Jizō-no-yu |
| 476 | 0.13 | — | 64.0° | aline common salt | Shuzenji | 修善寺(静岡) | Andes. and Tuff | Ma-yu |
| 477 | 0.13 | 103.0° | alkaline sulphur | Naruko | 鳴子(宮城) | Andesite | Andesite | Unagi-no-yu |
| 478 | 0.13 | — | 60.0° | acid vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Washi-no-yu |
| 479 | 0.11 | 180 | 59.0° | carbonated | Kamegawa | 龜川(大分) | Andesite | Hamada-Onsen |
| 480 | 0.10 | — | 60.0° | sulphated bitter | Izusan | 伊豆山(静岡) | " | Hashiri-yu |
| 481 | 0.10 | — | 45.0° | simple | Nagaoka | 長岡(" | Tertiary | Nagaoka-Onsen No. 1 |
| 482 | 0.09 | — | 48.0° | sulphur | Tsubame | 燕(新潟) | Andesite | Tsubame-Onsen Jet No. 1 |
| 483 | 0.08 | 140 | 48.0° | earth-muriated common salt | Atami | 熱海(静岡) | " | Me-no-yu |
| 484 | 0.08 | 241 | 93.0° | " | " | " | " | Komatsu-no-yu |
| 485 | 0.07 | — | 43.0° | common salt | Yugawara | 湯河原(神奈川) | " | Ōkura-no-yu |
| 486 | 0.04 | 1296 | 108.0° | earth-muriated common salt | Atami | 熱海(静岡) | " | Ō-yu |
| 487 | 0.02 | — | 81.0° | " | " | " | " | Kawara-no-yu |
| Emanation per litre of gas at 0° C. | | | | | | | | |
| 1 | 28.44 | — | — | — | Tamatsukuri | 玉造(島根) | Tertiary | River bed C |
| 2 | 25.98 | — | 95.0° | alkaline carbon-dioxated | Yumura (Hyōgo-ken) | 湯村(兵庫) | Granite | Ara-yu |
| 3 | 24.44 | — | — | — | Tamatsukuri (Shimane-ken) | 玉造(島根) | Tertiary | River bed B |
| 4 | 14.60 | — | 39.5° | carbonated | Deyu | 出湯(新潟) | Granite | Chōsen-kutsu-no-yu |
| 5 | 13.37 | — | 51.0° | — | Kachimi | 勝見(鳥取) | " | Private bath Vent No. 2 (Kinosaki-kyō) |
| 6 | 9.63 | — | 42.5° | sulphur | Yoshioka | 吉岡(" | Tertiary | Shimo-no-yu |
| 7 | 8.73 | — | 59.5° | saline common salt | Shuzenji | 修善寺(静岡) | Andes. and Tuff | Ishi-yu |
| 8 | 7.07 | — | 70.0° | " | " | " | " | Kikuya-no-yu No. 2 |
| 9 | 4.99 | — | 76.0° | acid vitriol | Kamegawa | 龜川(大分) | Andesite | Chinoike-Jigoku Vent No. 2 |
| 10 | 4.82 | — | 76.0° | " | " | " | " | Vent No. 1 |
| 11 | 4.56 | — | 37.7° | earth-muriated common salt | Yunogō | 湯郷(岡山) | Liparite | Shin-Sagi-no-yu |
| 12 | 4.11 | — | 37.0° | " | " | " | " | Sagi-no-yu (Onna-yu) |
| 13 | 4.11 | — | 69.0° | saline common salt | Shuzenji | 修善寺(静岡) | Andes. and Tuff | Dokko-no-yu |
| 14 | 3.34 | — | 48.0° | simple | Nagaoka | 長岡(" | Tertiary | Tamon-yu No. 3 |
| 15 | 3.31 | — | 50.0° | sulphated common salt | Yunotsu | 温泉津(島根) | " | Kyū-yu |
| 16 | 1.91 | — | 69.0° | iron carbonate | Shibasaki | 芝石(大分) | Andesite | Hijiri-yu |
| 17 | 1.77 | — | 47.0° | saline bitter | Funabana | 船原(静岡) | " | Kōgyoku-no-yu No. 1 |
| 18 | 1.56 | — | 46.5° | common salt | Shigaku | 志學(島根) | " | Shigaku-Onsen |
| 19 | 1.37 | — | 60.0° | acid vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Washi-no-yu |
| 20 | 1.29 | — | 53.0° | simple | Nagaoka | 長岡(静岡) | Tertiary | Nagaoka-Onsen No. 4 |
| 21 | 1.19 | — | 47.8° | " | Ito | 伊東(" | Quaternary | Matsubara-Moto-yu |
| 22 | 1.05 | — | 52.0° | earthy alkaline | Shirahone | 白骨(長野) | Chichibu Series | Ō-yu |
| 23 | 0.83 | — | 58.0° | acid hydrogen sulphide alum vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Yubatake-no-yu |
| 24 | 0.71 | — | 56.0° | — | " | " | " | Chiyo-no-yu |
| 25 | 0.70 | — | 56.0° | acid alum vitriol | " | " | " | Netsu-no-yu |
| 26 | 0.66 | — | 58.0° | — | " | " | " | Shirahata-no-yu |
| 27 | 0.65 | — | 56.5° | acid alum vitriol | " | " | " | Jizō-no-yu |

Gas evolving from:

(B) Cold Springs.

| No. | Examination per litre of water in Maché's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|---|-----------------------|--------------------|------------|---------|----------|------------------------------|
| 1 | 617.59 | 22 | 21.5° (23.5°) | earthy common salt | Masutomi | 増富(山梨) | Granite | Kanigawara No. 1 |
| 2 | 575.02 | — | 14.0° | " | " | " | " | " No. 2 |
| 3 | 409.31 | 13 | 21.5° (41°) | common salt | " | " | " | Kuridaira No. 1 (Part C) |
| 4 | 388.14 | 100 | 20.0° (41°) | " | " | " | " | " No. 1 (Part A) |
| 5 | 312.54 | 77 | 22.5° (40°) | earthy common salt | " | " | " | Nibuzawa No. 1 |
| 6 | 298.33 | — | 10.0° (41°) | — | " | " | " | Oshiba No. 1 |
| 7 | 298.22 | — | 18.0° | — | " | " | " | Yunokubo No. 1 |
| 8 | 291.93 | 100 | 20.0° | common salt | " | " | " | Kuridaira No. 1 (Part B) |
| 9 | 258.50 | 13 | 20.0° (10.5°) | — | " | " | " | " No. 3 |
| 10 | 257.42 | — | 10.0° (11°) | simple | Takayama | 高増山(岐阜) | " | Ena-Kōsen (Yamaguchi) |
| 11 | 240.20 | 45 | 17.5° (7°) | earthy common salt | Masutomi | 増富(山梨) | " | Tsugane-yu No. 1 (A) |
| 12 | 239.22 | — | 15.0° (10.5°) | " | " | " | " | Yunokubo-gawara-no-yu |
| 13 | 160.34 | — | 17.0° (12°) | carbonated (?) | Ikeda | 池田(島根) | " | Nobata-shin-yu |
| 14 | 146.97 | — | 18.5° | — | Masutomi | 増富(山梨) | " | Hatchō-jyaya-shita No. 1 |
| 15 | 109.96 | — | 30.0° (14°) | — | " | " | " | Ginsen-tō |
| 16 | 86.27 | 13 | 19.0° (7°) | earthy common salt | " | " | " | Tsugane-yu No. 1 (B) |
| 17 | 83.35 | — | 16.0° (7°) | " | " | " | " | " No. 1 (C) |
| 18 | 75.62 | — | 10.0° | — | " | " | " | Yunomukai No. 1 |
| 19 | 72.26 | — | 12.0° (20°) | simple | Hirukawa | 蛭川(岐阜) | " | Ena-Kōsen (Dairen-Fukushima) |
| 20 | 63.92 | — | 13.0° (1.0°) | " | Takayama | 高山(山梨) | " | " (Shimizu) |
| 21 | 56.41 | — | 28.5° (24.5°) | " | Tochiomata | 栃尾又(新潟) | " | Tochiomata-no-yu No. 4 |
| 22 | 46.04 | 530 | 25.0° (30°) | " | Murasugi | 村杉(山梨) | " | Murasugi-Kōsen No. 1 |
| 23 | 42.59 | — | 26.0° (31.5°) | " | " | " | " | " No. 2 |
| 24 | 40.28 | — | 13.5° (30°) | " | " | " | " | " No. 3 |
| 25 | 39.51 | 236 | 36.0° (27.5°) | " | Tochiomata | 栃尾又(山梨) | " | Jizai-kwan-uchi-yu No. 1 |
| 26 | 31.90 | — | 13.0° (21°) | " | Takayama | 高山(岐阜) | " | Okuhora |
| 27 | 30.36 | — | 13.0° (19°) | " | " | " | " | Inari-no-yu |
| 28 | 27.18 | — | 35.5° (26°) | " | Tochiomata | 栃尾又(新潟) | " | Tochiomata-no-yu No. 3 |
| 29 | 25.77 | — | 12.0° (20°) | " | Hirukawa | 蛭川(岐阜) | " | Shimo-issiki |
| 30 | 25.16 | — | 11.0° (20°) | " | Takayama | 高山(山梨) | " | Yamaki-no-yu |
| 31 | 23.70 | — | 10.0° (21.5°) | " | " | " | " | Ichi-no-sawa No. 1 |
| 32 | 19.42 | — | 12.0° (21°) | " | " | " | " | Shimoyama No. 1 |
| 33 | 18.48 | — | — | " | " | " | " | Kami-no-ido |
| 34 | 17.13 | — | — | " | " | " | " | Bunsaku-ido |
| 35 | 17.08 | — | 9.5° (21°) | " | " | " | " | Ichi-no-sawa No. 3 |
| 36 | * 15.77 | — | — | carbonated | Kasubuchi | 粕淵(島根) | Liparite | Kasubuchi-no-yu |
| 37 | * 15.38 | — | — | simple | Takayama | 高山(岐阜) | Granite | Spring A |
| 38 | 15.18 | — | 33.5° (20°) | — | Misasa | 三朝(鳥取) | " | Spring C |
| 39 | 14.20 | — | — | simple | Takayama | 高山(岐阜) | " | Shishi-no-neya |
| 40 | 13.74 | — | 22.5° (20°) | " | Murasugi | 村杉(新潟) | " | Shiroyama-Kōsen |
| 41 | 13.32 | — | 10.0° (7°) | " | Komono | 菰野(三重) | " | Spring A |
| 42 | 13.00 | — | 14.5° (9.2°) | " | Kaidani | 栢谷(岡山) | " | Tomada-no-yu |
| 43 | 11.57 | — | 33.0° (27°) | carbonated | Deyu | 出湯(新潟) | " | Tōshundai-no-yu No. 1 |
| 44 | 11.31 | — | 31.0° (28°) | " | " | " | " | " No. 2 |
| 45 | * 9.69 | 108 | 15.7° | " | Tombara | 樽原(島根) | " | Kamiyasumi Kōsen |
| 46 | 9.69 | — | — | simple | Bobata | 母畑(福島) | Gneiss | Spring B (K. Watanabe) |
| 47 | 8.77 | — | 8.2° | — | Masutomi | 増富(山梨) | Granite | Taikarizawa No. 2 |
| 48 | 8.42 | — | 4.5° | — | " | " | " | " No. 3 |
| 49 | 8.18 | — | 29.0° (13°) | simple | Komono | 菰野(三重) | " | Shika-no-yu |
| 50 | 7.45 | — | 18.7° (32°) | " | Imaita | 板新湯(山梨) | " | Imaita-Kōsen No. 1 |
| 51 | 6.99 | — | 33.5° (29.5°) | carbonated | Deyu | 出湯(山梨) | Granite | Shiraneya-no-yu |
| 52 | 6.54 | — | 8.2° | — | Masutomi | 増富(山梨) | " | Taikarizawa No. 1 |

| No. | Emanation per
litre of water
in Maché's units | Flow of water
in 24 hrs.
in hectolitres | Temp. of spring
in C. | Classification | Location | 地 名(府縣名) | Geology | Spring |
|-----|---|---|--------------------------|-------------------------|--------------------|----------|-------------------------|-------------------------------|
| 53 | 5.87 | ca. — | 36.0° | salt | Matsuzaki | 松崎(鳥取) | Andesite | Matsuzaki-kwan-no-yu |
| 54 | 5.83 | 25 | 24.4° (15°) | — | Yoshikata | 吉方(„) | Alluvium | Kinka-Onsen |
| 55 | 5.46 | — | 9.0° (3.5°) | simple | Shinji | 宍道(鳥根) | Granite | Ippai-shimizu |
| 56 | 5.30 | — | 34.0° | saline bitter | Higashiyama | 東山(福島) | Andesite or
Tertiary | Nuru-yu |
| 57 | 4.93 | — | 19.0° (1°) | simple | Yumura Shimane-ken | 湯村(鳥根) | Granite | Spring C |
| 58 | 4.55 | — | 32.0° | salt | Matsuzaki | 松崎(鳥取) | Andesite | Spring (in Matsuzaki Station) |
| 59 | 4.32 | 300 | 28.0° (16°) | simple | Takebe | 建部(岡山) | Granite | Yawata-Onsen |
| 60 | 4.19 | — | 16.0° (32°) | „ | Imaita | 今板(新潟) | — | Imaita-Kōsen No. 2 |
| 61 | 4.19 | — | — | — | Misasa | 三朝(鳥取) | Granite | Drinking water (N. Okazaki) |
| 62 | 4.08 | — | — | simple | Nekonaki | 猫啼(福島) | Gneiss | Spring A (Chōraku-en) |
| 63 | 4.03 | — | 9.0° | — | Masutomi | 増富(山梨) | Granite | Toyanosawa No. 1 |
| 64 | 3.87 | — | 35.0° (14°) | — | Yoshikata | 吉方(鳥取) | Alluvium | Shiratama-Onsen |
| 65 | 3.59 | — | 14.0° (16.5°) | — | Shimo-sase | 下佐世(鳥根) | Granite | Drinking water |
| 66 | 3.49 | 90 | 32.0° (11°) | alkaline common
salt | Yudani | 湯谷(鳥取) | Tertiary | Kabu-yu |
| 67 | 3.33 | 15 | — | — | Yoshikata | 吉方(„) | Alluvium | Kinoe-Onsen No. 1 |
| 68 | 3.26 | — | 14.0° (28°) | simple | Haguro | 羽黒(新潟) | — | Uba-yu |
| 69 | 3.18 | — | — | „ | Nekonaki | 猫啼(福島) | Gneiss | Spring C (Yakōji temple) |
| 70 | 3.15 | — | 10.5° | — | Masutomi | 増富(山梨) | Chichibu Series ? | Akashibu No. 3 |
| 71 | 3.14 | — | 24.0° (12°) | carbonated (?) | Ikeda | 池田(鳥根) | Granite | Nobata-no-yu |
| 72 | * 3.09 | — | — | — | Yumura (Hyōgo-ken) | 湯村(兵庫) | „ | Drinking water at Chimizu |
| 73 | 3.07 | — | — | simple | Nekonaki | 猫啼(福島) | Gneiss | Spring E |
| 74 | 3.04 | — | 31.0° | „ | Tōgō | 東郷(鳥取) | Andesite | Yōjyō-kwan-no-yu No. 5 |
| 75 | 3.02 | — | 11.5° (14°) | „ | Misasa | 三朝(„) | Granite | Spring A (K. Tsugawa) |
| 76 | 2.97 | — | 12.0° | — | Masutomi | 増富(山梨) | Chichibu Series ? | Akashibu No. 2 |
| 77 | 2.95 | — | 12.5° (21°) | simple | Takayama | 高山(岐阜) | Granite | Ichinosawa No. 2 |
| 78 | 2.95 | — | 10.5° | — | Kami-sekisuiji | 上積翠寺(山梨) | „ | Yōsō-Onsen |
| 79 | 2.94 | — | — | simple | Bobata | 母畑(福島) | Gneiss | Naka-no-yu No. 2 |
| 80 | 2.92 | — | 14.0° | — | Masutomi | 増富(山梨) | Chichibu Series ? | Itaya No. 1 |
| 81 | 2.86 | — | 12.0° | — | „ | „ | Granite | Umamichizawa No. 1 |
| 82 | 2.64 | — | 28.0° | simple | Sekine-Yunosawa | 関根湯澤(山形) | Gneiss | Yunosawa-Onsen |
| 83 | 2.61 | — | 13.0° | — | Masutomi | 増富(山梨) | Chichibu Series ? | Akashibu No. 1 |
| 84 | 2.36 | — | — | simple | Bobata | 母畑(福島) | Gneiss | Shimo-no-yu No. 2 |
| 85 | 2.29 | 362 | 33.0° | — | Masutomi | 増富(山梨) | Granite | Kinsen-tō |
| 86 | 2.27 | — | 36.5° (29°) | simple | Asama | 浅間(長野) | Tertiary | Me-no-yu |
| 87 | 2.24 | 15 | 26.0° (17°) | — | Masutomi | 増富(山梨) | Granite | Kuridaira No. 2 (A) |
| 88 | 2.18 | — | — | simple | Nekonaki | 猫啼(福島) | Gneiss | Tomaki-furu-yu |
| 89 | 2.13 | — | 12.0° | — | Masutomi | 増富(山梨) | Chichibu Series ? | Itaya No. 2 |
| 90 | 2.12 | — | 3.9° | — | „ | „ | Granite | Matsudaira No. 1 |
| 91 | * 2.02 | — | — | — | Yumura (Hyōgo-ken) | 湯村(兵庫) | „ | Drinking water at Shin-machi |
| 92 | 2.02 | — | 35.5° | salt | „ (Yamanashi-ken) | „ (山梨) | Alluvium | Tani-Onsen |
| 93 | 1.95 | — | — | simple | Nekonaki | 猫啼(福島) | Gneiss | Spring D |
| 94 | * 1.95 | — | — | — | Shimo-oritate | 下折立(新潟) | Granite | Ibo-no-yu |
| 95 | 1.92 | 11 | 20.0° (17°) | — | Masutomi | 増富(山梨) | „ | Kuridaira No. 2 (C) |
| 96 | 1.91 | — | — | — | Kami-sase | 上佐世(鳥根) | „ | Drinking water |
| 97 | 1.89 | — | — | simple | Bobata | 母畑(福島) | Gneiss | Toi-no-kuchi |
| 98 | * 1.88 | — | — | — | Sugatani | 菅谷(新潟) | Alluvium | Sugatani-Kōsen |
| 99 | 1.86 | — | 22.5° (15°) | — | Shigaku | 志賀(鳥根) | Andesite | Spring A |
| 100 | 1.86 | — | 33.8° | salt | Yumura | 湯村(山梨) | „ | Kōbōzue-Onsen |
| 101 | 1.85 | — | 14.5° (11°) | simple | Masutomi | 増富(„) | Granite | Oshiba No. 2 |
| 102 | 1.82 | — | — | „ | Nekonaki | 猫啼(福島) | Gneiss | Spring B (Chayū) |
| 103 | 1.55 | 30 | 11.0° | „ | Kami-sekisuiji | 上積翠寺(山梨) | Andesite | Yōgai-Onsen |
| 104 | 1.52 | — | — | „ | Bobata | 母畑(福島) | Gneiss | Spring A (C. Yabe) |
| 105 | 1.35 | — | — | — | Yoshikata | 吉方(鳥取) | Alluvium | Kinoe-Onsen No. 2 |

| No. | Emanation per litre of water in Maché's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|---|-----------------------|--|----------------------|----------|------------|---------------------------------|
| 106 | 1.33 | — | 12.5° | — | Isobe (Gumma-ken) | 磯部(群馬) | Tertiary | Ji-no-yu |
| 107 | 1.24 | — | 34.0 | simple | Oyuzawa | 大湯澤(秋田) | Diluvium | Shimo-naizawa-no-yu |
| 108 | * 1.22 | — | — | — | Hōgijiku | 寶木宿(鳥取) | Granite | Drinking water (T. Minamimatsu) |
| 109 | 1.19 | — | 28.0° (11) | — | Yamabe | 山邊(長野) | Tertiary | Mitarashi-no-yu |
| 110 | 1.18 | 20 | 13.0° (12) | carbonated common salt | Kawai | 川合(島根) | Granite | Urisaka-no-yu |
| 111 | 1.14 | — | 9.0° (2) | simple | Yumura (Shimane-ken) | 湯村(島根) | — | Spring B |
| 112 | 1.10 | — | 10.0° (11) | — | Ishioka | 石岡(茨城) | Diluvium | Well (K. Higashi) |
| 113 | * 1.10 | — | — | — | Komatsu | 小松(石川) | Alluvium | Komatsu Kōsen (A. K. Ito) |
| 114 | * 1.09 | — | — | — | Hōgijiku | 寶木宿(鳥取) | Granite | Well (K. Minamimatsu) |
| 115 | 1.08 | — | 28.0° | hydrogen sulphid | Nasu-Yumoto | 那須湯本(栃木) | Andesite | Kiraku-no-yu |
| 116 | 1.06 | — | — | — | Ushio | 海潮(島根) | — | Drinking water |
| 117 | * 1.02 | — | — | — | Utsuno | 宇津野(新潟) | — | Sakamoto-no-yu |
| 118 | 1.01 | 29 | 14.0° (12) | earth-maturated iron carbon-dioxated common salt | Kawai | 川合(島根) | Granite | Tōkōji-no-yu |
| 119 | 1.01 | — | 13.6° (13°) | muriated alkaline | Kasagi | 笠置(京都) | — | Kasagi-Kōsen No. 1 |
| 120 | 1.00 | — | 25.0° (10°) | sulphur (?) | Yunogō | 湯郷(岡山) | Liparite | Me-no-yu |
| 121 | 0.98 | 180 | 33.5° | hydrogen sulphid common salt | Oyuzawa | 大湯澤(秋田) | Diluvium | Well (S. Minamimatsu) |
| 122 | 0.98 | — | 20.0° (13°) | — | Tera-machi [Tottori] | 寺町(鳥取) | Alluvium | Elbisu-Onsen |
| 123 | 0.97 | — | 17.5° (15°) | common salt | Shinga | 新宮(石川) | Granite | Shingu Kōsen (A. V. Ito) |
| 124 | 0.96 | — | — | — | Yunogō | 湯郷(岡山) | Liparite | Well (M. Minamimatsu) |
| 125 | 0.95 | — | 16.0° (14) | — | Tadate | 田館(茨城) | Tertiary | Shibu-ido (O-sen) |
| 126 | 0.94 | — | 16.0° (15°) | common salt | Shingū | 新宮(石川) | Granite | Shingu-Kōsen No. 2 |
| 127 | * 0.92 | — | 21.1° | — | Komatsu | 小松(石川) | Alluvium | Komatsu Kōsen (B. V. Ito) |
| 128 | 0.90 | 650 | 32.0° | iron carbonate | Akayu (Akita-ken) | 赤湯(秋田) | Diluvium | Akayu-Onsen |
| 129 | 0.89 | — | 12.0° (4°) | simple | Hinobori | 日登(島根) | Granite | Spring A |
| 130 | 0.84 | — | 13.6° (13°) | muriated alkaline | Kasagi | 笠置(京都) | — | Kasagi-Kōsen No. 2 |
| 131 | 0.78 | — | — | — | Tokyo—Azabu-ku | 麻布(東京) | Diluvium | Well (S. Okubo) |
| 132 | 0.77 | — | — | — | Yunogō | 湯郷(岡山) | Liparite | Well (T. Okubo) |
| 133 | 0.75 | — | — | — | Tokyo—Azabu-ku | 麻布(東京) | Diluvium | Well (K. Kōjima) |
| 134 | 0.72 | — | — | salt | Gakutō | 學頭(島根) | Tertiary | Sudani-yakutō |
| 135 | 0.67 | — | — | — | Yunogō | 湯郷(岡山) | Liparite | Well (G. Minamimatsu) |
| 136 | * 0.66 | — | 22.0° | — | Kōbu | 構武(島根) | Alluvium | Myōbun-Kōsen |
| 137 | * 0.64 | — | — | — | Hōgijiku | 寶木宿(鳥取) | Granite | Well (S. Hori) |
| 138 | 0.64 | — | 6.0° (11) | simple | Yumura (Shimane-ken) | 湯村(島根) | — | Spring D |
| 139 | * 0.63 | — | — | — | Koyabara | 小屋原(石川) | Andesite | Spring B |
| 140 | 0.61 | — | 12.5° (29°) | — | Futami | 二見(富山) | Granite | Spring A |
| 141 | 0.58 | — | 29.0° (30°) | — | Yamabe | 山邊(長野) | Tertiary | Oboke-no-yu |
| 142 | 0.55 | — | 14.0° (13.5°) | — | Ishioka | 石岡(茨城) | Diluvium | Well (O. Nakazawa) |
| 143 | * 0.51 | — | — | — | Tokyo—Azabu-ku | 麻布(東京) | — | Well (T. Minamimatsu) |
| 144 | 0.49 | — | 17.0° (11) | — | Ishioka | 石岡(茨城) | — | Well (T. Higashi) |
| 145 | 0.49 | — | — | simple | Bobata | 母畑(福島) | Gneiss | Shimo-no-kyū-yu |
| 146 | 0.44 | — | 18.5° (15°) | — | Ōmori | 大森(東京) | Alluvium | Well (S. Minamimatsu) |
| 147 | 0.42 | 864 | 27.0° | salt | Gakutō | 學頭(島根) | Tertiary | Yunokawa-Kōsen |
| 148 | * 0.42 | — | — | — | Tokyo—Azabu-ku | 麻布(東京) | Diluvium | Well (H. Miyahara) |
| 149 | 0.42 | — | 34.0° | — | Ito | 伊東(静岡) | Quaternary | Private (H. S. A.) |
| 150 | 0.41 | — | 16.0° (16.5°) | — | Kawasaki | 川崎(神奈川) | Alluvium | Dokumizu |
| 151 | 0.40 | 3370 | 17.0° (18°) | earth-maturated common salt | Ōmori (Kanazawa-ken) | 大森(東京) | — | Morigasaki-Kōsen |
| 152 | 0.35 | — | 29.0° | — | Shimoda | 下田(静岡) | — | Atama Kōsen (A. V. Ito) |
| 153 | 0.33 | — | 22.0° | — | — | — | — | Well (A. V. Ito) |
| 154 | 0.28 | — | 14.0° (14°) | simple | Ishioka | 石岡(茨城) | Granite | Spring A (K. Higashi) |
| 155 | 0.28 | — | 36.0 | sulphated bitter | Tohi | 土肥(静岡) | Tertiary | Takasagoya-no-yu |
| 156 | * 0.27 | — | — | — | Saizu | 草薙(石川) | — | Yagizawa-Kōsen (A) |
| 157 | * 0.26 | — | — | — | Atami | 熱海(静岡) | Andesite | Drinking water (S. A.) |
| 158 | 0.25 | — | 31.0° | — | Shimoda | 下田(静岡) | — | Fuji-no-yu |

| No. | Emanation per litre of water in Mache's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-------------------------------------|---|---|-----------------------|---|----------------------|---------|-------------|----------------------|
| 159 | 0.25 | ca. | — | — | Baba-machi [Tottori] | 馬場町(鳥取) | Alluvium | Yōrō-no-yu |
| 160 | 0.23 | — | 35.0° | saline bitter | Funabara | 船原(静岡) | Andesite | Kōgyoku-no-yu No. 2 |
| 161 | 0.19 | — | 15.0° | acid vitriol | Noroshi | 狼煙(石川) | Alluvium | Noroshi-Kōsen |
| 162 | 0.19 | — | — | — | Osaki | 大崎(東京) | Diluvium | Well (T. Kikuchi) |
| 163 | 0.18 | — | — | simple | Bobata | 母畑(福島) | Gneiss | Spring C |
| 164 | 0.14 | — | 15.0° | sulphur | Nakaishida | 中石田(静岡) | Volc. Detr. | Kametsuru-Onsen |
| 165 | 0.12 | 77 | 34.5° (16°) | carbonated common salt | Fukumitsu | 福光(島根) | Tertiary | Yunohara-Onsen |
| 166 | 0.11 | — | 33.5 | salt | Ito | 伊東(静岡) | Quaternary | Yuda Onsen |
| 167 | 0.03 | — | — | — | Kusatsu | 草津(群馬) | Volc. Detr. | Drinking water |
| Emanation per litre of gas at 0° C. | | | | | | | | |
| 1 | 539.99 | — | 15.0° (10.5°) | earthy common salt | Masutomi | 増富(山梨) | Granite | Yunokubogawara-no-yu |
| 2 | 169.98 | — | 26.0° (31.5°) | simple | Murasugi | 村杉(新潟) | " | Murasugi-Kōsen No. 2 |
| 3 | 18.16 | — | 24.0° (12°) | carbonated (?) | Ikeda | 池田(島根) | " | Nobata-no-yu |
| 4 | 4.74 | — | 25.0° (10°) | sulphur (?) | Yunogō | 湯郷(岡山) | Liparite | Me-no-yu |
| 5 | 2.21 | — | 13.0° (1°) | carbonated common salt | Kawai | 川合(島根) | Granite | Urisaka-no-yu |
| 6 | 2.03 | — | 14.0° (1°) | earth-muriated iron carbon-dioxated common salt | " | " | " | Tōkōji-no-yu |
| 7 | 0.84 | — | 16.0° (16.5°) | — | Kawasaki | 川崎(神奈川) | Alluvium | Doku-mizu |
| 8 | 0.03 | — | 34.5° (10°) | carbonated common salt | Fukumitsu | 福光(島根) | Tertiary | Yunohara-Onsen |
| Gas evolving from: | | | | | | | | |

Table 3.

The determination of radium emanation was carried on with the fontactoscope of C. Engler and H. Sieveking.

I. Determined according to the authors' original direction.

II. Correction for the absorption of the radiation by the wall of the ionisation chamber was made.

III. Correction for the loss of emanation by inserting dissipation cylinder in the ionisation chamber was also made.

| No. | Emanation per litre of water in Mache's units | | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|-------|-------|---|-----------------------|-------------------------------|------------|---------|----------|----------------------------|
| | I† | II‡ | III‡ | | | | | | | |
| 1 | 9.67 | 11.65 | 11.88 | ca. | 50.0° | earthy common salt | Yunokawa | 湯川(北海道) | Liparite | Shin-yu No. 2 (Kinshōkwan) |
| 2 | 9.42 | 10.87 | 11.09 | 216 | 50.0° | earth-muriated common salt | " | " | " | Taki-no-yu (") |
| 3 | 7.05 | 8.13 | 8.29 | — | 66.0° | earthy common salt | Nezaki | 根崎(") | " | Rikuzō-no-yu |
| 4 | 2.72 | 3.15 | 3.21 | 152 | 76.0° | earth-muriated common salt | Awara | 蘆原(福井) | Alluvium | Funatsu No. 10 |
| 5 | 2.36 | 2.73 | 2.79 | 152 | 70.5° | " | " | " | " | " No. 21 |
| 6 | 2.26 | 2.61 | 2.66 | 152 | 75.0° | " | " | " | " | Nimen No. 23 |
| 7 | 2.06 | 2.38 | 2.43 | 3944 | 62.0° | bitter | Kaminoyama | 上ノ山(山形) | Liparite | Hie-no-yu |
| 8 | 1.85 | 2.14 | 2.18 | 152 | 68.2° | earth-muriated common salt | Awara | 蘆原(福井) | Alluvium | Tanaka-naka No. 6 |
| 9 | 1.83 | 2.12 | 2.16 | 152 | 70.0° | " | " | " | " | Funatsu No. 9 |
| 10 | 1.54 | 1.77 | 1.81 | 152 | 63.0° | " | " | " | " | Tanaka-naka No. 22 |
| 11 | 1.51 | 1.75 | 1.79 | 152 | 53.0° | " | " | " | " | Nimen No. 21 |
| 12 | 1.08 | 1.25 | 1.28 | 152 | 74.0° | " | " | " | " | Funatsu No. 6 |
| 13 | 1.04 | 1.20 | 1.22 | 1530 | 42.0° | earthy common salt | Miyagawa | 宮川(北海道) | Andesite | Miyagawa-no-yu |
| 14 | 0.94 | 1.09 | 1.11 | 270 | 45.5° | hydrogen sulphide common salt | Ōyuzawa | 大湯澤(秋田) | Diluvium | No. 33 (Ō-yu) |
| 15 | 0.91 | 1.05 | 1.07 | — | 8.0° | — | Matsuida | 松井田(群馬) | Tertiary | Ushio-Kōsen |
| 16 | 0.82 | 0.95 | 0.97 | — | 7.5° | — | " | " | " | Yunosawa-Kōsen (Kōngō-ji) |

| No. | Emanation per litre of water in Mache's units | | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-------------------------------------|---|-----------------|--------|---|-----------------------|-------------------------------|------------|---------|----------|-----------------------------------|
| | I [†] | II [†] | III | | | | | | | |
| 17 | 0.65 | 0.75 | 0.77 | ca. | 37.0° | common salt | Irinoyu | 入之湯(群馬) | Andesite | Iri-no-yu |
| 18 | 0.64 | 0.74 | 0.75 | 7 | 9.5° | alkaline | Matsuida | 松井田() | Tertiary | Rokutanda-Kōsen |
| 19 | 0.63 | 0.73 | 0.74 | — | 31.5° | common salt | Irinoyu | 入之湯() | Andesite | Kaede-no-yu |
| 20 | 0.61 | 0.71 | 0.72 | — | 33.5° | " | " | " | " | Me-no-yu |
| Emanation per litre of gas at 0° C. | | | | | | | | | | |
| 1 | 85.97 | 99.14 | 101.03 | — | 66.0° | earthy common salt | Nezaki | 根崎(北海道) | Liparite | Rikuzō-no-yu |
| 2 | 70.04 | 80.76 | 82.31 | — | 50.0° | earth-muriated common salt | Yunokawa | 湯川() | " | Taki-no-yu (<i>Rinchō-kwan</i>) |
| 3 | 63.14 | 72.78 | 74.17 | — | 50.0° | earthy common salt | " | " | " | Shin-yu (<i>M. 2</i> ()) |
| 4 | 33.55 | 38.69 | 39.43 | — | 76.0° | earth-muriated common salt | Awara | 蘆原(福井) | Alluvium | Funatsu <i>No. 10</i> |
| 5 | 22.45 | 25.88 | 26.37 | — | 70.0° | " | " | " | " | " <i>No. 9</i> |
| 6 | 13.45 | 15.51 | 15.80 | — | 62.0° | bitter | Kaminoyama | 上ノ山(山形) | Liparite | Hie-no-yu |
| 7 | 4.55 | 5.25 | 5.35 | — | 45.5° | hydrogen sulphide common salt | Oyuzawa | 大湯澤(秋田) | Diluvium | <i>No. 33</i> (<i>Ōyuzawa</i>) |
| 8 | 3.54 | 4.08 | 4.16 | — | 42.0° | earthy common salt | Miyagawa | 宮川(北海道) | Andesite | Miyagawa-no-yu |

Table 4.

The quantity of radium emanation and the temperature of springs in the following table were determined by Den'itirō Ishitani, Prof. of Physics, Imperial Peers' College, co-operated partly with Kaichirō Manabe, partly with Ichirō Yamakawa (both being assistants of Medical Faculty of Tōkyō Imperial University). As for particulars refer to Proceeding of the Tōkyō Mathematico-Physical Society (Tōkyō Sūgaku-Buturigakkwai Kizi), 2nd Ser.:

Vol. VI, No. 12, p. 182-183; No. 15, p. 221-222; No. 18, p. 276, 279; No. 20, p. 293; No. 21, p. 308-309; Vol.

VII, No. 1, p. 10; No. 3, p. 32, 33, 36; No. 9, p. 151; No. 12, p. 222-224; Vol. VIII, No. 1, p. 16-19, 22, 25.

For recalculation of Mache's units from the international unit following factor was used:

$$1 \times 10^{-10} \text{ curie} = 0.275 \text{ Mache's units.}$$

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|------------------------------|---------------------------------|---|-----------------------|--|----------|---------|---------------|-------------------------------------|
| | in 10 ⁻¹⁰ curies | in Mache's units (recalculated) | | | | | | | |
| 1 | 138.00 | 37.95 | — | > 31.0° | — | Arima | 有馬(兵庫) | Quartz Porph. | Zuihōji-Onsen |
| 2 | 42.00 | 11.55 | — | > 52.6° | salt | Tōgatta | 遠刈田(宮城) | Liparite | Taki-no-yu |
| 3 | 38.90 | 10.70 | 335 | 55.9° | carbonated | " | " | " | Kami-no-yu |
| 4 | 31.40 | 8.64 | — | 19.0° | earth-muriated iron, carbonate common salt | Arima | 有馬(兵庫) | Quartz Porph. | Maruyama-Kōsen |
| 5 | 26.70 | 7.34 | — | 63.1° | earth-muriated common salt | Kinosaki | 城崎() | Tertiary | Gesho-no-yu (<i>Font. 1</i>) |
| 6 | 18.35 | 5.05 | 204 | 63.0° | " | " | " | " | Kō-no-yu |
| 7 | 14.70 | 4.04 | — | 20.6° (3.5°) | muriated sulphur | Takedao | 武田尾() | Quartz Porph. | Yumoto |
| 8 | 13.60 | 3.74 | — | 46.0° | simple | Dogo | 後愛媛 | Granite | Yōjō-yu |
| 9 | 13.10 | 3.60 | — | 17.5° (24.) | — | Shioyu | 瀬湯(宮城) | " | Shio-yu |
| 10 | 12.95 | 3.56 | — | 46.0° | simple | Dogo | 道後(愛媛) | Granite | Kami-no-yu |
| 11 | 12.90 | 3.55 | — | 91.0° | earth-muriated common salt | Wakura | 和倉(石川) | Tertiary | Artes. Well of Wakazaki |
| 12 | 12.50 | 3.41 | 270 | 51.0° | salt | Sakunami | 作並(宮城) | " | Tsuru-no-yu |
| 13 | 12.00 | 3.30 | 216 | 59.3° | earth-muriated common salt | Kinosaki | 城崎(兵庫) | " | Yakagi-yu |
| 14 | 11.00 | 3.03 | — | 51.3° | salt | Tōgatta | 遠刈田(宮城) | Liparite | (<i>Font. 2</i>) (<i>N. 10</i>) |
| 15 | 10.40 | 2.86 | — | 24.6° (3.7°) | muriated sulphur | Takedao | 武田尾(兵庫) | Quartz Porph. | Moto-yu |
| 16 | 9.98 | 2.75 | — | 60.0° | earth-muriated common salt | Kinosaki | 城崎() | Tertiary | Ichi-no-yu (<i>Font. 3</i>) |

| No. | Emergence per
litre of water
to be consumed | Mod. salt
(included) | Flow of water
in 24 hr.
in hectolitres | Temp. of spring
in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|-------------------------|--|--------------------------|-----------------------------------|-------------------|-----------|---------------|--|
| 17 | 9.68 | 2.66 | — | 57.0° | saline bitter | Shiobara—Ōami | 鹽原—大網(栃木) | Tertiary | Onna-yu |
| 18 | 9.40 | 2.59 | — | 61.5° | earth-muriated
carbonated salt | Kinosaki | 城崎(兵庫) | " | Gosho-no-yu <i>Vent B</i> |
| 19 | 8.92 | 2.45 | — | 44.5° | " | " | " | " | Mandara-no-yu <i>Vent E</i> |
| 20 | 8.86 | 2.44 | — | 48.8° | salt | Sakunami | 作並(宮城) | " | Iwamatsu-Shin-yu |
| 21 | 8.72 | 2.40 | — | 46.6° | " | " | " | " | Me-no-yu |
| 22 | 8.58 | 2.36 | 270 | 50.6° | " | " | " | " | Kame-no-yu |
| 23 | 8.24 | 2.27 | — | 58.1° | carbonated | Namaze | 生瀬(兵庫) | Granite | Namaze-Kōsen |
| 24 | 8.16 | 2.24 | — | 55.8° | earth-muriated
carbonated salt | Kinosaki | 城崎(") | Tertiary | Mandara-no-yu <i>Vent B</i> |
| 25 | 8.04 | 2.21 | — | 15.0° (9.4) | simple carbon-
dioxated | Takarazuka | 寶塚(") | Granite | Tansan-sen <i>C</i> |
| 26 | 7.97 | 2.19 | — | 45.0° | common salt | Shiobara—Fukuwat. | 鹽原—福渡(栃木) | Tertiary | Hadaka-yu |
| 27 | 7.91 | 2.18 | — | 21.5° | simple carbon-
dioxated | Kōbe | 神戸(兵庫) | Granite | Jyareyama Tansan-sen <i>A</i> |
| 28 | 7.85 | 2.16 | — | 14.7° (9.2) | " | Takarazuka | 寶塚(") | " | Tansan-sen <i>B</i> |
| 29 | 7.83 | 2.15 | — | 21.5° | " | Kōbe | 神戸(") | " | Jyareyama Tansan-sen <i>B</i> |
| 30 | 7.78 | 2.14 | — | 18.8° | — | Arima | 有馬(") | Quartz Porph. | Shōtō-en Tansan-sen |
| 31 | 7.32 | 2.01 | — | 44.2° | earth-muriated
common salt | Kinosaki | 城崎(") | Tertiary | Ichi-no-yu <i>Vent D (Kase-yu)</i> |
| 32 | 7.04 | 1.94 | — | 15.4° (8.6) | simple carbon-
dioxated | Takarazuka | 寶塚(") | Granite | Tansan-sen <i>A</i> |
| 33 | 6.70 | 1.84 | — | 59.2° | earth-muriated
common salt | Kinosaki | 城崎(") | Tertiary | Ichi-no-yu <i>Vent C</i>
(<i>Sawano-yu</i>) |
| 34 | 6.45 | 1.77 | — | 51.1° | salt | Sakunami | 作並(宮城) | " | Taka-no-yu |
| 35 | 6.02? | 1.66 | 1400 | 56.0° | simple | Ao-ne | 青根(") | " | Shin-yu |
| 36 | 5.91 | 1.63 | — | 49.0° | sulphated bitter
sulphur | Yamanaka | 山中(石川) | Granite | Kiku-no-yu |
| 37 | 5.60 | 1.54 | 432 | 24.5° | carbonated | Kōbe | 神戸(兵庫) | Granite | Tennō-Kōsen |
| 38 | 5.51 | 1.52 | — | 50.5° | earth-muriated
common salt | Kinosaki | 城崎(") | Tertiary | Ichi-no-yu <i>Vent A</i> |
| 39 | 5.47 | 1.50 | — | 76.0° | " | Wakura | 和倉(石川) | " | Sō-yu |
| 40 | 4.92 | 1.45 | — | 54.2° | salt | Tōgatta | 遠刈田(宮城) | Liparite | Kamasaki-no-yu |
| 41 | 4.85 | 1.33 | 720 | 19.0° | carbonated | Kōbe | 神戸(兵庫) | Granite | Suwayama Tansan-sen |
| 42 | 4.47 | 1.23 | — | 17.4° | " | Arima | 有馬(") | Quartz Porph. | Sugigatani Tansan-sen |
| 43 | 4.44 | 1.22 | — | 46.0° | salt | Tōgatta | 遠刈田(宮城) | Liparite | Special bath in Naka-no-yu |
| 44 | 4.34 | 1.19 | 900 | 18.5° | carbonated | Kōbe | 神戸(兵庫) | Granite | Nunobiki Tansan-sen |
| 45 | 4.28 | 1.18 | 207 | 48.6° | earth-muriated
common salt | Kinosaki | 城崎(") | Tertiary | Jizō-no-yu |
| 46 | 4.17 | 1.15 | 450 | 69.1° | salt | Sakunami | 作並(宮城) | " | Taki-no-yu |
| 47 | 4.11 | 1.13 | 152 | 76.0° | earth-muriated
common salt | Awara | 蘆原(福井) | Alluvium | Artes. Well of Haiya-no-yu |
| 48 | 4.03 | 1.11 | — | 50.6° | " | Kinosaki | 城崎(兵庫) | Tertiary | Mandara-no-yu <i>Vent A</i> |
| 49 | 3.99 | 1.10 | — | 40.5° | saline common salt | Kamasaki | 鎌先(宮城) | Andesite | Mogamiya-no-yu |
| 50 | 3.97 | 1.09 | — | 47.0° | sulphated bitter
sulphur | Yamanaka | 山中(石川) | Tertiary | Ashi-no-yu |
| 51 | 3.84 | 1.06 | — | 48.0° | salt | Tōgatta | 遠刈田(宮城) | Liparite | Me-no-yu |
| 52 | 3.69 | 1.02 | — | — | " | Takarazuka | 寶塚(兵庫) | Granite | Drinking water |
| 53 | 3.60 | 0.99 | 504 | 25.0° | salt | Tatsunokuchi | 辰之口(石川) | Tertiary | Tatsunokuchi Min. Spr. |
| 54 | 3.41 | 0.94 | — | 13.0° | " | Fukatani | 深谷(") | " | Moto-yu |
| 55 | 3.31 | 0.91 | — | 50.5° | earth-muriated
common salt | Kinosaki | 城崎(兵庫) | " | Mandara-no-yu <i>Vent D</i> |
| 56 | 3.02 | 0.83 | — | 32.0° (25) | salt | Yugawa | 湯川(宮城) | — | Yugawa-Onsen |
| 57 | 2.77 | 0.76 | 450 | 53.2° | " | Sakunami | 作並(") | Tertiary | Kawara-no-yu |
| 58 | 2.75 | 0.76 | — | 41.2° | common salt | Akyu | 秋保(") | " | " |
| 59 | 2.74 | 0.75 | — | 13.0° | salt | Fukatani | 深谷(石川) | " | Oku-no-yu |
| 60 | 2.71 | 0.75 | 380 | 74.2° | acid alum vitriol | Myōdan | 明礬(大分) | Andesite | Jizō-no-yu (<i>Utsunoyu</i>) |
| 61 | 2.66 | 0.73 | — | 39.0° | sulphated com-
mon salt | Yuwaku | 湯涌(石川) | Tertiary | Shin-yu |
| 62 | 2.61 | 0.72 | — | 55.0° | sulphur | Awazu | 粟津(") | Liparite | Artes. Well of Kamiya |
| 63 | 2.58 | 0.71 | 1433 | 57.1° | salt | Gaga | 峨宮(宮城) | Gneiss | Gaga-Onsen |
| 64 | 2.48 | 0.68 | — | 15.0° (29°) | " | Kawasaki | 川崎(") | — | Kawasaki-Onsen |
| 65 | 2.38 | 0.66 | — | 10.5° | simple carbon-
dioxated | Arima | 有馬(兵庫) | Quartz Porph. | Jigokudani Tansan-sen |
| 66 | 2.37 | 0.65 | — | 62.1° | simple | Hokuto | 北投(臺北) | Tertiary | Tetsu-no-yu |
| 67 | 2.03 | 0.56 | — | 54.5° | earth-muriated
common salt | Kinosaki | 城崎(兵庫) | " | Mandara-no-yu <i>Vent C</i> |
| 68 | 2.02 | 0.56 | 300 | 17.1° (9.9°) | " | Takarazuka | 寶塚(") | Granite | Takarazuka-Kōsen |
| 69 | 1.98 | 0.55 | 1400 | 51.4° | simple | Ao-ne | 青根(宮城) | Tertiary | Myōgō-no-yu |

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|------------------------------|--------------------------------|---|-----------------------|---|---------------------|---------|---------------|-----------------------------|
| | in 10 ⁻⁹ curies | in Mach's units (recalculated) | | | | | | | |
| 70 | 1.98 | 0.55 | 1584 | 52.8° | common salt | Akya | 秋保(宮城) | Tertiary | Mi-yu |
| 71 | 1.91 | 0.53 | — | 51.2° | — | Beppu | 別府(大分) | Andesite | Sunayyu at sea coast |
| 72 | 1.79 | 0.49 | — | 43.0° | saline common salt | Kamasaki | 鎌先(宮城) | " | Kimuraya-no-yu |
| 73 | 1.73 | 0.48 | — | 41.5° | — | Arima | 有馬(兵庫) | Quartz Porph. | Valley water, Yubuna |
| 74 | 1.72 | 0.47 | — | — | — | Nashimotoya | 西宮() | Granite | Well water for sake |
| 75 | 1.70 | 0.47 | — | 71.0° | saline bitter sulphur | Yamashiro | 山代(石川) | Tertiary | Source in Araya |
| 76 | 1.54 | 0.42 | — | 43.6° | earth-minerated non-carbonate common salt | Arima | 有馬(兵庫) | Quartz Porph. | Hana-no-yu |
| 77 | 1.54 | 0.42 | 162 | 47.5° | simple | Beppu | 別府(大分) | Andesite | Azenashi no ya (V. 100-10) |
| 78 | 1.54 | 0.42 | — | 48.5° | " | Hokuto | 北投(臺北) | Tertiary | Kaikōsha Yūdaki-no-yu |
| 79 | 1.52 | 0.42 | 520 | 57.0° | simple carbon-dioxated | Beppu | 別府(大分) | Andesite | Reichō-sen |
| 80 | 1.52 | 0.42 | 3150 | 48.0° | saline common salt | Kamasaki | 鎌先(宮城) | " | Ichijō-no-yu |
| 81 | 1.50 | 0.41 | — | 15.5° | carbon-dioxated sulphur | Arima | 有馬(兵庫) | Quartz Porph. | Hana-no-bō cold Min. W. |
| 82 | 1.47 | 0.40 | 648 | 53.0° | iron carbonate | Beppu | 別府(大分) | Andesite | Furō-sen, Artes. Well |
| 83 | 1.43 | 0.39 | — | 22.0° | common salt | Daigamari | 臺森(宮城) | — | Daigamori-Kōsen |
| 84 | 1.41 | 0.39 | — | 57.1° | sulphur | Naruko | 鳴子() | Andesite | Taga-no-yu |
| 85 | 1.39 | 0.38 | — | 70.5° | " | Hokutō | 北投(臺北) | Volc. Detr. | Spring G |
| 86 | 1.36 | 0.37 | — | 55.0° | simple carbon-dioxated | Myōban | 明礬(大分) | Andesite | Tobi-yu |
| 87 | 1.34 | 0.37 | — | 40.0° | simple | Kimpōri | 金包里(臺北) | Tertiary | Satsumaya Uchi-yu |
| 88 | 1.34 | 0.37 | 7258 | 42.0° | " | Ao-ne | 青根(宮城) | " | Ō-yu |
| 89 | 1.30 | 0.36 | — | 45.0° | " (?) | Kannawa | 鐵輪(大分) | Andesite | Kumi-yu? |
| 90 | 1.28 | 0.35 | — | 56.0° | " | Hokutō | 北投(臺北) | Tertiary | Kaikōsha-waki-no-yu |
| 91 | 1.26 | 0.35 | — | 35.7° | alkaline common salt | Kwanshinei | 關仔嶺(嘉義) | " | Saka-no-yu |
| 92 | 1.25 | 0.34 | — | 32.5° | — | Arima | 有馬(兵庫) | Quartz Porph. | Hana-no-bō Uchi-yu |
| 93 | 1.21 | 0.33 | 180 | 63.0° | simple | Iizaka | 飯坂(福島) | Tertiary | Taki-no-yu |
| 94 | 1.19 | 0.33 | — | 34.5° | earth-minerated non-carbonate common salt | Arima | 有馬(兵庫) | Quartz Porph. | Mearai-yu |
| 95 | 1.19 | 0.33 | 225 | 33.0°? | sulphur | Hotta | 堀田(大分) | Andesite | Hotta-no-yu |
| 96 | 1.19 | 0.33 | — | 64.1° | salt | Shinkuruma | 新車(宮城) | " | A pump in Takashige Hotel |
| 97 | 1.15 | 0.32 | — | 41.0° | simple (?) | Beppu | 別府(大分) | " | Asami-Onsen |
| 98 | 1.11 | 0.31 | 108 | 46.0° | " | Iizaka | 飯坂(福島) | Tertiary | Kintaki-no-yu |
| 99 | 1.03 | 0.28 | — | 80.3° | salt | Urai | ウライ(臺北) | " | Spring B (on the left bank) |
| 100 | 1.01 | 0.28 | — | 23.0° | simple carbon-dioxated | Su-ō | 蘇澳(宜蘭) | Clay Slate | Tansan-sen |
| 101 | 1.01 | 0.28 | 270 | 49.1° | simple | Onikobe | 鬼首(宮城) | Tertiary | Todoroki-Onsen |
| 102 | 1.00 | 0.28 | 216 | 50.0° | iron carbonate | Kwankaigi | 觀海寺(大分) | Andesite | Ō-yu (Kwanakōji Onsen) |
| 103 | 1.00 | 0.27 | undant | — | simple carbon-dioxated | Izunayama | 飯網山(長野) | Volc. Detr. | Carb. Spr. N. 1 |
| 104 | 0.99 | 0.27 | — | 28.0° | — | Komatsugura | 小松倉(宮城) | — | Kinzan-Onsen |
| 105 | 0.97 | 0.27 | — | 47.0° | simple | Beppu | 別府(大分) | Andesite | Waki-yu in Hinago |
| 106 | 0.95 | 0.26 | undant | 41.0° | sulphur | Naruko | 鳴子(宮城) | " | Taki-no-yu |
| 107 | 0.95 | 0.26 | — | — | — | Arima | 有馬(兵庫) | Quartz Porph. | Drinking W., Zaihōji |
| 108 | 0.94 | 0.26 | — | 53.4° | earth-minerated non-carbonate common salt | " | " | " | Ni-no-yu |
| 109 | 0.94 | 0.26 | 198 | 57.0° | common salt | Kamegawa | 龜川(大分) | Andesite | Shi-no-yu |
| 110 | 0.93 | 0.26 | — | 14.0° | — | Miyatoko | 宮床(宮城) | — | Yamada-Onsen |
| 111 | 0.90 | 0.25 | 270 | 53.1° | simple | Obara | 小原() | — | Shin-yu |
| 112 | 0.90 | 0.25 | 648 | 47.0° | carbonated | Kawatabi | 川渡() | Tertiary | Mayu-no-yu |
| 113 | 0.87 | 0.24 | 54 | 38.0° | simple | Jōgi | 定義() | Andesite | Jōgi-Onsen |
| 114 | 0.84 | 0.23 | — | 32.5° | — | Kōnosu | 鴻巣() | — | Kōnosu-Onsen |
| 115 | 0.82 | 0.23 | — | 95.0° | sulphur | Hokutō | 北投(臺北) | Tertiary | Spring A |
| 116 | 0.81 | 0.22 | — | 53.4° | earth-minerated non-carbonate common salt | Arima | 有馬(兵庫) | Quartz Porph. | Ichino-yu |
| 117 | 0.81 | 0.22 | — | 27.2° | — | Kangane | 神峯(宮城) | — | Kangane-Onsen |
| 118 | 0.80 | 0.22 | — | 49.2° | carbonated | Beppu | 別府(大分) | Andesite | Takegawara-no-yu |
| 119 | 0.76 | 0.21 | — | 13.0° | salt | Fukatani | 深谷(石川) | Tertiary | Shin-yu |
| 120 | 0.76 | 0.21 | — | 61.5° | carbonated | Alayu Matsuzaki-ken | 赤湯(宮城) | Andesite | Shita-no-yu |

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|------------------------------|----------------------------------|---|-----------------------|--|-------------------|---------|-------------|----------------------------|
| | in cubic | in Maché's units (re-calculated) | | | | | | | |
| 121 | 0.76 | 0.21 | ca. 126 | 48.5° | saline bitter | Yoshina | 吉奈(静岡) | Andesite | Tōfuya-no-yu |
| 122 | 0.75? | 0.21 | — | 13.5° | — | Iizaka | 飯坂(福島) | Tertiary | Mitarase-Well |
| 123 | 0.73 | 0.20 | 270 | 54.0° | common salt | Kamegawa | 龜川(大分) | Andesite | Source in Kiyō-sen |
| 124 | 0.71? | 0.19 | 1,320 | 46.0° | iron carbonate | Beppu | 別府()) | " | Kusunoki-Onsen |
| 125 | 0.70 | 0.19 | — | 56.2° | salt | Onikōbe | 鬼首(宮城) | — | Shin-Mitaki-yu |
| 126 | 0.70 | 0.19 | 265 | 45.0° | saline bitter | Yoshina | 吉奈(静岡) | Andesite | Ō-yu |
| 127 | 0.70 | 0.19 | — | 26.0° (11°) | saline bitter earthy alkaline iron carbonate common salt | Hirano | 平野(兵庫) | Granite | Hirano Tansan-sen |
| 128 | 0.70 | 0.19 | 216 | 75.0° | iron carbonate | Shibaseki | 芝石(大分) | Andesite | Hijiri-yu |
| 129 | 0.70 | 0.19 | 270 | 58.0° | sulphur | Naruko | 鳴子(宮城) | " | Genzō-yu |
| 130 | 0.69 | 0.19 | — | 30.0° (28°) | — | Yunoheda | 湯邊田()) | — | Yunoheda-Onsen |
| 131 | 0.68 | 0.19 | — | 48.3° | — | Kawatabi | 川渡()) | Tertiary | Kagaya-Shin-yu |
| 132 | 0.66 | 0.18 | — | 93.5° | sulphur | Hokutō | 北投(臺北) | " | Spring D |
| 133 | 0.64 | 0.18 | — | 79.3° | simple | Tōi | 頭圍(宜蘭) | Alluvium | Spring in field |
| 134 | 0.64 | 0.18 | — | 48.0° | — | Kawatabi | 川渡(宮城) | Tertiary | Fujishimaya-Shin-yu |
| 135 | 0.62 | 0.17 | — | 49.7° | — | Hokutō | 北投(臺北) | — | Spring B |
| 136 | 0.58 | 0.16 | 144 | 58.0° | simple | Iizaka | 飯坂(福島) | Tertiary | Akagawa-no-yu |
| 137 | 0.55 | 0.15 | abundant | — | simple carbon-dioxated | Izunayama | 飯網山(長野) | Volc. Detr. | Carb. Spr. No. 7 |
| 138 | 0.54 | 0.15 | — | 57.0° | carbonated | Akayu, Miyagi-ken | 赤湯(宮城) | Andesite | Goten-no-yu (Ue-no-yu) |
| 139 | 0.49? | 0.14 | 108 | 59.0° | simple | Iizaka | 飯坂(福島) | Tertiary | Hako-yu |
| 140 | 0.48 | 0.13 | 1167 | 57.0° | common salt | Beppu | 別府(大分) | Andesite | Higashi-Onsen, Artes. Well |
| 141 | 0.48 | 0.13 | — | 44.0° | simple | Iizaka | 飯坂(福島) | Tertiary | Kotaki no-yu |
| 142 | 0.48 | 0.13 | — | 16.0° (27°) | common salt | Daigamori | 臺ヶ森(宮城) | — | Daigamori-Shin-yu |
| 143 | 0.47 | 0.13 | 54 | 55.0° | simple | Iizaka | 飯坂(福島) | Tertiary | Tōza-yu |
| 144 | 0.47 | 0.13 | — | 42.4° | — | Kawatabi | 川渡(宮城) | " | Me-no-yu (Musaka-no-yu) |
| 145 | 0.46 | 0.13 | — | 91.5° | sulphur | Hokutō | 北投(臺北) | " | Spring C |
| 146 | 0.44 | 0.12 | 51 | 54.0° | simple | Iizaka | 飯坂(福島) | " | Akagawabata-no-yu |
| 147 | 0.42 | 0.12 | 54 | 56.4° | salt | Naruko | 鳴子(宮城) | Andesite | Kawara-no-yu |
| 148 | 0.40 | 0.11 | 50 | 54.0° | simple | Iizaka | 飯坂(福島) | Tertiary | Sabako-no-yu |
| 149 | 0.40 | 0.11 | — | 90.0° | sulphur | Hokutō | 北投(臺北) | " (?) | Spring A |
| 150 | 0.38? | 0.11 | — | 53.0° | simple | Tōi | 頭圍(宜蘭) | Alluvium | Nishiyama's Artes. Well |
| 151 | 0.35 | 0.10 | — | 52.5° | — | Niizeki | 新關(宮城) | — | Niizeki-Onsen |
| 152 | 0.34 | 0.09 | — | 73.5° | salt | Urai | ウライ(臺北) | Tertiary | Spring (on the right bank) |
| 153 | 0.26 | 0.07 | — | 64.5° | alkaline common salt | Kwanshirei | 關仔嶺(嘉義) | " | Ue-no-yu |
| 154 | 0.21 | 0.06 | — | 53.2° | salt | Onikōbe | 鬼首(宮城) | — | Kyū-Mitaki-yu |
| 155 | 0.19 | 0.05 | — | 40-36.5° | " | Kimpōri | 金包里(臺北) | Tertiary | Honkura |
| 156 | 0.16 | 0.05 | abundant | 46.8° | sulphated bitter | Izusan | 伊豆山(静岡) | Andesite | Hashiri-yu |
| 157 | 0.14 | 0.04 | — | ca. 100.0° | salt | Onikōbe | 鬼首(宮城) | Tertiary | Miyazawa Shita-no-yu |
| 158 | 0.14 | 0.04 | — | 63.5° | simple | Obara | 小原()) | — | Shita-no-yu of Furu-yu |
| 159 | 0.13 | 0.04 | — | 74.0° | common salt | Yugawara | 湯河原(静岡) | Andesite | Hakoneya-Uchi-yu |
| 160 | 0.11 | 0.03 | — | 57.8° | — | Atami | 熱海()) | " | Muen-no-yu |
| 161 | 0.11 | 0.03 | — | 84.5° | simple | Shinkuruma | 新車(宮城) | " | Chiyo-no-yu |
| 162 | 0.10 | 0.03 | — | 84.0° | earth-mixed common salt | Atami | 熱海(静岡) | " | Kawara-no-yu |
| 163 | 0.08? | 0.02 | — | 93.0° | acid vitriol | Kannawa | 鐵輪(大分) | " | Umi-Jigoku |
| 164 | 0.08 | 0.02 | — | 79.5° | sulphur | Hokutō | 北投(臺北) | Tertiary | Spring F |
| 165 | 0.08 | 0.02 | — | ca. 100.0° | salt | Onikōbe | 鬼首(宮城) | " | Water from Fukage Geyser |
| 166 | 0.05 | 0.01 | — | 65.0° | — | Hokutō | 北投(臺北) | Volc. Detr. | Sekimon, Yukawa |
| 167 | 0.03 | 0.01 | — | 29.0° | — | " | " | " | Valley water |
| 168 | 0.02 | 0.01 | — | 52.0° | — | " | " | " | Third Fall, Yukawa |
| 169 | undetectable-smell | — | — | 27.0° | carbonated | Ayashi | 愛子(宮城) | — | Ayashi Tansan-sen |
| 170 | 0 | 0 | — | 100.0° | sulphur | Nakayama | 中山()) | Andesite | Hoshi-no-yu (Ihbi-yu) |
| 171 | 0 | 0 | — | ca. 100.0° | salt | Onikōbe | 鬼首()) | Tertiary | Miyazawa Taki-no-yu |
| 172 | 0 | 0 | — | 100.0° | alkaline sulphur earth-mixed common salt | Naruko | 鳴子()) | Andesite | Unagi-no-yu |
| 173 | 0 | 0 | 1296 | ca. 108.0° | — | Atami | 熱海(静岡) | " | Water from Ō-yu Geyser |

| No. | Emanation per litre of gas at 0° C. | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Gas evolving from: |
|-----|-------------------------------------|---------------------------------|---|-----------------------|---|-------------|---------|---------------|-----------------------|
| | in 10 curies | in Mache's units (recalculated) | | | | | | | |
| 1 | 24.30 | 6.68 | — | 71.0 | saline bitter sulphur | Yama-hiro | 山代(石川) | Tertiary | Source in Araya |
| 2 | 26.90 | 7.40 | — | 95.0 | — sulphur | Hokuto | 北投(臺北) | " | Spring 4 |
| 3 | 17.70 | 4.87 | — | 17.4 | carbon dioxide sulphur | Arima | 有馬(兵庫) | Quartz Porph. | Sugigatani Tansan-sen |
| 4 | 17.25 | 4.74 | — | 52.8° | common salt | Akyu | 秋保(宮城) | Tertiary | Mi-yu |
| 5 | 15.70 | 4.32 | — | 53.4° | earth-muriated non-carbonate common salt | Arima | 有馬(兵庫) | Quartz Porph. | Ichi-no-yu |
| 6 | 13.50 | 3.71 | — | 10.5 | simple carbon dioxide sulphur | " | " | " | Jigokudani Tansan-sen |
| 7 | 12.50 | 3.44 | — | 93.5° | — sulphur | Hokuto | 北投(臺北) | Tertiary | Spring 1) |
| 8 | 6.65 | 1.83 | — | 17.1° | carbon dioxide common salt | Takarazuka | 寶塚(兵庫) | Granite | Takarazuka-Kōsen |
| 9 | 6.46 | 1.78 | — | 33.0° | — sulphur | Hotta | 堀田(大分) | Andesite | Hotta-no-yu |
| 10 | 6.33 | 1.74 | — | 48.5° | saline bitter | Yoshina | 吉奈(静岡) | " | Tōfuya-no-yu |
| 11 | 2.63 | 0.72 | — | 23.0° | simple carbon dioxide | Suō | 蘇澳(宜蘭) | Clay Slate | Tansan-sen |
| 12 | 2.31 | 0.64 | — | 40° 30.5° | — | Kimpōri | 金包里(臺北) | Tertiary | Honkura |
| 13 | 2.12 | 0.58 | — | 20.0° 11° | earthly alkaline non-carbon dioxide common salt | Hirano | 平野(兵庫) | Granite | Hirano Tansan-sen |
| 14 | 1.96 | 0.54 | — | 64.5° | alkaline common salt | Kwan-shūrei | 關仔嶺(嘉義) | Tertiary | Ue-no-yu |
| 15 | 0.53 | 0.15 | — | — | — | Hokuto | 北投(臺北) | Volc. Rock | A Spring in Solfatara |
| 16 | † small quantity | | — | ca. 100° | earth-muriated common salt | Atami | 熱海(静岡) | Andesite | Ō-yu Geyser |
| 17 | † small quantity | | — | ca. 100° | — salt | Onikobe | 鬼首(宮城) | Tertiary | Fukiage Geyser |

† The determination was made with steam ejected from springs.

Table 5.

The quantity of radium emanation, the temperature of springs, the flow of water, and the amount of gas† produced from springs in the following table were determined by Hazime Ikeuti (Physical Faculty of Tōkyō Imperial University). As for particulars refer to Proceeding of the Tōkyō Mathematico-Physical Society (Tōkyō Sūgaku-Buturiggakkai Kizi), 2nd Ser., Vol. VII, No. 10, p. 177-180; No. 21, p. 423-426.

For recalculation of Mache's unit from the international unit following factor was used:

$$1 \times 10^{-10} \text{ curie} = 0.275 \text{ Mache's units.}$$

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|------------------------------|---------------------------------|---|-----------------------|----------------------------|---------------|-----------|----------|--------------------|
| | in 10 curies | in Mache's units (recalculated) | | | | | | | |
| 1 | 173.00 | 47.58 | ca. | 25.5° | simple | Murasugi | 村杉(新潟) | Granite | Murasugi-Kōsen |
| 2 | 2.57 | 0.71 | — | 72.0° | earth-muriated common salt | Matsunoyama | 松之山(新潟) | Tertiary | Netsu-no-yu |
| 3 | 2.56 | 0.70 | — | 65.0° | " | " | " | " | Ihii-no-yu |
| 4 | 2.47 | 0.68 | — | 73.0° | " | " | " | " | Washi-no-yu |
| 5 | 1.76 | 0.48 | — | 44.0° | " | " | " | " | Fukushimaya-no-yu |
| 6 | 1.59 | 0.44 | 111 | 42.0° | saline bitter | Yoshina | 吉奈(静岡) | Andesite | Tōfuya-no-yu No. 2 |
| 7 | 1.58 | 0.44 | — | 65.0° | sulphated common salt | Shibu [Hirao] | 澁[平穩](長野) | Diluvium | Ō-yu No. 2 |
| 8 | 1.37 | 0.37 | — | 13.0° | — | Matsunoyama | 松之山(新潟) | Tertiary | Ushiyashiki-no-yu |
| 9 | 1.26 | 0.35 | — | 49.0° | saline bitter | Yoshina | 吉奈(静岡) | Andesite | Ō-yu |
| 10 | 1.22 | 0.34 | 288 | 54.0° | earthly alkaline sulphur | Akakura | 赤倉(新潟) | " | Shin-yu |
| 11 | 1.12 | 0.31 | — | 48.0° | sulphated common salt | Shibu [Hirao] | 澁[平穩](長野) | Diluvium | Solfatara uchi-yu |
| 12 | 1.12 | 0.31 | — | 25.0° | — | Matsunoyama | 松之山(新潟) | Tertiary | Yuziri-no-yu |
| 13 | 1.03 | 0.28 | — | 36.0° | sulphur | Tsubane | 燕(新潟) | Andesite | Chi-no-ike-no-yu |

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|------------------------------------|---------------------------------|---|-----------------------|---------------------------|---------------|-------------|----------|---------------------|
| | in 10 ¹⁰ curies | in Mache's units (recalculated) | | | | | | | |
| 14 | 0.85 | 0.23 | 93 | 46.5° | saline bitter | Yoshina | 吉 奈(静岡) | Andesite | Kikusuitei-no-yu |
| 15 | 0.78 | 0.22 | — | 74.0° | sulphated common salt | Shibu [Hirao] | 澁 [平穩] (長野) | Diluvium | Ō-yu No. 1 |
| 16 | 0.68 | 0.19 | 100 | 52.5° | saline bitter | Yoshina | 吉 奈(静岡) | Andesite | Tofuya-no-yu No. 1 |
| 17 | 0.43 | 0.12 | — | — | sulphated common salt | Shibu [Hirao] | 澁 [平穩] (長野) | Diluvium | Wata-no-yu |
| 18 | 0.24 | 0.07 | — | 45.0° | sulphur | Tsubame | 燕 (新潟) | Andesite | Iwa-no-yu |
| | Emanation per litre of gas at 0°C. | | | | | | | | Gas evolving from : |
| 1 | 11.25 | 3.09 | 14.4 | 51.0° | entirely alkaline sulphur | Akakura | 赤 倉(新潟) | Andesite | Shin-yu |
| 2 | 2.29 | 0.63 | — | — | sulphur | Tsubame | 燕 () | " | Kawara-yu |

Table 6.

The quantity of radium emanation, the temperature of springs, the flow of water, and the amount of gas² produced from springs in the following table were determined by Suminosuke Ono, Prof. of Physics, Tōkyō Higher Normal School. As for particulars refer to Proceeding of the Tōkyō Mathematico-Physical Society (Tōkyō Sūgaku-Buturigakkwai Kizi), 2nd Ser., Vol. VII, No. 21, p. 421-422.

For recalculation of Mache's unit from the international unit following factor was used :

$$1 \times 10^{-10} \text{ curie} = 0.275 \text{ Mache's units.}$$

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|------------------------------------|---------------------------------|---|-----------------------|----------------|----------|------------|-----------------|-------------------------------|
| | in 10 ¹⁰ curies | in Mache's units (recalculated) | | | | | | | |
| 1 | 37.1 | 10.20 | 916 | 40.0° | — | Agune | 阿 久 根(鹿兒島) | Alluvium | Agune-Onsen No. 1 |
| 2 | 34.5 | 9.49 | — | 40.2° | — | " | " | " | " No. 2 |
| 3 | 34.3 | 9.43 | 1166 | 41.0° | — | " | " | " | " No. 3 |
| 4 | 33.4 | 9.19 | — | 40.0° | — | " | " | " | " No. 4 |
| 5 | 14.2 | 3.91 | 233 | 47.5° | carbonated | Hinagu | 日 奈 久(熊本) | Cretaceous | Gozen-yu |
| 6 | 13.8 | 3.80 | 389 | 48.5° | " | " | " | " | Otsugi-yu (Otoke-yu) |
| 7 | 10.5 | 2.89 | 147 | 47.0° | " | " | " | " | " (Onna-yu) |
| 8 | 4.6 | 1.27 | 864 | 50.0° | salt | Hayashi | 林 () | Alluvium | Moto-yu |
| 9 | 4.2 | 1.16 | 188 | 48.4° | " | " | " | " | Suiranrō-no-yu 1st Well |
| 10 | 3.2 | 0.88 | 156 | 45.2° | carbonated | Hinagu | 日 奈 久() | Cretaceous | Gata-yu (Onna-yu) |
| 11 | 3.2 | 0.88 | 475 | 48.8° | salt | Hayashi | 林 () | Alluvium | Suiranrō-no-yu 2nd Well |
| 12 | 3.1 | 0.85 | 205 | 45.2° | carbonated | Hinagu | 日 奈 久() | Cretaceous | Gata-yu (Otoke-yu) |
| 13 | 2.8 | 0.77 | 340 | 48.5° | " | " | " | " | Izumiya-no-yu |
| 14 | 2.1 | 0.58 | 97 | 46.5° | " | " | " | " | Kimparō-no-yu 1st Well |
| 15 | 1.4 | 0.39 | 180 | 45.0° | " | " | " | " | " mixture of 2nd and 3rd Well |
| 16 | 0.4 | 0.11 | ? | 39.0° | " | Yoshio | 吉 尾() | Chichibu Series | Yoshio-Onsen |
| | Emanation per litre of gas at 0°C. | | | | | | | | Gas evolving from : |
| 1 | 27.8 | 7.65 | 153 | 48.4° | " | Hayashi | 林 (熊本) | Alluvium | Suiranrō-no-yu 1st Well |
| 2 | 21.2 | 5.83 | 156 | 50.0° | " | " | " | " | Moto-yu |
| 3 | 20.5 | 5.64 | 333 | 48.8° | " | " | " | " | Suiranrō-no-yu 2nd Well |

Table 7.

The quantity of radium emanation and the temperature of springs in the following table were determined by Dr. H. Kondō, [vide Official Report of Kagoshima Prefecture ("Kagoshima-ken Kōhō"), Aug., 1913].

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in °C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|------------------------------|-----------------|---|------------------------|-------------------|--------------|----------|------------------------|----------------------------|
| | in 10 ⁻¹⁰ curies | in Mach's units | | | | | | | |
| 1 | 7.05 | 1.91 | ca. | 51.0° | hydrogen sulphide | Iwō-lani | 硫黄谷(鹿兒島) | Volc. Ash and Mud-Lava | Iwō-lani Onsen No. 1 |
| 2 | 3.69 | 1.00 | — | 65.0° | salt | Suri no-hama | 睨之濱(") | " | Surinohama Onsen (No. 100) |
| 3 | 3.35 | 0.90 | — | 66.5° | hydrogen sulphide | Eino-o | 榮之尾(") | " | Eino-o Onsen No. 1 |
| 4 | 3.33 | 0.90 | — | 46.0° | salt | Myōban-yama | 明礬山(") | " | Myōban-yama Onsen No. 5 |
| 5 | 3.21 | 0.87 | — | 59.5° | " | Iwōdani | 硫黄谷(") | " | Iwōdani Onsen No. 5 |
| 6 | 2.72 | 0.74 | — | 39.8° | iron carbonate | Anraku | 安楽(") | " | Spring A |
| 7 | 2.63 | 0.71 | — | 64.2° | salt | Myōban-yama | 明礬山(") | " | Myōban-yama Onsen No. 3 |
| 8 | 2.24 | 0.60 | — | 30.0° | carbonated | Shiose | 鹽浸(") | " | Shimo-no-yu No. 3 |
| 9 | 2.01 | 0.54 | — | 54.0° | salt | Yaji | 彌次(") | " | Yaji-ga-Onsen No. 1 |
| 10 | 1.60 | 0.53 | — | 68.0° | hydrogen sulphide | Myōban-yama | 明礬山(") | " | Myōban-yama-Onsen No. 2 |
| 11 | 1.54 | 0.42 | — | 48.7° | " | Iwōdani | 硫黄谷(") | " | Iwōdani-Onsen No. 4 |
| 12 | 1.50 | 0.41 | 000 | 47.0° | carbonated | Myōken | 妙見(") | " | Myōken-yu |
| 13 | 1.25 | 0.34 | — | 53.5° | iron carbonate | Anraku | 安楽(") | " | Shin-yu |
| 14 | 1.22 | 0.33 | — | 48.0° | " | " | " | " | Kyū-yu No. 1 |
| 15 | 1.14 | 0.31 | 64.50 | 42.0° | common salt | Shibadate | 芝立(") | " | Shibadate-Onsen |
| | Emanation per litre of gas | | | | | | | | Gas evolving from : |
| 1 | 7.33 | 1.98 | — | 47.0° | carbonated | Myōken | 妙見(鹿兒島) | " | Myōken-yu |

Table 8.

Radium Emanation in Petroleum from Nishiyama Oil Field in the Province of Etōgo.

(Tōkyō Sūgaku-Bunrō-shū, Kei, 2nd Ser., Vol. VIII, No. 1, p. 1.)

| Date of exp. | Well | Radium emanation per litre | | Observer |
|------------------|--------------------|---------------------------------|---------------------------------|--------------|
| | | of petroleum | of gas | |
| 27-29 Nov., 1914 | R. No. 22, Takiya | 1.66 × 10 ⁻¹⁰ curies | 0.35 × 10 ⁻¹⁰ curies | Kyōtoku Tōjō |
| 30 Nov., 1914 | R. No. 7, Nagamine | 1.44 " | — | " |

Radio-activity of Notable Spring Waters in Europe and in the United States of America.

The following tables give the results of measurements of the radio-activity of several well-known European mineral waters and of a few American waters. Many of these waters also contain radio-active gases. Reports of different observations on the same spring indicate that the waters and the gases in them vary greatly in radio-activity. Because of differences in the methods of measurement and in allowance for disintegration products and also because of the uncertain or arbitrary nature of some of the units of expression the figures have not been reduced to a common unit for direct comparison.

We have therefore recorded in these tables only the numbers from authentic origins which enable direct comparison. In Remarks are given geology, flow of water in 24 hours, and literature.

Abbreviations used in the tables :

Sommer.....Über die Radioaktivitätsverhältnisse der natürlichen Heilquellen des deutschen Sprachgebietes. Von Prof. Dr. Ernst Sommer.
Lazarus Handbuch der Radium-Biologie und Therapie von Prof. Dr. Paul Lazarus, 1913.
Curie Die Radioaktivität von M^{me} P. Curie, 1912. Band II.
Weidig Radioaktive Wässer in Sachsen von Dr.-Ing. M. Weidig, 1912. IV. Teil.
Ch. Z.....Chemiker Zeitung.
Ph. Z.....Physikalische Zeitschrift.
Bot. Gaz. National Bottler's Gazette.

Table 9.

(A) Results of measurements expressed in Mache's units.

| No. | Emanation per litre of water in Mache's units | Temp. of spring in C. | Classification | Location | | Altitude in ft. | Spring | Observer | Remarks |
|-----|---|-----------------------|--------------------------------|------------------------|-------------|-----------------|-----------------------------------|--------------------|--|
| 1 | 2400-1200 | — | — | Joachimsthal | Austria | 2100 | Mine water (Depth 500 ft.) | J. Stép | Ch. Z. 1912, p. 1470 |
| 2 | 2050.00 | — | — | " | " | " | Mine water | Schmidt | Ph. Z. 1907, p. 4 |
| 3 | 1964.40 | 7.0° | iron carbon-dioxide | Brambach | Germany | — | Neue Quelle | Weidig 1911 | 215 ft.; Contact Zone of Granite and Slate; Weidig, p. 102 |
| 4 | 600.00 | — | — | Joachimsthal | Austria | 2100 | Wernerlaufquelle | J. Stép 1905 | Sommer, p. 25 |
| 5 | 486.90 | 7.4° | alkaline | Brambach | Germany | — | Schillerquelle | Weidig 1912 | 18 ft.; Contact Zone of Granite and Slate; Weidig, p. 201 |
| 6 | 372.00 | 57.0° | mineral alkaline | Laevo Anneno in Ischia | Italy | — | Altrömische Quelle | Engler | Ch. Z. 1907, p. 813 |
| 7 | 366.90 | 8.5° | iron carbonate | Brambach | Germany | — | Grenzquelle | Weidig 1911 | Several litres per min.; Contact Zone of Granite and Slate; Weidig, p. 301 |
| 8 | 206.00 | 29.0° | sulphur | Landeck | " | 1500 | Georgenquelle | Schäfer 1909 | Gneiss; Sommer, p. 29 |
| 9 | 155.00 | 30.3° | simple | Gastein | Austria | 3400 | Grabenbäckerquelle | Mache | Ch. Z. 1907, p. 813 |
| 10 | 149.00 | 30.3° | " | " | " | " | " | Engler & Sieveking | " |
| 11 | 140.20 | 46.0° | " | " | " | " | Elisabethquelle (S) [†] | Sieveking | Ph. Z. 1905, p. 702 |
| 12 | 133.00 | 46.8° | " | " | " | " | " (H) [‡] | Mache | Ch. Z. 1907, p. 813 |
| 13 | 126.00 | 23.5° | common salt | Baden-Baden | Germany | 650 | Büttquelle | Engler & Sieveking | 645 ft.; Lower Rotliegende; Ch. Z. 1907, p. 814 |
| 14 | 122.40 | 46.8° | simple | Gastein | Austria | 3400 | Elisabethstollen (H) [‡] | " | Ch. Z. 1907 p. 813 |
| 15 | 121.90 | 41.9° | " | " | " | " | Chorinskyquelle (H) [‡] | " | " |
| 16 | 119.80 | — | sulphur | Landeck | Germany | 1500 | Friedrichsquelle | Schäfer 1909 | Gneiss; Sommer, p. 29 |
| 17 | 106.00 | — | simple | Gastein | Austria | 3400 | Wasserfallquelle | Mache & Meyer | Sommer, p. 21 |
| 18 | 73.70 | — | " | " | " | " | Elisabethstollen (S) [†] | " | " |
| 19 | 54.50 | 8.4° | alkaline sulphated | Karlsbad | " | 1230 | Eisenquelle | Sieveking | Ph. Z. 1905, p. 702 |
| 20 | 52.00 | — | " | Tannbach | " | — | Schlossteichquelle | Bamberger 1908 | Sommer, p. 43 |
| 21 | 51.00 | — | iron | Impos in Villanderthal | Tyrol | 3780 | Magenquelle ober Aunkaut | " 1907 | " p. 19 |
| 22 | 47.70 | — | iron carbon-dioxide | Disentis | Switzerland | 3770 | Placidusquelle unter | Schweitzer 1908-9 | " p. 17 |
| 23 | 47.00 | 8.4° | alkaline sulphated | Karlsbad | Austria | 1230 | Eisenquelle | Engler & Sieveking | Ch. Z. 1907, p. 813 |
| 24 | 46.70 | — | iron carbon-dioxide | Disentis | Switzerland | 3770 | Placidusquelle ober | Schweitzer 1908-9 | Sommer, p. 17 |
| 25 | 38.40 | 8.4° | alkaline sulphated | Karlsbad | Austria | 1230 | Eisenquelle | Mache | Ch. Z. 1907, p. 813 |
| 26 | 32.80 | 12.6° | alkaline sulphated common salt | Kreuznach | Germany | 340 | Inselquelle | Aschoff 1905 | quartz Porph. Sommer, p. 27 |

S. 1907, p. 115

Hauptquelle

| No. | Emission per litre of water in Maché's units | Temp. of spring in C. | Classification | Location | | Altitude in ft. | Spring | Observer | Remarks |
|-----|--|-----------------------|--|---------------------------|-------------|-----------------|-----------------------------------|---------------------------|---|
| 27 | 31.50 | 39.2° (?) | alkaline sulphated | Karlshad | Austria | 1250 | Mühlbach bei Vord. Quell. | Mache & Mexas 1905 | Sommer, p. 27 |
| 28 | 28.60 | 15.0° | carbon dioxide common salt | Nauheim | Germany | 470 | Karlsbrunnen | Schmidt & Kurz | Tert. Clay; Ch. Z. 1907, p. 81 |
| 29 | 27.90 | | earth-muriated common salt | Kreuznach | " | 340 | Quelle am Gassenhaus I | Ascholt 1905 | Quartz Porph. Schmidt & Kurz |
| 30 | 27.60 | 24.0° | " | " | " | " | Quelle am Haus Nr. | Schmidt & Kurz 1905 | Quartz Porph. Ch. Z. 1907, p. 81 |
| 31 | 26.00 | 8.3° (?) | earthy saline non-carbon dioxide | Griesbach | " | 1070 | Badquelle | Engler & Sieveking | Quartz Porph. Ch. Z. 1907, p. 81 |
| 32 | 24.00 | 59.0° | common salt | Baden Baden | " | 650 | Murquelle | " | Quartz Porph. Ch. Z. 1907, p. 81 |
| 33 | 23.49 | — | sulphur | Pistyan | Hungary | 530 | Brunnenschacht | Alexander & Weiss 1905 | Sommer, p. 37 |
| 34 | 23.40 | 31.2° | earth-muriated common salt | Münster a. Stein | Germany | 380 | Hauptbrunnen | Schmidt & Kurz 1905 | 180 hl.; Quartz Porph.; Sommer, p. 37 |
| 35 | 22.60 | 13.3° | muriated alkaline | Castellmare | Italy | — | Acidola | Engler | Ch. Z. 1907, p. 813 |
| 36 | 22.50 | — | iron | Kudowa | Germany | 1310 | — | " | Gravel bed; Lazarus, p. 200 |
| 37 | 22.10 | — | acid | Yellowstone National Park | U. S. A. | 7000 | Apollinaris Spring | Schmidt & Moore | Bot. Gaz. 1914, p. 82 |
| 38 | 21.90 | 11.3° | earthy carbon dioxide common salt | Söden in Taunus | Germany | 460 | Crampersbrunnen | Schmidt & Kurz | Taunus Phyllite; Ch. Z. 1907, p. 81 |
| 39 | 20.40 | 12.6° | earth-muriated common salt | Kreuznach | " | 340 | Inselquelle | " 1905 | Quartz Porph. Ch. Z. 1907, p. 81 |
| 40 | 20.00 | 11.1° | sulphated hydrogen sulphide | Nenndorf | " | 230 | Gewölbequelle | " | Tert. Clay, slate with Sph. Lazarus, p. 200 |
| 41 | 19.80 | cold | — | Fuengi near Rome | Italy | — | Badquelle | Engler | Ch. Z. 1907, p. 813 |
| 42 | 17.50 | — | — | Bründl | Austria | — | Quelle in der Kapelle | Bamberger 1908 | Sommer, p. 15 |
| 43 | 16.30 | — | sulphur | Talheim | Germany | — | — | " | Lazarus, p. 200 |
| 44 | 16.20 | 10.6° | — | Wetterau | " | — | Sauerbrunnen | Schmidt & Kurz | 501 hl.; Tert. Quartz Sand; Ph. Z. 1906, p. 217 |
| 45 | 16.00 | 9.2° | earthy alkaline saline non-carbon dioxide | Antegast | " | 1640 | Antoniusquelle | Frommel | 144 hl.; Gneiss; Ch. Z. 1907, p. 813 |
| 46 | 16.00 | cold | alkaline sulphated carbon dioxide | Marienthal | Austria | 2090 | Sauerbrunnen Markgraben | Zockelbauer & Dietel 1905 | Sommer, p. 31 |
| 47 | 14.50 | " | " | " | " | " | Ortssäuerling | " | " " |
| 48 | 14.20 | — | — | Kefermarkt | " | — | Dorfbrunnen | Bamberger 1908 | " p. 27 |
| 49 | 14.00 | — | — | Hundslof | " | — | Brunnenquelle | " | " p. 25 |
| 50 | 12.50 | 10.3° | — | Odenwald | Germany | — | Hirtenbrunnen | Schmidt & Kurz | Gerard, Ph. Z. 1906, p. 217 |
| 51 | 11.95 | 42.0° | common salt | Wisladen | " | 380 | Dr. Kurz's Quelle | Herrlich | Ch. Z. 1907, p. 813 |
| 52 | 11.44 1.00 | — | — | Stalnowick in Caracis | Russia | — | — | " | Ch. Z. 1910, p. 231 |
| 53 | 11.00 | 33°-48° | sulphur common salt | Lavey | Switzerland | 1350 | — | Suasso, Goy & Maché | Sommer, p. 29 |
| 54 | 10.10 | 22.5° | simple | Badenweiler | Germany | 1450 | Siegelsche Quelle | Engler & Sieveking | Ch. Z. 1907, p. 813 |
| 55 | 9.90 | — | — | Westerwald | " | — | — | Schmidt & Kurz | Tert. Clay; Ph. Z. 1906, p. 217 |
| 56 | 8.73 | — | alkaline saline non-carbon dioxide | Teplitz | Austria | 730 | Besenquelle bei Dux | Mache & Mexas 1905 | Sommer, p. 45 |
| 57 | 8.16 | cold | — | Franzensbad | " | 1500 | Franzensquelle | Sauberhann 1908 | " p. 17 |
| 58 | 8.00 | 10.6° | earthy carbon dioxide common salt | Homburg | Germany | 630 | Elisabethbrunnen | Schmidt & Kurz | Pre-Devon, Slate & Quartz Vein; Ch. Z. 1907, p. 813 |
| 59 | 7.88 | 22°-35° | sulphur | Ischl near Vienna | Austria | 700 | Franzensbad | Mache | Sommer, p. 13 |
| 60 | 7.80 | cold | iron carbon dioxide | Petersthal | Germany | 1330 | Karl Boschert | Frommel | Gran. Dyke in Gneiss; Ch. Z. 1906, p. 213 |
| 61 | 7.40 | " | earthy saline non-carbon dioxide | Freyersbach | " | 1475 | Gasquelle | " | Ch. Z. 1907, p. 813 |
| 62 | 7.10 | 9.8° | muriated bitter | Mergentheim | " | 60 | Karlquelle | " | 3000 hl.; Lower Muschelkalk; Schmidt, p. 1905 |
| 63 | 7.10 | 13.0° | earth-muriated common salt | Spessart | " | 470 | Albertquelle | Schmidt & Kurz | Zechstein; Ph. Z. 1906, p. 210 |
| 64 | 7.00 | 27.5° | simple | Badenweiler | " | 1450 | Hauptbadquelle | Engler & Sieveking | Ch. Z. 1907, p. 813 |
| 65 | 6.56 | 32.5° | alkaline saline alkaline muriated carbon dioxide | Teplitz | Austria | 730 | Steinbadquelle | Mache | " |
| 66 | 5.70 | — | — | Gleichenberg | " | 980 | Constantinquelle | Bamberger 1907 | Sommer, p. 21 |
| 67 | 5.70 | 74.0° | weak salt | Battaglia near Padua | Italy | 1900 | Surgone Grotta | Engler | Ch. Z. 1907, p. 813 |
| 68 | 5.15 | — | concentrated common salt | Sulza | Germany | 485 | Konstantinquelle | Wieprecht | Sommer, p. 43 |
| 69 | 5.00 | 87.0° | weak salt (?) | Alano near Padua | Italy | 100 | Surgone Montano centrale | Engler | Ch. Z. 1907, p. 813 |
| 70 | 4.70 | 65.0° | muriated alkaline | Ischia | " | — | Porto d'Ischia still concentrated | " | " |
| 71 | 4.50 | cold | alkaline sulphated | Reichelsch | Austria | 750 | Fempelquelle | Mache 1905 | Sommer, p. 37 |
| 72 | 4.50 | — | — | Gieszen | Germany | — | Schwache Quelle | Schmidt & Kurz | Tert. Clay; Ph. Z. 1906, p. 210 |
| 73 | 4.33 | — | — | Kislowick in Caucasus | Russia | — | — | " | Ch. Z. 1910, p. 231 |
| 74 | 4.33 | 10.4° | common salt | Kissinzen | Germany | 650 | Maxquelle | F. Jentzsch | Upper Permian; Ph. Z. 1906, p. 217 |
| 75 | 4.08 | cold | iron carbonate | Spa | Belgium | 1000 | Tonnelet Spring | Gerard & Chazy 1905 | Bot. Gaz. 1914, p. 82 |
| 76 | 3.78 | 21.2° | alkaline carbonate dioxide | Apollinaris | Germany | — | — | T. Kyll 1900 | Sommer, p. 13 |
| 77 | 3.77 | — | sulphur | Alvaneu | Switzerland | 3150 | Untere Quelle | Schweitzer | " |

| No. | Em. in c.c. per litre of water in Maché's units | Temp. of spring in C. | Classification | Location | | Altitude in ft. | Spring | Observer | Remarks |
|-----|---|-----------------------|---|------------------------|-------------|-----------------|--------------------------|------------------------------------|---|
| 78 | 3.40 | 12.5° | alkaline-mineralized hydrocarbon sulphate | Wendisch in Tannus | Germany | 440 | Natron-Lithiumquelle | Schmidt & Kurz | 46.7 hl.; Tertiary Limestone; Sommer, p. 47 |
| 79 | 3.2-1.6 | 30°-38° | simple | Wildbad | " | 1410 | Bohrlöcher im Badehaus | Engler & Sieveking | Granite; Ch. Z. 1907, p. 813 |
| 80 | 3.10 | — | — | Württemberg | " | — | Göppinger Sauerbrunnen | K. R. Koch | Ph. Z. 1907, p. 806 |
| 81 | ca. 3.00 | — | earthly sulphate of common salt | Salzschlirf | " | 820 | — | — | Lazarus, p. 199 |
| 82 | 2.60 | — | salt alkaline | Ragnoli near Naples | Italy | — | Manganello | Engler | Ch. Z. 1907, p. 813 |
| 83 | ca. 2.50 | 11.5° | earthly mineralized carbon dioxide | Wildungen | Germany | 980 | Helenenquelle | — | 76 hl.; Clayslate; Lazarus, p. 200 |
| 84 | 2.48 | — | earthly carbon dioxide | Altheide | " | 1310 | Charlottenbrunnen | Fresenius 1909 | Alluvium; Sommer, p. 13 |
| 85 | 2.20 | 72.0° | mineralized alkaline | Casamicciola in Ischia | Italy | — | Manzi II | Engler | Ch. Z. 1907, p. 813 |
| 86 | 2.10 | 0.8° | earthly saline non carbon dioxide | Rippoldsau | Germany | 1870 | Wenzelsquelle | Engler & Sieveking | Gneiss; Ch. Z. 1907, p. 813 |
| 87 | 1.90 | 90.0° | mineralized alkaline | Agnone near Naples | Italy | — | Purgativo | Engler | Ch. Z. 1907, p. 813 |
| 88 | 1.80 | cold | " | Pozzuoli near Naples | " | — | Aqua media | " | " |
| 89 | 1.52 | " | non carbon dioxide | St. Moritz | Switzerland | 5820 | Surpunt | Schweitzer | Sommer, p. 43 |
| 90 | 1.50 | " | iron | Roncegno | Austria | 1750 | Trinkquelle | Bamberger 1907 | " p. 37 |
| 91 | 1.40 | — | alkaline mineralized saline | Salzig | Germany | — | Quelle II | Schmidt | Low. Devon. Quartzite & Clay-slate; Sommer, p. 39 |
| 92 | 1.4-0.55 | — | — | Platonsk in Caucasus | Russia | — | — | — | Ch. Z. 1910, p. 231 |
| 93 | 1.35 | — | — | French Lick, Indiana | U. S. A. | — | Pluto Spring | Schlundt | Bot. Gaz. 1914, p. 82 |
| 94 | 1.31 | — | carbon dioxide | Passug | Switzerland | 2925 | Churwald. Quelle | Schweitzer | Sommer, p. 35 |
| 95 | 1.13 | cold | alkaline carbon dioxide | Tarasap | " | 3900 | Carolaquelle | " 1909 | " p. 43 |
| 96 | 1.13-0.41 | — | — | Essentuki in Caucasus | Russia | — | — | — | Ch. Z. 1910, p. 231 |
| 97 | 1.02 | 47°-52° | sulphur | Tronezin-Teplitz | Hungary | 850 | Quelle I | — | Lazarus, p. 200 |
| 98 | 0.84 | — | iron | Levico | Austria | 1640 | Starkquelle | Bamberger 1907 | Sommer, p. 29 |
| 99 | 0.76 | 31.7°-34° | simple | Ragaz-Pfäfers | Switzerland | 2000 | Alte Quelle | Schweitzer | " p. 37 |
| 100 | 0.71 | 23.3° | " | Vöslau | Austria | 810 | Hauptquelle | Mache & Meyer 1905 | " p. 47 |
| 101 | 0.58 | 36.7°-52° | mineralized sulphur | Baden | Switzerland | 1230 | Kesselquelle | Sury | " p. 13 |
| 102 | 0.31 | 14.0° | common salt | Reichenhall | Germany | 1540 | Kais. Karlquelle | Bender & Hobein 1909 | Muschelkalk; Sommer, p. 37 |
| 103 | 0.26 | 33.9°-51.1° | sulphated iron | Leukerbad | Switzerland | 4600 | Lorenzquelle | Bamberger 1907 | Sommer, p. 29 |
| 104 | 0.02 | — | concentrated common salt | Rheinfelden | " | 866 | — | Sommer 1908 | " p. 37 |
| | | | | | | | Gas evolving from: | | |
| 1 | 94.29 | 39.2° (?) | alkaline sulphated | Karlbad | Austria | 1230 | Mühlbrunnen vord. Quelle | Mache & Meyer 1905 | Sommer, p. 27 |
| 2 | 50.20 | — | " | " | " | " | Schlossbrunnen | " " | " " |
| 3 | 21.90 | — | alkaline saline | Teplitz | " | 730 | Urquelle | " " | " p. 45 |
| 4 | 15.00 | — | alkaline mineralized carbon dioxide | Gleichenberg | " | 980 | Constantinquelle | Bendorf 1907 | " p. 21 |
| 5 | 10.90 | cold | alkaline sulphated carbon dioxide | Marienbad | " | 2090 | Waldquelle | Mache, Maier and Zörkelbrüder 1905 | " p. 33 |
| 6 | 2.60 | 23.3° | simple | Vöslau | " | 810 | Hauptquelle | Mache & Meyer 1905 | " p. 47 |

(B) Results of measurements expressed in curies.

The quantity of radium emanation expressed in milligramme radium per minute in 10 litres of gas or water quoted from Curie's "Radioaktivität" and that expressed in milligramme radium bromide (Ra Br₂) per minute in 1 litre of gas or water quoted from Lazarus' "Handbuch der Radium-Biologie und Therapie" are recalculated into curies and Mache's units by using the following factors:

- 1 millig. min. emanation of radium = 1250×10^{10} curies.
 1 millig. min. emanation of radium bromide (RaBr₂) = 734×10^{10} curies.
 1×10^{-10} curie = 0.275 Mache's units.

The numbers in brackets denote the quantity of radium emanation calculated for reference under the supposition that the original numbers given in Curie's work refer to milligramme radium bromide (RaBr₂) per minute, and not to milligramme radium per minute.

| No. | Emanation per litre of water | | Temp. of spring in C. | Classification | Location | | Altitude in ft. | Spring | Observer | Remarks |
|-----|------------------------------|---------------------------------|-----------------------|---------------------|---------------|---------|-----------------|-------------------------|--------------------|----------------|
| | in 10 ⁻¹⁰ curies | in Mache's units (recalculated) | | | | | | | | |
| 1 | 649.59 | 178.64 | — | — | Agnas Lerez | Spain | — | Aceñas | Munoz del Castillo | Lazarus, p. 44 |
| 2 | 229.01 | 62.98 | 60.0° | mineral alkaline | La Bourboule | France | 2780 | Puits Choussy | Laborde | " |
| 3 | 162.50
(95.42) | 44.69
26.24 | 13.0° | iron carbon dioxide | Bussang | " | 2200 | Grande Salmade | " | Curie, p. 504 |
| 4 | 161.48 | 44.41 | 43.0° | sulphur | Luchon | " | 2600 | Grande source
Bardon | Moureu | Lazarus, p. 44 |
| 5 | 59.45 | 16.35 | 46.0° | simple (?) | Plombières | " | 1300 | Capucins | Curie & Laborde | " |
| 6 | 52.50
(30.83) | 14.44
8.48 | 12.0° | iron carbon dioxide | Dirza Corsica | " | 1960? | Source Dirza | Laborde | Curie, p. 504 |
| 7 | 47.50
(27.89) | 13.06
7.67 | 69.0° | simple (?) | Plombières | " | 1300 | Vauquelin | Curie & Laborde | " |
| 8 | 40.37 | 11.10 | 47.0° | salt sulphur | Aix-les-Bains | " | 860 | Source d'Alun | " | Lazarus, p. 44 |
| 9 | 33.47 | 9.20 | — | — | St. Lucasbad | Hungary | — | — | — | " |
| 10 | 21.25
(12.48) | 5.84
3.43 | 58.0° | saline | Bourbon-Lancy | France | 1780 | Le Lymbe | Curie & Laborde | Curie, p. 504 |
| 11 | 1.10 | 0.30 | 27.2° | saline sulphur | Uriage | " | 1350 | Source principale | Besson | Lazarus, p. 44 |
| | Emanation per litre of gas | | | | | | | Gas evolving from: | | |
| 1 | 2936.00 | 807.40 | 36.3° | simple | Gastein | Austria | 3400 | Grabenackerquelle | Mache & Meyer | Lazarus, p. 44 |
| 2 | 1416.62 | 389.57 | 60.0° | mineral alkaline | La Bourboule | France | 2780 | Puits Choussy | Laborde | " |
| 3 | 1347.62 | 370.60 | 43.0° | sulphur | Luchon | " | 2600 | Grande source
Bardon | Moureu | " |
| 4 | 736.94 | 202.66 | 69.0° | simple (?) | Plombières | " | 1300 | Vauquelin | Curie & Laborde | " |
| 5 | 507.50
(298.00) | 139.56
81.95 | 46.0° | " | " | " | " | Capucins | " | Curie, p. 504 |
| 6 | 226.07 | 62.17 | 47.0° | salt sulphur | Aix-les-Bains | " | 860 | Source d'Alun | " | Lazarus, p. 44 |
| 7 | 222.50
(130.65) | 61.19
35.93 | 58.0° | saline | Bourbon-Lancy | " | 1780 | Le Lymbe | Curie & Laborde | Curie, p. 504 |
| 8 | 90.28 | 24.83 | — | — | St. Lucasbad | Hungary | — | — | " | Lazarus, p. 44 |

Radio-activity of Mineral Springs in Miyagi and Nagano Prefectures.

(Supplement to Tables 1 and 2)

Table 10.

(A) Determined with Schmidt's electrometer.

| No. | Emanation per litre of water | | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-------------------------------------|------------------------------|---------------------------------|---|-----------------------|----------------|-------------|---------|----------|-------------------|
| | in 10 ¹⁰ curies | in Maché's units (recalculated) | | | | | | | |
| 1 | 237.31 | 65.26 | — | 9.0° | carbonated | Ōkuwa | 大桑(長野) | Granite | Shika-no-yu No. 2 |
| 2 | 224.25 | 61.67 | — | 9.5° | " | " | " | " | " No. 1 |
| 3 | 53.01 | 14.58 | 335 | 56.0° | " | Tōgatta | 遠刈田(宮城) | Liparite | Kami-no-yu |
| 4 | 21.20 | 5.83 | — | 9.2° | " | Ōkuwa | 大桑(長野) | Granite | Shika-no-yu No. 3 |
| 5 | 13.41 | 3.69 | — | 49.5° | salt | Sakunami | 作並(宮城) | Tertiary | Shin-yu |
| 6 | 11.42 | 3.14 | — | 51.0° | " | Tōgatta | 遠刈田(宮城) | Liparite | Naka-no-yu |
| 7 | 10.64 | 2.93 | — | 46.0° | " | Sakunami | 作並(宮城) | Tertiary | Me-no-yu |
| 8 | 1.55 | 0.43 | — | 83.5° | " | Shin-kuruma | 新車(長野) | Andesite | Moto-yu |
| Emanation per litre of gas at 0° C. | | | | | | | | | |
| 1 | 287.54 | 79.07 | — | 9.0° | carbonated | Ōkuwa | 大桑(長野) | Granite | Shika-no-yu No. 2 |
| 2 | 62.49 | 17.18 | — | 9.2° | " | " | " | " | " No. 3 |

(B) Determined with the fontactoscope of Kohlrausch and Löwenthal.

| No. | Emanation per litre of water in Maché's units | Flow of water in 24 hrs. in hectolitres | Temp. of spring in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|---|-----------------------|----------------|-------------|---------|----------|-----------------------------------|
| 1 | 5.04 | — | 51.5° | salt | Tōgatta | 遠刈田(宮城) | Liparite | Miuraya-no-yu (<i>Ōtoko-yu</i>) |
| 2 | 3.69 | — | 51.0° | " | " | " (宮城) | " | Naka-no-yu |
| 3 | 3.50 | 270 | 51.0° | " | Sakunami | 作並(宮城) | Tertiary | Tsuru-no-yu (<i>D. Koike</i>) |
| 4 | 3.20 | — | 51.0° | " | " | " (宮城) | " | Taka-no-yu |
| 5 | 2.62 | — | 52.0° | " | " | " (宮城) | " | Tsuru-no-yu (<i>Moritani</i>) |
| 6 | 2.20 | 450 | 63.0° | " | " | " (宮城) | " | Kawara-yu |
| 7 | 2.13 | — | 52.0° | " | Tōgatta | 遠刈田(宮城) | Liparite | Kamasaki-no-yu |
| 8 | 2.05 | 1400 | 57.0° | simple | Ao-ne | 青根(長野) | Tertiary | Shin-yu |
| 9 | 2.01 | — | 51.5° | salt | Tōgatta | 遠刈田(宮城) | Liparite | Miuraya-no-yu (<i>Ōtoko-yu</i>) |
| 10 | 1.86 | 450 | 69.5° | " | Sakunami | 作並(宮城) | Tertiary | Taki-no-yu |
| 11 | 1.64 | — | 55.0° | " | Tōgatta | 遠刈田(宮城) | Liparite | Ōkomuro-uchi-yu |
| 12 | 1.63 | 1400 | 52.0° | simple | Ao-ne | 青根(長野) | Tertiary | Myōgō-no-yu |
| 13 | 1.43 | — | 50.0° | salt | Tōgatta | 遠刈田(宮城) | Liparite | Jōtō-yu |
| 14 | 1.42 | 1584 | 43.0° | " | Akyū | 秋保(長野) | Tertiary | Sawa-no-yu |
| 15 | 1.42 | — | 74.0° | — | Moto-kuruma | 元車(長野) | Andesite | Moto-kuruma-yu |
| 16 | 1.29 | 7258 | 42.5° | simple | Ao-ne | 青根(長野) | Tertiary | Ō-yu |
| 17 | 0.86 | 1584 | 53.0° | common salt | Akyū | 秋保(長野) | " | Mi-yu |
| 18 | 0.75 | — | 44.0° | carbonated | Kawatabi | 川渡(長野) | " | Suzuki-no-yu |
| 19 | 0.74 | 648 | 48.5° | " | " | " | " | Ō-yu |
| 20 | 0.73 | — | 59.5° | " | Akayu | 赤湯(秋田) | Andesite | Ue-no-yu |
| 21 | 0.36 | — | 48.0° | " | Kawatabi | 川渡(長野) | Tertiary | Sukashi-yu |

II. Temperature.

Table 11.

| No. | Temp. of spring
in C. | Total residue
in grams | Flow of water
in 24 hrs.
in hectolitres | Emission per
litre of water
in Maché's units | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|--------------------------|---------------------------|---|--|------------------------------|--------------------|----------|-------------------------|--|
| 1 | 108.0° | 9.24 | 1296 | 0.04 | earth-mixed
common salt | Atami | 熱海(静岡) | Andesite | Ô-yu |
| 2 | 103.0° | 3.52 | — | 0.13 | alkaline sulphur | Naruko | 鳴子(宮城) | " | Unagi-yu |
| 3 | 102.0° | 4.19 | 9000 | 0.06 | common salt | Senami | 瀬沢(新潟) | — | Senami-Funtô |
| 4 | 100.0° | 1.12 | 270 | — | salt | Onikôbe | 鬼首(宮城) | Tertiary | Fukiage-yu |
| 5 | 100.0° | — | — | 0.27 | common salt | Shimogamo | 下賀茂(静岡) | " | Spring A
<small>near Jûmyô temple</small> |
| 6 | 99.0° | — | — | 0.22 | — | Kamegawa | 亀川(大分) | Andesite | Kamado-no-yu |
| 7 | 97.0° | — | 324 | — | vitriol | Noboribetsu | 登別(北海道) | Tertiary | Memo-yu |
| 8 | 96.0° | — | 1800 | — | " | Myanoshita | 宮下(神奈川) | Andesite | Spring A |
| 9 | 96.0 | — | — | 1.14 | alkaline sulphur | Nakabusa | 中房(長野) | Granite | Myôken-no-yu |
| 10 | 95.0° | — | abundant | 1.76 | alkaline carbon-
dioxated | Yumura (Hyôgo-ken) | 湯村(兵庫) | " | Ara-yu |
| 11 | 95.0° | — | — | 1.11 | sulphur | Futami | 二見(富山) | " | Dai-san-no-yu |
| 12 | 95.0° | 4.06 | 210 | — | acid alum | Osoreyama | 恐山(青森) | Andesite | Shin-taki-no-yu |
| 13 | 95.0° | — | — | — | — | Hokuto | 北投(臺北) | Tertiary | Spring A |
| 14 | 94.0° | — | — | 0.64 | — | Obama | 小濱(長崎) | Andesite | Gorin-yu |
| 15 | 93.0° | — | — | 0.93 | earth-mixed
common salt | Wakura | 和倉(石川) | Tertiary | Wakazaki no-yu No. 1 |
| 16 | 91.0° | — | 3060 | — | iron | Furô-sen | 不老泉(秋田) | " | Shirakumo-no-taki |
| 17 | 91.0° | 2.52 | 389 | 0.13 | acid alum vitriol | Myôkan | 明礬(大分) | Andesite | Jizô-no-yu |
| 18 | 90.0° | 1.40 | 1555 | — | saline bitter sulphur | Yunomine | 湯ノ峯(和歌山) | Tertiary | Oguri-no-yu |
| 19 | 90.0° | — | 3000 | — | salt | Hiraochi | 平落(鹿児島) | Volc. Ash &
Mud Lava | Naka-no-yu |
| 20 | 88.5° | — | — | 0.18 | common salt | Yugawara | 湯河原(神奈川) | Andesite | Uenoya-no-yu |
| 21 | 88.0° | — | — | 1.42 | sulphur (?) | Kazu | 葛(長野) | Granite | Spring A |
| 22 | 88.0° | 0.50 | — | 1.97 | simple | Yubiso | 湯槍(群馬) | " | Yubiso-no-yu No. 2 |
| 23 | 87.0° | — | 4200 | — | alum | Shiose | 鹽波(鹿児島) | Volc. Ash &
Mud Lava | Shimo-no-yu |
| 24 | 84.0° | — | 270 | — | simple | Dai | 臺(岩手) | Diluvium | Taki-no-yu |
| 25 | 83.0° | — | 130 | 1.13 | " | Kami-Suwa | 上諏訪(長野) | Alluvium | Tajiku-no-yu |
| 26 | 82.0° | 0.98 | 5148 | — | sulphur | Nozawa | 野澤(新潟) | Volc. Detr. | Asagama-no-yu |
| 27 | 82.0° | 0.40 | 207 | — | saline sulphated
bitter | Hokonagi | 鉾投(鹿児島) | Andesite | Hokonagi-no-yu |
| 28 | 80.3 | — | — | 0.28 | salt | Urai | ウライ(臺北) | Tertiary | Spring B
<small>on the foot bank</small> |
| 29 | 80.0° | 0.15 | 4680 | — | acid | Ôra | 大良(鹿児島) | Volc. Ash &
Mud Lava | Ôradani-no-yu |
| 30 | 80.0° | — | 135 | — | common salt | Masan (Chôsen) | 馬山(朝鮮) | — | Masan-Chên |
| 31 | 80.0° | — | — | — | salt | Shiriuchi | 尻内(北海道) | — | Tono-no-yu |
| 32 | 79.3° | — | — | 0.18 | simple | Toi | 頭圍(宜蘭) | Quaternary | Spring in field |
| 33 | 79.0° | 1.23 | — | 1.13 | sulphated bitter | Asamushi | 淺虫(青森) | Andesite | Tsubaki-no-yu No. 1 |
| 34 | 79.0° | — | 648 | 0.70 | saline bitter | Yujiku | 湯宿(群馬) | Tert. Tuff | Kubo-yu |
| 35 | 79.0° | — | — | 1.28 | common salt | Karayamazu | 片山津(石川) | Tertiary | Shiotsu-no-Tsubo |
| 36 | 78.0° | 2.39 | — | 1.87 | " | Kuradate | 蔵館(青森) | Alluvium | Hoyôen-no-yu |
| 37 | 77.0° | — | 1400 | 0.36 | saline common salt | Shuzenji | 修善寺(静岡) | Andes. & Tuff | Ayame-no-yu |
| 38 | 76.0° | — | — | 0.74 | salt | Ôwari | 大鰐(青森) | Liparite | Kwara-no-yu |
| 39 | 76.0° | 1.20 | 380 | 0.64 | bitter | Shido-daira | 志戸平(岩手) | Tertiary | Warabi-no-yu |
| 40 | 76.0° | — | — | 1.40 | common salt | Shirai (Hiro) | 流平(長野) | Diluvium | Ô-yu No. 3 & 2 |
| 41 | 76.0° | — | — | 1.33 | common salt | Yudanaka | 湯田中(新潟) | Misaka Series | Ô-yu No. 1 & 2 |
| 42 | 76.0° | 10.03 | 152 | 2.17 | earth-mixed
common salt | Awara | 蘆原(福井) | Alluvium | Furutsu-no-yu |
| 43 | 76.0° | — | — | 0.30 | " | Yunotani | 湯谷(熊本) | Aug. Andes. | Yunotani-Jigoku |
| 44 | 75.0° | — | — | 58.18 | " | Misasa | 三朝(鳥取) | Granite | Akazakiya-no-yu |

III. Flow of Water.

Table 12.

| No. | Flow of water
in 24 hrs.
in hectolitres | Temp. of spring
in C. | Total residue
in grams | Discharge of solid
materials
in 24 hrs.
in kilograms | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---|--------------------------|---------------------------|---|--------------------------------|------------------------------|-----------|-------------------------|------------------------|
| 1 | ca. 63000 | 40.0° | ca. 2.72 | ca. 17136 | acid alum vitriol | Takayu
(Yamagata-ken) | 高湯(山形) | Andesite | Sugawa-no-yu |
| 2 | 59530 | 48.5° | — | — | .. | Hokutō | 北投(臺北) | Tertiary | Taki-no-yu |
| 3 | 57802 | 37.5° | — | — | salt | Itō | 伊東(静岡) | Quaternary | Wada-Onsen |
| 4 | 54000 | 71.5° | 0.81 | 4374 | acid vitriol | Noboribetsu | 登別(北海道) | Tertiary | Taki-no-yu |
| 5 | 54000 | — | — | — | salt | Renge | 蓮華(新潟) | Andesite | Yakushi-no-yu |
| 6 | 36000 | 37° 38' | 1.03 | 3708 | alkaline | Kambara | 蒲原(..) | Chichibu Series | Kambara-Onsen |
| 7 | 24624 | 40.5° | 0.13 | 320 | simple | Innai-Yunosawa | 院内湯之澤(山形) | Tertiary | Yunosawa-Onsen |
| 8 | 21312 | 55.5° | 0.18 | 384 | .. | Shu-otsu (Chōsen) | 朱乙(朝鮮) | Granite | Shu-otsu-Onsen |
| 9 | 12600 | 60.0° | — | — | alum | Shibi | 紫尾(鹿兒島) | Volc. Ash &
Mud Lava | Shibi-Onsen |
| 10 | 10800 | 39.0° | 0.89 | 961 | simple | Onsei-in (Chōsen) | 溫井院(朝鮮) | Granite | Onsei-in-Onsen |
| 11 | 9540 | 53.9° | 1.44 | 1374 | iron carbonate | Anraku | 安樂(鹿兒島) | Volc. Ash &
Mud Lava | Anraku-Onsen |
| 12 | 9450 | 30.1° | 1.00 | 945 | simple | Yumoto
(Wakayama-ken) | 湯本(和歌山) | Mesozoic. | Tatsukami-Onsen |
| 13 | 9252 | 46.7° | 0.35 | 324 | sulphur | Myōban-yama | 明礬山(鹿兒島) | Volc. Ash &
Mud Lava | Iwō-yu |
| 14 | 9000 | 102.0° | 4.19 | 3771 | common salt | Senami | 瀬波(新潟) | — | Senami-Funtō |
| 15 | 7776 | 34.0° | 0.30 | 233 | simple | Imagami | 今神(山形) | Tertiary | Imakumano-Kōsen |
| 16 | 7200 | 62.0° | 1.19 | 857 | earthy alkaline
sulphur | Akakura | 赤倉(新潟) | Andesite | Akakura-Onsen |
| 17 | 6012 | 61.0° | 2.28 | 1371 | sulphur | Ōtaki | 大滝(秋田) | Alluv. Tal. Dep. | Tsuru-no-yu |
| 18 | 5400 | 60.0° | 0.57 | 308 | hydrogen sulphide | Eino-o | 榮之尾(鹿兒島) | Volc. Ash &
Mud Lava | Eino-o-Onsen |
| 19 | 4585 | 55.0° | 1.90 | 871 | acid muriated | Dake (Aomori-ken) | 岳(青森) | Volc. Detr. | Dake-no-yu |
| 20 | 4500 | 48.0° | 0.22 | 10 | simple | Yubara
(Okayama-ken) | 湯原(岡山) | Granite | Yakuō-tō |
| 21 | 4464 | 63.0° | 1.30 | 621 | hydrogen sulphide | Tateyama | 立山(富山) | Andesite | Tateyama-Onsen |
| 22 | 4140 | 38.0° | — | — | sulphated bitter | Tochinoki | 栃ノ木(熊本) | .. | Kwan-on-yu |
| 23 | 4066 | 58.5° | 1.91 | 777 | .. | Iwai | 岩井(鳥取) | Tertiary | Moto-yu |
| 24 | 3944 | 62.0° | 0.37 | 146 | simple | Kaminoyama | 上山(山形) | Liparite | Hie-no-yu |
| 25 | 3888 | 43.0° | 0.32 | 124 | .. | Yumura
(Shimane-ken) | 湯村(島根) | Granite | Shitsuni-no-yu |
| 26 | 3455 | 39.0° | 1.17 | 404 | sulphur | Dake (Fukushima-ken) | 岳(福島) | — | Dake-no-yu |
| 27 | 3230 | 43.0° | 0.29 | 94 | simple | Arifuku | 有福(島根) | Diorite | Arifuku-Onsen |
| 28 | 3150 | 48.0° | 5.03 | 1585 | saline common salt | Kamasaki | 鎌先(宮城) | Andesite | Ichijō-no-yu |
| 29 | 3060 | 50.0° | — | — | salt | Yufuin | 由布院(大分) | .. | Dakeshita-no-yu |
| 30 | 3060 | 91.1° | — | — | iron | Furosen | 不老泉(秋田) | Tertiary | Shirakumo-no-taki |
| 31 | 2772 | 44.0° | 2.97 | 768 | earthy common salt | Aoyama | 青山(北海道) | Andesite | Furōkaku-no-yu No. 1 |
| 32 | 2700 | 37.5° | 0.17 | 46 | simple | Maga | 真賀(岡山) | Granite | Kin-yu |
| 33 | 2592 | 50.5° | 0.86 | 223 | .. | Yoshioka | 吉岡(鳥取) | Tertiary | Kabu-yu |
| 34 | 2592 | 95.0° | 1.86 | 482 | alkaline | Ureshino | 嬉野(佐賀) | Alluvium | Ureshino-Onsen |
| 35 | 2074 | 100.0° | — | — | common salt | Atami | 熱海(静岡) | Andesite | Aoki-no-yu |
| 36 | 1953 | 64.0° | 1.58 | 309 | .. | Tamatsukuri
(Shimane-ken) | 玉造(島根) | Tertiary | Kami-no-yu |
| 37 | 1800 | 96.0° | — | — | acid alum (?) | Miyanoshita | 宮ノ下(神奈川) | Andesite | Spring A |
| 38 | 1800 | 52.0° | 0.59 | 106 | simple | Sandogoya | 三斗小屋(栃木) | .. | Sandogoya-Onsen |
| 39 | 1800 | 39.0° | 0.28 | 50 | .. | Tochiomata | 栃尾(新潟) | Granite | Tochiomata-no-yu No. 1 |
| 40 | 1730 | — | 0.93 | 161 | .. | Togo | 東郷(鳥取) | Andesite | Yōjō-kwan-no-yu |
| 41 | 1512 | 49.0° | 0.77 | 116 | .. | Takeo | 武雄(佐賀) | .. | Takeo-Onsen |
| 42 | 1320 | 51.0° | 0.65 | 86 | iron carbonate | Bappu | 別府(大分) | .. | Kusunoki-Onsen |
| 43 | 1296 | 38.0° | 2.25 | 292 | earthy muriated
common salt | Yunogo | 湯郷(岡山) | Liparite | Sagi-no-yu |

IV. The Quantity of Solid Ingredients.

Table 13.

| No. | Total residue
in grams | Flow of water
in 24 hrs. in lils. | Discharge of solid
ingredients
in 24 hrs.
in kilograms | Temp. of spring
in C. | Classification | Location | 地名(府縣名) | Geology | Spring |
|-----|---------------------------|--------------------------------------|---|--------------------------|---|--------------------|----------|-----------------|-----------------------|
| 1 | ca.
65,70 | ca. | ca. | — | earth-muriated iron carbonate
concentrated common salt | Arima | 有馬(兵庫) | Quartz Porph. | Ikeno-Ito-Kōsen |
| 2 | 56,58 | — | — | — | „ | Karato | 唐櫃(„) | „ | Karato-Kōsen |
| 3 | 42,00 | — | — | — | acid concentrated alum vitriol | Isobe Toyama ken | 磯邊(富山) | Tertiary | Isobe-Kōsen |
| 4 | 30,73 | 405 | 1245 | 80,0° | earthy iron carbonate
concentrated common salt | Shiriuchi | 知内(北海道) | — | Shiriuchi-Onsen |
| 5 | 30,28 | 2 | 6 | — | alkaline carbondioxated
concentrated common salt | Onishi | 鬼石(群馬) | Crystal. Schist | Yashio-Kōsen |
| 6 | 29,79 | 2700 | 8043 | 70,0° | sulphated common salt | Ryuko | 龍岡(朝鮮) | Gneiss | Ryūkō-Kōsen |
| 7 | 29,54 | 720 | 2127 | 15,5° | alkaline carbondioxated
concentrated common salt | Isobe Gunma ken | 磯部(群馬) | Tertiary | Isobe-Kōsen |
| 8 | 28,57 | — | — | — | concentrated common salt | Kashio | 鹿嶋(長野) | Crystal. Schist | Kashio-Kōsen |
| 9 | 27,86 | — | — | — | alkaline carbondioxated
concentrated common salt | Haraichi | 原市(群馬) | Volc. Detr. | Mine-Kōsen |
| 10 | 25,49 | — | — | — | acid sulphated concentrated
alum vitriol | Taku | 多久(佐賀) | Tertiary | Tentoku-Kōsen |
| 11 | 25,15 | — | — | — | muriated sulphur | Aoki | 碓(宮崎) | Alluvium | Aokigahara-Kōsen |
| 12 | 23,31 | 1800 | 4196 | 78,0° | acid alum vitriol | Tsukawara | 塚原(大分) | Andesite | Iwōyama-Onsen |
| 13 | 23,07 | — | — | — | earthy iron carbondioxated
concentrated common salt | Masuyama | 増山(大阪) | Tertiary | Masuyama-Kōsen |
| 14 | 21,64 | 36 | 78 | 43,0° | earth-muriated vitriol
common salt | Yunomoto Iki Prov. | 湯之本(長崎) | Liparite | Yunomoto-Onsen No. 2 |
| 15 | 21,09 | — | — | — | earthy alkaline concentrated
common salt | Muko | 武庫(兵庫) | Granite | Mukosan-Kōsen |
| 16 | 20,93 | 2160 | 4521 | 82,0° | earth-muriated common salt | Wakura | 和倉(石川) | Tertiary | Wakura-Onsen |
| 17 | 20,40 | — | — | — | earth-muriated concentrated
common salt | Ōshio | 大鹽(福島) | Andesite | Ōshio-Kōsen |
| 18 | 19,66 | — | — | 53,4° | earth-muriated iron carbonate
common salt | Arima | 有馬(兵庫) | Quartz Porph. | Ichi-no-yu |
| 19 | 15,79 | 630 | 995 | 74,0° | common salt | Katayamazui | 片山津(石川) | Tertiary | Katayamazui-Onsen |
| 20 | 15,64 | — | — | — | earthy iron carbonate
common salt | Daitō | 大塔(大阪) | „ | Daitō-Kōsen |
| 21 | 15,08 | — | — | 63,0° | earth-muriated common salt | Matsunoyama | 松之山(新潟) | „ | Netsu-no-yu |
| 22 | 14,78 | 270 | 399 | 44,5° | hydrogen sulphide common salt | Ōyuzawa | 大湯澤(秋田) | Diluvium | No. 38 (O. J. H.) |
| 23 | 13,76 | 360 | 495 | 18,7° | carbondioxated common salt | Takarazuka | 寶塚(兵庫) | Granite | Takarazuka-Kōsen |
| 24 | 11,50 | — | — | 46,0° | earth-muriated common salt | Atsushio | 熱鹽(福島) | Alluvium | Atsushio-Onsen |
| 25 | 11,32 | — | — | 60,0° | „ | Shinogamo | 下賀茂(静岡) | Tertiary | Shinogamo-Onsen |
| 26 | 10,20 | 29 | 30 | 13,0° | carbondioxated common salt | Kawai | 川合(島根) | Granite | Urisaka-no-yu |
| 27 | 10,03 | 152 | 153 | 76,0° | earth-muriated common salt | Awara | 蘆原(福井) | Alluvium | Funatsu No. 10 |
| 28 | 9,86 | — | — | 15,0° | earthy common salt | Masutomi | 増富(山梨) | Granite | Yunokubo-gawara no-yu |
| 29 | 9,74 | — | — | — | hydrogen sulphide common salt | Hiyanezaki | 比屋根岬(沖縄) | „ | Hiyanezaki-Kōsen |
| 30 | 9,38 | — | — | 17,0° | iron carbonate common salt | Anamori | 穴守(東京) | Alluvium | Anamori-Kōsen |
| 31 | 9,24 | 1296 | 1198 | 108,0° | earth-muriated common salt | Atami | 熱海(静岡) | Andesite | Ō-yu |
| 32 | 8,84 | — | — | 65,0° | „ | Obama | 小濱(長崎) | „ | Fukiage-yu |
| 33 | 7,81 | 1107 | 911 | 60,0° | common salt | Beppu | 別府(大分) | „ | Higashi-Onsen |
| 34 | 7,25 | — | — | — | acid alum vitriol | Kusatsu | 草津(群馬) | Volc. Detr. | Shiranesan-no-yu |

V. Hot Spring Spas arranged according to Altitude.

Abbreviations used in the table :

auto. automobiles
avail. available
betw. between
ft. foot or feet

jinr. jinrikisha
L. Lake
m. miles
mid. middle

omn. omnibuses
R. River
st. station
volc. volcano

Table 14.

| No. | Altitude
in ft. | Geology | Spa | 地名(府縣名) | Situation | Means of access |
|-----|--------------------|---------------------------|--------------------------|----------------|---|---|
| 1 | 6950 | Andesite | Shibu [Suwa] | 澁 [諏訪] (長野) | N. slope of Mt. Yatsugatake (9676 ft.). | { 12.4 m. from Chino st. (Chino—7.4 m. by
omn.—Sasahara—5 m. on ft.—spa). |
| 2 | 6000 | " | Mciji ["] | 明治 ["] (") | Near Shibu. | { 9.8 m. from Chino st. via Sasahara.
2.5 m. W. from Shibu [Suwa]. |
| 3 | 5468 | " | Sakawa | 酢 川(岩手) | Near the summit of Mt. Sukawa-dake (5468 ft.). | { 27 m. W. from Ichinoseki st. (Ichinoseki—
13.6 m. by omn.—Mayu—7.4 m. on ft.—spa). |
| 4 | 5300 | Granite | Nakadusa | 中 房(長野) | On the mid. slope of Mt. Atake (5300 ft.) of Japan Alps | { 14 m. from Akashina st. (Akashina—5 m. by
omn.—Ariake—9 m. on ft. or horse-back
—spa). |
| 5 | 5180 | Andesite | Manza | 萬 座(群馬) | At the base of Mt. Manza (494 ft.) near Mt. Shirane. | { 10 m. from Kusatsu; 24.5 m. from Kawara-
yu; only on ft. |
| 6 | 5088 | Quartz Porphy. | Nikkō-Yumoto | 日光湯本(栃木) | By L. Yumoto; E. of Mt. Nikkō-Shirane (7544 ft.). | 17 m. from Nikkō by jinr. |
| 7 | 5000 | Andesite | Sandogoya | 三斗小屋(") | Near the summit of Nasu volc. (5000 ft.). | 7.5 m. up from Nasu-Yumoto on horse-back. |
| 8 | 4720 | Granite | Kamikochi | 上 高 地(長野) | Betw. Mt. Yake and Hotaka of Japan Alps. | { 27 m. from Matsumoto st. (Matsumoto—
11 m. by omn.—Shimajima—10 m. on ft.—
spa). |
| 9 | 4640 | Volc. Detr. | Karawa | 鹿 澤(群馬) | Betw. Mt. Asama and Azumaya; S. W. of Manza spa. | 9.8 m. from Tanaka st. on horse-back. |
| 10 | 4627 | Andesite | Tateyama | 立 山(富山) | On the Mt. Tate-yama (9689 ft.). | { 32 m. from Toyama (Toyama—8 m. by
jinr.—Kamitaki machi—23.4 m. on ft. by
steep path—spa). |
| 11 | 4500 | " | Renge | 蓮 華(新潟) | Mid. slope of Mt. Renge (4003 ft.). | 23.3 m. from Itoigawa st. (steep path). |
| 12 | 4500 | Volc. Detr. | Kusatsu | 草 津(群馬) | E. of Mt. Shirane (7500 ft.). | { 26.2 m. from Karuizawa. 50.8 m. from
Takasaki st. via Nakanojō (whence to spa
25 m. on horse-back). |
| 13 | 4500 | Andesite | Hoppo | 發 補(長野) | E. of Shibu [Hirao], on the way to Kusatsu from Shibu. | { 15.7 m. from Toyono st. (Toyono—10 m.
by auto.—Shibu—5.7 m. on ft.—spa). |
| 14 | 4000 | " | Sogayu | 酸 湯(青森) | S. W. of Mt. Hakkōda (5184 ft.). | 25 m. from Aomori by steep path. |
| 15 | 3785 | Tertiary | Otari | 小 谷(長野) | Ft. of Mt. Amakazari (6732 ft.). | { 39 m. from Akashina st. via Omachi, omn.
avail. |
| 16 | 3750 | Chichibu Series | Shirahone | 白 骨(") | N. E. of Mt. Norikura-gadake (10448 ft.) of Japan Alps. | { 29.5 m. from Matsumoto st. (Matsumoto—
14.5 m. by omn.—Inokuni—15 m. on ft. or
horse-back—spa). |
| 17 | 3464 | Granite | Kawamata | 川 俣(栃木) | N. of Mt. Nantai (8194 ft.). | { 12.3 m. from Nikkō-Yumoto, near Nishi-
zawa gold mine. |
| 18 | 3300 | Andesite | Yunohanazawa
Hakone | 湯之花澤(神奈川) | At the ft. of Mt. Koma-ga-dake of Hakone. | 8 m. from Yumoto (Hakone). |
| 19 | 3200 | Jurassic | Kawachi | 河 内(石川) | W. slope of Mt. Hakusan (8867 ft.). | { 30.2 m. from Komatsu st. (Komatsu—19.6 m.
by omn.—Onnabara). |
| 20 | 3000 | Granite | Kashi | 甲 子(福島) | At the base of Mt. Asahi (6498 ft.). | { 31.5 m. from Ikuu st. for 21.3 m. jinr. avail. |
| 21 | 3000 | Tertiary | Gosunki | 五 色(山形) | On the Itaya-tōge, near Mt. Azuma (volc., 6511 ft.). | { 14.3 m. from Shirakawa st. (Shirakawa—
2.4 m. by jinr.—Mafune). |
| 22 | 3000 | Andesite | Nasu-Yumoto | 那須湯本(栃木) | S. E. ft. of Nasu volc. (6310 ft.). | 2 m. from Itaya st., on ft. |
| 23 | 2944 | " | Takayu
(Yamagata-ken) | 高 湯(山形) | N. ft. of Mt. Zaō-dake (6034 ft.). | 10.4 m. from Kuroiso st., omn. avail. |
| 24 | 2760 | " | Ashino
Hakone | 蘆 之 湯(神奈川) | By L. Ashi; at the base of Mt. Koma-ga-dake of Hakone. | { 7.3 m. from Kaminoyama st., omn. avail.
8.5 m. from Yumoto (Hakone). |
| 25 | 2709 | Volc. Ash &
And. Detr. | Kirishima | 霧 島(鹿児島) | S. W. skirt of Mt. Karakuni-dake (or Kirishima), volc. (5610 ft.). | { 4.5 m. from Miyamoshita.
From Makizono st. to Kami-nakatsu 4 m. by
jinr., then 6 m. by kago or on horse-back. |
| 26 | 2700 | Andesite | Irinoyu | 入 之 湯(群馬) | E. slope of Usui-tōge (3038 ft.). | 6 m. down from Karuizawa, on ft. |
| 27 | 2700-2500 | " | Ihara | 伊 香 保(") | E. slope of Mt. Haruna (4808 ft.). | { 10.6 m. from Maebashi st. each by elec. tram.
20.3 m. from Takasaki st. } |
| 28 | 2600 | Alluvium | Suwa | 上 下 諏 訪(長野) | By L. Suwa, W. skirt of Mt. Yatsugatake (9676 ft.). | 120 m. from Tadamaichi st. (Tokyo). |
| 29 | 2510 | Liparite | Jozankei | 定 山 溪(北海道) | On the R. Toyohira. | { 17.7 m. from Sapporo (Sapporo—0.5 m. by
horse-tram.—Ishiyama). |
| 30 | 2500 | Andesite | Akakura | 赤 倉(新潟) | E. slope of Mt. Myōkō (8098 ft.). | 4.3 m. from Taguchi st., jinr. avail. |
| 31 | 2500 | Metals Series | Shirane | 四 萬(群馬) | Upper course of R. Agatsuma. | 11.3 m. from Nakanojō, omn. avail. |
| 32 | 2400 | Andesite | Unzen | 温 泉(長崎) | On the mid. slope of Mt. Unzen (4300 ft.). | 7 m. from Obama by chair. |
| 33 | 1223 | " | Miyamoshita | 宮 下(神奈川) | On the upper reach of R. Hayakawa, at the base of Mt. Myōjin and Myōjō. | 4 m. from Yumoto (Hakone). |
| 34 | 1150 | Quartz Porphy. | Arima | 有 馬(兵庫) | N. ft. of Mt. Rokkōzan (1550 ft.). | { 14 m. from Kōbe by kago.
6 m. from Sanda st., auto. avail. |

VI. Politico-economic Statistics with regard to Hot Spring Spas.

Table 15.

(A) Arranged according to Number of Visitors (1909).

| No. | Number of visitors per annum | Their total stays in days | Date of discovery | Number of inns | Number of inns with private baths | Number of public baths | Total number of baths | Spa | 地名 府縣名 | Population | Number of hotels |
|-----|------------------------------|---------------------------|-------------------|----------------|-----------------------------------|------------------------|-----------------------|--------------------------|--------------|------------|------------------|
| 1 | 108 192 | 549 947 | mytholog. age | 202 | 202 | 21 | ca. 1000 | Beppu | 別府 (大分) | 14045 | 3120 |
| 2 | 91 755 | 92 098 | — | 7 | 0 | 1 | 1 | Kojo | 神戶 (兵庫) | 2600 | — |
| 3 | 65 000 | 195 000 | 270-310 | 13 | 13 | 2 | 16 | Yugawara | 湯河原 (神奈川) | 500 | 80 |
| 4 | 64 370 | 126 910 | — | 33 | 33 | 10 | 43 | Kami Suwa | 上諏訪 (長野) | 13445 | 2887 |
| 5 | 56 787 | — | 81-113 | 18 | 8 | 5 | 13 | Iioka | 飯坂 (福島) | 4800 | 1000 |
| 6 | 52 500 | 325 200 | 1648-52 | — | — | 3 | — | Ichiku | 市來 (鹿兒島) | 729 | 127 |
| 7 | 45 565 | 203 152 | mytholog. age | 47 | 0 | 4 | 4 | D-go | 道後 (愛媛) | 1786 | 432 |
| 8 | 43 731 | — | 1104 | 9 | 9 | 3 | 12 | Kwankaiji | 觀音寺 (大分) | 150 | 30 |
| 9 | 39 297 | — | — | 19 | 4 | 4 | 8 | Yuno | 湯野 (福島) | — | 616 |
| 10 | 35 795 | 73 136 | 947-67 | 55 | 45 | 13 | 58 | Asama | 淺間 (長野) | 10000 | 300 |
| 11 | 35 443 | 55 325 | 1458 | 26 | 17 | 6 | 26 | Kominoyama | 上之山 (山形) | 8600 | 1000 |
| 12 | 35 000 | 150 000 | 850 | 17 | 17 | 1 | 18 | Takayu Yamazota-kei | 高湯 (山形) | 440 | 50 |
| 13 | 34 153 | 1 039 847 | 1156-59 | 20 | — | 5 | — | Yamaga | 山鹿 (熊本) | 7541 | 1588 |
| 14 | 33 557 | 193 217 | 28 B.C. | 38 | 12 | 15 | 94 | Ikao | 伊香保 (群馬) | 772 | 373 |
| 15 | 32 349 | 101 261 | bef. 270 | 35 | — | 1 | — | Takeo | 武雄 (佐賀) | 3200 | 570 |
| 16 | 31 451 | 165 670 | 806-10 | 41 | — | 23 | — | Shiobara | 鹽原 (栃木) | 2000 | 350 |
| 17 | 30 499 | 109 894 | 673-86 | 17 | 7 | 3 | 10 | Musashi | 武藏 (福岡) | 733 | 132 |
| 18 | 28 832 | 147 489 | 729-49 | 22 | 0 | 3 | 3 | Yamanaka | 山中 (石川) | 3300 | 760 |
| 19 | 28 704 | 96 879 | 718 | 11 | 11 | 1 | 12 | Awazu | 粟津 (津) | 580 | 150 |
| 20 | 27 082 | 181 604 | 1596-1615 | 38 | 28 | 4 | 32 | Owani | 大鰐 (青森) | 3779 | 615 |
| 21 | 26 875 | 58 471 | 1312 | 10 | 7 | 3 | 10 | Akayu Yamagata-kei | 赤湯 (山形) | 4135 | 801 |
| 22 | 26 528 | 60 395 | 1598 | 10 | 10 | 2 | — | Yumoto and Tenosawa | 湯本及雲之澤 (神奈川) | 1853 | 300 |
| 23 | 25 351 | 219 165 | 720 | 39 | 0 | 6 | 6 | Kinosaki | 城崎 (兵庫) | 1023 | 421 |
| 24 | 25 146 | 262 494 | 1409 | 34 | 34 | 4 | 38 | Hinagu | 日奈久 (熊本) | 2493 | 442 |
| 25 | 24 969 | 111 411 | 806-10 | 20 | 25 | 7 | 32 | Shuzenji | 修善寺 (靜岡) | 2252 | 355 |
| 26 | 23 638 | 174 306 | 668-71 | 45 | 45 | 3 | 51 | Atami | 熱海 (静岡) | 5866 | 803 |
| 27 | 22 808 | — | — | 19 | 0 | 7 | 13 | Yufuin | 由布院 (大分) | — | 300 |
| 28 | 22 523 | 94 168 | 724-29 | 33 | 17 | 24 | 41 | Hirao | 平穏 (長野) | 3031 | 747 |
| 29 | 22 000 | — | 717-24 | 17 | 17 | 1 | 22 | Higashi-yama | 東山 (福島) | — | 103 |
| 30 | 21 267 | 79 237 | 1573-92 | 20 | 14 | 1 | 15 | Wakura | 和倉 (石川) | 570 | 111 |
| 31 | 20 286 | 170 907 | bef. 1624 | 8 | 3 | 5 | 8 | Obama | 小濱 (長崎) | 2500 | 500 |
| 32 | 20 083 | — | 1884 | 25 | 25 | 3 | 30 | Awara | 蘆原 (福井) | 1000 | 180 |
| 33 | 19 375 | 96 759 | 725 | 18 | 18 | 2 | 20 | Yamashiro | 山手 (石川) | 2200 | 420 |
| 34 | 18 764 | 37 875 | 673-86 | 22 | 5 | 5 | 10 | Iessho | 別所 (長野) | 1002 | 222 |
| 35 | 16 560 | 115 920 | 1736-41 | 18 | 5 | 2 | 7 | Hijiori | 肘折 (山形) | 500 | 80 |
| 36 | 16 324 | 56 340 | 1025 | 10 | 10 | 1 | — | Miyanoshita | 宮下 (神奈川) | 617 | 103 |
| 37 | 16 055 | 45 590 | 947-57 | 20 | 9 | 7 | 16 | No-awa | 野澤 (長野) | 1059 | 288 |
| 38 | 15 033 | 45 454 | 1346-70 | 8 | — | 3 | — | Matsumeyama | 松之山 (新潟) | 240 | 30 |
| 39 | 13 263 | 132 636 | 821 | 25 | 25 | 2 | 27 | Atsumi | 濃海 (山形) | — | 100 |
| 40 | 12 968 | 55 356 | bef. 1270 | 15 | — | 5 | — | Kanmawa | 巖輪 (大分) | 1000 | — |
| 41 | 12 455 | 52 948 | 270 | 19 | 1 | 1 | 2 | Ureshino | 嬉野 (佐賀) | 1200 | 200 |
| 42 | 12 441 | 53 226 | 835 | 18 | 13 | 2 | — | Tamatsukuri Miyazaki-kei | 玉造 (宮城) | — | 380 |
| 43 | 12 378 | 32 833 | 1652-55 | 7 | 7 | 1 | 8 | Katayama-ku | 片山津 (石川) | 500 | 140 |
| 44 | 11 365 | 79 598 | 1624-44 | 33 | 33 | 10 | — | Ito | 伊東 (靜岡) | 8363 | 1379 |
| 45 | 11 361 | 50 595 | — | 0 | 6 | — | — | Tatsunokuchi | 長之口 (石川) | 270 | 45 |
| 46 | 11 288 | 22 450 | 834-48 | 15 | 15 | 2 | 17 | Onogawa | 小野川 (山形) | 175 | 60 |
| 47 | 11 018 | 70 462 | — | 9 | — | 6 | — | Soraku | 副田 (鹿兒島) | 800 | 200 |

* before.

| No. | Number of visitors per annum | Their total stays in days | Date of discovery | Number of inns | Number of inns with private baths | Number of public baths | Total number of baths | Spa | 地名 (府縣名) | Population | Number of houses |
|-----|------------------------------|---------------------------|-------------------|----------------|-----------------------------------|------------------------|-----------------------|----------------------|-----------|------------|------------------|
| 48 | 10 151 | 73 524 | 1387 | 13 | 13 | 7 | 20 | Dai | 臺 (岩手) | 170 | 27 |
| 49 | 10 132 | 64 605 | noolog. age | 27 | — | 2 | — | Arima | 有馬 (兵庫) | 1778 | 329 |
| 50 | 10 093 | 47 007 | — | 11 | 1 | 2 | 3 | Dake (Fukushima-ken) | 岳 (福島) | 230 | 22 |
| 51 | 10 039 | 34 415 | 1825 | 19 | — | 2 | — | Funagoya | 船小屋 (福岡) | 357 | 55 |
| 52 | 9 994 | 23 601 | — | 24 | 12 | 4 | 16 | Shimo-Suwa | 下諏訪 (長野) | 7000 | 1300 |
| 53 | 9 805 | 40 653 | 1573-92 | 4 | — | 3 | — | Hotta | 堀田 (大分) | 100 | 20 |
| 54 | 9 334 | 67 999 | 806-10 | 28 | — | 3 | — | Tawarayama | 俵山 (山口) | 716 | 112 |
| 55 | 8 637 | 65 718 | 1428 | 4 | 3 | 1 | 4 | Kamasaki | 鎌先 (宮城) | 141 | 14 |
| 56 | 8 300 | 50 000 | 1335 | 10 | — | 1 | — | Murasugi | 村杉 (新潟) | 250 | 33 |
| 57 | 7 603 | 86 447 | 540-71 | 14 | — | 5 | — | Nasu-Yumoto | 那須湯本 (栃木) | 700 | 60 |
| 58 | 7 133 | 138 285 | 1521-46 | 10 | — | 2 | — | Takarazuka | 寶塚 (兵庫) | 600 | 100 |
| 59 | 6 846 | 17 879 | 1661-73 | 8 | 8 | 2 | — | Unzen | 溫泉 (長崎) | 527 | 108 |
| 60 | 6 740 | 140 200 | bef.* 642 | 70 | — | 7 | — | Kusatsu | 草津 (群馬) | 1313 | 300 |
| 61 | 6 617 | 57 613 | 784 | 6 | 6 | 0 | 14 | Shima | 四萬 () | 320 | 58 |
| 62 | 5 984 | 25 650 | — | 2 | — | — | 2 | Ashinoyu (Hakone) | 蘆之湯 (神奈川) | 81 | 9 |
| 63 | 5 558 | 56 937 | — | 17 | 0 | 1 | 1 | Myōban | 明礬 (大分) | 280 | — |

(B) Arranged according to administrative district.

| No. | Number of visitors (1909) | Their total stays in days | Total stays in days per visitor | | | Number of spas† | Prefecture (Ken) | 府縣名 |
|-----|---------------------------|---------------------------|---------------------------------|------|------|-----------------|------------------|-----|
| | | | Average | Max. | Min. | | | |
| 1 | 19 891 | 138 627 | 7 | 27 | 1 | 48 | Hokkai-dō | 北海道 |
| 2 | 62 798 | 372 529 | 6 | 36 | 1 | 32 | Aomori | 青森 |
| 3 | 38 247 | 224 387 | 6 | 14 | 3 | 18 | Iwate | 岩手 |
| 4 | 28 367 | 230 415 | 8 | 26 | 1 | 39 | Akita | 秋田 |
| 5 | 65 926 | 369 737 | 6 | 17 | 4 | 27 | Miyagi | 宮城 |
| 6 | 229 618 | 920 822 | 4 | 13 | 1 | 30 | Yamagata | 山形 |
| 7 | 105 046 | 355 073 | 3 | 7 | 1 | 17 | Fukushima | 福島 |
| 8 | 128 236 | 403 833 | 3 | 19 | 1 | 45 | Niigata | 新潟 |
| 9 | 43 376 | 292 516 | 7 | 11 | 2 | 6 | Tochigi | 栃木 |
| 10 | 87 981 | 378 741 | 4 | 21 | 1 | 35 | Gumma | 群馬 |
| 11 | 7 995 | 34 884 | 4 | 8 | 1 | 11 | Ibaraki | 茨城 |
| 12 | 2 530 | 11 300 | 5 | 10 | 2 | 6 | Chiba | 千葉 |
| 13 | 3 156 | 10 426 | 3 | 8 | 1 | 7 | Saitama | 埼玉 |
| 14 | 4 158 | 20 605 | 5 | 10 | 2 | 3 | Tokyo | 東京 |
| 15 | 119 520 | 360 496 | 3 | 6 | 2 | 8 | Kanagawa | 神奈川 |
| 16 | 109 875 | 515 829 | 5 | 19 | 1 | 21 | Shizuoka | 静岡 |
| 17 | 31 332 | 130 979 | 4 | 30 | 1 | 11 | Yamanashi | 山梨 |
| 18 | 262 433 | 723 305 | 3 | 13 | 1 | 36 | Nagano | 長野 |
| 19 | 26 081 | 89 833 | 3 | 10 | 1 | 21 | Gifu | 岐阜 |
| 20 | 31 876 | 139 451 | 4 | 7 | 2 | 11 | Toyama | 富山 |
| 21 | 142 666 | 558 156 | 4 | 5 | 1 | 19 | Ishikawa | 石川 |
| 22 | 1 000 | 3 355 | 3 | 4 | 2 | 2 | Fukui | 福井 |
| 23 | 536 | 1 179 | 2 | 3 | 2 | 2 | Shiga | 滋賀 |
| 24 | 560 | 910 | 2 | 3 | 1 | 3 | Aichi | 愛知 |
| 25 | 4 389 | 8 768 | 2 | 5 | 2 | 5 | Mie | 三重 |
| 26 | 2 651 | 1 076 | — | 4 | — | 5 | Nara | 奈良 |
| 27 | 22 650 | 105 041 | 5 | 9 | 2 | 14 | Wakayama | 和歌山 |

† Spas refer only to those taken in calculation.

| No. | Number of
visitors
(1909) | Their total
stays in days | Total stays in days per visitor | | | Number
of
spas† | Prefecture (Ken) | 府 縣 名 | |
|------------|---------------------------------|------------------------------|---------------------------------|------|------|-----------------------|------------------|-------|---|
| | | | Average | Max. | Min. | | | | |
| 28 | 6 523 | 10 339 | 2 | 6 | 1 | 4 | Kyōto | 京 | 都 |
| 29 | 56 462 | 509 675 | 9 | 19 | 4 | 8 | Hyogo | 兵 | 庫 |
| 30 | 18 233 | 66 327 | 4 | 10 | 1 | 10 | Tottori | 島 | 取 |
| 31 | 30 519 | 86 289 | 3 | 7 | 1 | 13 | Shimane | 島 | 根 |
| 32 | 19 616 | 65 793 | 3 | 9 | 1 | 10 | Okayama | 岡 | 山 |
| 33 | 6 945 | 11 150 | 2 | 11 | 1 | 7 | Hiroshima | 廣 | 島 |
| 34 | 39 438 | 227 672 | 6 | 14 | 1 | 12 | Yamaguchi | 山 | 口 |
| 35 | 11 252 | 11 171 | — | — | — | 1 | Tokushima | 徳 | 島 |
| 36 | 404 | 12 091 | 30 | 30 | 30 | 1 | Kagawa | 香 | 川 |
| 37 | 49 165 | 204 952 | 4 | 5 | 2 | 2 | Ehime | 愛 | 媛 |
| 38 | 2 135 | 4 371 | 2 | 2 | 2 | 3 | Kōchi | 高 | 知 |
| 39 | 40 963 | 145 614 | 4 | 3 | 3 | 3 | Fukuoka | 福 | 岡 |
| 40 | 420 720 | 1 448 205 | 3 | 10 | 1 | 24 | Ōita | 大 | 分 |
| 41 | 93 366 | 1 496 486 | 16 | 30 | 3 | 4 | Kumamoto | 熊 | 本 |
| 42 | 29 123 | 193 928 | 7 | 8 | 1 | 7 | Nagasaki | 熊 | 本 |
| 43 | 46 980 | 192 630 | 4 | 18 | 3 | 4 | Saga | 長 | 崎 |
| 44 | 10 474 | 47 831 | 5 | 5 | 5 | 3 | Miyazaki | 佐 | 賀 |
| 45 | 142 164 | 931 374 | 7 | 19 | 2 | 41 | Kagoshima | 宮 | 崎 |
| 46 | 70 088 | 182 175 | 3 | 8 | 1 | 37 | Chosen | 鹿 | 島 |
| Total | | | Absolute | | | Total | | | |
| 2 677 364 | | | Average | Max. | Min. | 676 | | | |
| 12 300 947 | | | 5 | 36 | 1 | | | | |

VII. Summer Resorts (without hot spring) arranged geographically.

Abbreviations used in the table are the same as those in Table 14.

Table 16.

| No. | Name | 地名(府縣名) | Altitude in ft. | Situation | Means of access | Number of visitors per annum* |
|-----|-----------------------|-------------|-----------------|---|--|-------------------------------|
| 1 | Ōnuma Park | 大沼公園(北海道) | 400 | S. slope of Mt. Koma-ga-take (3627 ft.). | Summer st., 17.5 m. from Hakodate, in 51 min. | 7353 |
| 2 | Lake Towada | 十和田湖(青森-秋田) | 1470 | S. of Aomori. | { From the Kosaka copper mine, via Oyu spa, 22 m. horse and jinr. avail. } | 646 |
| 3 | „ Chūzenji | 中禪寺湖(栃木) | 4194 | S. ft. of Mt. Nantai-san (8194 ft.). | 10 m. W. from Nikkō, jinr. avail. | 16174 |
| 4 | „ Haruna | 榛名湖(群馬) | 4000 | W. ft. of Mt. Haruna-Fuji (4808 ft.). | 5 m. W. from Ikao spa., on ft. | 35 |
| 5 | Mt. Tsukuba | 筑波山(茨城) | 2892 | N. W. of Lagoon Kasumi-ga-ura. | { From Tsukuba to the ft. of the mount. 10 m., jinr. or omn. avail. } | 4311 |
| 6 | „ Kano | 鹿野山(千葉) | 1218 | On the borders of Kazusa and Awa. | { From Kasarazu st. to the ft. of the mount. 12 m., jinr. avail. } | — |
| 7 | „ Mitake | 御嶽(東京) | 2851 | W. of Tōkyō. | 8 m. W. from Ōme st., on ft. | — |
| 8 | Lake Ashi | 蘆ノ湖(神奈川) | 2380 | 11 m. S. W. from Yumoto (Hakone). | jinr. avail. | 1069 |
| 9 | Gotemba | 御殿場(静岡) | 1589 | E. ft. of Mt. Fuji. | 69 m. from Tōkyō, in 3 hrs. 48 min. | — |
| 10 | Sano Waterfall Garden | 佐野瀑園(„) | — | E. ft. of Mt. Ashidaka-yama (2966 ft.), near Mt. Fuji. | 0.8 m. N. W. from Sano st., jinr. avail. | 186 |
| 11 | Mt. Fuji | 富士山(静岡-山梨) | 12395 | On the boundary between Kai and Suruga Provinces. | { 7 m. from Yoshida-guchi to Fushibō, 5 m. from Gotemba-guchi to Fushibō, 5.5 m. from Su-bashiri guch. to Umagasa, 7.5 m. from Omiya-guchi to Hakomori dō. } | 25500 |
| 12 | Lake Shoji | 精進湖(山梨) | 4000 | N. W. ft. of Mt. Fuji. | { From Kōfu 17 m. (on ft.). From Gotemba or Ōtsuki to Yoshida by horse train, then 17 m. on ft. } | 283 |
| 13 | Fujimi | 富士見(長野) | 3200 | S. W. ft. of Mt. Yatsu-ga-dake (9676 ft.). | 109.9 m. from Iidamachi st., Tōkyō, in 8 hrs. 35 min. | — |
| 14 | Karuizawa | 軽井澤(„) | 3770 | S. of Usui toge (3088 ft.). | 88.3 m. from Ueno st., Tōkyō, in 4 hrs. 54 min. | 8599 |
| 15 | Togakushi | 戸隠(„) | — | E. ft. of Mt. Togakushi-yama (8000 ft.). | 12 m. N. W. from Nagano, on ft. | 1200 |
| 16 | Lake Nojiri | 野尻湖(„) | — | E. ft. of Mt. Madarao-san (5062 ft.). | 24 m. N. E. from Kashiwabara st., jinr. avail. | 890 |
| 17 | Yoro | 養老(岐阜) | — | N. E. ft. of Mt. Yōrō-san. | 0.5 m. S. from Ōgaki (by light railway). | 1795 |
| 18 | Ōwagayama | 大岩山(富山) | 700 | N. extremity of the ridge adjacent to Mt. Tate-yama (2989 ft.). | { 13.7 m. E. from Toyama, jinr. avail. From Namerigawa st. to Kami-ichi by light railway, thence 5 m. by jinr. } | 4734 |
| 19 | Mt. Kōya | 高野山(和歌山) | 2858 | 2 m. S. from R. Kinokawa. E. of Wakayama. | { 11 m. from Hashimoto st., of wh. h. to Kane (5 m.) jinr. avail., the rest on ft. or by kago. } | — |
| 20 | „ Hiei | 比叡山(京都) | 2800 | N. E. of Kyōto. | { 12 m. from Kōyaguchi st., thence 3.7 m. by jinr., the rest on ft. or by kago. } | — |
| 21 | „ Atago | 愛宕山(„) | 3043 | W. of Kyōto. | About 3 hrs. from Kyōto on ft. | — |
| 22 | Mino-o | 箕面(大阪) | — | N. of Ōsaka. | 6 m. from Saga st. to the summit. | — |
| 23 | Mt. Maya | 摩耶山(兵庫) | 2290 | N. E. of Kōbe. | { From Ōsaka by electric tram, in 30 m. 4.9 m. from Ikeda st. (11.5 m. from Ōsaka) } | 495 |
| | | | | | { 3 m. from Sannomiya; 6 m. from Sumiyoshi; kago avail. } | — |

*According to the statistics in 1909.

VIII. List of Watering Places (with the number of visitors in 1909) arranged geographically.

Table 17.

| No. | Geographical division | Number of
visitors
per annum | Number of
watering
places | Name of notable places | |
|-----|---|------------------------------------|---------------------------------|---|--------------------------------|
| 1 | S. W. coast of Hokkaido | 19 000 | 3 | Zenibako | |
| 2 | N. part of Pacific coast { | from C. Oma to C. Kurosaki | 5 951 | 5 | Same-minato |
| 3 | | from C. Kurosaki to C. Inubo | 81 807 | 22 | Shōbuda, Oarai, Hira-iso |
| 4 | | from C. Inubō to C. Nojima | 70 078 | 18 | Ohara, Ichi-no-miya, Katsu-ura |
| 5 | Bay of Tokyo (from C. Nojima to Misaki) | 10 416 | 13 | Hōjō, Tateyama | |
| 6 | Bay of Sagami (from Misaki to C. Iro) | 28 471 | 7 | Hayama, Zushi, Kama-kura, Katase, Oiso | |
| 7 | Bay of Suruga (from C. Iro to C. Ommae) | 2 495 | 3 | Shizu-ura, Numazu | |
| 8 | Sea of Tōtōmi (from C. Ommae to C. Irako) | 3 030 | 1 | Benten-jima | |
| 9 | Bays of Atsumi, Chita and Ise (from C. Irako to C. Daiō) | 105 231 | 12 | Gamagōri, Ōno, Morozaki | |
| 10 | Coast of Kii (from C. Daiō to Str. Yura) | 3 551 | 1 | Myōkō-ura | |
| 11 | Coast around the Bay of Osaka | 22 613 | 8 | Sakai, Hamadera, Suma, Maiko, Akashi, Iwaya | |
| 12 | E. part of Inland Sea (coast around the Bay of Harima) | 402 | 3 | Ushimado, Hoden | |
| 13 | Middle part of Inland Sea (coast around the Bay of Bingo) | 1 219 | 7 | Sensui-jima (Tomo), Itozaki | |
| 14 | W. part of Inland Sea (coast around the Bays of Hiroshima, Iyo and Suwo) | 5 921 | 5 | Baishinji | |
| 15 | S. coast of Shikoku | 2 000 | 1 | | |
| 16 | Coast along the Seas of Hibiki-nada, Genkai and Amakusa (from C. Kawajiri to Shimo-no-seki and N. and W. coast of Kyushu) | 69 775 | 7 | Mutsure-jima, Karatsu, Nezumi-jima | |
| 17 | S. part of the Sea of Japan (from C. Kawajiri to C. Suzu) | 6 123 | 13 | Inasahama, Uratomi, Mikuni | |
| 18 | N. part of the Sea of Japan (from C. Suzu to C. Ōma) | 10 973 | 7 | Shimao, Higashi-iwase, Kujira-nan-i, Koto-ura | |
| | Total | 449 656 | 136 | | |

* Watering places refer only to those taken in calculation.

PART III

NOTES

on

Prominent Spas, Seaside Resorts,

and

Summer Retreats.

HOKKAI-DŌ

Although hot springs in *Hokkai-dō* are generally believed not to be small in number, those already known are situated mostly in the south-western portion of the main island. Of these only such ones, as located in the places accessible by some means of transportation, are utilized but not a few are neglected owing to their remote and inconvenient positions. Hot springs gushing out in volcanic regions are, as a rule, sulphur springs of high temperature. In non-volcanic districts hot springs are comparatively seldom met with and are inclined to be carbonated, saline, or weak saline springs. We are going to describe some of these hot springs that are comparatively abundant and properly taken care of.

Noboribetsu. These hot springs are the most famous in Hokkai-dō and lie 4,3 m. N. W. of the *Noboribetsu* station, *jinrikisha* or coaches being available over the distance. When these springs were first discovered many years ago, no path was found to go there. It was only in the Ansei Era (1854-1860) that officials were despatched by the Shōgunate Government to lay out a road there-to. In the course of the Meiji Era (1868-1912), the road was much improved. The place is 600 ft. above sea-level and very rich in scenic charms in all seasons, especially in spring with various blossoms and in autumn with tinted leaves; nevertheless it is delightfully cool in summer. Baths and inns offer satisfactory accommodation and make the place an ideal holiday resort. Springs gush out from a crater on the foot of a hill, 0,2-0,3 m. from the inns up the *Noboribetsu* River. This place is the origin of the *Kusuri-e-san-betsu*, a branch of the *Noboribetsu*, and forms a basin surrounded by precipices on all sides except in the S. Abundant steam with sulphurous gases is constantly issuing out with resounding noise from two or three places on its bottom and white vapour fills the whole valley. Rocks acted upon by the steam especially on the precipices on the N. and the E. side are turned to gray and loose masses sometimes with yellow ocher and sulphur. Only in some spots, there is comparatively fresh andesite. Near the crater there are several other spots, where springs of different temperature and composition occur. see Plate 12

Some 3,5 m. distant from the crater, we find an open-air bath of a hot spring called "*Karurusu*." About 0,2 m. to the N. W. of the crater, there is an old craterlet with a geyser, which was throwing up water tens of feet high till 1880. Stepping a little distance further uphill we come across an oblong pond of gray-coloured warm water extending to the N. E., with the circumference of 165 ft. and surrounded by rather steep hills. At the S. E. corner of this pond, there are cold and hot springs. The scenery of this place is at its best in autumn when maples turn crimson, and no holiday-maker can enjoy their stay too much. The *Shimofuri Waterfall*, sometimes called "*Kachidoki-no-taki*," is in the upper course of the *Noboribetsu*, and there are many scenic wonders in its vicinity. *Mt. Noboribetsu* may be climbed from this place. The hot spring village has more than fifty houses inhabited by some two hundred people and also a post office. Two inns out of ten have their own baths and besides there are five public baths. There is a billiard room for visitors' amusement. Bathers were numbered 4,975 in 1909 and their total stays 33,250 days long. In recent years foreign visitors have been gradually increasing and the average number is ten a day. Although visitors come here all the year round, this place is most prosperous in summer. see Plate 14

Karurusu. 8 m. N. W. of the *Noboribetsu* station; 800 ft. above sea-level. A branch of *Mt. Noboribetsu* slopes towards the place and surrounds the place on all sides, a clear stream flowing across the valley and hot springs gushing out at its eastern extremity. The whole view is very lovely. Temperature indoors neither runs above 85° F. even in dog days nor falls below 33° in winter when snow covers the grounds 4 ft. high. Five houses. Although these springs were recognized by ancient inhabitants, they came to the knowledge of people from Japan Proper about in September, 1888, when surveyers accidentally discovered them during their engagement in the survey for colonial troop barracks. see Plate 7, fig. 1 and 4

Omnibuses available over the distance of 4,2 m. from the *Noboribetsu* station (fare: 35 *sen* a passenger), and farther stage of journey, 3,7 m. by mountain road (if on horse-back, charge 50 *sen*). 8,3 m. from the *Horobetsu* station (horse, fare 60 *sen*). Three inns, all without private baths. 298 bathers in 1909, most bathers come from neighbouring places for recuperation.

There are two springs, one *Kame-no-yu* or "tortoise spring" and the other *Tsuru-no-yu* or "crane spring." The springs, it is said, are named after the famous *Karl'sbad* of Austria on account

of similar property. (*Karurusu* is the Japanised pronunciation of "*Karls*").

Jōzankei. 17.6 m. from Sapporo up the *Toyohira* River, 2,510 ft. above sea-level. *Jinrikisha* and coaches available. *Jōzankei* is an important place on the road to *Abuta*. Although natives knew their existence long ago, they were recorded in history for the first time in the Ansei Era (1854-1860). In August, 1869 an itinerant monk *Jōzan* opened a bath, hence the present name. The public bath in front of the place now called "*Moto-yu*" is the very one set up by the monk. Hot springs gush out among a thickly wooded valley surrounded by high mountains and penetrated by a beautiful stream of the *Toyohira* with wonderful rocks. *Naka-no-yu* or "middle spring" was accidentally discovered by a visitor to "*Moto-yu*" while sauntering in the neighbourhood some time in 1674. *Shika-no-yu* or "deer spring" was discovered in 1984-5. Three inns; baths are dug into natural rocks and cemented all over. The bathers were numbered 2,080 in 1909 and their total stays 37,276 days long.

Sea-Bathing at Zenibako. 0.5 m. N. E. of the town of *Zenibako*. The seashore is where the wide expanse of *Ishikari* Plain meets the Japan Sea to the N. W., and the range of view is broad and wide. The whole shore and the bottom of the sea 360 ft. off are of white sand. Sand dunes stand side by side on the shore and there is a salt-manufactory. There are six rest-huts for the use of bathers, besides two ordinary inns with hot sea-water bath. The place is patronized by people from *Sapporo*, *Otaru*, etc. The number of bathers was 10,000 in 1912.

Aoyama and Miyagawa. Both the hot springs are on the River *Yuno-kawa*, 3 m. N. E. of the *Kombu* station and 4.3 m. N. W. of the *Kariputo* station, and less than 0.2 m. distant from each other. The district is a wide highland, forming a colony dotted with 75 farm-houses and now under cultivation.

Ōnuma Park. There are two lakes at the N. extremity of *Kameda-gōri* in Oshima province, that is *O-numa* and *Ko-numa*, both 400 ft. above sea-level. The whole shape is like the numeral 8, the narrowest portion being called "*sebatto*." The part of the lakes to the W. of *Sebatto* is named "*Ko-numa*" and that to its E. "*Ō-numa*," the entire water is known again as "*Ō-numa*." This was certainly a crater of an ancient volcano. *Ko-numa* is 1.7 m. N. to S., the whole lakes 3.8 m. E. to W., and their circumference 21.2 m. There are three villages on the lake shore. The whole surface of water looks like a large sheet of mirror, and water is clear and drinkable, its depth being 2 or 3 to 10 fathoms. The bottom is generally of slate covered by grits, five streams contributing to the lakes on all sides; *Mt. Koma-ga-dake* rears its sword-like peaks towards the sky, and adds a great deal to the quaintness of the sight. There are islets of various shapes and sizes, more than 140 in number, the biggest 180,000 square ft. and the smallest less than 1 square ft., all of them being thickly wooded, mostly with cherry, yew, maple, etc. The lake shore has many indentations which form bays, and the water is very calm. *O-numa* is famous for carp, gibel, etc., their annual production amounting to ¥ 3,000 worth. Thus the landscape is beautiful and air is pure and invigorating. Temperature: 20° to 30° F. in winter, 40° to 60° in spring, 70° to 80° in summer, and 50° to 60° in autumn.

For the convenience of visitors a temporary station *Onuma-Kōen* is opened every year from May 1st till October 31st, 18.3 m. from *Hakodate*.

Inns: *Kōyō-kyūan* (Maple House), *Hyakkwa-en* (Gardens of Hundred Flowers), etc. The former faces *O-numa* towards N. E., with its garden planted with maples, and commands a wide view of cloud-capped *Koma-ga-dake* and islets in the lake. The latter stands on *Ko-numa* with its fine view of white-winged vessels sailing past among many islands. Boat to hire on the lakes. Foreign and Japanese visitors are numbered more than 2,000 in August and September.

Yunokawa lies 3.5 m. N. E. of *Hakodate*, and not far from the volcano *Koma-ga-dake*. The district is flat, being located at the mouth of the *Matsukura* and facing the strait of *Tsugaru*. Its seashore is of white and beautiful sand, whence one can descry distant mountain ranges in *Mutsu* Province over the sea. The place is flanked by hills on the N. and the E. side and forms a truly quiet pleasure resort. The scenery is best in spring, although of course it is attractive through the whole year. Its climate is mild and not subject to sudden changes, thus making it a good health resort. Houses 124 and population 593. Village, post, police offices and primary school. Many shops standing side by side form a regular street. Its discovery dates back to not later than 1654, but the place was abandoned till 1886, in which year the citizens of *Hakodate*, convinced of the

existence of a hot spring vein in the neighbourhood, tried to bore the earth with a machine for making a well. Their effort was in vain even when they reached 500 ft. to 1,900 ft., but overcoming extreme financial difficulty, was awarded with ample success. As the quantity of this spring amounted to 8,5 hectolitres per hour, next year they began to run the place as a public resort and since then have contrived to bore more springs, lay out a park and repair roads leading there-to, thus bringing about to-day's prosperity. Electric cars from *Hakodate* over the distance of 4,5 m. The road is good and post, telegraph, and telephone communications are sufficient. Convenient communications also with other places.

Eight hotels; the best are *Senshin-kwan*, *Rinchō-kwan*, etc., all provided with private baths and three inns also with special baths for foreign visitors. Two doctors in the town and one shampooer. The room charge for foreigners about ¥25 a month. Bathers come chiefly from *Hakodate*. The average visitors are 2,500 per annum; total stays 600 days long. Not a few visitors from among the crews of foreign men-of-war anchoring at *Hakodate* and foreign summer tourists to *Hokkai-dō*. The season is from April to June.

see Plate 7,
fig. 3

AOMORI-KEN

Asamushi is at the S. head of *Aomori Bay* and 7,3 m. from *Aomori*. A short distance from the station of the same name. Broad seascape with small islands dotted here and there and the abundance of pine trees add much to the charms of the surroundings. There are eight sources of hot springs. History tells us that the famous monk *Enkō Daishi* on his N. E. itinerancy happened to come here and noticed a deer bathing in the sea. Convinced of the existence of a hot spring, he persuaded the villagers to establish baths. This is generally believed to be the inception of the present baths. Naturally early inhabitants religiously adored the springs and dared not to bathe in them but used them only for the purpose of dipping and steaming hemp for clothings. Thus the springs obtained the name of "*Asa-mushi-no-yu*" or 'Spring for steaming hemp,' gradually corrupted into the present characters. 210 houses; population 1,050. Salt-manufactories. Communications also maintained by sea. The famous *Tsubaki-yama* is 12,2 m. N. to the village. 26 inns; 2 doctors. Bathers come from the neighbouring places but an appreciable number of visitors from distant countries by railway, etc. Visitors 3,373 in 1909 and their total stays 11,997 days long.

Ōwani. A few minutes' walk from *Owani* station on the Ō-u Line, about 7 m. from *Hirosaki*, and 270 m. from *Sendai*.

The hot springs are situated to the S. W. of *Aomori-ken*, near the boundary of *Akita-ken*, surrounded by mountains, of which *Akara-yama* to the S. and *Iwaki-san* or the "*Tsugaru-Fuji*," (4,600 ft. high) so called on account of its similarity in shape to the famous Mt. Fuji, rises to the W. of the springs. The small stream *Hira-kawa* runs through the village which has 615 houses and 3,800 inhabitants. The spa lies at an elevation of 248 ft. above sea-level. Excellent apples are grown in the neighbourhood (the yearly production ca. ¥15,000).

In the environs of Ōwani bath there are many hot springs, namely *Nuru-yu* on the N. E., *Yumoto* at the foot of *Iwaki-san* and *Ikariga-seki* on the S. E. and others. *Kuradate* hot springs lie on the opposite side of the river from Ōwani. One of the hot springs, *Yamabuki-no-yu*, was found before the Keichō Era (1596-1615).

No less than 32 hot springs are found in Ōwani bath in the space of half a mile. There are two resident doctors, 38 inns with private baths and 4 public baths at Ōwani Spa, in 1909 the visitors numbered 27,082, the total stays being 181,604 days.

Kuradate. The bath lies on the opposite side of the river *Hira-kawa* to *Owani*. The village has 469 houses and 3,400 inhabitants, 5 public baths. Each inn is provided with its own bath. The inhabitants of the villages *Ōwani* and *Kuradate* built greenhouses near the source of springs, utilizing the heat in the germination of *soja* beans and other vegetations to advantage. *Kuradate* has one resident doctor. One of the hot springs, "*Shimo-no-yu*," was known in the Kenkyū Era (1190-1199). Number of visitors: 1,755, their total stays being 6,157 days (1909).

Ikariga-seki. About 10 minutes' walk from the *Ikariga-seki* station on the Ō-u Line, 5 m. S. from *Ōwani*, 36 m. S. from *Aomori*. *Jinrikisha* and omnibuses available.

The springs occur in a valley which runs N. and S. between the mountains. The village is situated at an elevation of 680 ft. above sea-level and has about 1,000 inhabitants.

Post, telegraph and telephone office. Local places of interest: *Senjō-ga-taki* ('Ten thousand feet Waterfall') 5 m., *Ito-taki* ('Thread Waterfall') 2 m., *Furu-ake-fudō* 2,5 m., ruins of an old castle 2 m., etc., are all worth visiting. The hot springs are here 5 in number, of which two belong to public baths.

The hot spring "*Nitsu-no-yu*" was known before 1592. Number of visitors: 3,212, their total stays being 33,304 days (1909).

IWATE-KEN

Dai. Passengers from the S. alight at *Hanamaki* station on the N. E. Main Line, 22 m. N. from *Morioka*, 91 m. N. from *Sendai*, those from the N. alighting at *Ishidoriya* station, both stations are about 6 m. distant from the springs, *jinrikisha* and omnibuses being available.

Dai is a small bathing-place with mountains on all sides, lying on the bank of the upper course of a tributary flowing into the River *Kitakami-gawa*, and at an altitude of some 1,500 ft. above sea-level. Pop. 170. A kind of porcelain called "*Dai-yaki*," produced in this village, is a speciality of this district. *Dai* Mine is about 5 m. from the place.

There are more than 14 hot springs in the village, of which seven belong to public baths.

Number of visitors: 10,151, the total stays being 73,524 days (1909). Most bathers come from neighbouring places for recuperation.

Shido-daira. About 6 m. W. from the *Hanamaki* station, 91 m. N. from *Sendai*, and 308 m. from *Tōkyō*, the journey from the capital taking about 13 hours. *Jinrikisha* and omnibuses available from *Hanamaki*.

The bathing-place lies on the N. bank of a small stream, the *Toyosawa-gawa*, with mountains to the E. and W., at an elevation of 700 ft. above the sea. Pop. 120.

In this neighbourhood there are *Osawa Hot Springs* (1,5 m. from *Shido-daira*) and *Namari Hot Springs* (5 m.). Report claims that the spring was already known to the people before the Enryaku Era (782-806 A.D.).

The place has only one spring and one inn. The new bath-house has recently been constructed by the keeper of the inn *Kubota Ryo-kwan*, and has no equal in its construction and appointments in the neighbouring spas. Number of visitors: 2,914 (1909).

Ōsawa. 8,5 m. W. from the *Hanamaki* station; 1,5 m. from *Shido-daira*; 2,5 m. from *Namari Hot Springs*.

The place is situated on both banks of the *Toyosawa-gawa*, with mountains on the E., the other three sides being open country communicating with the towns. Pop. 280. Post-office in the village. 2 public baths and a playground on the hillside. The eight scenic views of Ōsawa are well known in this place for their beauty. Number of visitors: 1,975 (1909).

AKITA-KEN

Ō-yuzawa, 1,3 m. N. W. from the *Jimba* station on the Ō-u Line, *jinrikisha* available over $\frac{2}{3}$ of the distance, is shut in on all sides by densely wooded mountains, and here being one of the three most beautiful forests of Japan. The bath-place lies at an elevation of 700 ft. above the sea, and is a quiet summer retreat. The existence of the hot springs is believed to be dated from the time of the eruption of *Bandai-zan* (an active volcano in Iwashiro Province) in 1889. Number of visitors: 3,000, the total stays being 21,700 days (1909).

Akayu is situated a little higher up in the mountain 0,5 m. from *Ō-yuzawa*, the water being a carbonated spring containing iron. Number of visitors: about 1,000, the total stays being about 10,000 days (1909).

Ō-yu. From Ō-date (251,5 m. from *Fukushima* on the Ō-u Line, in 12 hrs. 27 min.) to *Kosaka* Mine (a mining town containing 14,975 inhabitants) by *Kosaka* Railway (14,1 m., in 1 hr. 25 min.), built specially for the better exploitation of the *Kosaka* Mine, thence 8,6 m. S. E. by *jinrikisha* or omnibuses.

The bath-place is a small village with 5,438 inhabitants (660 households). Electricity, generated by the waterfalls near the village, supplies both motive power and light to Kosaka Mine. The hot spring "*Kami-no-yu*" is said to have been found in the Bummei Era (1469-1487 A.D.), but the bath-houses were first constructed in the Era of Eihō (1673-1681). There are four springs in the village, not more than 0,5 m. distant from each other, the water being all weak common salt springs. Number of visitors: 2,125, the total stays being 14,875 days (1909). see Plate 15

An interesting excursion may be made from the spas in this region to the beautiful mountain lake *Towada* with its wonderful landscape view. **Lake Towada**, literally "Ten Inlets Lake," 5,6 m. wide and 6,2 m. long, circumference 31 m., is 1,500 ft. above sea-level, is surrounded by the Mt. *Hanabe-yama* (3,200 ft.), *Towada-dake* and *Herai-dake* (3,250 ft.). The Hakkōda mountain range towers high on the horizon to the north, giving a fine view from the springs. The shore of the lake is picturesquely studded with grotesque rock formations, and its irregular contour and many windings, from which come the principal inlets, give an interest to all lovers of nature, the inlets in number giving us the origin and meaning of the name "Ten Inlets Lake," while another attraction is found at the southern end in a number of charming little islands (*Ebisu-jima*, *Kabuto-jima*, *Yoroi-jima*, *Tane-jima*, *Hōrai-jima*, etc.) and the famous old Shintō temple "*Towada-jinja*" in its setting of greenery on the shore but a short distance from *Ebisu-jima*. In autumn the glories of the wonderful foliage are shown in all their splendour, as the maples turn their leaves into brilliant scarlets and reds, these being reflected in the placid waters present a sight never to be forgotten. Boating facilities give opportunity for pleasure and enjoyment while the angler will find a sporting satisfaction, the lake being abundant in trout. The outlet forms another interesting feature in the cascade "*Chōshi-no-taki*," which tumbles in a foaming spray to pour itself into the River *Oirase*. Refreshment places near the lake are open during the summer months.

Ōdaki (8,6 m. S. E. from the *Odate* station) lies by the S. bank of the *Yoneshiro-gawa*, Ōdaki having 300 inhabitants. The stream itself adds a beauty and distinction to the place. In the public bath there is a hot waterfall for massage. Each inn is provided with private bath. Number of visitors: 2,023, the total stays being 10,482 days (1909).

Innai-Yunosawa. 1,8 m. from *Innai* station (122 m. from *Fukushima*, in 7 hrs.) on foot.

The bath-place (679 ft. above the sea) is on the E. and W. enclosed by mountains, the stream *Yunosawa-gawa* with clear water running close by the village. Pop. 800. Several waterfalls (*Yuno-no-taki*, *Tōnaru-no-taki*, *Ō-daki*, *Mitsuai-no-taki*, etc.) each a wonderful sight. In 1907 the Garrison Hospital of Akita built here a sanatorium. There are two hot springs in this spa, the flow of water being very generous. Number of visitors: 2,488, the total stays being 12,858 days (1909). see Plate 10

Shibukuro. The valley of the upper course of the *Tama-gawa*, which originates from high mountains, such as *Yake-yama*, *Eboshi-dake* (5,356 ft.), *Koma-ga-dake* (5,264 ft.), etc., bounding two prefectures Iwate and Akita, is generally called "*Shibukuro-zawa*" or simply "*Shibukuro*." In this valley occur a number of hot springs, of which the most noted are *Hato-no-yu*, *Tsuru-no-yu*, *Kuro-yu*, and *Kani-no-yu*.

These springs, more than 2,700 ft. above the sea, are located in Sempoku-gun, Akita Prefecture, about 34-45 m. N. E. from the *Ōmagari* station on the Ō-u Main Line, not far from Lake Tazawa, all being very difficult to be reached. Waters coming from the springs are all received by the *Tama-gawa*, and on its bed is found sinter-deposit in crust of crystalline structure. It has been lately ascertained that the crust is identical with the so-called "*Hokutōlite*," a kind of new mineral found at Hokutō Hot Springs in Taiwan. see Plate 3

MIYAGI-KEN

The new Riku-u Line (94,6 m.) starts from *Kogota* and is to terminate at *Sakata* Harbour on the Japan Sea. It crosses the Ō-u Line at *Shinjō* and runs along the bank of the *Mogami-gawa*. The line is now under construction and only the section to *Kawatabi* (20 m. from *Kogota*) has been opened to traffic. In the vicinity of *Kawatabi* and *Naruko* many hot springs are found, of which the following are noted:

Yuizumi and Onikobe. The hilly district N. W. of the town of *Iwade-yama* has been noted for horse-breeding for centuries, and several of the favourite mounts of the late Emperor were

raised here. The neighbourhood being volcanic, hot springs are found issuing almost everywhere, especially along the upper course of the *Arao-gawa*. These springs are in two clusters, one called *Yuizumi Hattō*, or the "Eight bath-resorts at Yuizumi," this latter term itself meaning "hot spring." The other is called *Onikōbe Gotō*, or the "Five bath-resorts at Onikōbe." The eight spas of Yuizumi are *Kawatabi*, *Tanaka*, *Aka-yu*, *Moto-kuruma*, *Shin-kuruma*, *Naruko*, *Kawara*, and *Nakayama*, and these are separated from each other by a distance varying from 100 yds. to 4 m., all being situated along the highway on the plain. Though lying in out of the way places, these resorts are reputed to be the best N. of Sendai in general arrangements and accommodation. *Kawatabi* (4 m. from *Ikezuki*) has five baths and is credited with being efficacious for leg-dropsy; *Tanaka* 1,3 m. W. is sheltered at the back by hills and has three baths; *Aka-yu* lies 0,5 m. off *Tanaka*, and its three springs yield carbonic acid waters; *Moto-kuruma* is 0,5 m. W. from the above and its waters belong to salt springs; *Shin-kuruma* occupies about the centre of the eight resorts and has five springs. *Naruko*, 3 m. from *Kawatabi*, is a prosperous village, containing more than one hundred houses; *Kawara* lies 0,3 m. further W., while the last resort, *Nakayama*, has two springs and is bounded on one side by a river, the other three sides being shut in by mountains. The vicinity abounds in natural beauty spots such as *Kokurosaki* with its dense pine woods variegated with maples, *Ikezuki-numa*, and *Mizuno-kojima*, the latter an islet in the *Tamatsukuri-gawa* with a shrine to Benten standing on it. There are also waterfalls, pools, high cliffs, etc. Of these eight springs above mentioned the two, *Tanaka* and *Nakayama*, were completely destroyed by a terrible flood of the *Arao-gawa* in 1910.

The Onikōbe Hot Springs, being located amidst hills and therefore less accessible than those of Yuizumi, are not so generally known. Of the five resorts, the first in order is *Sabusawa*, which is 5 m. distant from *Naruko*; then *Mitaki*, separated by 0,8 m. from *Sabusawa*; *Todoroki*, 2,4 m. from *Sabusawa*, and lastly *Ara-yu*. The waters of all these springs except *Ara-yu* (sulphur spring) are alkaline salt springs. At *Todoroki* is a geyser, or '*fuki-age*.' Formerly there were two holes from which intermittent hot springs gushed out with a loud detonation, but in 1901 the larger of the two ceased to be active, and at present the geyser comes out only from the smaller hole, at intervals of about 2 hrs., the column shooting up about 20 ft. The water contains bicarbonate of calcium, sulphate of potash, etc., the geyser having the property of encrusting and transforming into a fossilized state any plants or objects coming within its range.

It is very interesting to see the geyser in operation. The hot liquid gushes out from a hollow (about 10 ft. long and 4 ft. wide) in rocks, which was formed by the action of the water in course of time. A few minutes before the discharge of the geyser, steam evolves this being followed by a fearful detonation, while bubbling water comes out from several spots, and at last the hot water column shoots out suddenly straight up into the air, reaching a height of about 20 ft. The evolution of bubbling water continues for about two or three minutes with the same force, forming a small pool in the hollow to the depth of about 1 ft. As soon as the evolution stops, the whole amount of hot water disappears entirely, so that the pool at once dries up, being probably received again by the same hole under-ground from which the water comes forth. This process is repeated about 12-15 times every day.

| | Naruko | Moto-kuruma | Shin-kuruma | Kawara | Kawatabi | Tanaka | Akayu | Todoroki |
|-------------------------|--------|-------------|-------------|---------|----------|---------|-------|----------|
| Number of visitors 1900 | 2,734 | 1,920 | 1,617 | 315 | 3,950 | 340 | 1,565 | 927 |
| The total stays in days | 12,926 | 7,760 | 5,940 | 1,610 | 16,020 | 1,610 | 7,360 | 5,695 |
| Found in | 835 | 1,781 | 1,844 | unknown | 901 | unknown | 1,781 | 1,615 |

Kamasaki. 3 m. from *Shiroishi* station on the N. E. Main Line, 189,3 m. from *Utsu*, Tokyo, in 8 hrs. 31 min. The spa lies on the upper course of the *Kosute-gawa* and is shut in on all sides by hills. The springs are said to be efficacious especially for wounds, bruises, *kakke*, skin diseases, etc. The spa is a small village, at an elevation of 600 ft. above the sea, inhabited by only 140 souls, the springs dating back to 1428 A.D. Number of visitors: 8,637, the total stays being 65,718 days (1909). Inns: *Ichijō-kwan*, *Kimura-ya*, *Mogami-ya*. The spa is visited mostly by patients to take the cure by the waters.

YAMAGATA-KEN

Atsumi is situated on the coast lying between Niigata and Sakata, about 85 m. N. E. from Niigata (in 8 hrs. by steamship, fare ¥ 1,40), on the N. bank of the *Atsumi-gawa*, and is 1 m. distant from the sea. The beach of *Nezumi-ga-seki*, noted for its beautiful scenery, is 7,5 m. S. W. from the spa. The springs were found in 811 A.D., and are believed to be efficacious for anaemia, wounds and disorders of the female genital organs. The *Yunohama* hot springs lie also on the coast 17 m. further N. from the spa. Atsumi has electric light and telephone. Each inn is provided with private bath. Number of visitors: 13,263, the total stays being 132,636 days (1909).

Yunohama. 2 m. N. W. from *Kamo-machi*, the steamship plying between Niigata and Sakata calls the town *Kamo*. A white sandy beach extends from the spa towards the N. E. to Sakata, and near Yunohama issue hot springs from the water's edge on the beach, so that the spa affords also facility for hot sand baths. Good for sea-bathing. Number of visitors: 4,203, the total stays being 52,461 days (1909). The result of meteorological observations made in Kamo-machi is as follows:

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity | |
|---------------------|-----------------|------|------|-------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|---------|
| | Mean | | | | Absolute | | | Total | % | Total | Mean | Dir. |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | | |
| For the year, 1912. | | | | | | | | | | | | |
| January | 2.0 | 4.3 | -0.5 | 4.9 | 11.6 | -4.7 | 72 | 49.8 | 16 | 176.3 | 7.1 | |
| February | 4.2 | 7.5 | 1.3 | 6.2 | 21.6 | -3.9 | 69 | 72.9 | 23 | 252.7 | 5.7 | |
| March | 4.6 | 8.3 | 1.3 | 7.0 | 19.0 | -1.2 | 70 | 122.0 | 33 | 182.5 | 5.5 | N 59° W |
| April | 10.2 | 14.5 | 5.6 | 8.9 | 23.0 | -1.2 | 68 | 175.0 | 44 | 118.3 | 5.2 | N 89° W |
| July | 22.1 | 25.5 | 19.2 | 6.3 | 29.1 | 15.1 | 90 | 142.9 | 32 | 316.4 | 2.7 | N 75° W |
| December | 3.6 | 6.6 | 0.7 | 5.9 | 11.4 | -2.7 | 77 | 15.5 | 5 | 227.7 | 6.1 | N 48° W |
| For the year, 1913. | | | | | | | | | | | | |
| January | 1.3 | 3.5 | -1.4 | 4.9 | 8.2 | -6.2 | 76 | 29.8 | 10 | 226.3 | 6.9 | N 49° W |
| February | 2.0 | 4.9 | -1.0 | 5.9 | 14.3 | 6.7 | 69 | 74.8 | 25 | 116.3 | 6.0 | N 52° W |
| March | 3.4 | 6.3 | 0.1 | 6.2 | 15.5 | -5.0 | 67 | 134.8 | 36 | 120.0 | 5.3 | N 64° W |
| April | 10.1 | 14.7 | 5.2 | 9.5 | 25.4 | -1.2 | 72 | 170.9 | 46 | 255.0 | 3.5 | |
| May | 13.5 | 17.6 | 9.5 | 8.1 | 25.2 | 2.9 | 72 | 177.0 | 40 | 65.3 | 4.2 | |
| June | 17.9 | 21.7 | 11.2 | 7.5 | 25.4 | 9.4 | 79 | 150.9 | 35 | 121.1 | 4.1 | |
| August | 22.0 | 26.3 | 18.2 | 8.1 | 29.5 | 15.5 | 82 | 222.8 | 52 | 273.0 | 3.1 | |
| September | 18.9 | 23.3 | 15.2 | 8.1 | 27.0 | 10.9 | 80 | 182.5 | 49 | 234.7 | 2.9 | |
| October | 14.7 | 18.5 | 11.1 | 7.4 | 24.7 | 5.2 | 79 | 133.9 | 39 | 212.7 | 3.9 | |
| November | 9.3 | 12.6 | 5.5 | 7.2 | 19.1 | 0.3 | 68 | 90.7 | 30 | 254.6 | 5.3 | |
| December | 4.8 | 7.2 | 1.8 | 5.4 | 18.0 | 1.7 | 76 | 43.1 | 15 | 305.0 | 5.8 | |
| For the year, 1914. | | | | | | | | | | | | |
| January | 4.0 | 6.9 | 0.8 | 6.0 | 14.8 | -2.7 | 71 | 68.3 | 23 | 204.4 | 5.5 | |
| February | 2.7 | 5.1 | -0.1 | 5.1 | 10.1 | 3.3 | 71 | 72.8 | 24 | 130.0 | 4.8 | |
| March | 5.7 | 10.3 | 2.8 | 7.6 | 17.0 | -2.1 | 60 | 132.0 | 36 | 127.5 | 4.7 | |
| April | 8.9 | 12.8 | 4.7 | 8.1 | 23.5 | -2.0 | 72 | 109.7 | 43 | 112.9 | 3.7 | |
| May | 16.1 | 20.3 | 11.7 | 8.7 | 28.4 | 6.0 | 76 | 175.7 | 40 | 125.0 | 2.6 | |
| June | 19.3 | 23.2 | 15.4 | 7.8 | 30.0 | 10.9 | 82 | 195.3 | 44 | 104.5 | 2.9 | |
| July | 23.1 | 26.4 | 19.8 | 6.6 | 31.5 | 14.4 | 86 | 147.9 | 33 | 349.4 | 3.0 | |

Kami-no-yama (47,8 m. from Fukushima, in 3 hrs. 28 min.) was formerly the residential seat of a small Daimyō and famed as one of the most flourishing pleasure resorts in Dewa, both on account of its hot springs and of the existence of its gay life. The spring issues near the Shintō shrine of *Hie*, the waters being conducted by pipes to the bath-inns. To the E. of the town rises the volcanic cone of *Zaō*. *Tsukioka Park*, on the W. elevation of the town, was formerly a residential site of the Daimyō and commands a fine view of the Zaō Range and the *Mae-kawa* flowing below. A Shintō shrine, dedicated to the memory of the feudal ancestors of the place, stands on the ground. The spa lies at an elevation of 574 ft. above sea-level, surrounded by mountains on three sides, viz. on the S., E., and W. Pop. 8,600. 10 resident doctors; post, telegraph, and telephone office. 6 public baths. Number of visitors: 35,443, the total stays being 55,325 days (1909). Inns: *Yone-ya*, *Kame-ya*, *Nakamura-ya*, *Yumoto-ya*.

Meteorological Observations.

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | Amount of Precipitation | Wind Velocity | |
|---------------------|-----------------|-------|-------|-------|----------|-------|-------------------|----------------------------|-------------------------|---------------|------|
| | Mean | | | | Absolute | | | | | Mean | Dir. |
| | Mean | Max | Min. | Range | Max. | Min. | | Total | % | Total | |
| For the year, 1911. | | | | | | | | | | | |
| December | 0.4 | 4.6 | 3.1 | 7.6 | 19.0 | -8.5 | 83 | 81.9 | — | | 1.33 |
| For the year, 1912. | | | | | | | | | | | |
| January | -1.04 | 2.81 | -5.19 | 8.00 | 9.5 | -9.0 | 81.74 | 74.8 | — | | 0.97 |
| February | 2.06 | 6.42 | -1.92 | 8.33 | 14.5 | -9.5 | 76.70 | 78.4 | — | | 8.21 |
| October | 11.80 | 16.90 | 7.60 | 9.30 | 25.0 | 1.0 | 81.70 | 112.1 | 26.9 | | 1.43 |
| November | 4.30 | 7.90 | 0.90 | 7.00 | 19.5 | -6.0 | 82.30 | 73.9 | 64.7 | | 1.47 |
| December | -0.46 | 2.93 | -3.64 | 6.58 | 9.0 | 10.0 | 85.30 | 32.7 | 110.2 | | 0.97 |
| For the year, 1913. | | | | | | | | | | | |
| January | -2.07 | 1.79 | -6.02 | 7.81 | 10.5 | -12.5 | 83.47 | 57.6 | 65.5 | | 0.62 |
| February | 1.88 | 2.39 | -5.95 | 8.12 | 8.0 | -12.0 | 79.23 | 76.7 | 15.0 | | 2.00 |
| March | 1.15 | 5.55 | 3.21 | 8.63 | 16.0 | -9.0 | 77.20 | 119.0 | 24.6 | | 2.40 |
| April | 9.83 | 16.22 | 3.45 | 12.77 | 27.5 | -4.0 | 71.22 | 138.9 | 37.9 | | 1.69 |
| May | 12.97 | 18.79 | 7.24 | 11.55 | 26.5 | -1.0 | 72.65 | 126.8 | 60.5 | | 2.63 |
| June | 18.00 | 23.57 | 12.90 | 10.68 | 29.0 | 8.0 | 76.29 | 132.4 | 64.3 | | 2.53 |
| July | 21.03 | 25.22 | 17.35 | 7.84 | 32.0 | 12.0 | 83.53 | 91.4 | 105.8 | | 1.23 |
| August | 21.03 | 26.55 | 16.44 | 10.11 | 32.2 | 13.0 | 80.95 | 182.3 | 185.5 | — | |
| September | 16.61 | 22.58 | 11.72 | 10.83 | 27.5 | 5.0 | 81.59 | 125.3 | 30.4 | — | |
| November | 5.66 | 11.02 | 8.00 | 10.22 | 18.0 | -4.0 | 80.79 | 77.5 | 37.1 | — | |
| December | 1.97 | 5.85 | -1.64 | 7.50 | 20.5 | -8.0 | 87.71 | 36.5 | 139.0 | — | |
| For the year, 1914. | | | | | | | | | | | |
| January | 0.25 | 4.34 | 3.52 | 7.56 | 10.5 | -9.0 | 80.51 | 78.1 | 19.2 | | 2.17 |
| February | -0.45 | 3.87 | 4.25 | 8.12 | 10.0 | -6.5 | 79.34 | 72.9 | 25.6 | | 2.03 |
| March | 4.39 | 9.40 | 0.22 | 9.63 | 15.5 | -6.0 | 74.83 | 116.0 | — | | 2.12 |
| April | 7.03 | 12.82 | 1.33 | 11.48 | 25.0 | -4.5 | 71.67 | 150.1 | — | | — |
| May | 15.39 | 22.00 | 9.14 | 12.85 | 29.5 | 2.0 | 72.92 | 165.9 | 63.9 | | 2.17 |

Aka-yu (36.1 m. from Fukushima, in 2 hrs. 53 min.) lies on the N. E. corner of the plain of Yonezawa, on a slope of *Eboshi-dake*, and at an elevation of 700 ft. above the sea. The spa is about a mile from the *Aka-yu* station and boasts as its sights the Eboshi Park (or *Kairaku-en*) where are planted those remarkable blossoming cherry-trees, and the *Hakuryō Pond*. It is a pleasure resort of the province, the healing waters having been used for bathing since 1312 A.D. Pop. 1,135; post, telegraph and telephone office; 2 resident doctors. Number of visitors: 26,875, the total stays being 58,471 days (1909). Inns: *Goten-mori*, *Tansan Hotel*.

Meteorological Observation.

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | Amount of Precipitation | Wind Velocity | | | | |
|---------------------|-----------------|------|-------|-------|----------|-------|-------------------|----------------------------|-------------------------|---------------|---|-------|------|------|
| | Mean | | | | Absolute | | | | | Total | % | Total | Mean | Dir. |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | | | | |
| For the year, 1912. | | | | | | | | | | | | | | |
| January | 0.1 | 3.1 | -3.7 | 7.0 | 7.8 | -9.8 | 80 | 94.8 | — | 3.1 | | | | |
| February | 2.1 | 5.2 | -1.2 | 6.4 | 9.5 | -10.4 | 75 | 108.2 | 75.7 | 3.1 | | | | |
| March | 4.1 | 7.9 | 1.1 | 6.7 | 18.2 | -2.0 | 73 | 146.0 | 115.5 | 4.2 | | | | |
| April | 10.1 | 15.9 | 5.1 | 10.8 | 28.0 | -2.0 | 65 | 100.6 | 55.9 | 5.1 | | | | |
| May | 14.4 | 20.4 | 9.7 | 10.5 | 29.8 | 3.2 | 67 | 220.2 | 58.6 | 3.7 | | | | |
| June | 19.6 | 23.3 | 15.9 | 7.0 | 29.7 | 9.4 | 74 | 140.1 | 58.5 | 2.1 | | | | |
| July | 22.9 | 26.5 | 19.9 | 6.4 | 31.0 | 12.6 | 78 | 113.5 | 139.3 | 2.2 | | | | |
| August | 24.8 | 29.3 | 21.3 | 7.9 | 33.6 | 16.0 | 74 | 194.0 | 61.6 | — | | | | |
| September | 18.5 | 22.4 | 14.7 | 7.8 | 20.0 | 8.0 | 77 | 102.6 | 96.5 | 2.8 | | | | |
| October | 12.9 | 18.1 | 8.8 | 9.3 | 25.5 | 3.0 | 74 | 128.5 | 35.2 | 2.2 | | | | |
| November | 6.1 | 8.9 | 3.2 | 6.7 | 21.0 | -3.0 | 74 | 73.6 | 56.2 | 3.1 | | | | |
| December | 0.0 | 2.8 | -2.5 | 4.1 | 9.0 | -8.0 | 84 | 35.8 | 128.2 | 2.1 | | | | |
| For the year, 1913. | | | | | | | | | | | | | | |
| January | -0.7 | 1.3 | -6.0 | 7.3 | 7.0 | -15.0 | 84 | 66.6 | 63.7 | 2.9 | | | | |
| February | -1.2 | 2.8 | -5.4 | 8.3 | 8.0 | -10.0 | 79 | 107.2 | 30.8 | 2.9 | | | | |
| March | 0.3 | 4.4 | -4.24 | 8.6 | 15.0 | -10.0 | 77 | 154.3 | 42.4 | 3.6 | | | | |
| April | 9.8 | 15.0 | 4.5 | 11.2 | 26.0 | -1.5 | 69 | 159.7 | 40.1 | 2.5 | | | | |
| May | 13.3 | 17.9 | 7.9 | 10.4 | 26.0 | 0.1 | 66 | 155.3 | 67.0 | — | | | | |
| June | 18.2 | 23.0 | 13.5 | 9.5 | 27.0 | 7.0 | 73 | 170.5 | 32.3 | 3.2 | | | | |
| July | 20.8 | 24.8 | 17.7 | 7.0 | 30.5 | 12.5 | 75 | 111.5 | 126.5 | — | | | | |
| August | 21.0 | 25.3 | 17.0 | 8.1 | 31.0 | 12.0 | 71 | 155.4 | 162.9 | - | | | | |
| September | 16.6 | 22.1 | 12.3 | 9.8 | 27.5 | 5.5 | 71 | 136.7 | 89.1 | 1.9 | | | | |
| October | 12.1 | 17.2 | 7.9 | 9.6 | 21.2 | 2.0 | 74 | 107.3 | 109.6 | 1.7 | | | | |
| November | 5.8 | 10.2 | 1.9 | 8.3 | — | — | 71 | 102.7 | 96.9 | 2.4 | | | | |
| December | 0.0 | 2.2 | -2.0 | 4.9 | 8.0 | -8.5 | 78 | 44.6 | 161.8 | - | | | | |
| For the year, 1914. | | | | | | | | | | | | | | |
| January | -0.2 | 3.7 | -4.0 | 7.9 | 8.0 | -10.0 | 73 | 99.9 | 59.5 | 2.3 | | | | |
| February | -0.4 | 3.7 | -4.2 | 7.8 | 7.0 | -9.2 | 74 | 94.7 | 71.3 | 2.0 | | | | |
| March | 4.2 | 8.3 | 1.0 | 7.9 | 14.5 | -9.0 | 70 | 135.1 | 32.7 | 3.8 | | | | |
| April | 7.2 | 13.1 | 2.6 | 10.5 | 24.0 | -3.0 | 65 | 174.6 | 99.0 | 4.1 | | | | |
| May | 15.4 | 21.6 | 9.6 | 11.6 | 28.5 | 4.5 | 63 | 182.9 | 53.0 | 2.5 | | | | |

see Plate 18 **Onogawa** (5 m. S. W. of Yonezawa, *jirrikisha* or omnibuses available) lies on the E. bank of the upper reaches of the *Kinomo-gawa*, which runs from the foot of *Azuma-san*, and has an altitude of 1,000 ft. above the sea. The spa is shut in by mountains to the E. and W., the N. being open country facilitating communication with the town. Pop. 175. The hot springs are reputed good for disorders of the digestive system, the water being saline, colourless, and slightly alkaline in reaction. 2 public baths; 15 inns with private baths. Here one can find everywhere hot springs by boring. The springs were found before 848 A.D. Number of visitors: 11,288, the total stays being 22,450 days (1909). Inns: *Ogi-ya*, *Onogawa Hotel*.

Meteorological Observations.

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity | | |
|---------------------|-----------------|------|------|-------|----------|-------|-------------------|----------------------------|---|-------------------------|---------------|------|------|
| | Mean | | | | Absolute | | | Total | % | | Total | Mean | Dir. |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | | | |
| For the year, 1911. | | | | | | | | | | | | | |
| December | 1.4 | 4.7 | -3.5 | 1.2 | 14.5 | -10.0 | 60.0 | 84.1 | | 171.3 | 1.9 | | |
| For the year, 1912. | | | | | | | | | | | | | |
| January | -1.4 | 2.1 | -4.6 | -2.2 | 7.1 | -9.0 | 59.3 | 63.1 | | 159.9 | 2.3 | | |
| February | 0.7 | 4.3 | -2.4 | 1.6 | 10.0 | -7.1 | 57.3 | 71.2 | | 115.4 | 2.1 | | |
| October | 11.4 | 16.8 | 6.7 | 10.1 | 23.3 | 1.5 | 82.9 | 108.0 | | 41.4 | 1.5 | | |
| November | 4.7 | 8.9 | 1.6 | 7.3 | 19.8 | -6.4 | 83.7 | 69.3 | | 112.4 | 1.5 | | |
| December | -0.2 | 3.0 | -3.3 | 6.3 | 8.2 | -9.5 | 85.5 | 46.9 | | 228.8 | 1.3 | | |
| For the year, 1913. | | | | | | | | | | | | | |
| January | -2.3 | 1.5 | -5.6 | 7.1 | 8.0 | -11.3 | 83.2 | 58.9 | | 155.2 | 1.3 | | |
| February | -1.6 | 2.2 | -5.6 | 7.8 | 7.0 | -13.7 | 78.0 | 86.3 | | 60.6 | 2.2 | | |
| March | 0.04 | 4.3 | -4.2 | 8.5 | 14.0 | -11.8 | 76.8 | 114.9 | | 98.1 | 2.1 | | |
| April | 8.2 | 13.6 | 2.8 | 10.8 | 24.0 | -3.1 | 71.9 | 112.6 | | 37.0 | 1.8 | | |
| May | 11.8 | 17.3 | 5.9 | 11.4 | 24.5 | -0.8 | 70.3 | 183.9 | | 81.5 | 2.1 | | |
| June | 16.8 | 22.2 | 11.2 | 10.9 | 27.0 | 4.8 | 76.0 | 149.3 | | 105.1 | 1.3 | | |
| July | 20.3 | 24.4 | 15.9 | 8.5 | 30.0 | 10.6 | 80.7 | 107.2 | | 120.7 | 0.9 | | |
| August | 20.9 | 26.1 | 16.1 | 10.0 | 31.4 | 10.4 | 76.4 | 181.6 | | 254.2 | 1.0 | | |
| September | 16.7 | 22.4 | 12.2 | 10.1 | 26.8 | 5.2 | 76.6 | 116.4 | | 102.7 | 1.2 | | |
| October | 11.3 | 16.6 | 6.8 | 9.8 | 20.2 | 0.2 | 74.7 | 108.0 | | 96.5 | 1.9 | | |
| November | 5.4 | 10.5 | 1.0 | 9.4 | 17.9 | -4.8 | 70.8 | 87.8 | | 141.8 | 2.2 | | |
| December | 1.7 | 5.0 | 1.7 | 6.7 | 15.8 | -7.2 | 77.6 | 45.2 | | 261.1 | 1.1 | | |
| For the year, 1914. | | | | | | | | | | | | | |
| January | 0.2 | 4.4 | -3.6 | 8.0 | 11.0 | -9.8 | 70.3 | 80.0 | | 155.7 | 1.1 | | |
| February | 0.2 | 4.1 | -4.6 | 8.7 | 9.7 | -9.0 | 67.7 | 73.5 | | 77.5 | 3.0 | | |
| March | 4.0 | 9.0 | -0.2 | 9.2 | 16.0 | -8.6 | 66.4 | 106.1 | | 64.2 | 1.9 | | |
| April | 7.3 | 13.1 | 1.7 | 11.4 | 24.1 | -3.8 | 66.1 | 162.0 | | 147.9 | 2.7 | | |
| May | 15.2 | 22.4 | 8.1 | 14.2 | 31.0 | 2.2 | 71.4 | 170.6 | | 92.1 | 1.6 | | |

Sekine-Yunosawa. About 1,3 m. S. from *Sekine* station on the Ō-u Line; also 1,3 m. W. from the next station *Osawa*, both an agreeable walking distance. There is only one spring, the temperature being 28° C., and one bath-house in the village.

Namegawa (2,4 m. S. W. from the *Tōge* station on the Ō-u Line and over 14,6 m. from *Yonezawa*) lies in a ravine on the slope of *Azuma-san* in Uzen Province. Waterfalls are the chief attraction of the spa. *O-taki*, the largest one, over 900 ft., is in a few minutes' walk, and *Takakura-yama*, wooded with old trees, affords a beautiful sight, especially in spring and in autumn, on account of rhododendrons and of maples. Number of visitors: 2,544, the total stays being 16,525 days (1909). Inn: *Fukushima-ya*. see Plate 19

Itaya (13,5 m. from Fukushima, in 18 min.) lies near the Itaya Pass through which the line is laid. From the valley of Fukushima the ground gradually ascends, and the line passes through many tunnels, 1 m. to 2 m. long, and peaks, mountain streams, and waterfalls greet the eyes of the traveller in an everchanging confusion as the train emerges from one tunnel only to enter another. The pass is about 2,460 ft. above the sea, and *Tōge* station, next to *Itaya*, bears a general resemblance to *Kuma-no-taira* in the Usui Pass. This section of the country is subject to a heavy snowfall, which not infrequently blocks the track in winter. The waterfalls popular as beauty spots in the neighbourhood are *Nunobiki*, *O-taki*, and *Sangai-daki*, the first descending over 600 ft.

Goshiki lies about 2 m. from *Itaya* and amidst the mountains of the same name. The distance from *Yonezawa* is about 14,6 m. The spa, lying at an elevation of 3,000 ft. above the sea, affords a fine view over the surrounding mountains. Excellent skiing in winter. Only one inn. It is really a lovely summer resort, where one can enjoy cool invigorating mountain air. The springs were found previous to 686 A.D. Number of visitors: 2,860, the total stays being 31,400 days (1909). Inn: *Munekawa-ryokwan*. see Plate 20

FUKUSHIMA-KEN

Iizaka is a favourite pleasure resort of the people of Fukushima, the distance from the town being 6 m., light railway and automobiles available. The village is situated on the bank of the *Surikami-gawa*, over which a peculiar suspension bridge, called *Totsuna-bashi*, 222 ft. in length, has been constructed. The place lies at an altitude of 550 ft. above the sea. The village on the opposite side of the river is called **Yuno** and is also a bath-resort. The houses in the two villages, being constructed on the terraced banks of the stream, present a peculiar appearance; for though they are three or even four-storied structures when viewed from the river side, they appear to have only one story as seen from the road. The ground rises towards the N. W., the highest portion being *Daisaku-yama*, while an extensive plain extends towards the S. E. It is said, that *Yamato Dake* or *Yamato-Dake-no-nikoto*, a famous hero and military genius of olden times (81-113 A.D.), used the spring for the first time for bathing purposes though it is believed that the presence of the spring was already known to the people before that time. There are 7 hot springs in Iizaka, the water being all weak saline, the temperature ranging from 50° to 70° C. The water is believed to be efficacious for wounds, swellings, etc. Number of visitors: 56,787 (1909). Houses 1,000. Pop. 4,800. Inns: *Kwasui-kwan*, *Kado-ya*, *Masu-ya*, *Senshū-kaku*, *Akagawa-ya*.

The spa *Yuno* has 616 houses. There are 7 hot springs in *Yuno*, all weak saline, the temperature ranging from 45° to 71° C. The springs are said to be efficacious for rheumatism and other complaints. Number of visitors: 39,275 (1909); 39,533 (1910).

Higashi-yama lies 2,2 m. to the S. E. of the *Wakamatsu* station (38,7 m. from *Kōri-yama*, in 2 hrs. 55 min.) on the Gan-etsu Line and is the most popular spa in this district. Coaches and automobiles are available. It is shut in by peaks on the N. and E., its W. side only being left open and communicating with the city. The bath-houses stand on both banks of the rock and boulder strewn little stream called *Yu-gawa*. The spa lies at an altitude of 850 ft. above the sea. There are 11 hot springs, all weak saline, the temperature ranging from 38° to 61° C. Number of visitors: 7,380 (1909); 22,000 (1910). Inns: *Shin-taki*, *Mukai-taki*, *Fudō-taki*, *Nihachi-ya*, etc.

Kashi 14,3 m. W. from the *Shirakawa* station on the N. E. Main Line (115,7 m. from *Ueno*, in 4 hrs. 49 min.); from the named station 4,9 m. W. to *Mafune-mura*, jinrikisha available, from *Mafune-mura* as far as to *Uma-gaeshi* 7,3 m. on horse-back, thence only on foot.

The bath-place lies by the banks of the upper flow of the *Abukuma-gawa* on the E. slope of *Asahi-zan*, which rises to the N. E. of the active volcano *Nasu-zan* (6,300 ft.). The place, being at an elevation of more than 3,000 ft. above the sea, affords a wide range of magnificent scenery. The stream, which is bedded with huge singularly shaped rocks, and its bank dotted with old pines, wistaria, rhododendron, etc., adds charm to the natural beauty of this place. The hot springs issue from the crevices of granite rocks, and some bath-houses are built directly on the rocks. Going up the valley a little more than 2.4 m. W. towards the *Asahi-zan*, are found two waterfalls, i. e. *O-daki* and *Me-daki* (the former descends over 300 ft., the latter over 200 ft.).

Owing to their low temperature (48.5–51° C.), the waters are used only in summer for bathing, while the bath is closed during the winter. This may be found a quiet summer resort though convenience is limited there being but one inn. Number of visitors: 4,367, the total stays being 18,633 days (1910).

Nekonaki. From *Shirakawa* station on the N. E. Main Line (115.7 m. from *Ueno*, in 4 hrs. 49 min.) 14.7 m. E. to *Ishikawa-machi* by omnibus; 0.8 m. W. from *Ishikawa-machi* is situated the cold spring. Two small streams, *Kitasu-gawa* and *Tano-gawa*, run on the N. of the place, the valley extending from E. to W. The cold springs here gush out from the base of the hill. The vicinity of *Ishikawa* is noted for the occurrence of diverse kinds of minerals, especially rare. In the granite (mostly pegmatite) are found: *Quartz*, *Smoky Quartz*, *Yellow Quartz*, *Smoky Crystal*, *Muscovite*, *Biotite*, *Vermiculite*, *Tourmaline*, *Almandine*, *Andalusite*, *Columbite*, *Monazite*, *Titanite*, etc. In the crystalline schist are found: *Rose Crystal*, *Fiedmontite*, etc. In the contact rock are found: *Garnet*, *Epidote*, *Zoisite*, etc. In veins of iron ore is found *Rhodonte*. *Monazite* sand may be here found also in the river-bed near the town, when the level of water becomes very low. The waters of *Nekonaki* are used for bathing, after they have been heated artificially. Number of visitors: 3,028, the total stays being 3,550 days (1909). Most visitors come from the neighbouring villages.

Bobata. About 4.9 m. N. from *Ishikawa-machi*, *jinrikisha* available. The village is shut in on two sides, W. and E., by hills. The cold springs issue from the crevice of granite rocks, the temperature being 14° C. and 27° C. respectively. On the banks of the stream there are two springs, the upper one being called "*Kami-no-yu*," the lower on the opposite side of the stream "*Shimo-no-yu*," about 0.7 m. distant from each other.

Number of visitors: 2,166, the total stays being 6,300 days (1909).

NIIGATA-KEN

Senami. Tourists coming from Tōkyō leave train at *Murakami*, the terminus of the *Murakami* Line, a branch of *Shin-etsu* Line (junction: *Niitsu*), and take *jinrikisha* from the station to the spa (about 2 m.).

During the period of oil mania, which swept this country some years ago, some capitalists of the village of *Iwafune* planned to bore the ground for oil at *Senami*. They selected a site on a sand hill near the coast, about 1.3 m. S. from the town of *Senami* and N. from *Iwafune* (sea port), and commenced the operations on July 9th, 1904. After laborious works the well attained the depth of 500 ft., and still no oil coming out, the venture was considered a failure, though the works went on. When they reached about 600 ft. in depth, the capital becoming exhausted, it was decided to give up the undertaking. One of the enterprisers made up his mind to continue boring as long as the remaining fuel could keep the boring machine going. When the well was 837 ft. deep, a small quantity of lukewarm liquid flowed from the tube, which was at first believed to be volatile oil. This was, however, not the case, but after a few hours a large amount of hot water began to shoot up into the air, even more than 90 ft. high.

Though they were disappointed, as no oil was forthcoming, their attention was now fixed upon the hot water that gushed out of the tube. It was soon discovered that the water contains mineral ingredients, of which common salt predominates. Thus the place commenced to be a thriving watering-place in this region. The spa stands on a hill, wooded with pine-trees, close by the sea. The *Minomo-gawa* is stocked with salmon, and the neighbouring sea abounds in *tai* (pagrus) and *iwashi* (sardine). An island called *Aō-jima* (about 23 m. from *Senami*) adds to the scenic beauty of the place. The sand hill, on which the spa lies, produces *matsu-take* (mushrooms), the opposite bank of the *Minomo* being noted for *shii-take* (*Cortinellus shiitake*). Boiling water gushes out of

[illegible]

an iron pipe (4 inches in diameter), so abundantly as about 9,000 hectolitres in 24 hours. The temperature of the water, measured at the upper end of the pipe, shows 104° C. Mineral sinter, which is deposited inside the pipe, should be removed every two months. There are 13 inns with private baths in the spa. The water is said to be efficacious for dyspepsia, rheumatism, neurasthenia, etc.

Yuzawa (at Onna-gawa-mura). About 15 m. N. E. from the *Nakajō* station on the Shin-etsu Line; from Nakajō to *Shimoseki-mura jinrikisha* available. The hot springs are 0,5 m. from the village, lying on the bank of the *Ara-kawa*. Pop. 200. There are two hot springs in the neighbourhood, i. e. **Takase-Onsen** (0,7 m. E.) and **Taka-no-su-Onsen** (2,8 m. E.) up the valley. Bath-houses are built direct upon the springs, which issue from the crevices in rocks close by a stream. Number of visitors: 5,327, the total stays being 10,554 days (1909).

Takase. 0,7 m. E. from *Yuzawa*, and 2,1 m. W. from *Taka-no-su*, situated on the bank of the *Ara-kawa*. Hot spring gushes out from the bottom of the river. Number of visitors: 8,377, the total stays being 16,727 days (1909).

Takanosu. 2,1 m. from *Takase* and 2,8 m. from *Yuzawa*. It stands on the bank of the *Ara-kawa*. A wooden suspension bridge across the stream is a fine sight. The stream abounds in *ayu* (*Plecoglossus altivelis*). One inn with baths in the spa. The water is said to be efficacious for diseases of stomach, syphilis, etc. Number of visitors: 3,060, the total stays being 8,730 days (1909).

Murasugi. 5,5 m. S. E. from the *Suibara* station on the Murakami Line; for the further stage of journey *jinrikisha* (fare 45 *sen*) and omnibuses available. It lies at the W. foot of *Hishiga-dake*, with mountains on the S. and the E., open on the N. and the W. towards the plain of Niigata, and 400 ft. above the sea. The village has 250 inhabitants (33 households). There is another hot spring "**De-yu**," 2 m. N. from the spa. Fish-shaped fossils called "*uo-iwa*" are found in the neighbourhood of Murasugi (0,5 m. W.).

The springs issue from the side of granite hill and the water is conducted to baths by pipes. The discovery of the springs dates back to the Era of Kembu (1334-38). The temperature of springs is 26° C. at 31° C. of air temperature, the water containing a small quantity of solid matters. The water is therefore heated for bathing.

Since the springs were examined last year to test their radio-activity, the place has become a thriving bathing-place in the province of Echigo. A number of big *cryptomerias* dot the spa, hence the name "*Mura-sugi*" or 'village cryptomeria.' Murasugi is also a pretty forest resort of the province (pine and cryptomerias abundant in the neighbourhood). The springs are said to be efficacious especially for rheumatism, wounds and eczems. The public baths here are divided into two classes, i. e. *Ichino-yu* (1st class) and *Ni-no-yu* (2nd class). Number of visitors: 8,300, the total stays being 50,000 days (1909).

Deyu. 5 m. S. E. from *Suibara* station on the Murakami Line; it lies at the W. foot of *Gozu-san*, 400 ft. above the sea, only 2 m. from Murasugi. The place has mountains on the E. and W. side. Pop. 388. Found in 808 by *Kōbō Daishi*. 9 inns. 1 public bath. The hot water issuing from granite formation is conducted to the basin of a public bath called "*Chōsen-kutsu*," built on the temple-grounds of *Kwanhō-ji*. Number of visitors: 2,699, the total stays being 17,354 days (1909); 23,200 bathers (1913).

The springs are said to be good for skin diseases, dyspepsia, metal poisoning, *kakke*, disorders of the female genital organs, haemorrhoids, rheumatism, etc. Near Murasugi and Deyu lies a cold spring "*Imaita*."

Tochiomata. Passengers coming from Tōkyō take a Shin-etsu Line train to the *Raikō-ji* station (220,2 m. from *Ueno*, in 12 hrs. 2 min.), which is the junction for the Uonuma Light Railway, leading to *Ojiya*, 18,1 m. in 45 min. *Ojiya* lies on the W. bank of the *Shinano-gawa* and is noted for silk and hempen fabrics for summer wear, *habutae*-tissue, etc. From *Raikō-ji* the line traverses an extensive and fertile plain of Echigo irrigated by the two large rivers of *Shinano* and *Agano* and a number of smaller streams. A highroad leads from *Ojiya* towards the S. E. to the town of *Koide-machi* (post, telegraph, and telephone office) up the stream *Uono-gawa*, the distance being 12,2 m., *jinrikisha* (fare, 75 *sen*) and omnibuses (fare, 50 *sen*) available; from *Koide-machi* to *Kami-ortate-mura* 8,8 m. by a steep path *Yunotan-kaido*, *jinrikisha* available (fare Y 1); from *Kami-ortate-mura* to the hot spring about 1,2 m. on foot. The bathing-place is located at the foot of

Yatsumi-yama, and on the right bank of the *Yunosawa-gawa*, a branch of the River *Sanashi*. The place, being shut in on all sides by high mountains, has a limited area and is 920 ft. above the sea. The spot, where the hot springs gush out, is of granite formation. Bath-houses for public use are built in a hollow of rocks, water issuing from their crevices.

As the temperature of the water is only a little higher (39° C.) than that of the human body, bathers remain in the baths usually for a very long time, sometimes even 5–6 hours (see “*Yu-no-gō*” Hot Springs in Okayama-ken). Healing property is said to be remarkable, especially for rheumatism, chronic eczems, disorders of the female genital organs in various forms, hysteria, scrofula, skin diseases, wounds, etc.

It must be noticed that the water contains a considerable amount of radium emanation, its radio-activity being the strongest among all mineral springs in the Northern and Eastern Japan. Murasugi Cold Springs are next Tochiomata in their radio-activity (see “*Murasugi*” Cold Springs in Echigo). Moreover, the temperature of the spring here is higher than that of Murasugi. With regard to the flow of water, the former is also superior to the latter. Number of visitors: 2,007, the total stays being 23,452 days (1909). 0.3 m. down the valley lies *Ōyu-Onsen* on the opposite side of the stream. Here hot springs are scattered all over the village. Bath can be taken in this spa in winter, as the temperature of water is higher than that of Tochiomata. But the spring is of weak radio-activity. Number of visitors: 831, the total stays being 2,798 days (1909).

Yuzawa. 24.2 m. S. from *Koide-machi* (12.2 m. from *Ojīya* station), *jūrikisha* available (6 hrs.). It is a poor mountain hamlet with 208 inhabitants, on the bank of the *Uono-gawa*, about 1,000 ft. above the sea. Number of visitors: 5,524, the total stays being 20,362 days (1909).

Akakura is situated at the E. base of *Myōkō-zan*, 4.3 m. W. of the *Taguchi* station (158 m. from *Ueno*, in 8 hrs. 12 min. by the Shin-etsu Line), whence *jūrikisha* available. Water is conducted from springs more than 4 m. up the mountain. The water shoots down so rapidly along a steep slope of the mountain that it keeps almost its original temperature during the descent, the loss of temperature being less than 10° C. even in winter. The inner surface of the wooden canals used for the conduct of water gets thickly covered with sediments within a few years, which, therefore, must be removed from time to time. The sediments of different colours and densities grow gradually thicker in the course of time, forming an apparent stratification, just as the year rings of trees. The thickness of the mineral sinter amounts to about half an inch in one year. The water is used for the treatment of uterus diseases, affections of stomach and intestines, diseases of the skin, dyspepsia, etc.

Inns stand 2,500 ft. above the sea and command a view over the extensive plain of Echigo and the Japan Sea, while in fine weather the islands of *Sado* is visible afar. A branch bath-house has recently been established near the station, the water being brought down from the spring by pipes and is called *Myōkō-Onsen*. There are two thermal springs *Tsubame-Onsen* and *Seki-Onsen*, on the N. W. slope of the mountain, about 2 m. from Akakura. The former springs are sulphurous and the latter ferruginous saline. Flow of water (Akakura): 7,200 hectolitres in 24 hrs. The spa Akakura is noted for its high location with an extensive view and also for its cool climate in summer. Fresh sea-fish is obtainable easily from *Naoetsu*, 22 m. distant. 3 public baths; all inns provided with their own baths. Number of visitors: 2,496; total stays: 11,925 days (1909). Inns: *Kōun-kwan*, *Kōgaku-rō*, *Akakura branch inn*, *Kashima-ya*.

Myōkō-zan (8,090 ft.) is called the “*Fuji of Echigo*” and stands opposite to Mt. *Kurohime*, from which it is separated by the valley of the *Seki-gawa*. It is best to climb the mountain by the pass leading to the hot spring which supplies water to the bath-houses at Akakura. There are two small solfataras where the hot spring gushes out. The pass becomes steeper as it ascends, iron chains being stretched in some places to aid climbers.

At its summit stands an *Amida-dō*, and is found a cold spring close by. The range of view comprises such mountains as *Asama*, *Fuji*, *Kurohime*, and *Izuna*, and also in the N. E. the Japan Sea, the plain of Echigo, and the island of *Sado* in fine weather. Starting from the inns at Akakura it takes six or seven hours both ways.

Matsunoyama. The highway leads from *Takada* (176.7 m. from *Ueno*, in 9 hrs. 15 min.) to *Oshima-mura* via *Yasuzuka*, thence a steep and narrow path reaches the spa village via *Muroo-mura*. The place with 210 inhabitants lies at the N. foot of the *Anunizu* Range, facing the stream

of the *Hifumi-gawa*, and is about 33 m. E. from Takada. There are two public baths and all the eight inns there are owned by one inn-keeper *Tanabe*.

The springs are suitable for the treatment of eczema, rheumatism, gout, affections of genital organs, urinary diseases, chronic hysteria, etc. Number of visitors: 15,035, the total stays being 45,454 days (1909).

TOCHIGI-KEN

Nasu. The village of *Yumoto* (about 3,000 ft.), which occupies a corner of the plain of *Nasu* and is the most thriving community in the neighbourhood, is really a small hamlet of little more than 30 houses, mostly bath-house inns. It is sheltered on the N. E. by the mountain range and opens towards the S. W. The springs issue from the E. bank of the *Yu-gawa* at the foot of *Nasu-dake*, and the waters are strongly sulphurous with some quantity of iron. At 0.8 m. from *Yumoto* is the **Takao-mata Hot Spring** (about 3,000 ft.), and, at about 2 m. from *Yumoto*, the **Benten Hot Spring** (about 4,000 ft.), encircled on the three sides by rocky elevations, and with a small temple for the God *Kwan-on* situated near the spring. 1.2 m. E. is found **Kita Hot Spring** (about 4,000 ft.), in a depression between hills, so that it is only between 9 a.m. and 3 p.m. that the sun shines on the place. 1 m. W., over a steep climb, is **Daimaru Hot Spring** (about 4,000 ft.) with the waters issuing from crevices in rocks, while **Sando-goya** (about 5,000 ft.) is situated 4.5 m. further on, by the new road constructed along the side of *Usu-ga-take* and commanding a wide panorama below. The waters here are partly simple thermals, partly sulphur springs. see Plate 25

Itamuro Hot Spring is 11 m. from *Sando-goya*, the road making a wide detour around the S. W. base of *Chausu-yama*. On the N. E. and W. it is shut in by the *Nasu Range* and only on the S. it is open towards the plain of *Nasu*. The bath-houses are shut up in winter, the keepers returning to their respective villages. A short cut leads from here to *Yumoto*, while there is a path leading S. to *Kuroiso*.

The water of *Nasu Hot Spring* being strongly sulphurous, a too prolonged immersion may cause skin eruption, headache, or dizziness. The sights near *Nasu*, besides the springs already mentioned, are *Sesshō-seki* and *Onsen-jinja*, etc.

Sesshō-seki, or "Death-stone," no longer exists, but its site is pointed out near the river-bed, about half a mile from the village. Though the stone no longer exists, the whole scene is extremely desolate.

Nasu-dake. The *Nasu* volcanic range extends over the three provinces of *Shimozuke*, *Iwashi* and *Iwaki*, and is a lofty chain running through the central part of N. E. Japan. The range is composed of *Chausu*, *Nangetsu-san* and *Sambon-yari*. The first is a helmet-shaped active volcano, comparable to *Asama* in *Shinano*, and its prominence and conspicuous shape cause it to eclipse all other peaks. The ascent, 7 m. from *Yumoto*, is very easy and follows the road leading to **Benten Hot Spring**, then to **Daimaru** and **Sando-goya** bath resorts. About 2.4 m. from the last mentioned place the path divides, one branch leading to *Sando-goya*, and the other to the top.

The air is strongly charged with sulphurous gas, but otherwise the ascent is very easy, for it is practicable even in palanquins. The other path, though shorter, is far steeper and requires a guide. Striking left at the *Onsen-jinja*, a wide panorama of the *Nasu Plain*, the *Hakkō-san Ridge*, *Tsukuba*, etc., is obtained. In scenic beauty *Takaomata* surpasses all other resorts in *Nasu*. Continuing the ascent the path again branches, the left one leading to the top, and the right one to the *Benten* and *Daimaru* bath resorts. From this parting place upward it is a rather laborious climb up a sharp ascent and through bamboo grass, till a narrow resting-place called *Fudō-awa* is reached. From here the rivers *Nasu* and *Shira-kawa* appear like little ribbons. Continuing the ascent we notice *Nangetsu* below us, while the side of *Chausu-yama* is brought almost alongside us. Rhododendrons, creeping pines, and other plants growing at high altitude make their appearance amidst volcanic rocks. The active volcano, with its top like an inverted helmet, is brought very near, and streaks of white smoke issuing from its fissure make a strange sight. The path leads across a small level tract called *Amida-ga-hara*, where some stone images of *Jizō* stand, and we reach *Chausu*. On its side there is a crescent-shaped cave, called "*Hakken-ishi no-omuro*," which is 5 ft. deep and exhales sulphurous smoke. Climbing still, we come to a big rock, and below it are two craters, "Male crater" and "Female crater," the former showing greater activity than the other. The two craters lie between the big rocks standing to E. and W. on the top, but small craters and solfataras are abundant. The spectacle is less awe-inspiring than that seen on *Asama* or *Aso*, but far more weird than that at *Ōjigoku* at *Hakone*.

5 public baths; some inns are provided with delightful hot water baths. A peculiar sort of bathing, called "*Kazoe-yu*" or 'Numbering bath,' nearly the same as the "Time bath" at *Kusatsu*, can be seen at some public baths at *Yumoto Spa*. Number of visitors: 7,603, the total stays being 86,447 days (1909). Inns: *Komatsu-ya*, *Matsukawa-ya*, *Matsu-ya*, *Tokitwa-kwan*.

Shiobara.

Nishi-nasuno (92.1 m. from *Ueno*, in 3 hrs. 30 min.) is a station where those bound for **Shiobara Hot Springs** have to leave the train, *Shiobara* being situated 13.5 m. to the N. W. of the

station. As far as *Sekiya*, 7.3 m., where the ground begins to ascend, tramcars are in operation, seven times a day, single fares 28 *sen* ordinary class, and 42 *sen* special class. From *Sekiya* to *Shiobara jūnikisha* and omnibuses are in service.

Shiobara consists of 4 hamlets, called *Shimo-Shiobara*, *Naka-Shiobara*, *Kami-Shiobara*, and *Yumoto-Shiobara*, with a total population of 1,863. Hot springs issue at *Ōami*, *Fukuwata*, *Shiogama*, *Shionoyu*, *Hataori*, *Monzen*, *Sumaki*, *Furu-machi*, *Furu-Yumoto*, and *Ara-yu*, all situated along the banks of the *Hōki-gawa* or its tributaries, and within easy reach from one another. Of the above, *Fukuwata*, *Monzen*, and *Furu-machi* are the most thriving and provide better accommodation than the others. *Shiobara* is one of the most picturesque and most delightful pleasure and health resorts within easy reach of *Tōkyō*, affording all the year round, especially in spring and autumn, nature's charming displays of flowers, with the crimson tints of foliage in the latter season, while the crystalline stream of the *Hōki-gawa* with its interesting rocks and cascade do not fail to delight those who are susceptible to what is beautiful and romantic. In summer too the elevated situation of the place makes it a delightfully cool retreat.

The discovery of these charming health resorts must have occurred in ancient times; at any rate the ubiquitous saint *Kōbō Daishi* is said to have first visited the place about the middle of the 9th century and bruited the efficacy of the waters. The local chieftain, *Shiobara Iyetada*, then opened roads and made the hot springs more accessible.

It was not till the Governorship of the late Viscount *Mishima*, however, that attempts were carried out on a large scale for building roads and otherwise facilitating the convenience of the journey. About 1884, Viscount *Mishima* as Governor of *Tochigi*, reconstructed the road, removed the rocks, and built bridges; and his far-sighted undertakings, though stoutly opposed at that time by the Local Assembly, have been amply justified by the result, for it was from that time that *Shiobara* gradually grew in popularity, and to-day it contains a large number of villas of wealthy people of *Tōkyō* and other places; there is also an Imperial villa in the place. Between *Sekiya* and *Ōami* (3.7 m.), the first hot springs in this series of spas, some romantic scenery begins, affording a delightful change after the journey over a flat, uninteresting plain. The *Nyū-shō-bashi*, over the small stream at the end of *Sekiya*, makes the beginning of the natural beauty of *Shiobara* proper, with the narrow valley of the mountain stream of *Hōki-gawa* shut in by long ridges of hills. A walk of about a mile brings us to the *Mikaeri-bashi*, spanning the stream about the cascade of the same name. Here various sights abound, chiefly falls and cascades.

Inn Tariff: 1st class ¥ 3, tiffin ¥ 1.5; 2nd class ¥ 2, tiffin ¥ 1; 3rd class ¥ 1.5, tiffin 80 *sen*; rooms and bath are extra. Local souvenirs of *Shiobara* are *Okina-ame* (a kind of sweet-meat), fossil leaves and other fossil objects, cotton fabric dyed with hot spring incrustations, *Shiobara* wood-work, etc.

Ōami (Inn, *Satō*). The springs issue from crevices in big rock on the N. bank of the *Hōki-gawa*. The sights near this place are mostly falls and rocks. There is also a tunnel bored through a rock measuring 102 ft. in length, 12 in height and 15 in width. The boring was done in 1884, when the new road to *Shiobara* was constructed. Passing through this tunnel, which is called *Hakuun-dō*, we notice on the right some basaltic columns called *Zaimoku-iwa*. The waters are saline bitter.

Fukuwata (Inns: *Masu-ya*, *Matsu-ya*, *Izumi-ya*, *Maru-ya*, *Kanō-ya*, *Yoshino-ya*, *Tama-ya*, *Makino-ya*, *Sakaguchi-ya*), 1.4 m. from *Ōami*, forms part of *Shimo-Shiobara* and is 1,150 ft. above the sea. The place is surrounded by peaks, while the *Hōki-gawa* flows past its N. W. boundary, the little stream of *Fudō-ga-sawa* joining the river on the opposite bank of *Fukuwata* proper. There are a number of hot springs, as *Fudō-no-yu*, *Iwa-no-yu*, *Same-no-yu*, *Atwa-no-yu*, *Yagen-no-yu*, *Hadaka-no-yu*, and others, the waters being all alkaline. The Imperial villa is situated at this place. The sights are *Fudōga-sawa*, *Fukuwata Park*, *Onsen-jinja*, *Tengu-iwa*, etc. The *Tengu-iwa*, or the 'Hobgoblin Rock' overhangs the road, about 0.3 m. from *Fukuwata* and on it grow a number of pines. The *Nodachi-iwa*, or 'Solitary Field Rock,' is a large flat boulder under the Hobgoblin Rock and lies in the bed of the small stream called *Sayo*. Hundreds of persons may stand on this boulder. From the top of *Torito-yama* or of *Ura-yama*, rising respectively on the N. W. and the S. W. of *Fukuwata*, the whole panorama of *Nasuno* may be observed.

Shiogama (Inn, *Ko-ume-ya*), 0.5 m. further inward from *Fukuwata*, has hot alkaline springs issuing from both banks of the *Hōki-gawa*. The tomb of *Takao*, who was born here, stands in this place. Noted sights are *Ani-ototo-no-taki*, *Tamasudare-no-se*, *Kotarō-ga-fuchi*, *Oyakake-no-matsu*, etc.

Shionoyu, 1,600 ft. above the sea, (Inns: *Kashiwa-ya*, *Tama-ya*, *Myōgi-ya*), lies in a secluded spot a mile from *Shiogama*, reached after crossing the *Shio-waki-bashi* spanning the *Hōki-gawa*. It lies on the bank of the another stream, called *Kanomata-gawa*, and is sheltered by mountains on E. and W. Azaleas in spring and maples in autumn make a specially gorgeous display. There are three common salt springs. Sights near this place are *Sennin-awa*, *Shiro-yama*, *Shiyū-no-taki*, *Raitei-no-taki*, *Soren-no-taki*, *Yūhi-no-taki*, etc. All these *taki* or cascades are among the grandest of the 70 falls and cascades in *Shiobara*. *Raitei* Fall is specially famous, its height being 150 ft. and width 200 ft.

Hataori (Inns: *Nuri-ya*, *Kami-ya*, *Yamato-ya*, *Sano-ya*) lies at a distance of 0.2 m. to the N. W. of *Shiogama* and at the bend of the *Hōki-gawa* and contains five springs, all muriated alkaline. On the S. rises *Fuji* and the range of *Kijūroku*. *Fumon-ga-fuchi*, *Hōshū-no-taki*, *Kiyomi-no-taki*, etc., are among the sights here. Tradition relates that the painter named *Fumon*, believed to have been a brother of *Takao*, was drowned in the pool, which is therefore called *Fumon-ga-fuchi*. His Buddhist picture is kept at the *Myōun-ji*.

Monzen (Inns: *Yamaguchi-ya*, *Matsumoto-ya*, *Fukui-ya*, *Aoki-ya*, *Sakamoto-ya*, *Kikuchi-ya*, *Miyata-ya*) lies N. of, and almost contiguous to, the last-named place, and about half a mile from *Shiogama*. Sheltered on the N. E. by *Tera-yama* and bounded on the S. W. by the *Hōki-gawa*, it is the most thriving village in *Shiobara*, possessing a village office, post-office, primary school and other public institutions. This place has five springs, all of an alkaline nature. The Buddhist temple of *Myōun-ji* is the largest and oldest temple in *Shiobara*, having been founded by *Myōun*, nurse of *Shigemori*, eldest son of *Kiyomori*. The temple preserves a long over-dress used by *Takao*; while on the premises stands the tomb of the founder.

Sumaki (Inn, *Nemoto-ya*) is 0.5 m. from *Monzen* and stands on the slopes of *Kijūroku* Range, the place being at an elevation of 1,580 ft. An interesting feature of the bath-room here is that several jets of hot water are brought down into the basin by pipes, these jets giving an excellent massaging effect to whatever part of the body is placed under them.

Furu-machi (Inns: *Kaji-ya*, *Kami-Aizu-ya*, *Yorozu-ya*, *Naka-Aizu-ya*, *Nasu-ya*, *Fūsen-rō*, *Kome-ya*, *Myōga-ya*, *Hitachi-ya*) stands opposite *Monzen* on the other bank of the *Hōki-gawa*, the two places being connected by an iron bridge called *Hōrai-bashi*. The hills *Ōkubo* and *Hazama* rise to the N. E., while *Kurashita* and *Kijūroku* stand opposite. *Furu-machi* is continuous with *Monzen*, and the houses in the two combined give the largest total in the *Shiobara* villages. There are six or seven springs of an alkaline nature. The sights in the neighbourhood are *Onsen-sha*, *Senshin-taki*, *Genzan-no-ana*, *Hachiman-jinja*, *Shōji-gawa*, etc. *Genzan-no-ana* is a cave at the N. E. base of *Goten-yama*, N. W. of *Furu-machi*, where *Aritsuna*, grandson of *Genzammī* *Yorimasa*, consealed himself. *Hachiman-jinja*, 360 yds. from the above, is dedicated to *Honda-wake-no-Mikoto*, and on the premises are found three of the "Seven Wonders of *Shiobara*," these being *Sakasa-sugi* ('cryptomeria growing upside down'), *Ichiiya-take* ('bamboo grown in one night') and *Fuyu-no-tade* ('polygonum growing in winter'). The *Shōji-gawa* is included among the wonders, because fish are entirely absent from it, though they come up to near the *Hachiman-bashi* where the *Shōji* joins the *Hōki*. On the banks of the river grow *Kataha-no-ashi* ('reeds with leaves growing on one side only,' and hence regarded as another wonder). The two remaining marvels are the *Fūfu-garasu* ('pair of ravens') at *Ara-yu* and the *Fuyu-no-momo* ('winter peach') at *Shiogama*, but this last no more exists.

Ara-yu (Inns: *Kame-ya*, *Kami-Fuji-ya*, *Daikoku-ya*, *Tsuta-ya*, *Gensen-kan*, *Kimijima-ya*, *Shimo-Fuji-ya*) is 7 m. from *Furu-machi*, up the river and via *Kami-Shiobara* and *Furu-Yumoto*, the road, which lies to the right, being fairly level. The left route is shorter, the distance being only 5 m., but the road is more laborious. There are two ponds *en route*. *Ara-yu* is at the greatest elevation of all the spas in *Shiobara* and even in midsummer the thermometer rarely goes up to 70° F. There are four springs, the waters being acid. The sights near here are *Onsen-jinja*, *Jisō* Temple, and the *Fūfu-garasu*. *Shiobara* is singularly free from ravens, and in this spot only a single pair is met with, which is hence regarded as a wonder.

Furu-Yumoto is found among the hills, 2 m. inward from *Ara-yu*, and 5 m. along the *Hōki* from *Furu-machi*. This is the pioneer hot spring in *Shiobara*; it is also called *Kajiwara-no-yu*, because *Kajiwara*, a follower of *Yoritomo*, is said to have bathed here when he accompanied his master on the hunting excursion at *Nasuno*. Before it was visited by a severe earthquake in 1659, the

place contained over 80 houses and a fine temple, but now only one inn is left. There are two falls near here.

There are 23 public baths, 3 doctors in Shiobara. Number of visitors: 31,451, total stays being 165,670 days (1909).

Nikkō.

The glory of nature at *Nikkō* is ideally complemented by the glory of art enshrined there, for Nature's bold essay in landscape gardening with mountains, cascades, crystalline streams, ancient trees, lovely lakes, and the brilliant chromatic display of leaves in autumn, is utilised to best advantage by the cunning hand of master designers and craftsmen, who have reared amidst such surroundings gorgeous piles profusely decorated. It is not to be wondered at that foreigners should regard *Nikkō* as by far the most noteworthy sight in all Japan, while the tribute which Japanese pay to the romantic beauty of the place is sufficiently evidenced by that popular saying, meaning, "Don't use the word 'splendour' till you have seen *Nikkō*," 'splendour' being in Japanese '*kekko*.'

Through trains run several times a day between Ueno and *Nikkō*. Passengers have to change at *Utsunomiya*, when going by non-through trains. Time required: about four hours. Fares: 1st class, ¥ 3.60; 2nd class, ¥ 2.14.

Nikkō (90.9 m. from *Ueno*, in 4 to 4¾ hrs.) is the terminus of the *Utsunomiya-Nikkō Branch Line*. The station (1,746 ft. above the sea-level) is situated on the right bank of the *Daiya-gawa*, along which a long town stretches on constantly rising ground from the station toward the N.W. —the distance from the station to the *Mihashi*, or 'Sacred Bridge,' which is at the centre of the most interesting section of the town, is about 1 m. There is an electric-tram service starting from the station and passing the various hotels, (and an automobile is available in summer).

Hotels: *Kanaya Hotel* (on a hill, close by the *Mihashi*, about 1 m. from the station), *Nikkō Hotel* (near the Park at *Nishi-machi*, a little over 1 m. from the station). Inns: *Konishi-Ryokwan*.

The Four Seasons in *Nikkō*.

Spring. In late spring, and in the month of May, *Nikkō* offers charming scenes by reason of the flowers of azalea-trees, purple, red, and white, and the delicate green shoots of larch-trees. The azaleas found here are somewhat different from those growing elsewhere. The best places for enjoying the flowers are in the neighbourhood of the street of *Nikkō*, *Nakimushi-yama* ridge, *Yashū-hara*, *Kujira-yama* on the upper course of the *Tamozawa*, the road from *Nikkō* to *Chūzenji* hill-sides facing *Uma-gaeshi*. Then *Chūzenji*, especially at *Tera-ga-saki* and *Aka-iwa* on the lake shore, and *Mae-Shirane* at *Yumoto* are also noted for the flowers. For the vivid green of the larch buds, the best places are *Senjō-ga-hara* near *Yumoto*, the S. foot of *Tarō-san*, the W. foot of *Nantai-san*, and also the W. side of *Yashū-hara* and *Nakimushi-yama*.

Summer. Refreshing cool climate, boating and fishing on *Chūzenji* and *Yumoto* Lakes, viewing the sunrise from the top of *Nantai*, gorgeous carpet of late azaleas and iris flowers at *Senjō-ga-hara* and neighbourhood in early July, and then warbling of Japanese nightingales and other singing birds may be mentioned as attractions of *Nikkō* in this season.

Autumn. *Nikkō* is superb in autumn for the scarlet tints of its foliage; in this respect it is considered far to excel other places noted for the autumn scarlet, such as *Usui* and *Shiobara*. By Japanese, therefore, this season is considered the best time for visiting *Nikkō*. The fascinating colouring begins from about the latter part of September, but the sight is generally at its best about the middle of October. The best places for seeing the autumn colouring are *Ogura-yama*, the vicinity of *Kirifuri* Fall, right bank of the *Daiya* as one ascends from the Sacred Bridge, and on the road between *Uma-gaeshi* and *Chūzenji*, *Misawa*, *Ken-ga-mine*, *Naka-no-chaya*, *Fudō-zaka*, *Kegon* Fall, etc. At *Chūzenji* the places are *Uta-ga-hama*, *Tera-ga-saki*, *Shōbu-ga-hama*, *Aka-iwa*, etc., and at *Yumoto* there are *Yu-daki*, *Tade-no-umi*, *Konsei-tōge*, etc. The view of the moon as reflected on the lake is a delight to Japanese artists and poets. The maples, which are the chief contributors to this chromatic splendour, are of divers varieties in *Nikkō*, some being peculiar to one place and others to other places. More than twenty four kinds are counted by Japanese lovers of this autumn sight.

Winter. Near *Chūzenji* the snow accumulates to a depth of about 3 ft. on the ground, and hence *Yumoto* and *Chūzenji* are practically deserted by holiday seekers during the cold season. The lake does not freeze thick enough for skating. However, there are great hopes that sleighing and skiing, for which excellent grounds are abundant, may become a popular winter attraction in *Nikkō*.

Places of interest: Nikkō Kōen, or Public Park, the Grand Cryptomeria Avenues, *Kirifuri-no-taki*, or 'Mistfalling cascade,' *Gamman-ga-fuchi*, *Sōmen-no-taki*, or 'Vermicelli cascade,' *Urami-no-taki*, or 'Back-viewing cascade,' *Jikwan-no-taki*, *Jakkwō-no-taki*, *Haguro-no-taki*, *Aioi-no-taki*, *Nana-taki*, for 'Seven waterfalls,' etc.

Nikkō to Chūzenji.

From the Mihashi to Lake Chūzenji, a distance of about 10 m., *jirikishas*, *kago* (chairs), and saddle-horses are available; while electric trams go as far as *Uma-gaeshi* (5 m.), which is halfway up. The waterfalls *Hōdō*, *Hannya*, *Kegon* and *Shirakumo* are worth visiting.

Chūzenji.

The hamlet of Chūzenji lies on the N. shore of Lake Chūzenji, which is about 10 m. from Nikkō and 4,194 ft. above the sea-level. The place acquired its name from the temple of Chūzenji, which was established by *Shōdō-Shōnin* over 1,100 years ago.

Hotel: *Lake-side Hotel* near the bridge, *Ōjiri-bashi*. Inns: *Kome-ya*, *Tsuta-ya*, near the shore (foreign meals to order).

Lake Chūzenji. This clear, fresh-water lake, is 15 m. in circumference (its greatest length from E. to W. is about 5 m., its breadth, 2,5 m.). It lies at the foot of *Nantai-san* and is surrounded by beautiful hills on the other sides. Its shores are rich in indentations, which add much to its scenic beauty. Towards the E. of *Oimatsu-ga-saki* are found the deeper parts of the lake,—soundings showing the greatest depth (i.e. off *Kōzuke-jima*) to be 172 metres (567 ft.). The lake, formerly devoid of fish, now abounds with trout, carp, etc., having been stocked by the Government in 1873 with *iwana* (a species of trout), in 1874 with carp, in 1875 with *hara-aka*, in 1879 with eels and lampreys, in 1882 with trout and *ame-no-urao*, in 1890 with *higai*, and in 1892 with American trout.

Highway to Yumoto.

The highway to Yumoto (7,5 m.) starts from Chūzenji, running along the N. shore. First we pass the bronze *torii* of the Chūgūshi Shrine, then come to *Osaki*, where in a forest of pines, cryptomerias, birches, and maples, stand many summer villas. *Shōbu-ga-hama* (3 m. from *Ōjiri-bashi*—boat available) is an important stage-town (where horses or *jirikisha* may be hired). Here the highway takes a sharp turn towards the N. and traverses a forest of large birches. On the way to *Senjō-ga-hara*, we pass *Akanuma*, a reddish-coloured marshy land, overgrown in many places with reeds.

Senjō-ga-hara is a plain, rendered very attractive in summer by the blossoms of *Yashū-bana* (a kind of azalea) and wild lilies. The place also affords a complete view of the surrounding mountains—*Nantai-san*, *Ōmanago*, *Komanago* and *Tarō-zan* towards the E., and *Shirane-san*, *Mae-Shirane-san*, and *Konsei-zan* to the W.

Yudaki or the 'Hot water cascade,' is formed by the waters from Yumoto Lake running down a steep rocky bed. By taking the highway towards Yumoto, we soon come to a spot whence we may look down upon the cascade, while the left-hand path brings one to a point from which one may view it from below.

One may then rejoin the Yumoto highway by a short cut which brings one out at the spot above mentioned. The cascade makes a fall of 270 ft., its waters flowing through a forest of maples and other fine trees.

Yumoto Hot Springs.

The Yumoto spa is 17 m. from *Nikkō* and 7,5 m. from *Chūzenji*,—*jirikisha* available throughout. Hotel: *Namma Hotel*. Inns: *Kama-ya*, *Ita-ya*, etc.; these each have detached houses; foreign meals to order. Yumoto is surrounded by mountains on all sides except to the S., where it is open towards the lake. The place is at the high altitude of 5,088 ft. above the sea-level, being higher than Chūzenji by 900 ft., and than the *Mihashi*, *Nikkō*, by 3,060 ft. In the hottest days of summer the thermometer never registers more than 82° Fahr. It is literally a summer resort, the hotel and inns being practically closed between the middle of November and the middle of March.

Lake Yumoto, called '*Yu-no-umi*,' is 1 m. long and 0,2 m. wide and receives its waters from the *Shirane-sawa*, the *Konsei-zawa*, etc., as well as from the thermal springs. On the E. shore of the lake (i.e. on the right-hand side of the road to Chūzenji) is a small peninsula, called *Usagi-jima*, which is covered by a forest of splendid conifers. In fact the lake is surrounded on all sides by a

beautiful forest, in which are found trees like *kome-tsuga*, *take-momi* (fir), *kara-matsu* (larch), *kara-hinoki*, and *asunaro*. There are boats for hire, either for rowing or for fishing. The lake abounds in *funa* (gibel) and eels.

There are many hot springs, which contain, as a rule, hydrogen sulphide with a slight trace of acid. For the most part they are colourless and transparent, but some are slightly turbid. In the hotel and inns there are private baths, but public baths, of which there are more than 10, are largely patronised, as they are believed to be more efficacious than the private baths. The public baths have names, which, taking them in order from the S., are as follows:—*Kawara-no-yu*, *Tsuru-no-yu*, *Donsu-no-yu*, *Naka-no-yu*, *Taki-yu*, etc. Of these the hottest are *Ara-yu* (156° Fahr.) and *Kawara-no-yu* and *Gosho-yu* (both 149°), while *Donsu-no-yu* (105°) has the lowest temperature. Number of visitors: 1,249 (1909).

GUMMA-KEN

Ikao, Kusatsu, and other Hot Springs.

The mountainous district situated northward from *Takasaki* and *Karuizawa* is noted for the abundance of its hot springs, the waters of which possess various medical qualities, while the localities where they issue form delightful summer resorts. These hot springs are found at *Ikao*, *Kusatsu*, *Shima*, and *Sawatari*.

Kusatsu and Neighbouring Hot Springs.

Kusatsu Spa may be reached by any of these six routes:—

(a) From *Ikao* (also from *Takasaki*, *Iizuka*, or *Maebashi*) by electric tram to *Shibukawa*; from *Shibukawa* to *Nakanōjō* by horse-tram; from *Nakanōjō*, via *Kawara-yu* (12.5 m. in 3 hrs.) and *Otsu*, to *Kusatsu* by *basha* (entire distance 25 m., in 7 hrs.; fare ¥ 1.40).

(b) To *Nakanōjō*, as stated above; from *Nakanōjō* to *Sawatari* (6 m.) partly by *basha* or *jīnrikisha*, and the rest on foot; from *Sawatari* to *Kusatsu*, across the *Kuresaka Ridge* (18 m.) on foot.

(c) From *Ikao* or *Haruna* on foot as far as *Hakojima*, near a station on the *Shibukawa-Nakanōjō* horse-tramway, via *Benten-daki*, *Ō-daki*, etc., the path lying along the stream *Numa-gawa*; the rest of the way to *Kusatsu* either as in (a) or (b). Between *Hakojima*, and *Nakanōjō*, there is an interesting pedestrian path along the *River Agatsuma-gawa*.

(d) From *Ikao* to *Haruna* either on foot or by chairs (or horse-back); from *Haruna* to *Kawara-yu* (20 m. from *Ikao*), via *Ōdo*, *Ōkashiwagi*, either on foot or horse-back,—the path lying along the W. slope of *Haruna*; the rest of the way to *Kusatsu* as in (a). Between *Haruna* and *Kawara-yu*, the path is rich in beautiful scenery, this part being known as the “*Yabakei*” of *Kwantō*.

(e) From *Karuizawa* or *Nagano* by rail to *Toyono*; from *Toyono* to the *Shibu Spa* (11 m.) by *basha* or *jīnrikisha*; from *Shibu Spa* to *Kusatsu* on foot or on horse-back.

(f) From *Karuizawa* or *Kutsukake* to *Kusatsu* (26.2 m.), via *Okuwa* (14 m.), *Haneo*, *Otsu*, etc. on foot or horse-back (fare ¥ 4.50).

Kusatsu Hot Springs.

Kusatsu has been celebrated from ancient times for the medical efficacy of its baths, as *Arima* has been in the neighbourhood of *Kyōto*. Local chronicles record the visits of many celebrated persons to *Kusatsu*, that of *Yoritomo* having especially contributed, it is said, to the spreading of the fame of the waters, which are chiefly sulphuric, the sulphur being present in the form of sulphuric acid. The other constituents are iron, alum, and arsenic. The temperature is exceedingly high, ranging from 113° to 128° Fahr. The waters have such a powerful effect on the skin that in the first stage sores are brought out on the tender parts. The chief public bath, called “*Netsu-no-yu*,” stands in the village square, and people suffering from obstinate skin-diseases take courses in it. The waters are so hot that the bathing is done under semi-military discipline, with a ‘bath-master’ directing the operation. The first process consists in stirring the waters thoroughly with boards to aerate and soften them (fig. 1); then the bathers pour 250 dippers of water on their heads to prevent congestion (fig. 2). At the word of command from the master they enter the bath and endure the ordeal for 3.5 to 4 min., remaining practically motionless, for even a slight agitation of the water makes the temperature doubly trying to those immersed in the bath (fig. 3). At intervals of one minute or half a minute the master cries out the time left, to which all the bathers answer in chorus. Of the

various baths, *Takino-moto-yu* is 148° F. and contains sulphuretted hydrogen, iron, and acid; *Netsu-no-yu* is 144° F.; *Washi-no-yu* and *Jizō-no-yu* 140° F. (fig. 3); *Goza-no-yu* 137° F.; *Nagi-no-yu* 127° F.; *Kakke-no-yu* 143° F.; *Kompira-no-yu* 130° F. Kusatsu stands 4,500 ft. above the sea and is a clean looking village, three-storied inns with gabled fronts occupying the square. This square is full of vapour from the spring of boiling sulphur water which is brought here by wooden troughs to collect sulphur deposit (fig. 1). Kusatsu lies in a depression in a table-land, and a small stream runs through it, from the bed of which hot springs gush out here and there.

At an extreme end of the village, and isolated from the rest, is a leper's quarter (called *Yuno-sawa*) with a special bath for the afflicted. Sights near Kusatsu are *Sai-no-kawara*, *Kōri-dani*, *Jōfu-no-taki*, *Kakomi-yama*, *Doku-mizu*, *Shirane-jinja*. From Kusatsu two roads lead N.W. to *Shinano*, one to *Suzaka* via the *Torii-tōge*, and the other to *Shibu-Onsen* over the 19,5 m. pass of the same name, a very laborious expedition. This path, however, secures one the advantage of visiting *Sirane-san* and of enjoying the milder waters of the hot springs at *Shibu*. It should be noted that those who have passed through the bathing course at Kusatsu generally make a week's stay at either *Sawatari* or *Shibu* to effect the complete after-cure of the irritated skin.

Water pipes laid from the hill, good drainage, no mosquito in summer. Number of visitors: 6,740, the total stays being 140,200 days (1909). Inns: *Ichii-Ryokwan*, *Shirane Hotel*, *Nisshun-kwan*, *Chōyō-kwan*, *Bōun-kwan*.

Shirane-san, called "*Kusatsu Shirane*" to distinguish it from *Nikkō Shirane*, is an active volcano, 7,500 ft. high, that has exploded four times during the last few decades, viz. in 1882, 1897, 1898 and 1902. It lies about 2,5 m. off the highway and requires an extra couple of hours or so to visit it. At the top three craters exist, the central one called *Yugama*, the W. one *Karc-gama*, and the E. one *Mizu-gama*. The last one contains yellowish-coloured water, while from the *Yugama*, which means "hot-water cauldron," dense vapour is ascending and jets of hot water containing sulphur are thrown up. The other one, "dried-up cauldron," is dry. The woods all round the mountain present a desolate appearance with their blasted skeleton trees, the effect of the recent eruptions, though the woods near *Shinano* have escaped destruction. Near *Shirane* stands *Moto-Shirane*, another cone-shaped volcano, while in front of it rises *Manza-yama*, at the base of which are the hot springs of *Manza* (5,180 ft. above sea-level).

Kawarayu, 11,3 m. S. W. of *Nakanojō* on the way to Kusatsu and 2,200 ft. above the sea, is situated on the high precipitous bank of the upper course of the River *Agatsuma*, opposite the peaks of Mts. *Tengu* and *Takama*. An extent of the valley in the vicinity of the spa being exceedingly beautiful in scenery, is regarded as '*Yabakei* of Kwantō (Central Japan),' as already mentioned. (*Yabakei* is a valley near *Beppu*, noted for its scenic beauty). The spa may be a quiet summer resort with hot springs.

Shima spa includes three hot springs, *Yamaguchi*, *Shin-yu* and *Hinatami*, 0,5 m. from each other, all located in *Sawada-mura*, 10 m. N. from *Nakanojō*, *Kōzuke* Province, *jirikisha* and omnibuses available. The spa, 2,500 ft. high above the sea, stands by a ravine with several waterfalls on its course. The main inns own private baths and are also provided with vapour baths. The best season for visit is from May to August. The springs are noted in efficacy next to Kusatsu among all the natural baths in this district. The water is said to be efficacious especially for dyspepsia, swellings, diseases of the skin, etc. Number of visitors: 6,617, the total stays being 57,613 days (1909). Inns: *Sairyō-kwan*, *Sekizen-kwan*.

Sawatari (Inn, *Taishō-kwan*) is 6 m. W. of *Nakanojō* and at an altitude of 2,200 ft. above the sea. Number of visitors: 2,000, the total stays being 30,000 days (1909).

Isobe (73,9 m. from *Ueno*, in 3 hrs. 29 min., 15 m. from *Karuizawa*) is noted for mineral springs used for baths after heating. The spa is situated in a sort of ravine, by the bank of a stream, the *Usui-gawa* at an elevation of 1,200 ft. above the sea. For a distance of 0,3 m. from the station, the road is lined by cherry-trees.

Shiro-yama, 0,7 m. S. E. of the station, is the site of the old castle of *Sasaki Moritsuna* (end of the 12th century). The place is a low hill, with flat summit, 100 ft. above the surrounding plains, and commands a wide prospect. From here are the bizarre, jagged mountain peaks of *Myōgi-san* visible on the S. W., and Mt. *Asama* (an active volcano) with its splendid view on the W. There have recently been planted a large number of cherry-trees, which attract numerous visitors in the flower season.

The springs, though cold, are highly saline, and impregnated with carbonic acid gas. Each of them is one of the most concentrated common salt springs in Japan, the total quantity of solid constituents in 1 kilogram being more than 30 grams. In *Hara-machi*, a small town, 2,5 m. distant from Isobe, is also a cold spring, which contains more than 34 grams of solid matter in 1 kilogram of water. Carbonic acid gas is utilized here for obtaining carbonate of lime for tooth-powder and water for the manufacture of aerated water. Salt, obtained by evaporating mineral water, is sold at this spa. The water is suitable for the treatment of dyspepsia, swellings, haemorrhoids, uterus diseases, etc. Number of visitors: 6,340, the total stays being 15,451 days (1909). Inns: *Hōrai-kwan*, *Taigaku-rō*, *Isobe-kwan*, etc

Ikao and Neighbourhood.

The spa town of Ikao may be reached by three routes: (1) From Tōkyō via *Takasaki*—Tōkyō to Takasaki by railway (63 m. in about 3 hrs.) and from Takasaki by electric car (20,3 m. in 2,3 hrs.); visitors from Yokohama may effect junction with the Tōkyō—Takasaki Line, by taking train or electric car from *Shinagawa* to *Akabane*; (2) From Karuizawa or Nagano, via *Iizuka* (immediately N. of Takasaki),—to Iizuka by rail and from Iizuka to Ikao by electric car (19 m. in 2 hrs.); (3) From Nikkō or Mito, via *Oyama* and *Maebashi*,—to Maebashi by rail and from Maebashi to Ikao by electric car (16,6 m. in 2 hrs.); between Ueno (Tōkyō) and Ikao, through tickets issued, via either Takasaki or Maebashi.

Shibukawa is a prosperous town of 7,046 inhabitants, being 9,1 m. from Maebashi, 12,8 m. from Takasaki, and 7,5 m. from Ikao, all by electric tramway. The town is picturesquely situated, being in full view of Akagi Range towards the N. E. and of *Haruna* towards the S. W., while the rivers *Tone* and *Agatsuma*, one flowing from the N. and the other from the W., join at its N. out-skirts. From Shibukawa to Ikao, electric cars available as already stated; to *Kusatsu*, *Shin'a*, *Sawatari*, *Kawarayu* (all spa towns), tramway (horse) as far as *Nakanojō* (13 m. in 2 hrs. 30 min.—fare 50 *sen*), from Nakanojō to the several hot springs named, either by *basha*, a rude kind of stage coach, or *jūrikisha*. There is a tramway service also between *Shibukawa* and *Numata* (13,2 m. in 2 hrs.—fare 45 *sen*); the latter town lying on the highway to Nikkō and Mt. *Shirane-san*.

Ikao is one of the most popular summer resorts within easy reach from Tōkyō and Yokohama and is situated on a mountain slope with an elevation of 2,500 to 2,700 ft. The chief attraction of this place for Japanese is the hot spring, which has the temperature of 113° F. and contains a small quantity of iron and sulphate of soda. The waters are believed to be specially efficacious for women's diseases, and hence Ikao finds more patrons among the fair sex than among the men. Ikao is essentially a spa, for the little village containing about one thousand souls consists of inns and houses selling local souvenirs, or otherwise purveying to the needs of guests staying at the inns. The principal street that divides the village into E. and W. halves makes a rather sharp ascent, graded by an almost continuous flight of stone steps. The W. half ends in a deep ravine where flows a small stream, while the other half is bounded in by a low mountain ridge. The hot waters gush out at the back of a small gully called *Yumoto*, or "Source of Hot Water," which lies about half a mile from the top of the street, and to which an excellent level road leads. At their source the waters are clear and are drunk by visitors, and from this place they are conveyed by bamboo pipes to all the bath-houses. When left to cool, the waters precipitate yellowish deposit of iron oxide. Ikao is rich in wild-flowers, lilies of various kinds being especially prominent from June to September, while in late spring the mountain sides echo with the songs of Japanese nightingales and cuckoos. For those prefer walking from *Shibukawa* as far as Ikao, a distance of 6 m., instead of travelling in the electric car, time 1 hr., it may be noted that about 3,5 m. up is a resting-station, called *Mikage-no-matsu*, named from the aged pine growing close by, where stands the stone tablet commemorating the short rest the late Empress-Dowager Eishō-Kōtaikō took when she visited Ikao. A little further on we come to the place where a wooden post is erected to mark the path leading to the *Fu-nyū* Fall. Ikao is a little over a mile from here. The scenery commanded from Ikao is extensive and beautiful and comprises the valleys of the two rivers *Tone* and *Agatsuma*, the mountain ranges marking the border of the vast plain of *Musashi* in which Tōkyō is situated, and then the Nikkō Range. Some of the larger hotels have suitable provision for the accommodation of foreign guests. 15 public baths, 1 spa physician and 1 dentist. Number of visitors: 33,557, the total

stays being 193,217 days (1900). Hotels: *Ikao Hotel* (semi-European style), *Lake Hotel*, *Hashimoto Hotel*. Inns: *Buibu* or *Nino-Kogure*, *Chigira*, *Horai-kean*, *Chitose-kean*, etc.

Meteorological Observations.

Summary.

| Month | Air temperature | Relative Humidity | No. of hours with Sunshine | Amount of Precipitation | Wind Velocity | |
|---------------------|-----------------|-------------------|----------------------------|-------------------------|---------------|-----------|
| | Mean | | Total | Total | Mean | Direction |
| For the year, 1913. | | | | | | |
| January | -2.03 | 61.0 | 209.3 | 31.2 | 3.19 | N 73° W |
| February | -0.62 | 60.4 | 191.4 | 23.6 | 3.23 | N 71° W |
| March | 1.09 | 58.4 | 228.1 | 52.9 | 3.48 | N 72° W |
| April | 9.98 | 73.3 | 186.2 | 68.2 | 3.10 | S 1° W |
| May | 11.83 | 71.3 | 230.1 | 169.4 | 3.39 | S 60° W |
| June | 16.47 | 80.2 | 168.2 | 230.9 | 2.64 | S 3° E |
| July | 19.99 | 87.5 | 157.9 | 196.6 | 2.38 | S 36° E |
| August | 20.08 | 83.3 | 184.4 | 237.9 | 2.58 | S 14° E |
| September | 16.04 | 81.3 | 144.4 | 182.3 | 2.30 | S 31° W |
| October | 11.85 | 80.1 | 141.9 | 219.0 | 2.55 | S 57° W |
| November | 5.68 | 67.2 | 170.4 | 70.7 | 2.98 | N 81° W |
| December | 1.72 | 61.0 | 173.0 | 36.5 | 2.99 | N 65° W |
| For the year, 1914. | | | | | | |
| January | 0.37 | 54.4 | 210.1 | 14.6 | 3.20 | N 61° W |
| February | -0.48 | 58.1 | 171.0 | 35.8 | 1.90 | N 54° W |
| March | 4.50 | 69.9 | 167.6 | 89.2 | 2.96 | N 88° W |
| April | 7.56 | 61.7 | 213.9 | 130.5 | 3.50 | N 55° W |
| May | 14.12 | 72.0 | 183.0 | 225.0 | 2.62 | S 30° W |
| June | 18.05 | 80.6 | 157.9 | 165.7 | 2.06 | S 5° E |
| July | 22.07 | 81.8 | 179.7 | 134.4 | 2.12 | S 28° W |
| August | 22.60 | 87.0 | 203.5 | 508.5 | 2.36 | S 0° W |
| September | 18.51 | 86.7 | 101.0 | 264.6 | 1.83 | S 18° W |
| October | 11.71 | 74.8 | 178.1 | 103.3 | 1.90 | S 82° W |

Nanae-no-taki and **Benten-daki** supply a pleasant walk, the former only about 0.5 m. from Ikao, the other side of the ravine, while the latter, formed by the water coming from Lake Haruna, is a little over 1.5 m. further on in the wood that contains *Nanae*. *Nanae* is so called because the little mountain torrent here divides into seven little cascades. Near the spot stand rest-houses where Japanese buckwheat vermicelli (*soba*) and other delicacies are served. *Benten* is a *bona fide* fall, some 40 ft. in height and 10 ft. wide. It is utilized for generating hydro-electricity to supply motor power to the trams from Shibukawa to Ikao. Visitors to Haruna shrine may reach *Benten* by returning by the path leading to it along the lake.

Garameki-Onsen is a lukewarm spring amidst hills near *Minowa*, at the S. E. foot of *Sōma-ga-dake*. It is about 2.4 m. from Ikao; the waters are alkaline and are heated before being used.

Haruna-san ranks with *Akagi* and *Myōgi* as one of the three celebrated mountains in Kōzuke Province. From Ikao to the shrine of *Haruna-jinja* the distance is about 6 m., of which the first stage of about 1.2 m. is a rough climb over the mountain slope, though practicable by "chairs" (fare ¥ 2) or on horseback. It is perhaps more advisable to walk. On reaching to the top of the

ascent we come to a plateau, and diverging left we come after a few minutes to the *mushi-yu* (vapour-bath), about 1,2 m. away. The place was much frequented by rheumatic and other patients in order to take advantage of the sulphurous gases that used to issue from the ground; but since of late these gases ceased to issue, the vapour-bath has been closed. The twin-peaks rising on the left are *Futatsu-dake*. Advancing further and over a little ascent called *Yasemune-tôge*, the road traverses for about 2,5 m. the descending moor with *Sôma-ga-dake* standing left and *Takane* on the N., and at last we reach Lake Haruna. *Sôma-ga-dake*, otherwise called *Kurokami-yama*, is 4,800 ft. in height; the routes to the summit are all steep, iron-chains being stretched in some places to aid climbing. On the summit stands a 6 ft. stone statue of Taira-no-Masakado. The panorama commanded extends over a wide region, including *Fuji* nearly due S., several of the lofty ranges of Kai and Shinano and all the peaks standing close by.

Ikao-Fuji, or **Haruna-Fuji** (4,808 ft. in height) rises at the N. E. part of Lake Haruna, and is so named from its shape. At the foot is found a cattle ranch in miniature. At the foot facing the lake is a small mound called *Hitomokko-yama*. Legend says that a genius having undertaken to build this Fuji in one night, the day began to dawn while he had yet to pile on one more "mokokful" of earth ("*mokko*" being a rope net-work mat for carrying earth and other things), and he was obliged to leave the earth where the mound stands. Hence the name, which means "one mokkoful mound."

Lake Haruna is a crater lake, about 5 m. from Ikao, and is 0,8 m. from E. to W., 1 m. from N. to S., and 2,3 m. round. The water is clear, and besides *Ikao Fuji*, there stand around the basin, *Eboshi-dake*, *Bingushi-dake*, *Suzuri-iwa*, and *Kamon-ga-dake*. The lake is stocked with salmon and other fish, a special company being organized for undertaking pisciculture here. The tea-house (Lake Hotel, branch of Ikao Hotel) on the shore serves lunch or provides accommodation for the night. The shore is noted for irises and fire-flies, while in winter the frozen surface of the lake affords excellent skating. Ice is also hewn here in winter. The outflow of the lake forms the Benten Fall. *Tenjin-tôge* is a short ascent on the road leading from the lake to the shrine. On the top stands a big *torii* for Haruna-jinja (Shintô shrine) and there are tea-houses to right and left of the road. The view of the lake from this point is really picturesque. The distance from here to the shrine is 1,2 m. Near the shrine, and beyond the small current flowing left of the road, stands a queer-shaped rock, called *Tsuzura-iwa*, because it looks like a set of gigantic trunks piled one upon another.

Yujiku hot spring (29,5 m. N. from Maebashi, and 20,4 m. N. W. from Shibukawa) is on the right bank of the *Akaya-gawa* and on a highway (*Mikuni-kaidô*), with mountains on the S. and N. Pop. 500. In the neighbourhood are several springs: **Hôshi-Onsen** (7,2 m.), **Yushima-Onsen** (2,5 m.), **Sasa-no-yu** (2,5 m.), etc. There are five springs in Yujiku, the temperature of water being widely different from one another. Number of visitors: 1,500, the total stays being 5,000 days (1909).

Yubiso. 15. m. from *Numata*, and 25,4 m. N. from *Shibukawa*; from Shibukawa to Numata (13,2 m.) horse-tramway available and the rest only on foot. The place (1,500 ft. above the sea) lies on the right bank of the *Yubiso-gawa*, the upper course of the *Tone-gawa*. Pop. 194. The water is said to be especially efficacious for rheumatism. **Yubara-Onsen** is about 2,5 m. distant. Number of visitors: 495, the total stays being 3,382 days (1909).

Yubara. 23 m. N. from Shibukawa, 2,5 m. S. from *Yubiso-Onsen*. Pop. 398. 3 springs. Number of visitors: 519, the total stays being 1,908 days (1909).

Irinoyu, 8,5 m. N. W. from *Yokokawa* station on the Shin-etsu Line, lies on the E. slope of *Usui-tôge*, about 2,700 ft. above the sea, 6 m. from *Karuizawa*. The place is shut in by hills on all sides. Maple trees are found in the surrounding hills. Only 1 inn with baths. Number of visitors: 121, the total stays being 966 days (1909).

Nashiki-Kôsen. 2,5 m. W. from the *Kami-kambai* station on the Ashio Line (junction: see Plate 28 *Kiryû*). The spa, 1,490 ft. high above the sea, is situated in a gorge at the foot of *Mt. Akagi*, a stream with clear water running near by. Thickly wooded hills surrounding the place keep the air always fresh and bracing. The garden of an inn *Nashiki-kwan* is planted with cherries, plums and peaches, and affords a beautiful sight in spring. It is a quiet summer retreat for inhabitants in the vicinity.

IBARAKI=KEN

Hiraiso, 2.4 m. N. of *Minato*, Pop. 7,856, is noted both as a watering-place and also as the foremost fishing centre in *Hitachi* Province. The view of the waves from the N. height of this place is very grand, and the height bears the name of *Kéantō-jō* ('Wave-viewing station') specially bestowed by *Rekkō*, the celebrated prince of Mito in the middle of the 19th century. see Plate 29

Ōarai lies to the S. of Hiraiso and is the most noted pleasure resort in this district. The coast is dotted with rocks and backed by a low ridge. This shore is noted for shell-fish, particularly ear-shell. see Plate 30 The place can be reached from Mito either by *jinrikisha*, 9 m., or by descending the *Naka-gawa* by steamer as far as *Iwai*, whence the distance is 1.4 m. and is covered by *jinrikisha*. Aged pines grow in the precincts of *Isozaki-jinja*, a Shintō shrine, from which a good view of the ocean is obtained.

KANAGAWA=KEN

Hayama lies, 3.7 m. of the S. of *Zushi*, (33 m. from Tōkyō, in 1 hr. 23 min.), and as it projects far more into the sea it commands a better view than does the latter. On the S. lies a pine-clad point called *Chōja-ga-saki* with an inn (*Chōja-en*) built on it; this is an excellent spot for bathing. see Plate 31 An Imperial villa and the retreats of many notable personages stand here. Pop. 6,620. see Plate 32 Number of visitors: 1,551, the total stays being 29,096 days (1909).

Ōiso (40.8 m. from Tōkyō, in 2 hrs. 5 min.). The Japanese people, though living so close to the sea, never appreciated the benefits of sea-bathing, till Dr. Matsumoto Jun, the surgeon general of the Army, taught them to resort to seaside places, by first inducing a friend of his to open a seaside hotel at Ōiso (1886). Ōiso is now of full of villas and has grown to be a flourishing town with a population of 8,000. The late Prince Itō had a villa here, and the town thus became a centre of reliance whither statesmen flocked at times of political calm or stress. Ōiso, one of the most popular watering-place, is also a delightful winter resort, the climate being mild in winter. see Plate 33 Number of visitors: 17,502, the total stays being 107,796 days (1909). Inns: *Shōsen-kaku*, *Tōryū-kan*.

Hakone and Neighbourhood.

Hakone, the name of a large mountainous district, justly famous on account of its hot springs, beautiful scenery, and salubrious climate, is easily reached from Tōkyō or Yokohama by taking the Tōkaidō Line to Kōzu (48 m. from Tōkyō), thence by electric car to Yumoto Spa, which lies at the entrance to the interior of Hakone.

Tramway from Kōzu: 39 trips each way between Kōzu and Yumoto (Hakone), the distance being covered in a little over an hour. Tariff: between Kōzu and Odawara, 1st class, 45 *sen*; 2nd class, 30 *sen*; 3rd class, 15 *sen*; between Kōzu and Yumoto, 1st class, 90 *sen*; 2nd class, 60 *sen*; 3rd class, 30 *sen*.

Hakone may also be reached from two other directions, viz. from *Mishima* and *Gotemba*, both being stations on the Tōkaidō Line. (1) *Mishima* to *Odawara*, via the Hakone Pass. This is a part of the old Tōkaidō Highway, traversed so frequently by *Daimyōs* with their vast retinues on their periodical visits to *Yedo* (Tōkyō). Between *Mishima* and *Odawara*, via the Hakone Pass, the distance is 19 m.—roughly 9.5 m. up and the same down. The strictly mountainous part consists of about 17 m. of narrow, slipperly and steep pathway. After gaining the summit and descending slightly, we come to *Hakone-shuku* (Hakone-machi)—a hamlet on the shore of the beautiful Lake of Hakone (*Ashi-no-ko*, lit. 'a sea of reeds'). In feudal times there stood at this spot a barrier gate, where all the travellers passing through the Tōkaidō were examined. (2) From *Gotemba* to *Yumoto*. This road leads to the ridge of *Otome-tōge* (5 m. from Gotemba). Whence in 0.7 m. to *Sengoku-bara*, after which via the famous *Miyanoshita* Spa to *Yumoto*. This path is entirely outside the old Tōkaidō.

Hakone as a Popular Resort. Hakone is the name of an extensive mountainous region, which separates middle Japan from the Eight provinces of *Kéantō*. With the beautiful lake of *Ashi-no-ko*, the numerous hot springs, and splendid mountain scenery, the place fairly rivals *Nikkō* as a popular resort. If *Nikkō* may boast of its matchless shrines, Hakone's chief distinction lies in its plentiful spas and its charming views of *Fuji*. The hot springs are 12 in number and of different composition, as will be shown later on. Hakone has also a beautiful climate: the annual average temperature at *Yumoto* being 58° Fahr. (min. 19° and max. 92°) and at *Hakone-machi* 50° Fahr. (min. 10° and max. 88°). In spring the cherry-blossoms of *Kowakidani*, in summer the luxuriant verdure of the mountains and valleys, as well as the lake *Ashi-no-ko* (with facilities for boating), in autumn glorious tints of manifold leaves, and in winter the superb snow scenes are all so many attractions of Hakone. To persons interested in science, Hakone offers many opportunities of investigation, with its two-fold volcanic craters, a rare species of salamander (*Sanshō-urwo*), and its varied forms of

vegetation, to one moreover interested in Japanese history, Hakone will be found interesting on account of its associations with *Yoritomo*, *Hōjō Sōun*, *Hideyoshi*, etc.

Itinerary in Hakone. Undoubtedly the best plan for visitors to Hakone is to make *Miyanoshita* their headquarters. *Miyanoshita* is 4 m. from *Yumoto* and may be reached by *jinrikisha* in 1 hr. or on foot in 1 hr. 15 min. (there is a good road, though steep, by the *Haya-kawa*). The chief recommendation of *Miyanoshita*, besides its excellent baths, pretty surroundings, and salubrious atmosphere, is the hotel *Fuji-ya* (European style), which provides excellent accommodation. From here short excursions may be made, either in 'sedan chairs,' *kago*, *jinrikisha*, automobile, or on foot, to neighbouring places of interest. For example, a day's excursion may be tried, taking in *Kowakidani*, *Ashi-no-yu*, *Moto-Hakone* (or *Hakone-machi*), and from either one of the last-named places boat may be taken across the lake to *Umijiri*, thence coming back to *Miyanoshita* by way of *Ubako*, *Owakidani*, and *Kiga*.

Twelve Hot Springs of Hakone. The old-time 'Seven hot springs of Hakone' have been increased to twelve by the recent openings of five new ones:

Yumoto, situated at the foot of *Yuzaka-yama* and on the right bank of the *Haya-kawa*,—waters almost pure, with little trace of minerals.

Tōnosawa, 0.3 m. higher up from *Yumoto*, on the new road,—waters also of the same quality as *Yumoto*.

Miyanoshita, 3.5 m. yet higher up on the same road,—waters contain salt and soda. The place is 1,300 ft. above sea-level.

Sokokura, situated close to *Miyanoshita*,—waters also of the same quality as the above mentioned.

Dōgashima, near *Sokokura*,—waters almost pure, containing scarcely a trace of minerals.

Kowakidani, 2,000 ft. above sea-level and at the base of *De-yama*,—waters contain acid and iron.

Kiga, 1,300 ft., situated at the base of *Ninotaira* plateau and by the *Haya-kawa*,—waters contain salt and soda.

Gōra, 2,600 ft., and at the base of *Sōunjigoku*,—waters contain acid and sulphur.

Sengokubara, close to *Gōra*,—waters also of the same quality as the above mentioned.

Ubako, 2,877 ft., and situated at the base of *Kami-yama*,—waters contain a small quantity of salt and soda.

Ashi-no-yu, 2,760 ft., on a hill at the base of *Koma-ga-take*,—waters contain much sulphur.

Yunohana-zawa, 3,300 ft., at the foot of *Koma-ga-take*,—waters contain much sulphur.

Geography of the Hakone District.

The mountains of Hakone, together with *Fuji*, the *Ashigara* group, and other mountains in the N., such as *On-take*, *Asama*, and the *Usui Ridge*, constitute a thick wall on the W. and N. W. side of the *Kwantō* District (*Kwantō* means 'E. of the Barrier,' referring to Hakone where the famous barrier gate stood). These mountains send their off-shoots into the little peninsula of *Izu*, as if to make the separation between the regions E. and W. of these mountains even more secure. Among the Hakone group of hills, the better known are *Kintoki-zan*, *Koma-ga-take*, *Myōjin-ga-take*, and *Kami-yama* (4,788 ft.)—the last named being the highest peak in the group. The Hakone and *Ashigara* groups form a watershed from which flow rivers either towards *Sagami-nada* or the Bay of *Suruga*.

These mountains are all cone-shaped, with a blunt apex, a sure evidence that they owe their existence to volcanic eruptions. As may be seen from the plains of *Suruga* (e. g. from *Sano* on *Tōkaidō* Line) they have a long sloping base, with an angle of 7° to 12°. Most of these cone-shaped hills show at their summits traces of both older and more recent volcanoes, having double crater ridges. For instance, *Kintoki*, *Myōjō*, *Kurakake*, *Yamabushi*, *Mikuni*, *Umijiri*, *Nagao* and *Otome* are remnants of the older crater ridges, while *Futago*, *Kami-yama*, *Koma-ga-take* are the remaining portions from the ridges of the newer craters. Lake Hakone, where in clear weather is reflected the peerless form of *Fuji*, *Sengokubara* over which graze hundreds of cattle, and *Miyagino*, which is covered in summer with plantations of Indian corn, are none other than the sites of once active volcanoes. There are two rivers which drain the E. side of this watershed—the *Haya-kawa* and the *Sukumo-gawa*. The *Haya-kawa* has its source at *Ashi-no-ko* and, after irrigating *Sengokubara*, forms a rapid at *Chōshi-no-kuchi*, then, after crossing the *Miyako* Plain, it is enriched at *Sokokura* by the waters of the *Jakotsu-gawa*; now winding through the valleys lying between the *Myōjō* and *Sengen* hills, the *Haya-kawa* is finally joined at *Yumoto* by the *Sukumo*, to flow further on to *Sagami-nada*. (The new road leading from *Yumoto* to the other hot springs lies mostly along the

course of the Haya-kawa). The Sukumo has its origin in *Kurakake-yama*, and, before it meets the Haya-kawa, is joined at *Hata-juku* by the *Taki-zaka*. On the banks of the streams grows *Yama-wasabi* (Eutrema), which is sold as a preserved article of food, and in the river are found *Sanshō-urwo*, a kind of salamander. (The old Tōkaidō road lies along the course of this stream).

Some of these old volcanic peaks will repay climbing on account of the panoramic view obtained from the top. The upper and lower peaks of *Futago-yama* ('Twin hills'), which are the southernmost of the Hakone group and close to the lake, are very pretty mamelons, though the ascent is somewhat steep. *Kami-yama* is the highest of all the peaks of Hakone, and from its summit may be enjoyed the grandest view of any in this district. *Koma-ga-take* will repay the climber by affording the completest view of Lake Hakone.

In Hakone there are many fissures or holes whence issue hot vapours or steam. Such are *Owakidani*, *Sōunjigoku*, *Iwō-yama* and *Yunohana-zawa*, which emit either sulphur dioxide or sulphuretted hydrogen; at *Kowakidani* the exhalation consists chiefly of steam. The emission takes place with the greatest force at Ōwakidani ('Valley of the Greater Boiling'), otherwise called *Ojigoku*, or 'the Big Hill.'

Hot Springs.

Yumoto (Inn: *Fukuzumi* on the right bank of the Haya-kawa, excellent accommodation) is the oldest known spa in Hakone, being situated at the gateway, so to speak, which leads to all the other spas. Here the old and new roads branch off to the right and left, the one along the *Sukumo-garwa*, leading finally to Hakone-machi and Mishima, and the other (new road) along the Haya-kawa, leading to Miyanoshita and other spas. *Jinrikisha* is available over the whole of the new road.

The hot waters issue at the S. foot of *Yuzaka-yama*, whence they are conveyed in pipes to Yumoto. Specialties: toys, lacquer wares, cabinet work, marquetry works (desks, book-cases, screens, etc.).

Tono-sawa (Inns: *Shin-tamano-yu*, *Suzuki*—famous on account of its third story flat being made of *Jindai-sugi*, cryptomeria wood long buried in the earth—*Ichino-yu* and *Fukuzumi*) is reached after an ascent of 0,3 m. on the new road from Yumoto. The spa houses are situated by the two bridges spanning the Haya-kawa, at a place among the mountains where it makes an S-shaped bend. Cool and closed in by mountains, the place makes a fine summer resort. Electric Power House, situated near Tōno-sawa, supplies power to the Yokohama Electric Light Company.

Miyanoshita. Hotel: *Fujiya Hotel* is a splendid establishment—the buildings being partly in European and partly in Japanese style (European Building: 15 special double rooms, with private bath—Tariff, American plan ¥18; 24 ordinary double rooms—Tariff, ¥13-15; 28 single rooms—Tariff, ¥6,50-8,00. Japanese Building: 27 single rooms fitted up in European style—Tariff, ¥6,50-7,00). The hotel can accommodate altogether 130 guests. *Nara-ya* is also a first-class establishment, in both Japanese and European style, with 73 rooms—Tariff, ¥3,50-7,00. The late Emperor once stayed here when he visited Hakone many years ago. *Ryōun-kan*, also a good Japanese inn, boasts of possessing a bath tank where *Toyotomi Hideyoshi* frequently bathed during his siege of Odawara. (Curio shop, Nakada Shōkwai).

Miyanoshita is 4 m. from Tōnosawa, the road making many turns as it follows the course of the Haya-kawa. In 0,8 m. from Tōno-sawa we come to a cascade, *Tokiwa-no-taki*, and in 2 m. more, to *Ohira-dai*, a level spot whence Mt. Fuji is visible, and also Miyanoshita itself, only 1 m. off across a dale. Miyanoshita is a wide, level tract or terrace in the valley of the Haya-kawa, and is 1,223 ft. above sea-level. This wide flat is surrounded on nearly all sides by mountains—*Myōjin-ga-take* and *Myōjō-ga-take* across the river to the N. E., *Koma-ga-take*, *Kamuri-ga-dake*, and *Hōrai-zan* towards the E., and a chain of hills running from the S. W. towards the E. Only in the E. is a little opening, whence the blue waters of Sagami-nada may be seen. Miyanoshita is pre-eminent throughout Hakone on account of its excellent hotel accommodation, fresh air, and convenient location for making short excursions.

Dōgashima (Inns: *Ōmi-ya*, *Yamato-ya*) is situated 0,2 m. below Miyanoshita, close to the Haya-kawa. Being in the bottom of a valley, it is very quiet and secluded. In its neighbourhood are two cascades—*Shirabe-no-taki* and *Shiraito-no-taki*.

Sokokura (Inns: *Tsuta-ya*, *Ume-ya*, *Sengoku-ya*) is practically a part of Miyanoshita. It is a cluster of houses lying on the bank of the *Jakotsu-garwa*, which pours its waters into the Haya-kawa in the form of a cascade.

Kiga (Inns: *Kame-ya*, *Sengoku-ya*) is situated 0.4 m. up from Miyanoshita, on the right bank of the Haya-kawa. The spa is situated on a level space, surrounded by hills on all sides, except towards the S. E. It is a very secluded spot, somewhat similar to Tōnosawa.

Miyagino is a cultivated plain extending on both sides of the Haya-kawa and reached in about 0.3 m. from Kiga. Now studded with farm-houses, it formed originally a part of a large volcano, the rest of the crater becoming filled with water and being known as the *Ashi-no-ko*. To the S. E. of the village are found many cryptomeria trunks long buried in the earth, which on account of the peculiar colouring of the wood are much prized as building material, being known as the *Jindai-sugi* or cryptomeria wood of the mythological age. Specialty: buckwheat-rolls, to be eaten with soy.

Gōra, the most recently discovered of the 12 hot springs, is situated on a level tract on a slope of *Sōun-zan* and is drained by two streams on the E. and the W. Its altitude is from 1,813 ft. to 2,322 ft. Gōra is favoured with natural advantages which should make it the very center of Hakone. It commands a wide prospect, which includes Miyanoshita and the larger part of the valley of the Haya-kawa, and also the sea beyond. The air is fresh and comparatively dry, with plenty of sunshine. Above all the mineral waters are practically inexhaustible, as they are conveyed from the stream flowing from the Ōwakidani springs. A scheme is on foot to extend as far as Gōra the electric tramway from *Kōzu*, now reaching to *Yumoto*, and large pleasure garden, with band-stand, swimming tank, club house, and recreation hall, has already been laid out. It is believed that the electric tramway will probably be completed in a few years.

Sengokubara (Inn, *Ishimura*) consists really of two springs, one lower down (1.2 m. from *Miyagino*) and the other about 0.3 m. higher up. The two springs are resorted to by the inhabitants of the neighbourhood.

Ōwakidani, or the 'Valley of the Greater Boiling,' is otherwise called Ōjigoku, or the 'Great hell,' (0.7 m. from the lower waters of Sengokubara and 1.3 m. by a round-about path from Gōra). It is a gorge full of solfataras, which from its middle part up to the top for about 0.8 m. literally reeks with sulphurous fumes. Wherever a little hole is dug—1 or 2 ft. deep—out start steam and fumes. The ground everywhere presents a light brownish colour and overlying crust is so weak and unreliable that no step is safe unless directed by a guide. There is very little vegetation. The hot mineral waters, which collect into a stream, are made somewhat lower down to form a reservoir, whence they are conveyed to Gōra and also to Sengokubara. At the head of the gorge one may obtain superb view of Fuji and other mountains, a singular contrast to the weird and desolate scene of the solfataras below.

Ubako (Inn, *Shūmei-kwan*), 0.9 m. from Ōwakidani and the innermost of the 12 baths, is situated on the S. slope of *Kamuri-ga-dake*, at an altitude of 2,877 ft. Towards the W. lie the pasture lands belonging to the village of Sengokubara, and beyond them the *Ashi-no-ko* and Mt. Fuji, while in the N. there looms up the hill, *Dai-ga-take*. The hot waters bubble out of rock crevices, and bath-tanks have been improvised by excavating rocks. In the neighbourhood of the tanks, fossil leaves or twigs are sometimes found.

Umijiri, at the N. end of Lake *Ashi-no-ko* and reached in about 1 m. from the Ubako Spa, is the outlet for *Ashi-no-ko*, where the Haya-kawa has its source. From Umijiri to Moto-Hakone and Hakone-machi there is a regular boat service.

Kowakidani (1 m. from Miyanoshita; Hotels: *Mikawa-ya*, *Kaikwa-tei*). Kowakidani is situated on a slope of *Kojigoku-yama*, and at an elevation of 2,000 ft. Opposite the spa, across the valley of the *Jakotsu-gawa*, there rise tall peaks of *Myōjō-ga-take* and *Myōjin-ga-take*. In the neighbourhood of the spa there have recently been planted 10,000 cherry-trees and a large number of maples and azaleas, so that the place is destined to be one of the most attractive places in Hakone. A little higher up (0.4 m.) the hill *Kojigoku-yama* is a cave which constantly emits sulphurous fumes.

Chisuji-no-taki, or the 'Waterfall of 1,000 threads,' is in the upper course of the *Jakotsu-gawa*, where the waters of the river dash down from innumerable crevices on a high cliff, giving the appearance of thousands of white threads hanging down from the height above.

Ashi-no-yu (4.5 m. from Miyanoshita by the new road or 2.7 m. by the old road).

Inns: *Matsuzaka-ya* (17 European rooms, besides 100 Japanese rooms); *Kinokuni-ya*.

The road winds through the valleys and round the base of a hill for about 2.5 m., until we reach a flat space called *Benten-yama*, whence a good view may be obtained of Sagami-nada, the peninsula of Misaki, and the distant hills of *Arwa* and *Kazusa*, while the nearer mountains in sight

are *Iwō-yama* to the left and *Futago-yama* and *Koma-ga-take* on the other side. *Ashi-no-yu*, a little below this elevation, is 2,760 ft. above sea-level and is delightfully cool in summer. In winter the place is well protected from the N. W. cold winds by *Koma-ga-take*. The only draw-backs are the occasional thick fogs of summer and the heavy snows of winter.

Yunohana-zawa (Inn: *Bosenkaku*), about 0.6 m. N. of *Ashi-no-yu* and 1 m. from *Kowakidani* by a short cut, is situated in a valley at the S. base of *Koma-ga-take*. The place is at an altitude of 3,300 ft. It lies on the road leading from *Ashi-no-yu* to the top of *Kami-yama*.

Moto-Hakone (Inns: *Matsuzaka-ya* and *Sakamoto-ya*, both facing the Lake *Ashi-no-ko*, on the surface of which on a clear day the reflected figure of *Fuji* may be seen—the famous *Sakasa-Fuji*, ‘inverted *Fuji*.’ *Matsuzaka-ya* has 13 rooms fitted up in European style, as well as Japanese rooms). *Moto-Hakone* is a village of 40 houses and is 2.2 m. (by the new road) from *Ashi-no-yu*.

Moto-Hakone (lit. ‘Original *Hakone*’) was the chief village of the mountain district of *Hakone*, before *Hakone-machi* was founded in 1618 as a barrier town. Though it has since been overshadowed by the new *Hakone*, there is no question that the place is most picturesquely situated. Shut in on one side by a thickly-wooded hill, on the other side a charming vista opens out showing one of the finest views of *Fuji*, as well as its inverted image reflected in the lake, and to the left the Imperial villa on a small headland projecting into the lake.

Ashi-no-ko, or Lake *Hakone* as it is generally called by foreigners, has a maximum length of 3.4 m. from N. to S. and is 1.4 m. from E. to W. in its greatest width; its circumference being about 9.8 m. It is in the form of a gourd, with a narrow middle part where the promontory *Tōga-shima* protrudes. The Imperial villa, *Hakone-Rikyū*,—a building partly in Japanese style and partly in European—stands on the edge of this promontory. The scenery of the lake, if inferior to that of *Chūzenji* near *Nikkō* or that of *Nishi-no-umi* at the N. base of *Fuji*, is charming enough to attract numerous visitors in summer. There are rowing-boats for hire, as well as Japanese house-boats with boatmen, and, as already remarked, there is a regular ferry service maintained between *Hakone-machi* (via *Moto-Hakone*) and *Umijiri* at the N. end of the lake.

Hakone-machi (0.6 m. from *Moto-Hakone*; Inns: *Ishiuchi Ryokwan*, *Enshū-ya*, *Hakone Hotel*) is reached from *Moto-Hakone* by a road which is overshadowed by an avenue of tall cryptomerias, round the ancient branches of which twine ivy and parasitic orchids, imparting a venerable aspect to their stately hosts. At a spot about 0.2 m. from this side of the town, there are the remains of the old barrier gate, where formerly all travellers through the *Tōkaidō*, from the greatest town to the humblest, were examined, the custom being only abolished with the Restoration of 1868. The barrier, it will be noticed, was most strategically situated, being across a narrow pathway with a high and steep hill on one side and the lake on the other.

The villagers mostly made their living as inn-keepers till about 20 years ago, when the *Hakone* tunnels of the *Tōkaidō* Railway were opened to traffic. Since then the village, which had been never too prosperous, rapidly declined. Within recent years, however, the place has become quite a popular summer resort for both Japanese and foreigners. There are many houses to let, where Europeans bringing their servants with them may set up housekeeping. Some of the inns, like the *Hakone Hotel*, have rooms fitted up with beds and furnish European food.

Summary.

| Spa | Altitude above sea-level in ft. | Population | Number of houses | Number of baths | Number of inns | Number of visitors in 1919 | Their total stays in days |
|---------------------|---------------------------------|------------|------------------|-----------------|----------------|----------------------------|---------------------------|
| Yumoto and Tonosawa | 150 | 1,853 | 309 | 6 | 10 | 26,528 | 60,395 |
| Miyanoshita | 1,300 | 917 | 163 | 13 | 10 | 16,324 | 50,345 |
| Kiga | 1,300 | 70 | 13 | 3 | 2 | 1,207 | 1,947 |
| Ashi-no-yu | 2,760 | 81 | 9 | 2 | 2 | 5,984 | 25,050 |
| Gora | 2,600 | 8 | 2 | 1 | 1 | 190 | 572 |
| Utsunomiya | 2,870 | 13 | 1 | 1 | 1 | 1,708 | 10,903 |
| Yunohana-zawa | 3,300 | 7 | 1 | 1 | 1 | 181 | 848 |

Yugawara, situated in a quiet, pretty spot by the *Fujiki-gawa*, is 350 ft. above the sea-level and is 7.2 m. from *Hakone-machi* (the path lying across the *Jikkoku-tōge*). It is walled in on three sides by mountains—on the N. by the *Hakone* hills, on the S. W. by *Higane-yama* (*Jikkoku-tōge*) and *Izusan*; being open only on the S. E. towards *Sagami-nada*.

Odawara-Atami Light Railway (15.8 m.) lies mostly along the coast, on rocky cliffs skirting the group of Hakone hills, excepting at one place—*Yoshihama*—where the tracks lie close to a sandy beach. All along the route there is a splendid view of the sea with its numerous isles. Nine trains run daily from each end, between 6 a.m. and 3 p.m.; the time required to cover the entire distance being 2.5 hrs., and the fare, 2nd class ¥ 1.20, 3rd class 70 *sen* (no 1st class cars).

Yugawara (also called *Mongawa*) is a station of the light railway (10.8 m. from *Odawara*, in 2 hrs.; fare 2nd class 90 *sen*), 1.9 m. from the hot springs of Yugawara; from the station to the spa *jinrikisha* (fare 20 *sen*), also *basha* (fare 20 *sen*, for exclusive hire 80 *sen*) available. Pop. 500; 2 public baths; 1 doctor. Number of visitors: 6,500, the total stays being 195,000 days (1909).

Inns: *Nakanishi-ya*, *Ueno-ya*, etc.

SHIZUOKA-KEN

Izusan (14.4 m. from *Odawara*, by light railway in 2 hrs. 20 min.; fare, 2nd class ¥ 1.17).

Izusan spa, near the station, is situated on a rocky cliff facing the sea, inns being built close to its edge. It is reached from the station by a steep path down. Walled in on three sides by hills and rocks, it is open towards the sea, with a full view of *Vries Island* and the headland of *Awa*. Hot water bubbles up from rocky crevices with violent force, 42 ft. above the sea. The water is called *Hashiri-yu* ("Running Hot Water"). Pop. 1,370. The springs are said to be good for brains. There are 8 inns, each provided with its own baths. In *Sagami-ya*, the best inn in *Izusan*, is a large swimming bath, 60 ft. in length and 18 ft. in width. Hot water massage by "*Yudaki*" (hot waterfall) can be taken in almost all inns. Number of visitors: 2,768, the total stays being 19,363 days (1909). Inns: *Sagami-ya*, *Enoshima-ya*, etc.

Atami (15.8 m. from *Odawara*, in 2 hrs. 30 min.; Inns: *Atami Hotel* or *Higuchi*, with both European and Japanese accommodation, *Fuji-ya*, *Suzuki*, *Sagami-ya*) is a fashionable watering-place, with a resident population of 5,866. Surrounded by hills on three sides and open towards the sea on the S. E., it is warm in winter (lowest temperature 35° Fahr.) and cool in summer (highest temperature 85° Fahr.). Its beach offers excellent sea-bathing, and the scenery is charming. The prospect takes in towards the left *Izusan* and *Manazuru-ga-saki* Promontories, and yet further off the mountains of *Awa*, while towards the right stand out *Uwomi-ga-saki* Promontory and *Vries Island* smoking in the distance, with the pretty little islet of *Hatsu-shima* in between. Atami has many attractive walks, either by the seashore or on the hillsides.

The hot waters, which contain a large quantity of salt, have their source in a powerful geyser. It is called *Ō-yu*, or the 'Great Hot Water.' The geyser breaks out once in about 4 hrs., when it shoots forth boiling water and steam with great force and noise. Originally the jet shot straight upward, but it is now partially smothered by placing a big stone over it. Sometimes the geyser continues active for more than 12 hrs., such cases occurring several times a year. The waters are transparent, with a temperature of 209–220° Fahr., and are carried through pipes to the various bath-houses. Near the geyser is an inhalation house (*Kyūki-kwan*), into which the steam from the geyser is conveyed for the benefit of patients suffering from bronchial and lung troubles. There are doctors connected with the house.

Formerly there were in Atami seven springs in all, but the number of the springs has considerably increased (about 20), since the people has begun to bore the ground for finding hot springs, in consequence of which the world-famous intermittent spring has never been in full play and the discharge of water decreased remarkably. The springs of Atami are believed to be efficacious especially for the treatment of rheumatism and eye diseases.

Geologically *Atami* forms part of an extinct crater, the hills behind it—*Iwato-zan*, *Higane-yama*, *Kamizawa-toge*, *Takichi-yama*—forming one-half of the old crater wall, while the other half has been submerged by the sea. The geyser is a faint reminder of the crater's former activity.

Atami Park is situated on a hillside 0.6 m. N. of the town. The Park contains more than 3,000 *ume*-trees, which blossom early in January, about two months before the flower season in Tokyo, also peach-trees, and several splendid old pines and cypresses (*hinoki*), with a pretty, silvery stream winding through the *ume*-grove.

Higane-yama, popularly known as *Jikkoku-tōge*, is a high ridge lying on the path between *Hakone-machi* and *Atami*. The ridge, which is 2,684 ft. high, is reached after a steep ascent of 3.6 m. from *Atami*. The upper portions are bare of trees, being covered only by low shrubs. The

view from the top is superb, embracing all the surrounding provinces, as its name of Jikkoku-tōge (lit. 'Ten Province Pass') indicates. On the summit is a stone monument giving the points of the compass. The first object to strike the eye is Fuji, which bursts upon one's gaze as one gets to the top, and is followed by other well-known views of the surrounding provinces—mountains of the Ashigara and Hakone groups, of Kai and Shinano, the *Fuji-kawa*, the classical beaches of *Miho-no-matsubara*, *Tago-no-ura*, and *Kiyomi-gata*, the towns of *Numazu* and *Mishima*, the Izu hills of Izu Province, such as *Amagi* and *Managi*, the town of Atami and its neighbouring sights, the islands of *Hatsu-shima*, *Oshima* (Vries Is.), *To-shima*, *Nii-jima*, etc., the mountains of Awa and Kazusa, etc.

Number of visitors: 23,638, the total stays being 174,306 days (1909).

Itō (13 m. S. from Atami). There is steamship service which connects Itō with Atami, Kōzu, Odawara, and Tōkyō (*Reigan-jima*). A road leads from Itō to Shūzenji (12 m.), *jinrikisha* and omnibuses available. The spa lies on the E. coast of Izu Peninsula, and is surrounded on three sides by hills, a branch range of *Amagi-san* and *Hakone-yama*, while, it is open to the sea on the E. A small stream of the *Matsu-kawa* runs through the spa town towards the E. It is divided into two parts, *Matsubara* and *Kusumi*, separated by the *Matsu-kawa*. The place is protected by the hills from the N. W. wind, which prevails in winter in this region, and warm and fresh sea breezes make it one of the best winter resorts with a mild climate. The island of *Hatsu-shima* lying near the coast and the mountain range of Awa and Kazusa far beyond present a very fine view. It is noticeable that the spa stands on a vein of hot springs. Hot springs are found almost everywhere. But those which are now open to public use, such as *Shishido-Onsen*, *Deki-yu*, *Wada-Onsen*, *Shin-Onsen*, *Yuda-Onsen*, *Me-no-yu*, etc., have been known to people from ancient times and are believed to be most efficacious, especially *Deki-yu* and *Shin-Onsen* for diseases of the skin, wounds, and disorders of the female genital organs, and *Wada-Onsen* for rheumatism.

In the whole town of Itō there are at present more than 300 baths, of which 10 are for public use; 33 inns with private baths. Close by *Shin-yu* is a special bath for oxen and horses. In the Era of the Tokugawa Shogunate government the water from the spring *Deki-yu* was transported in wooden barrels to Tōkyō for bath. The spa is also an excellent place for sea-bathing in summer. Pop. 8,563 (1,379 households). Lake "*Yoshida-no-Oike*" with beautiful scenery, is near Yoshida-mura (S. of Ito). The lake is stocked with carps and *funa* and a great place for fishing. Number of visitors: 11,365, the total stays being 79,568 days (1909). Inns: *Denko-en*, *Tōkyō-kwan*, *Masu-ya*.

Shuzenji (3.1 m. from Ōhito; *jinrikisha* fare 25 *sen*; omnibus 15 *sen*) claims with Atami the honour of being the most popular spas in Izu. A temple founded here in the Daidō Era (806–9) by Kūkai or Kōbō-Daishi has given rise to the name of this place. It is bounded on the N. and S. by mountain ranges, which leave a narrow strip of land extending from E. to W., watered by the *Katsura-gawa* which joins the *Kano-gawa*. Several hot springs issue from the bed of the Katsura, and of these the one called *Dokko* is left to the common use of the public from ancient times, and is most noted for its remarkable efficacy for rheumatism. The spring "*Dokko-no-yu*" is said to have been discovered in the Era of Daidō (806–10) by the famous priest Kūkai. Of the six other public baths the most noted are *Sugi-no-yu* or "Cryptomeria hot spring," *Ishi-yu* or "Stone hot spring," *Chigo-no-yu* or "Child hot spring," and *Ayame-no-yu* or "Iris hot spring." The spa lies at an altitude of 250 ft. above the sea. Pop. 2,250; 355 houses; 14 hot springs. Number of visitors: 24,969, the total stays being 111,414 days (1909). Inns: *Arai*, *Kiku-ya*, *Asaba*, *Yomo-rō*, etc.

Asahi Waterfall is found to the W. of Shuzenji and measures over 300 ft. high and 15 ft. wide, the water being divided into two parts in the middle, and beautifully reflecting the morning light. Its water flows into the *Kano-gawa*.

Summary.

| Spa | For the year, 1909 | Spring
(March, Apr., May) | Summer
(June, July, Aug.) | Autumn
(Sept., Oct., Nov.) | Winter
(Dec., Jan., Feb.) | Total |
|----------|-----------------------------|------------------------------|------------------------------|-------------------------------|------------------------------|---------|
| Izusan | Number of visitors | 973 | 616 | 610 | 500 | 2,700 |
| | Their total stays in days | 6,813 | 4,429 | 4,355 | 3,780 | 19,362 |
| Atami | Number of visitors | 6,804 | 3,986 | 4,434 | 3,111 | 18,335 |
| | Their total stays (in days) | 54,631 | 27,904 | 32,867 | 58,904 | 174,306 |
| Shuzenji | Number of visitors | 9,104 | 6,405 | 5,673 | 3,787 | 24,969 |
| | Their total stays in days | 38,959 | 34,273 | 20,787 | 17,402 | 111,411 |

Nagaoka. 0,6 m. W. from the *Nanjō* station on the Sunzu Electric Tramway which leads from *Mishima* to *Ōhito* (10,6 m. in 1 hr.); *jīnrikisha* (fare, 20 *sen*) and omnibuses (fare, 15 *sen*) available. Pop. 358. The springs were found in 1907, and are said to be efficacious for rheumatism and eczema. Mt. Fuji forms a sublime view from the spa. On account of the mild climate of this place, a number of wealthy people have built villas here. 1 public bath, 14 inns, 1 doctor. Number of visitors: 8,229, the total stays being 27,966 days (1909). Inn: *Yamato-kwan*.

Kona, 0,5 m. W. from the *Nanjō* station, has 5 baths, of which 1 is for public use. The water is said to be efficacious for scald. Pop. 586. Number of visitors: 7,331, the total stays being 12,757 days (1909).

Hatage. 2 m. W. from the *Daiba* station on the Sunzu Electric Tramway; *jīnrikisha* (fare, 35 *sen*) and omnibuses (fare, 12 *sen*) available. Pop. 360. 5 inns with private baths. There are 8 hot springs in the spa, efficacious for diseases of the skin and venereal diseases. Number of visitors: 1,514, the total stays being 28,382 days (1909).

Funabara. 7 m. S. W. from *Ōhito*, *jīnrikisha* (fare, 56 *sen*) and omnibuses (fare, 30 *sen*) available. The spa is 7,5 m. S. W. from *Shuzenji*, 1,3 m. N. from *Yoshina*, and 7 m. E. from *Tōhi*. It is penetrated by a stream called *Funabara-gawa*. It is a quiet summer retreat and the cold is not very intense even in winter. Number of visitors: 1,630, the total stays being 6,160 days (1909). Inn: *Shūfū-kwan* (*Suzuki*). The private bath of the inn is called *Kōgyoku-yu* and believed to be efficacious for skin diseases, haemorrhoids, gonorrhoea, uterus diseases, dyspepsia, etc.

Yoshina. 8 m. S. from *Ōhito*, *jīnrikisha* (fare, 56 *sen*) and omnibuses (fare, 32 *sen*) available. The place lies on the bank of the *Yoshina-gawa* at the N. foot of *Amagi-san*, at an altitude of 360 ft. above the sea. The spa is shut in on two sides by mountains densely wooded with pines and cryptomeria. The springs were found in the Era of *Eihō* (1081-84) and said to be efficacious for rheumatism, uterus diseases, haemorrhoids, dyspepsia, etc. Pop. 240. Indeed it is a quiet spa. Number of visitors: 2,345, the total stays being 14,319 days (1909). Inn: *Tōfu-ya*.

Mt. Amagi, 4,981 ft. high, forms the backbone of the peninsula, being part of the Fuji volcanic chain. The whole mountain extends for 14,6 m. from S. to N. and 2,4 m. from E. to W. and its summit is 7,3 m. uphill from *Yugashima*. It is densely wooded and partly an Imperial game preserve. The southern foot of *Amagi-san* is also rich in hot springs.

Yugashima is found about 7,3 m. to the S. of *Shuzenji*, sheltered on the S. by the range of *Amagi*. The inns with baths stand close by the banks of the river *Kano-gawa*. The spa (625 ft. above the sea-level) is a quiet summer resort in *Izu Province*. 7 springs, 2 public baths, 1 doctor. Pop. 1,630. Number of visitors: 2,732, the total stays being 6,493 days (1909). Inns: *Yumoto-kwan*, *Ochiai-rō*.

The road from here to *Shimoda Port* at the southernmost extremity of the peninsula, about 24,4 m., leads over the *Amagi Pass*, but is negotiable by *jīnrikisha* or omnibuses.

Yugano (26,3 m. S. from *Ōhito*) lies at the S. foot of *Amagi-san* on a highway leading to *Shimoda* (sea-port), and on the bank of the *Kawazu-gawa*. The water is said to be efficacious for eye diseases. Pop. 340. Number of visitors: 50, the total stays being 106 days (1910).

Konabe. Near *Yugano* there is another hot spring called *Konabe*, with 205 inhabitants, 1 public bath, and 2 inns. The water is believed to be efficacious for diseases of the skin, swellings and wounds.

Yatsu (30 m. S. from *Ōhito* and 9,3 m. N. from *Shimoda*) stands on the E. coast of the peninsula, facing the stream *Kawazu-gawa*. Pop. 335. 1 public bath. 2 inns. Efficacious for syphilis, haemorrhoids, uterus diseases, eye diseases and dyspepsia. Number of visitors: 176, the total stays being 344 days (1910).

Rendaiji. 2,3 m. N. from *Shimoda*, *jīnrikisha* available. The place, lying on the bank of the *Inōzawa-gawa*, is bounded on the S. and N. by hills and situated along a highway leading to *Shimoda*. 2 public baths. 6 inns with private baths. Efficacious for swellings, diseases of the skin, uterus diseases, syphilis, rheumatism, etc. Number of visitors: 1,852, the total stays being 2,369 days (1909).

Kōchi. 2,5 m. N. from *Shimoda* by the highway of *Shimoda-kaidō*. The place is penetrated by the *Inōzawa-gawa*. The spring was found in 1839. The efficacy of the water is said to be the same as that of *Rendaiji*. There are 132 houses and 4 inns in the hamlet. Number of visitors: 431, the total stays being 468 days (1909).

Shimoda has a hot spring called *Akama-Kōsen* near the sea coast. It is 6-7 ft. above the sea-level and close to the *Inōzawa-gawa*.

Shimogamo. 7.5 m. S. W. from Shimoda, *jinrikisha* available. From the bank of the *Aonogawa* issue several hot springs, the water being all alkaline common salt springs. Efficacious for swellings and uterus diseases. Pop. 2,834. 5 inns. Number of visitors: 4,030, the total stays being 4,093 days (1900).

Tohi. 20 m. S. W. from Numazu (in 2 hrs. by steam-boat, fare, 25 *sen*). The place (30 ft. above the sea) lies on the W. coast of Izu Province, facing Suruga Bay. Pop. 3,627. *Mt. Fuji* and *Miho-no-matsubara* are visible from the spa. 3 public baths. 11 inns. 2 doctors. The water is said to be efficacious for rheumatism, dyspepsia, diarrhoea, catarrhs of intestines, disorders of the female genital organs, eczema, etc. Number of visitors: 17,148 (1913). Inns: *Asaka*, *Meiji-kwan*.

YAMANASHI-KEN

Yumura, 0.8 m. S. from *Kōfu* (80.3 m. from *Iidamachi*, in 6 hrs. 13 min.), lies on a path leading to *Mitake*, on a flat ground at the foot of *Yumura-yama*. Pop. 200. The springs are said to have been found in 809 by Daigaku Zenshi. On account of easy access from *Kōfu*, the spa is mostly patronized by its citizens. Number of visitors: 14,088 (1909).

Masutomi. 12.8 m. N. from *Nirazaki* station (88.3 m. from *Iidamachi*, in 6 hrs. 52 min.) on the Central Main Line (*Iidamachi-Nagoya*), and 15.5 m. from *Hinoharu* station (7 hrs. 39 min.). The place is 23.5 m. distant from *Kōfu*. There are two routes to reach the village of *Masutomi*, where are cold springs of the strongest radio-activity in Japan.

(a). Starting from *Nirazaki* station and going N. by a highway, which leads to *Iwamurada* near *Komoro* (*Shinano* Province) along the valley of the *Shio-kawa*, we reach *Mamyōda* via *Shimo-jō*, *Naka-jō*, and *Odagawa* (*jinrikisha* available as far as *Mamyōda*); from *Mamyōda* the way diverges into two, the left one leading to *Wakamiko* (Pop. 2,818) and the right to *Kami-kandori* via *Fujita*. Proceeding farther up the valley of the *Shio-kawa*, with the two mountains *Kana-ga-dake* (5,544 ft.) and *Kaya-ga-dake* (5,874 ft.) on the right, we come to *Higashi-muki*.

(b). One who starts from *Hinoharu* station goes at first to *Wakamiko* and then across the *Sudama-gawa* going N. E. the road meets with the route (a) near *Higashi-muki*.

A small distance to the hamlet the way ascends by degrees, the banks of the *Shio-kawa* becoming higher and steeper. Here hills stand direct from the brinks of the stream on both the sides, agitated water forming abysses and cascades here and there, while old pine-trees grown on quaint-shaped rocks add to the charms of the sight. The abysses are called "*Sen-ga-fuchi*" or 'one thousand abysses.' A little up the stream, there is a poor bathing-place "*Azuma-Onsen*" at the foot of a rocky and precipitous hill. The way now leads across the stream to *Negoya*. The two big *keyaki*-trees (*Zelkova acuminata*), which stand on the ground of *Negoya-jinja* (*Shintō* shrine), are said to be more than one thousand years old. A little farther we cross again the *Shio-kawa* and soon come to a small hamlet called *Taira*.

After passing the place we descend to the stream, on which stands the village of *Egusa* (Pop. 2,423). From *Egusa* we must climb *Torii-zaka*, one of the steepest passes in *Kai* Province, the ascent being about 0.8 m. A good example of the erosion of rocks by water can be seen from the village on the S. slope of the hill, which may interest geologists. Though the ascent is not more than one mile, it is indeed a laborious work, the path being very steep and stony. When we reach the top, we can get a picturesque view over the valley of the *Shio-kawa*, the hamlet *Hishi* lying below on its bank. The *Hishi-jinja*, one of the places of historical interest in the vicinity, stands close by the stream.

Passing through the hamlet, and turning right, we begin again to climb over "*Nagasaka-tōge*" or 'long pass,' the ascent being longer than over *Torii-tōge*, but not so steep. From the top of the pass a splendid, panoramic view of the surrounding mountains can be had—*Kimbu-san* (8,647 ft.), *Mizugaki-yama*, a branch peak of the *Kimbu-san* Range on the N. E., and *Koma-ga-dake* (9,900 ft.) of *Kai* Province on the S. W.

Coming down the pass, the road leads to a hill with two hamlets at its foot. The one lying on the S. side is called "*Hinata*" meaning 'sunshine,' and the other on the N. side "*Hikage*" meaning 'shade.' In front of the latter hamlet flows a small stream called "*Hontani-gawa*," which is an upper course of the *Shio-kawa*. A small hamlet *Oshiba-mura* with a dozen houses, lying on the opposite side of the stream forms an entrance to the cold springs of *Masutomi*. While crossing the

bridge, we see gas rising from the bed of the stream, and at Ōshiba-mura we come across several springs for the first time. One spring, issuing from the hill-side near Ōshiba-mura, is called *Ginsen-tō* or "Silver Spring," the water being carbonated and calcareous. The bath-house, having been destroyed, is now under construction. After passing through an extensive region belonging to the paleozoic system, our eyes are greeted by granite rocks here.

About 0.4 m. up the stream there is on the right bank another hamlet with some 30 households, called *Higashi-obi*. It is a poor mountain hamlet, most of whose inhabitants are farmers and some miners for rock crystals or tungsten at the foot of Kimbu-san. At a little distance from the hamlet, is found a tract of land, which is famous for a large number of cold springs of strong radio-activity. These springs, issuing from hill-sides, gorges, and the bed of the stream, are equally located on a granite formation, not very distant from one another, and the water contains as its main constituents common salt and bicarbonates of the alkaline-earth metals, impregnated with a large amount of carbonic acid gas. The water is moreover, endowed with strong radio-activity. The water of various sources, which shows strong radio-activity, is of low temperature, so that it must be previously heated for bath. *Kinsen-tō* or "Gold Spring" seems to have been known to people for a long time and is used for bath, on account of its comparatively higher temperature (33° C.), but the water shows the weakest radio-activity of all the springs in the neighbourhood. This is, however, only one spring used for bath, though its equipments are very poor, while others remain even now in a state of *laissez-faire*.

Several springs produce a considerable quantity of gas, rich with radium emanation. Both water and gas can undoubtedly be utilized with good results for medical purposes, internally or externally, but no arrangement has yet been made for that end. The place is delightfully situated amidst scenic surroundings. The upper course of the Hontani-gawa is a rival in the beauty of natural scenery with *Mitake* of Kai province. The scenery consists in densely wooded peaks, fantastically shaped rocks, narrow ravines, and blue meandering streams, all these intermingled with trees. Rhododendrons in late spring and tinged foliage in autumn are also attractive.

Mt. Kimbu is noted for its crystals and scheelite. There is also the site of an old gold mine, carried on by *Takeda Shingen*. On the valley, into which the spring water flows, near *Kinsen-tō*, leaves of some trees are covered with mineral sinter. The water is barrelled here and conveyed on horse-back to Hinoharu station to be again sent to Tōkyō. In the N. of the Hontani-gawa there is another branch called *Kamase-gawa* of the Shiokawa, nearly paralled with the former. This stream supplies water to a group of hamlets, such as *Shiokawa*, *Gōdo*, *Mikado*, *Wada*, and *Kuromori*, the last lying on the uppermost bank of the stream. This group is generally called "*Nishi-obi*" and possesses a large number of cold springs producing carbonic acid gas on the bank of the *Kamase-gawa*. The surrounding scenery is also attractive. Going farther up the valley, we come to a pasture called "*Matsudaira Bokujō*," from which *Mizugaki-yama* with ragged peaks is in full view. The ascent of this mountain is very hard, but a trip even to its foot does not fail to interest one. On account of their inconvenient location, Masutomi cold springs are still left to their own devices. The springs were first examined a few years ago for their radio-activity by M. Minagawa, an expert of Yamanashi-ken, but it was Dr. Ishizu, an expert of the Tōkyō Imperial Hygienic Laboratory, that determined at their sources the quantity of radium emanation contained in the water and brought the fact to public notice.

Kurobira is a watering-place, situated on the N. slope of *Mitake*, about 14 m. distant from Kōfu. The springs are found in the 3rd year of Genroku (1619) by a Shintō priest of *Mitake-jinja*. Though the baths are poor, the road leading from Kōfu via *Mitake* is beautiful and worth travelling.

Mitake is noted for the best natural scenery in the neighbourhood of Kōfu, from which to *Kin-ō-jinja* on the *Shimmichi* ('New Road') is 11 m. To the foot of *Wada-tōge*, about 1.4 m., *jirikisha* are available. After reaching the foot of *Wada-tōge*, an ascent of about a mile brings us to the top, from which a fine view can be obtained. On descending the pass we come to a cultivated plain, then proceed for about 2.4 m. round the foot of *Takasago-yama*, and find ourselves on the bank of the *Ara-kawa*, where the grove of *Tenjin-no-mori* stands. Here begins sight No. 1 of the beauties of *Mitake*. From the grove we follow the valley of the mountain stream for about 5 m. N. The foaming torrent as it dashes against the fantastic rocks that intercept the current, the cliffs rising here and there on the banks, the waterfalls tumbling down by the road-sides, all combine to attract the attention of those who are susceptible to what is beautiful and sublime.

see Plate 10,
fig. 2 and 4

see Plate 33,
fig. 2 and 3

see Plate 34,
fig. 1

see Plate 35,
fig. 1

At times the path leaves the valley and the sound of the dashing torrent is heard far below our feet. The grove at this place is called *Tenke-rin* ('Heaven's Drum Grove'), and leaving the grove we come to an open place, but soon rejoin the valley and find ourselves confronted with a column-like gigantic rock rising perpendicularly several hundred feet. This is called *Kakuen-hō* and is considered one of the best sights in Mitake. Near this column there is the *Kinkei-kwan Hotel*, which occupies a site of great scenic beauty. Still climbing upward, we come to statue erected in memory of the famous road-maker, and near it stands a stone monument with an inscription giving the history of this singular exploit. Continuing the ascent a stone gate is met with, and at a short distance from it the bridge *Shōsen* spans the mountain torrent. The views commanded from the bridge are really romantic. Then comes the *Senga-daki* cataract, the most interesting of all in Mitake, and next the second stone gate, after which we reach the somewhat open space occupied by the village of *Ikari*, where the road-maker lived and died, and at last, after a mile's progress, the Shintō shrine of *Kin-ō* is reached, 10 m. from Kōfu, quite impressive in its structure and solemn surroundings. The best time to visit Mitake is in autumn, when the valley is ablaze with the crimson tints of maples and other leaves.

Mt. Kimbu-san can be climbed from this place. The crystals obtained from the mountain are noted for their quality, and their yearly output amounts to ¥100,000, then come those obtained from *Takemori-yama*. Tungsten is extracted from the crystal-bearing veins. The crystal produced in Kai are purple, black, *mizuri* (containing water), *kusairi* (containing leaves), and brown, and these are worked into balls, rosaries, rings, stamp stones, stationary articles, hair-ornaments for women, and other decorative wares.

Kami-Sekisuiji (2 m. N. from Kōfu) lies on the slope of *Sekisuiji-yama*, on which extends a dense state-owned forests, commanding a fine view over the plain of Kōfu and *Mt. Fuji* beyond. The spring is said to have been found in the Tenshō Era (1573-92). Number of visitors: 350, the total stays being 9,450 days (1909).

Shimobe (22 m. S. from Kōfu, and 10 m. S. from *Kajikazawa*) lies on the E. bank of the *Fuji-kawa*, about 9 m. lower than *Kajikazawa*, at the foot of *Amaga-take*. Near by the spa runs a small ravine, from whose bank the cold springs gush out. The baths are said to have been used by Takeda Shingen for the treatment of wounds. People believe that the water is efficacious especially for bruises, wounds, and itches. Number of visitors: 4,979, the total stays being 26,248 days (1909).

NAGANO=KEN

Suwa (121.7 m. to Kami-Suwa from *Iidamachi*, in 11 hrs. 5 min.; 130.9 m. from *Nagano*; 70 m. from *Nagano*). The two Suwas (*Kami-Suwa* and *Shimo-Suwa*) are almost continuous along the shores of the Lake Suwa, and seem to lie on subterranean beds of hot springs, as hot water issues wherever a bore is made. In the depth of winter the lake is alive with skaters and a grand meet is held at the Suwas. Pop.: 13,405 at Kami-Suwa; 6,630 at Shimo-Suwa. Takashima Castle Grounds in Kami-Suwa mark the site where in the feudal days the Daimyō named Suwa resided; now the grounds constitute a public park.

Lake Suwa-ko is at the greatest elevation of all the large lakes in Japan, it being 2,600 ft. above the sea. In circumference it is about 10 m. and has lofty ridges surrounding it. At its N. W. end is an outlet, which forms the headwaters of the *Tenryū-gawa*. From another corner the lovely shape of *Fuji* is seen far away rising above the shoulder of the nearer ranges. The sight of the lake from the top of *Shiojiri-tōge* is extremely pretty. The lake produces carp, gibel, eel, and other fresh-water fishes.

Geologists say that the lake is gradually silting up, and that the old castle grounds at Kami-Suwa, which are now found at an considerable distance from the shore, were formerly at the edge of the lake. The appearance of cracks on the surface of the frozen lake is generally regarded as a sign that the ice can be walked upon with safety, this phenomenon arising from the fact that the increase of volume of the water when thoroughly frozen causes the central part to rise up and finally to crack. The lake does not appear to be of volcanic origin.

Shrines of Kami-Suwa and Shimo-Suwa. The one at Kami-Suwa is called *Ichino-miya* and is situated in grounds rendered very impressive by patriarchal cryptomerias; it is noted also for its beautiful architecture. The shrine at Shimo-Suwa is somewhat inferior in general appearance to the other. 10 public baths, 27 doctors, 1 hospital (Kami-Suwa). Number of visitors: 64,370, the total

stays being 126,910 days (1909). Inns: *Botan-ya, Nunohan, Suwa Hotel, Kogetsu-kwan.*

4 public baths (Shimo-Suwa). Number of visitors: 9,994, the total stays being 23,601 days (1909). Inns: *Kame-ya, Maru-ya.*

Meteorological Observations.

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation |
|---------------------|-----------------|------|-------|-------|----------|-------|-------------------|----------------------------|----|-------------------------|
| | Mean | | | | Absolute | | | Total | % | Total |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | |
| For the year, 1911. | | | | | | | | | | |
| January | -2.6 | 2.6 | -7.9 | 10.5 | 9.0 | -14.7 | 82 | — | — | 70.8 |
| February | 0.0 | 5.8 | -4.0 | 9.8 | 14.3 | -10.1 | 60 | — | — | 39.5 |
| March | 4.0 | 9.7 | -0.5 | 10.2 | 19.9 | -11.4 | 69 | — | — | 87.5 |
| April | 9.9 | 15.5 | 4.3 | 11.2 | 25.5 | -5.8 | 63 | — | — | 199.2 |
| May | 13.8 | 20.4 | 7.1 | 13.2 | 26.4 | 0.0 | 57 | 243.9 | 56 | 61.8 |
| June | 19.3 | 24.2 | 14.5 | 9.7 | 29.4 | 10.5 | 68 | 174.9 | 41 | 361.2 |
| July | 22.4 | 27.0 | 17.9 | 9.1 | 33.5 | 13.4 | 71 | 179.7 | 41 | 231.3 |
| August | 22.8 | 27.1 | 18.5 | 8.6 | 30.9 | 15.9 | 71 | 191.4 | 46 | 306.7 |
| September | 20.4 | 24.7 | 10.2 | 8.6 | 28.4 | 8.5 | 73 | 161.6 | 43 | 160.3 |
| October | 12.0 | 16.6 | 7.5 | 9.1 | 24.1 | 1.8 | 71 | 159.2 | 46 | 71.4 |
| November | 8.5 | 13.2 | 3.8 | 9.4 | 18.5 | -2.0 | 68 | 177.6 | 58 | 68.6 |
| December | 1.1 | 5.8 | -3.6 | 9.4 | 1.33 | -8.6 | 69 | 167.8 | 56 | — |
| For the year, 1912. | | | | | | | | | | |
| January | -1.8 | 3.3 | -6.8 | 10.1 | 7.4 | -15.6 | 73 | 190.9 | 62 | 50.1 |
| April | 10.3 | 16.0 | 3.9 | 12.7 | 27.9 | -5.4 | 51 | 234.0 | 60 | 123.0 |
| May | 14.3 | 20.1 | 8.6 | 11.4 | 28.4 | 2.5 | 57 | 229.3 | 53 | 91.0 |
| June | 18.2 | 23.4 | 13.0 | 10.4 | 27.3 | 6.9 | 65 | 196.6 | 45 | 110.8 |
| July | 22.3 | 26.7 | 18.0 | 8.7 | 31.9 | 12.6 | 68 | 109.2 | 38 | 235.9 |
| August | 23.8 | 29.3 | 18.3 | 11.0 | 32.5 | 15.0 | 66 | 246.6 | 60 | 119.5 |
| September | 17.9 | 21.8 | 13.9 | 7.8 | 30.5 | 8.0 | 74 | 118.7 | 32 | 241.7 |
| For the year, 1913. | | | | | | | | | | |
| January | -4.8 | 1.9 | -11.6 | 13.5 | 7.2 | -20.6 | 83 | 191.3 | 62 | 26.1 |
| February | -1.8 | 4.2 | -7.7 | 11.9 | 10.4 | -17.4 | 72 | 193.8 | 64 | 38.0 |
| March | 1.6 | 7.6 | -4.3 | 11.9 | 19.8 | -12.0 | 57 | 233.7 | 63 | 43.5 |
| April | 11.0 | 16.7 | 5.3 | 11.4 | 27.2 | -2.0 | 61 | 177.6 | 45 | 107.2 |
| May | 13.2 | 19.3 | 7.3 | 12.1 | 26.0 | 0.3 | 59 | 225.4 | 52 | 162.1 |
| June | 18.0 | 22.9 | 13.0 | 9.9 | 28.1 | 6.3 | 66 | 169.0 | 39 | 167.2 |
| July | 22.5 | 27.9 | 17.1 | 10.8 | 32.6 | 11.6 | 66 | 211.6 | 48 | 129.6 |
| August | 22.5 | 27.8 | 17.1 | 10.7 | 32.1 | 11.2 | 65 | 204.1 | 49 | 117.0 |
| September | 17.3 | 22.3 | 12.4 | 9.9 | 26.2 | 6.1 | 66 | 162.1 | 44 | 73.6 |
| October | 13.0 | 17.8 | 8.2 | 9.6 | 22.7 | 2.3 | 69 | 162.7 | 47 | 109.5 |
| November | 6.1 | 11.6 | 0.7 | 10.9 | 16.9 | -3.9 | 69 | 174.5 | 57 | 99.3 |
| December | 1.5 | 6.3 | -3.2 | 9.5 | 14.1 | -14.1 | 68 | 170.6 | 57 | 51.8 |
| For the year, 1914. | | | | | | | | | | |
| January | 0.1 | 5.2 | -5.0 | 10.2 | 13.5 | -10.0 | 63 | 198.1 | 61 | 32.9 |
| February | -0.6 | 4.8 | -6.0 | 10.8 | 10.1 | -13.4 | 67 | 179.0 | 59 | 45.9 |
| March | 5.1 | 10.3 | 0.0 | 10.3 | 17.3 | -6.4 | 68 | 178.1 | 48 | 113.0 |
| April | 8.4 | 14.8 | 2.0 | 12.8 | 25.5 | -6.7 | 54 | 214.4 | 55 | 48.8 |
| May | 15.2 | 20.7 | 9.7 | 11.0 | 25.8 | 0.3 | 62 | 170.3 | 39 | 229.5 |

Asama (2,4 m. to the N. E. of *Matsumoto*) is a popular resort (1,600 ft. above the sea) for Matsumoto residents. Here hot water issues everywhere, but great inconvenience is occasioned by the scanty of ordinary water. The place is provided with many inns where comfortable accommodation may be obtained. *Hino-ru* and *Nishi Isukawa-ya* are among the first-rate establishments. Pop. 1,600. Number of visitors: 35,795, the total stays being 73,136 days (1909).

Meteorological Observations.

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation |
|---------------------|-----------------|------|-------|-------|----------|-------|-------------------|----------------------------|----|-------------------------|
| | Mean | | | | Absolute | | | Total | % | Total |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | |
| For the year, 1911. | | | | | | | | | | |
| January | -2.3 | 2.5 | -7.1 | 9.6 | 8.7 | -13.5 | 77 | 140.0 | — | 54.5 |
| February | 1.1 | 6.6 | -4.4 | 11.1 | 15.5 | -10.5 | 73 | 179.8 | 59 | 31.4 |
| March | 4.8 | 10.4 | -0.8 | 11.2 | 20.5 | 13.2 | 67 | 172.2 | 44 | 80.4 |
| April | 9.0 | 16.3 | 3.5 | 12.8 | 25.0 | -5.5 | 62 | 195.5 | 50 | 172.6 |
| May | 14.1 | 21.7 | 6.5 | 15.3 | 27.0 | -2.0 | 59 | 253.8 | 58 | — |
| June | 19.8 | 26.0 | 13.7 | 12.3 | 31.0 | 9.6 | 70 | 170.0 | 39 | 345.4 |
| July | 22.9 | 28.1 | 17.7 | 10.4 | 33.9 | 13.0 | 70 | 178.8 | 40 | 243.9 |
| August | 23.6 | 28.8 | 18.5 | 10.3 | 31.8 | 15.8 | 74 | 195.1 | 47 | 174.0 |
| September | 21.3 | 26.5 | 16.1 | 10.4 | 31.5 | 8.6 | 74 | 192.3 | 44 | 123.1 |
| October | 12.4 | 17.6 | 7.2 | 10.4 | 25.5 | 1.5 | 60 | 150.7 | 43 | 71.1 |
| November | 9.1 | 14.0 | 4.1 | 9.9 | 20.4 | -3.4 | 64 | 153.1 | 50 | 68.1 |
| December | 1.4 | 6.2 | -3.4 | 9.6 | 12.9 | -8.8 | 77 | 142.9 | 48 | 31.0 |
| For the year, 1912. | | | | | | | | | | |
| January | -1.9 | 3.3 | -7.0 | 10.4 | 8.5 | -15.9 | 61 | 157.1 | 51 | 33.0 |
| April | 11.0 | 17.9 | 4.1 | 13.8 | 29.0 | -3.4 | 57 | 227.8 | 58 | 69.6 |
| May | 14.5 | 21.3 | 7.8 | 13.5 | 29.5 | 1.0 | 62 | 230.2 | 53 | 72.3 |
| June | 11.5 | 24.5 | 13.0 | 11.5 | 29.0 | 4.7 | 67 | 178.1 | 41 | 138.5 |
| July | 23.3 | 28.1 | 18.5 | 9.6 | 32.6 | 12.4 | 68 | 170.0 | 38 | 142.1 |
| August | 24.2 | 29.8 | 18.5 | 11.2 | 32.6 | 14.0 | 77 | 223.4 | 54 | 160.6 |
| September | — | — | — | — | — | — | — | 103.2 | 28 | — |
| For the year, 1913. | | | | | | | | | | |
| January | -4.9 | 1.0 | -10.9 | 11.9 | 5.9 | -18.7 | 74 | 146.8 | 47 | 40.3 |
| February | -1.2 | 4.4 | -6.9 | 11.3 | 13.4 | -17.8 | 64 | 156.6 | 51 | 32.0 |
| March | 2.0 | 8.4 | -4.4 | 12.7 | 21.4 | -12.3 | 51 | 196.5 | 53 | 15.8 |
| April | 10.9 | 18.0 | 3.8 | 14.1 | 28.5 | -9.9 | 58 | 150.1 | 40 | 101.5 |
| May | 13.7 | 20.6 | 6.8 | 13.7 | 26.6 | 0.1 | 59 | 206.3 | 47 | 121.4 |
| June | 18.7 | 23.9 | 13.4 | 10.5 | 28.5 | 6.1 | 63 | 161.6 | 37 | 167.2 |
| July | 23.0 | 28.5 | 17.5 | 11.0 | 33.0 | 11.9 | 63 | 192.8 | 47 | 97.7 |
| August | 22.9 | 28.2 | 17.5 | 10.7 | 33.4 | 11.9 | 67 | 187.8 | 45 | 147.7 |
| September | 17.4 | 22.7 | 12.2 | 10.5 | 26.2 | 4.5 | 70 | 165.8 | 45 | 52.4 |
| October | 13.0 | 18.3 | 7.9 | 10.7 | 22.1 | -1.0 | 79 | 147.5 | 48 | 97.6 |
| November | 6.3 | 12.1 | 0.5 | 11.6 | 18.9 | -4.6 | 78 | 156.3 | 51 | 71.8 |
| December | 1.9 | 6.3 | 2.5 | 8.8 | 15.1 | -17.8 | 70 | 151.8 | 59 | 35.4 |
| For the year, 1914. | | | | | | | | | | |
| January | 0.3 | 5.9 | -5.2 | 11.0 | 13.5 | -13.8 | 70 | 156.9 | 51 | 18.1 |
| February | -0.7 | 4.9 | 6.3 | 11.3 | 11.0 | -15.6 | 72 | 100.1 | 53 | 24.1 |
| March | 5.8 | 11.5 | 0.2 | 11.3 | 17.9 | -6.3 | 76 | 172.9 | 58 | 60.7 |
| April | 8.6 | 15.9 | 1.4 | 14.5 | 26.0 | -7.1 | 52 | 198.8 | 51 | 30.1 |
| May | 15.5 | 22.1 | 8.9 | 13.2 | 28.0 | 1.5 | 66 | 156.1 | 30 | 100.0 |

Yamabe (2.4 m. E. of the *Matsumoto* station) offers only poor accommodation. 2,000 ft. above the sea. Pop. 300. Number of visitors: 9,119, the total stays being 34,057 days (1909). Inns: *Azumi-ya*, *Marunaka-ryokwan*.

Tenryū-kyō. As the shortest natural route connecting *Nakasen-dō* region with the *Tōkai-dō*, the descent down the rapids of the *Tenryū-gawa* formerly played an important part in the economic relations and transportation facilities between these two highways in the middle part of the main island of Japan Proper. For one who starts from either from *Tōkyō* or *Yokohama* and has to return to the same place after enjoying the adventure, four days will be required. The best plan is to stop overnight at either *Kami-Suwa* or *Shimo-Suwa*; to halt at either *Jida* or *Tokimata* on the 2nd day; the 3rd day is devoted to the passage down from *Tokimata*, the night being spent at *Hamamatsu*, on the *Tōkai-dō* Line; and on the 4th day one returns to one's starting-point.

From *Tokimata* to *Mitsujima*, 19.5 m. in about 4 hrs. In about 30 min. from *Tokimata* we come to the picturesque gorge called *Tenryū-kyō*, which marks the entrance to the romantic rapids and races that are to follow.

Nakabusa. There are two routes to reach the mountain spa *Nakabusa*:

(a) *Hotaka-guchi.* From *Akashina* station, crossing the *Kawate-Kaidō*, we reach *Hotaka-machi*, then across the *Karasu-gawa*, come to *Nishi-Hotaka-mura*, and going over the Pass *O-tōge* and up the stream *Nakabusa-gawa*, reach the place, the whole distance being over 15 m. From *Akashina* 2.5 m. to *Hotaka*, *jirikisha* scarcely available, thence 3.7 m. only on foot.

(b) *Ariake-guchi.* Going from *Akashina* 2.5 m. N. along the *Kawate-Kaidō*, we come to *Nanaki-mura*, then turning left towards the S. foot of *Ariake-san*, reach *Furu-umaya* of *Ariake-mura* (2.5 m.), *jirikisha* available so far (fare, 55 *sen*); from *Furu-umaya*, passing through the plain at the foot of *Ariake-san*, *Miyagi* is reached, where porters for baggages and guides can be hired. From *Miyagi* we go 5 m. up the stream to *Shinano-zaka* by a road maintained by the Forestry Department (*Nakabusa-Kindō*), and cross the gorge and over the peak, around the middle slope of *Ariake-san*, the ascent being very steep. The road coming from *Hotaka* meets that coming from *Ariake-guchi* at this point. Ascending 2.5 m. farther from *Shinano-zaka*, we reach at last the bathing-place of *Nakabusa*. Fee for carrying baggage from *Miyagi* to *Nakabusa*: 10 *sen* per *kwanme* (8.3 pounds) as a rule; for fragile articles 50 % more, and for articles longer than 3 ft. and wider than 2 ft. double the ordinary rate. Guide 50 *sen*; horse ¥ 1.50; *kago* ¥ 2.50.

Nakabusa lies at the middle of the slope of *Ariake-san*, at the foot of the *Hakuba Range*, the place being the first step for those who want to climb the Japanese Alps. The spa is surrounded on every side by lofty mountains. The eminent peaks, such as *Hakuba-san*, *Ō-rence*, *Yari-ga-take*, *Shakushi-dake* and *Eboshi-dake*, rise high in the sky far beyond, while the stream *Nakabusa-gawa* with clear water runs near the spa. A dense forest with big and old trees is in its neighbourhood. The place is 5,300 ft. above the sea-level. Everywhere within about 0.3 m. in the gorge issue hot springs, the flow of water being abundant. There are three bath-tanks, of which the one called '*Kashikiri-yū*' is for reservation. Besides the usual baths, special arrangements for hot water massage (*Takino-yū*) and for vapour bath, as well as for swimming, are provided outside. The water is alkaline, containing hydrogen sulphide, iron, alumina, etc., and claims to be efficacious for dyspepsia, disorders of the brain, wounds, syphilis, rheumatism, *kakke*, disorders of the female genital organs, etc. The discovery of the springs dates back to the Era of Enryaku (782-806). Number of visitors: 22,000 (1912).

Shirahone. The spa is situated at the N. E. foot of *Norikura-ga-take*, which rises on the boundary of the two provinces, *Shinano* and *Hida*. *Yake-ga-take* stands to the N. of the spa by which a small stream runs; on the opposite side of the stream there is a high and steep bank, forming a precipice hanging over the water. The springs are said to be efficacious for diseases of the stomach.

Norikura-ga-take may be best climbed from the spa (in 6-7 hrs.).

Number of visitors: 3,000 (natives) and 50 foreigners (1912).

Kamikōchi. 27 m. W. from *Matsumoto*; from *Matsumoto* to *Shimajima* in *Azumi-mura*, 11 m., *jirikisha* available (fare 66 *sen*), thence on foot. The spa lies on a flat plateau enclosed by high mountains, at an elevation of 4,720 ft. above the sea. *Kasumi-dake* rises to the S. E., *Yari-ga-take* to the N. W., *Hotaka-ga-take* to the N. W., and *Yake-ga-take* to the S. of the spa.

The two mountains, Yari and Yake, can be climbed from this place. *Shirahone* hot springs are 12 m. N. from here. The air being bracing and the heat moderate in summer, it is fast growing popular as a summer resort and has begun to be noticed by foreigners. At present there is only one inn in the spa. Number of visitors: 397, the total stays being 786 days (1909).

Kuzu. 21 m. N. W. from *Akashina* station, *jūrikishū* available throughout. The spa lies at the foot of the mountains *Takinosawa-dake* and *Nanakura-ga-take*, on the bank of the upper course of the *Takase-gawa*, at an elevation of 2,847 ft. above the sea. The three mountain lakes, *Kizaki*, *Naka-ami*, and *Aoki*, are about 7 m. N. E. There are 6 hot springs now in use and only one inn in the spa. Number of visitors: 2,417, the total stays being 10,585 days (1909).

Shino-noi Line. This Line leads from *Shiwojiri* and connects with the Shin-Etsu Line at *Shino-noi*, the total extension being 42.1 m. via this Line, through trains run between Tōkyō (*Ida-machi*) and *Nagano* on the one hand and between *Nagoya* and *Nagano* on the other, the former occupying 13 hrs. 41 min. and the latter 10 hrs. 52 min. The presence of the mighty range recently called the 'Northern Alps of Japan' is prominent feature of the district traversed by this line.

Japan Alps.

This is the title of a volcanic range extending through the central part of *Honshū* or the Main Island of Japan, and forming the boundary between Suruga, Kai, Tōtōmi, Shinano, Hida, Etchū, and Echigo. It consists of three lofty ridges called respectively Southern, Middle, and Northern.

(1) The 'Southern Alps of Japan' are made up of the Akaishi Ridge, which, rising in the Atsumi Peninsula in Mikawa, passes to the N. of Hamana-ko Lagoon, winds between the valleys of the rivers *Kamanashi* and *Tenryū*, and ends at the shore of Lake Suwa-ko. This range furnishes a good example of upheavals caused by the sliding of strata. Of the spurs or clusters of peaks in this range, those that attain an elevation of approximately 10,200 ft. above the sea are *Akaishi* (10,145 ft.), *Shirane-san* (loftiest point 10,332 ft. and including *Kita-dake*, *Aino-take*, *Notori-san*), and *Koma-ga-take* of Kai Province, in which *Jisō*, *Hō-dō*, and *Koma-ga-take* itself all rise about 10,000 ft. above the sea.

(2) The 'Middle Alps of Japan' are constituted by the Kiso Range, which rises in the Chita Peninsula in Owari, enters Shinano, where it sends up *Ena* and *Koma-ga-take* and forms a watershed for the Kiso and the Tenryū, finally terminating at the S. end of the Matsumoto Plain. The cluster of peaks with *Koma-ga-take* of Shinano as their monarch forms the highest elevation in this chain.

(3) The 'Northern Alps of Japan' is the title applied to the Hida Range, which forms the majestic chain that extends from N. to S. on the E. boundary separating Hida and Etchū from Shinano. This comprises *On-take* 10,447 ft., *Norikura* 10,142 ft., *Jōnen-take* and *Otenjō-dake* 10,447 ft., and *Shiratori-dake*. Other noted peaks slightly lower than the above are *Tsubame-dake*, *Kasa-ga-take*, *Goroku-take*, *Washiha-dake*, and *Tsurugi-ga-dake*. Running further N. it sends up *Tate-yama*, and finally descends into the Japan Sea in the steep precipices of *Oyashirazu*. The range extends for a distance of 98 m. with a breadth of 37 m.

Prof. Shiga, in his 'Essay on Japanese Scenery,' writes that the 'Japan Alps' are a medley of volcanic upheavals that have burst through granite strata form the wildest and grandest solitude of rocky pinnacles and wooded peaks in Japan. It is such a solitude, that in summer at some hot springs human faces are rarely seen for weeks at a time, the only living creatures that meet the eyes of solitary travellers or mountaineers in this wild tract being the golden-coloured eagles that soar above the mountains, and pretty deers that stare stolidly at the strange figures. A peculiar species of swallow, haunting cascades or falls, and sluggish salamanders, hiding in mountain creeks, characterize this region as compared with the original Alps. Then in their comparatively primitive aspect the Japanese Alps compare favourably with their somewhat vulgarized prototype, at the Rev. Walter Weston, who is probably the highest authority on the 'Alps of Japan,' writes in his excellent work on this particular subject.

Those in Tōkyō or Yokohama who wish to explore this region should take the first train at Ueno and stop a night at *Matsumoto*. The next day *Norikura-ga-dake* is climbed, then we descend to *Shirahone* Hot Springs, and, after attacking the *Abō-tōge* (6,396 ft. above the sea) on the boundary between Shinano and Hida, *Iwō-ga-take*, *Hotaka-ga-take*, *Yari-ga-take* (11,578 ft.) and *Kasa-ga-take* are climbed in succession; to be followed by the ascent of the peaks on the boundary between Shinano and Etchū, such as *Shishi-ga-take*, *Harinoki-tōge* (8,200 ft.), *Daikoku-dake*, *Yari-ga-take*, and so forth. *Shirouma-dake* and *Dairenge-san* (9,610 ft.), between Etchū and Echigo, are next

attempted, and after a rest at **Renge Hot Springs**, 5,445 ft., where a single inn stands, we descend to *Kotaki-mura* on the bank of the *Itoi-gawa* and finally emerge on the town of the same name after a walk of 11 m. Thence we return to our starting-point by train via *Naoetsu*.

For those starting from either Ōsaka or Kōbe the first objective is *Takayama*, the principal city in Hida. Then proceeding along the road for Matsumoto, the first real ascent is tried at *Norikura-ga-take*, after which, following the itinerary mentioned above, one enters Etchū via *Harinoki-tōge*, to end the adventure with the ascent of the Tateyama Range. Tateyama commands one of the grandest and most magnificent views in Japan, and, though inferior in height, is said to surpass the European Alps in so far as regards the number of lofty peaks that can be descried from the top. Descending to *Ashikura-ji*, one then proceeds to *Toyama*, not quite 17 m. distant, and then back home by train. The expedition requires about one month, whether one starts from Tōkyō or Ōsaka.

Ascent of Norikura-ga-dake.

Norikura-ga-dake stands over 10,400 ft. high above the sea and is the sixth highest peak in Japan Proper. There are six ascents, of which, two are from Shinano, viz. from *Ōnogawa-mura* and **Shirahone Hot Springs**, and the other four from Hida, viz. from *Nonugi*, *Aoya*, *Iwaidani*, and *Hirayu Hot Springs*.

(1) From **Shirahone Hot Springs** the path leads in about 5 m. to the cultivation plot at the headwaters of the *Ōno-gawa*, and 1.4 m. farther on to *Kanayama-taira*, where galena ores are said to have been extensively mined by *Takeda Shingen* (1521-73), and the working of which is still being carried on. Ascending the slope, a little further on the path comes to a small stream strongly smelling of sulphuretted hydrogen, issuing from a crevice a little higher up. This place is 8,500 ft. above the sea. We soon enter the crawling pine zone, with rhododendron shrubs flowering here and there. At an altitude of 10,000 ft. the path leads over hardened lava and scoriae, making the climbing very laborious, and soon the summit is reached. Near it is an old crater 984 ft. in diameter, the wall of which rises steeply at its E. side, but is much lower on the N. N. W. On the elevated side a small shrine dedicated to *Asahi-Gongen* and also a stone post for triangular survey are found. No less than 47 small tarns also exist about this place, most of them dry. From the cultivated plot to the summit the distance is about 7 m.

(2) From **Hirayu Hot Springs** the path leads in about half a mile to the smelting works, then to a magnificent cascade 600 ft. in height tumbling down rocky sides. The path makes plenty of ups and downs, leads past the mine office, through a pine wood, and after an hour's climb over places strangely devoid of rocks and water, enters the crawling pine zone. Passing over *Yutsu-ga-take* which forms an ante-peak, as it were, to Norikura, we find *Ebiko-dake* rising on the right. Going up still higher we find, when we reach Lake Ōnifu, that *Ebiko* is now below us, and from three sides of the lake rugged peaks shoot up. The water is yellowish and the whole scenery strikes one as uncanny. Then begins a scramble over sharp slopes, the path passing along a narrow ridge like a horse's back, while now and then even such a path is absent. The crawling pines become scarce, but alpine plants take their place. Past *Tsuru-ga-ike* tarn and in sight of another tarn called *Kame-ga-ike*, an altitude of over 10,000 ft. is registered. The upward journey over volcanic rocks is a tedious process, the path making ups and downs. No. 5 tarn is seen to the left. Then there is another high elevation to be attacked, the path to it being strewn with large lava blocks, and finally we reach the summit of Norikura-ga-dake. The twin peaks bound in a lake, the largest tarn existing at such an altitude in Japan, for in diameter it measures 1,000 ft. On the right peak stands the Shintō shrine of *Asahi-jinja*, while the left one is sacred in honour of *Ontake-jinja*, and it is this latter peak that forms the highest point in Norikura-ga-dake, its summit, according to the latest calculation, rising 10,630 ft. above the sea. The views obtained from this vantage point vie in grandeur with those from the top of On-take.

Ascent of Yari-ga-take.

Yari-ga-take ('Spear peak') is, next to Fuji, the highest mountain in Japan, being 11,600 ft. above the sea. The following description of the ascent of this peak is based on the '*Yari-ga-take Tankenki*' (Exploration of Yari-ga-take) of Mr. *Kojima Ōsui*, who is an acknowledged Japanese authority on mountaineering.

At Shimajima Village a guide is hired. Four miles on there is a mineral spring called *Furohira*, and thence the track leads, over the *Tokumoto-tōge* (7,100 ft.) between *Nabekaburi-yama* and *Kasumi-dake*. The path then makes a descent and leads to a hut in a dense forest on the banks of the *Azusa-gawa*, 7 hrs. from the starting point. In the river trout abounds, and beyond towers up the sublime granite ridge of *Hotaka* resembling in its situation and outline a mountain near Chamounir in the European Alps. From the hut to the top 9 to 10 hrs. are taken; then down to the hunter's cave at *Akasaka* (6,400 ft.) 2 hrs. 30 min. are required even by the hardest climbers. From the cave the path is led along a mountain stream which has to be forded here and there, and in 3 hrs. the snow-streaked granite giant of *Hotaka* rises on the left, while on the right stand well-wooded and milder ranges. The peaks rising about are all of granite and in their spiry shapes remind the beholder of the wild mountain landscapes that are found in Chinese paintings. Nowhere in Japan, except at remote places in *Yamato*, can such primitive grandeur and sublime solitude be found as in this neighbourhood. From the hunter's cave the forest zone is left, and the path begins to ascend over snow-fields. Continuing the upward journey, the presence of huge boulders indicates the nearer approach to the summit. Some are piled up, others threaten to tumble down, while here and there they form natural arches, and stepping from one stone to another, or clambering over boulders or across snow-fields, and sometimes crawling over risky spots the summit is reached at last. At this spot a single rocky pinnacle is seen to shoot up perpendicularly, its sides falling away sheer all round like a spear-point, the S. E. corner alone exhibiting a slight slope. On the descent a sturdy man can reach the hunter's cave in 2 hrs. 30 min., hence 12 hrs. 30 min. to Shimajima.

The views from the summit embrace to the N. the innumerable peaks, many of them nameless, of Shinano and Etchū; to the W., *Kasa-ga-take*; to the S., the ridge of *Hotaka* and *Norikura*, and beyond, the peaks of *On-take* and to its E. *Koma-ga-take* of Shinano; and to the S. W., the lofty range forming the boundary between Shinano and Kai, and far away, for the distance is 85 m. as the crow flies, Fuji lifts up her noble head. The view of Japan Sea to the N. W. is generally obscured owing to mist.

Ascent of Hotaka-ga-take. This is also called *Myōjin-dake* and ranks third on the list of high mountains, it standing 11,500 ft. above the sea. The ascent is made from *Higashi-Hotaka-mura*, 6 m. to the S. W. of the station, *Tokugō* being the usual starting-point. From the village to the summit it is an ascent of a little over 17 m., and though involving hard strain the journey can be accomplished in one day, viz. 6 hrs. in ascent and 5 in descent. Hotaka stands before us soon after the start as a granite cliff shooting up 5,000 ft. from the ravine of the *Azusa-gawa*. The trouble with this mountain is that no track has yet been made to facilitate the ascent, so that the climber has to force his way through tall bamboo-grass and to scramble up rocks by dint of holding on to shrubs or creepers. At 8,500 ft. above the sea there is a snow-covered slope making an angle of about 40°, and, when this is safely climbed, large masses of smooth rock bar the passage. Scrambling over it we at length reach the ridge from which rise several peaks, of which the one on the extreme left is the highest, with a summit consisting of broken blocks of granitic rock. The panorama viewed from here is practically identical with that seen from *Yari-ga-take*.

Nakabusa Hot Springs lie to the W. of Hotaka and is 14.5 m. distant from Matsumoto. The spa occupies a solitary spot surrounded by high mountains and deep valleys and is resorted to by students who are anxious to utilise their summer holidays to the best advantage for the body and the mind.

Ascent of Ariake-yama, 8,075 ft. is best made from the hot springs, the ascent being divided into 9 stages, each covering about 0.5 to 0.6 m. and indicated by a post. At the 6th stage the twin mountains of *Chō-ga-take* and *Amakazari* rise, the S. one connected with *Kasumisawa-san* and the N. one with *Ariake*. At the 7th stage the road becomes steeper and here the *Ōjigoku-dani* ('Big Hell Valley') is found. At this spot one part of the road has been washed away by heavy rain, leaving a huge rock overhanging above, while a dense mist always obscures the abyss. From the opening at the 8th stage, the town of *Ōmachi*, the two lakes of *Kozaki* and *Nakatsuna*, and then the noble ridge of *Iida* are seen. At the top stands a *torii*, a little beyond which we come to the small Shintō shrine of *Ariake-jinja*. The panorama of the Matsumoto Plain unfolded below and of the rivers and surrounding mountains well rewards one for the toil of the ascent.

Hirao. There are five hot springs noted for their efficacy in *Hirao-mura*, viz. *Kambayashi*, *Kutsuno*, *Shibu*, *Andai* and *Yudanaka*. These springs, not distant from one another, can be easily reached from the station *Toyono* (141.5 m. from Ueno, in 7 hrs. 35 min.), the distance being 10 m. Automobiles, omnibuses, and *jinrikisha* are available. Post, telephone and telegraph office. The five springs are situated on the bank of the *Yomase-gawa*, *Kambayashi* lying on the uppermost portion and *Yudanaka* on the lowest.

| | Kambayashi | Kutsuno | Shibu | Andai | Yudanaka |
|----------------------------|------------|---------|-----------------------|--|--|
| Population | 9 | 907 | 915 | 200 | 1,900 |
| Houses | 3 | 194 | 200 | 50 | 300 |
| Inns | 1 | 2 | 13 | 7 | 10 |
| Springs | — | — | common salt | simple | bitter |
| Diseases that may be cured | — | — | rheumatism, paralysis | disorders of the brain, diseases of the stomach and intestines | venereal diseases, disorders of the female genital organs, dyspepsia |
| Private baths | 2 | 1 | 8 | 3 | 2 |
| Public baths | 1 | 3 | 7 | 3 | 10 |
| Doctors | 0 | 0 | 1 | 0 | 2 |
| No. of visitors | 222 | 297 | 10,063 | 4,601 | 7,340 |
| Total stays (in days) | 723 | 509 | 52,984 | 18,155 | 21,707 |

Meteorological Observations. Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation |
|---------------------|-----------------|------|------|-------|----------|-------|-------------------|----------------------------|-------|-------------------------|
| | Mean | | | | Absolute | | | Total | Total | |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | |
| For the year, 1911. | | | | | | | | | | |
| January | -2.4 | 1.9 | -6.6 | 8.5 | 9.3 | -12.3 | 70 | — | — | — |
| February | 0.1 | 4.0 | -3.8 | 7.8 | 10.6 | -10.3 | 71 | — | — | 41.1 |
| March | 3.5 | 7.8 | -0.9 | 8.7 | 15.9 | -9.3 | 74 | — | — | 69.0 |
| April | 9.3 | 14.5 | 4.1 | 10.4 | 27.0 | -6.6 | 68 | 187.8 | 48 | 120.6 |
| May | 13.9 | 20.0 | 7.8 | 12.2 | 25.3 | 0.1 | 59 | 263.4 | 60 | 49.1 |
| June | 18.6 | 23.3 | 14.0 | 9.3 | 28.1 | 9.0 | 70 | 206.8 | 47 | 252.5 |
| July | 22.2 | 26.5 | 17.9 | 8.6 | 31.7 | 12.2 | 74 | 197.3 | 44 | 183.1 |
| August | 23.3 | 28.1 | 18.6 | 9.5 | 31.7 | 16.0 | 73 | 214.1 | 51 | 224.1 |
| September | 20.6 | 25.6 | 15.6 | 10.0 | 31.8 | 0.6 | 73 | 180.9 | 49 | 130.1 |
| October | 12.5 | 16.9 | 8.1 | 8.7 | 26.0 | 2.4 | 71 | 154.4 | 44 | 90.4 |
| November | 9.0 | 13.7 | 4.4 | 9.3 | 20.6 | -1.6 | 70 | 164.8 | 54 | 76.0 |
| December | 1.3 | 5.2 | -2.5 | 7.6 | 14.1 | -8.1 | 77 | 155.8 | 52 | 41.7 |

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation |
|---------------------|-----------------|------|------|-------|----------|-------|-------------------|----------------------------|----|-------------------------|
| | Mean | | | | Absolute | | | Total | % | Total |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | |
| For the year, 1912. | | | | | | | | | | |
| January | 1.9 | 1.9 | -5.8 | 7.7 | 5.9 | -11.3 | 70 | 132.6 | 43 | 70.3 |
| April | 9.4 | 15.0 | 3.8 | 11.2 | 25.0 | -6.1 | 59 | 234.5 | 60 | 44.3 |
| May | 13.4 | 18.7 | 8.1 | 10.6 | 28.5 | 3.0 | 65 | 224.1 | 51 | 93.3 |
| June | 17.4 | 21.8 | 13.0 | 8.8 | 27.0 | 6.5 | 69 | 212.3 | 48 | 82.0 |
| July | 21.5 | 25.6 | 17.5 | 8.0 | 30.5 | 11.0 | 74 | 205.3 | 46 | 256.6 |
| August | 23.4 | 28.2 | 18.6 | 9.5 | 31.4 | 14.0 | 74 | 270.8 | 65 | 155.9 |
| September | 17.4 | 21.5 | 13.3 | 8.2 | 28.1 | 7.0 | 78 | 136.2 | 37 | 154.9 |
| For the year, 1913. | | | | | | | | | | |
| January | -3.5 | 0.7 | -7.7 | 8.5 | 5.9 | -12.4 | 75 | 127.0 | 41 | 60.0 |
| February | -1.2 | 3.4 | -5.7 | 9.0 | 14.1 | -12.3 | 73 | 165.0 | 54 | 60.4 |
| March | 0.6 | 5.3 | -4.0 | 9.3 | 15.8 | -11.8 | 74 | 218.4 | 59 | 34.9 |
| April | 10.1 | 15.4 | 4.7 | 10.8 | 28.1 | -4.6 | 68 | 214.9 | 55 | 86.8 |
| May | 12.5 | 18.0 | 6.9 | 11.0 | 23.7 | 0.6 | 68 | 224.9 | 51 | 105.0 |
| June | 17.3 | 21.6 | 13.0 | 8.7 | 29.0 | 5.5 | 71 | 202.5 | 46 | 128.9 |
| July | 21.3 | 25.6 | 16.9 | 8.7 | 30.5 | 10.7 | 75 | 204.1 | 46 | 82.2 |
| August | 21.3 | 26.0 | 16.7 | 9.2 | 31.0 | 13.0 | 72 | 230.5 | — | 171.0 |
| September | 16.5 | 21.1 | 11.9 | 9.2 | 25.1 | 5.2 | 74 | 175.9 | 47 | 33.1 |
| November | 5.9 | 10.0 | 1.7 | 8.3 | 18.4 | -1.7 | 75 | 136.6 | 45 | 103.4 |
| December | 1.4 | 5.2 | -2.5 | 7.7 | 13.8 | -10.9 | 76 | 150.9 | 52 | 59.8 |
| For the year, 1914. | | | | | | | | | | |
| January | 0.1 | 4.6 | -4.5 | 9.1 | 13.3 | -10.4 | 75 | 168.1 | 55 | 39.4 |
| February | -1.0 | 2.9 | -4.8 | 7.7 | 8.6 | -10.8 | 75 | 168.8 | 56 | 39.9 |
| March | 4.6 | 9.1 | 0.2 | 8.8 | 19.6 | -5.6 | 72 | 195.0 | 53 | 59.2 |
| April | 7.8 | 13.3 | 2.3 | 11.0 | 24.0 | -5.3 | 68 | 217.9 | 55 | 75.9 |
| May | 15.2 | 20.4 | 9.9 | 10.6 | 29.0 | 0.4 | 70 | 211.3 | 48 | 104.5 |

Kakuma. 1.2 m. S. from Shibu hot springs. Pop. 280. 3 springs, 6 public baths, 6 inns, of which two are provided with private baths. The water is said to be good for *kakke*, brain, and diseases of the skin. Number of visitors: 2,110, the total stays being 8,511 days (1909).

Bessho. 7 m. S.W. from Ueda (113.4 m. from Ueno, in 6 hrs.) on the Shin-etsu Line, *jinrikisha* and omnibuses available. The spa is shut in on the two sides, S. and W., by hills and penetrated by the *Yu-kawa*, hot springs issuing from its S. bank. Pop. 1,396 (294 households), 22 inns, 5 public baths, 5 inns with private baths, and 2 resident doctors in the village. Number of visitors: 18,764, the total stays being 37,875 days (1909).

Meteorological Observations.

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation |
|---------------------|-----------------|------|------|-------|----------|-------|-------------------|----------------------------|----|-------------------------|
| | Mean | | | | Absolute | | | Total | % | Total |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | |
| For the year, 1911. | | | | | | | | | | |
| February | 1,8 | 6,3 | -2,8 | 9,2 | 14,0 | -8,8 | 68 | — | — | 46,7 |
| March | 5,6 | 11,1 | 0,1 | 11,0 | 20,0 | -8,8 | 69 | — | — | 62,0 |
| April | 10,8 | 16,5 | 5,1 | 11,4 | 30,4 | -5,8 | 66 | 185,4 | 47 | 164,2 |
| May | 14,7 | 21,7 | 7,6 | 14,1 | 27,9 | 0,8 | 57 | 103,4 | 44 | 44,4 |
| June | 20,1 | 25,6 | 14,5 | 11,0 | 30,9 | 10,0 | 78 | 117,5 | 27 | 335,0 |
| July | 23,9 | 28,7 | 19,0 | 9,8 | 33,9 | 13,0 | 88 | 118,8 | 27 | 224,4 |
| August | 24,6 | 29,6 | 19,6 | 10,0 | 33,4 | 16,5 | 72 | 164,7 | 39 | 260,1 |
| September | 21,5 | 26,7 | 16,4 | 10,3 | 31,9 | 9,9 | 88 | 113,6 | 31 | 145,3 |
| October | 14,0 | 18,7 | 9,2 | 9,5 | 28,9 | 1,8 | 60 | 144,5 | 42 | 74,3 |
| November | 10,2 | 15,4 | 5,0 | 10,4 | 25,5 | 0,0 | 70 | 109,0 | 36 | 65,9 |
| December | 1,9 | 6,5 | -2,7 | 9,2 | 12,5 | -7,8 | 77 | 91,7 | 31 | 35,2 |
| For the year, 1912. | | | | | | | | | | |
| January | -0,3 | 4,6 | -5,2 | 9,8 | 10,5 | -10,3 | 75 | 138,9 | 45 | 35,7 |
| April | 11,1 | 17,9 | 4,4 | 13,5 | 27,9 | -4,8 | 50 | 239,9 | 61 | 75,2 |
| May | 15,2 | 21,6 | 8,8 | 12,8 | 31,4 | 2,9 | 61 | 212,8 | 49 | 87,1 |
| June | 18,9 | 24,5 | 13,4 | 11,1 | 29,4 | 5,9 | 70 | 163,0 | 37 | 201,8 |
| July | 23,5 | 28,3 | 18,8 | 9,6 | 34,1 | 12,5 | 70 | 144,7 | 33 | 192,0 |
| August | 24,9 | 30,7 | 19,1 | 11,5 | 33,4 | 15,0 | 69 | 236,5 | 57 | 82,7 |
| September | 18,9 | 23,3 | 14,5 | 8,8 | 29,9 | 9,5 | 79 | 92,0 | 30 | 182,8 |
| For the year, 1913. | | | | | | | | | | |
| January | -3,3 | 1,6 | -8,2 | 9,8 | 5,4 | -13,3 | 73 | 152,1 | 49 | 37,1 |
| February | -0,5 | 4,5 | -5,4 | 10,0 | 15,0 | -11,2 | 68 | 102,3 | 53 | 37,8 |
| March | 2,3 | 8,2 | -3,7 | 11,9 | 19,5 | -11,2 | 61 | 206,3 | 56 | 37,1 |
| April | 11,6 | 18,0 | 5,1 | 12,9 | 28,9 | -2,2 | 70 | 193,2 | 49 | 108,2 |
| May | 13,6 | 20,0 | 7,1 | 12,9 | 27,0 | 0,8 | 65 | 235,1 | 54 | 111,3 |
| June | 17,8 | 22,7 | 12,8 | 9,9 | 27,9 | 7,4 | 67 | 161,9 | 37 | 158,5 |
| July | 22,2 | 27,2 | 17,3 | 9,9 | 31,9 | 11,5 | 71 | 187,0 | 42 | 66,2 |
| August | 21,9 | 26,6 | 17,2 | 9,4 | 30,9 | 12,5 | 73 | 171,9 | 41 | 198,0 |
| September | 17,0 | 21,7 | 12,2 | 9,5 | 27,0 | 5,4 | 75 | 154,9 | 42 | 41,2 |
| October | 12,8 | 17,6 | 8,0 | 9,6 | 21,0 | 0,5 | 77 | 125,9 | 30 | |
| November | 6,1 | 11,5 | 0,7 | 10,8 | 18,2 | -3,8 | 71 | 154,1 | 50 | 85,5 |
| December | 1,7 | 6,6 | -3,2 | 9,8 | 13,3 | -10,0 | 72 | 161,3 | 54 | 44,0 |
| For the year, 1914. | | | | | | | | | | |
| January | 0,9 | 6,8 | -4,9 | 11,7 | 13,9 | -10,2 | 68 | 178,8 | 58 | 21,0 |
| February | -0,6 | 4,4 | -5,6 | 10,0 | 11,4 | -13,2 | 70 | 155,3 | 51 | 32,0 |
| March | 5,8 | 10,9 | 0,6 | 10,2 | 19,2 | -6,4 | 69 | 164,3 | 44 | 84,4 |
| April | 8,4 | 14,7 | 2,1 | 12,6 | 26,1 | -3,7 | 55 | 196,6 | 50 | 50,6 |
| May | 15,1 | 21,0 | 9,1 | 11,9 | 29,2 | -0,1 | 63 | 174,7 | 40 | 135,1 |

Nozawa, 24 m. N. E. from *Toyono* station, *jirikisha* available, lies on the right bank of the *Chikuma-gawa*, with hills on the N., S. and E., at an elevation of 993 ft. above the sea. The springs were found in the Tenryaku Era (947-57). Pop. 1,659, 7 public baths, 12 inns, 1 doctor. The spa may become a lovely summer retreat, the climate being moderate during the summer months. Number of visitors: 16,055, the total stays being 45,590 days (1909).

Meteorological Observations.

Summary.

| Month | Air temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation |
|---------------------|-----------------|------|------|-------|----------|-------|-------------------|----------------------------|----|-------------------------|
| | Mean | | | | Absolute | | | Total | % | Total |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | |
| For the year, 1913. | | | | | | | | | | |
| April | 8.9 | 14.7 | 3.1 | 11.6 | 25.7 | -3.7 | 70 | 183.9 | 47 | 75.7 |
| May | 12.5 | 18.0 | 7.0 | 11.0 | 25.4 | 0.3 | 65 | 199.2 | 46 | 110.7 |
| June | 17.3 | 22.1 | 12.5 | 9.6 | 28.0 | 4.4 | 73 | 157.6 | 36 | 111.8 |
| July | 21.2 | 26.0 | 16.5 | 9.5 | 31.5 | 9.8 | 75 | 148.8 | 33 | 96.0 |
| August | 21.5 | 26.4 | 16.6 | 9.8 | 29.6 | 12.7 | 71 | 184.8 | 44 | 262.4 |
| September | 16.6 | 21.3 | 11.9 | 9.4 | 25.1 | 5.2 | 70 | 128.1 | 34 | 39.3 |
| October | 12.5 | 17.0 | 7.9 | 9.1 | 23.3 | 1.4 | 76 | 113.4 | 33 | 129.4 |
| November | 5.7 | 9.7 | 1.7 | 7.9 | 17.0 | -1.4 | 80 | 110.3 | 36 | 185.9 |
| December | 1.5 | 5.4 | -2.3 | 7.8 | 14.5 | -9.2 | 75 | 87.8 | 31 | 164.5 |
| For the year, 1914. | | | | | | | | | | |
| January | -0.1 | 4.5 | -4.8 | 9.3 | 12.5 | -11.0 | 66 | 117.8 | 38 | 286.0 |
| February | -0.6 | 3.4 | -4.5 | 7.9 | 8.4 | 10.5 | 64 | 105.6 | 35 | 187.7 |
| March | 4.4 | 8.7 | 0.0 | 8.4 | 19.5 | -6.5 | 68 | 147.8 | 40 | 94.8 |
| April | 7.8 | 13.4 | 2.2 | 11.1 | 24.0 | -5.5 | 65 | 151.4 | 38 | 111.2 |
| May | 15.1 | 20.8 | 9.4 | 11.4 | 28.5 | 0.0 | 65 | 169.1 | 39 | 112.7 |

Karuizawa (88.3. m. from Ueno, in 4 hrs. 54 min.) being situated at an elevation of 3,270 ft. above the sea, has recently risen to be a highly popular summer resort, especially for foreigners, as the place is easily accessible from Tōkyō. It extends 4.6 m. from E. to W. and 6 m. from N. to S. and is shut in on three sides—N., S., and E.—by mountains, with an opening only in the S. W. direction. The grassy moors in the S. W., the only open direction, are bounded at their W. extremity by the sharp ridge of *Yakaze*. These moors occupy about two-thirds of the whole area of Karuizawa. The land is generally elevated towards the N. E. and slopes towards the S. W., and the streams, of which the *Kawagoe-ishi-kawa* is relatively the largest, all rise in the N. and flow W. It was by Archdeacon Shaw of the S. P. G. Mission and Prof. Dixon, then of the Imperial Tōkyō University, that Karuizawa as a summer resort was first introduced to the foreign public. They visited the place in 1886, and spent the summer at the houses of certain of the inhabitants. The villa built in 1888 by the Archdeacon on the top of *Ōkatsuka* was the first foreigner's house erected here. Several other foreigners followed his example, until before long Karuizawa, which was otherwise doomed to decay, blossomed forth as a prosperous village. The grateful villagers have erected a monument in memory of the Archdeacon. Visitors to Karuizawa in 1911 numbered 6,597, i. e. 5,406 Japanese and 1,191 foreigners, and the number of days they stayed there aggregated 121,644.

The meteorological records for the years 1911 and 1913 are as follows:

Meteorological Observations.

Summary.

| Month | Air Temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation |
|---------------------|-----------------|------|-------|-------|----------|-------|-------------------|----------------------------|----|-------------------------|
| | Mean | | | | Absolute | | | Total | % | Total |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | |
| For the year, 1911. | | | | | | | | | | |
| January | -2.6 | 3.3 | -8.5 | 11.8 | 10.2 | -13.7 | 65 | — | — | 32.0 |
| February | 1.6 | 8.2 | -4.9 | 13.0 | 14.0 | -12.1 | 59 | — | — | 11.7 |
| March | 4.4 | 11.4 | -2.7 | 14.1 | 17.5 | -11.7 | 61 | — | — | — |
| June | 16.0 | 20.3 | 11.6 | 8.7 | 22.1 | 7.7 | 73 | 67.5 | 15 | 419.8 |
| July | 19.1 | 23.4 | 14.8 | 8.5 | 29.7 | 11.3 | 84 | 124.8 | 28 | 375.2 |
| August | 20.8 | 25.3 | 16.3 | 9.0 | 29.6 | 12.3 | 82 | 116.4 | 28 | 405.1 |
| September | 18.5 | 23.4 | 13.6 | 9.8 | 30.1 | 7.6 | 72 | 118.2 | 32 | 178.3 |
| November | 7.4 | 14.1 | 0.8 | 13.2 | 22.4 | -6.5 | 60 | 108.0 | 35 | 44.8 |
| December | -0.3 | 6.5 | -6.6 | 13.2 | 15.0 | -11.0 | 61 | 146.1 | 49 | 2.7 |
| For the year, 1913. | | | | | | | | | | |
| January | -6.7 | -0.7 | -12.7 | 12.0 | 4.0 | -21.5 | 84 | 180.9 | 59 | 35.2 |
| February | -3.9 | 2.0 | -9.7 | 11.7 | 9.4 | -20.8 | 76 | 197.1 | 65 | 31.5 |
| March | -1.3 | 5.3 | -7.9 | 13.2 | 17.4 | -15.3 | 81 | 198.7 | 54 | 44.7 |
| April | 8.0 | 18.4 | 1.2 | 13.6 | 27.5 | -8.3 | 81 | 170.0 | 43 | 94.9 |
| May | 9.7 | 16.3 | 3.0 | 13.4 | 20.6 | -5.0 | 64 | 196.4 | 45 | 153.6 |
| June | 14.7 | 20.3 | 9.1 | 11.2 | 26.2 | 0.8 | 70 | 129.0 | 30 | 177.3 |
| July | 18.4 | 23.5 | 13.2 | 10.2 | 29.7 | 7.6 | 78 | 147.1 | 33 | 95.1 |
| August | 17.9 | 23.8 | 12.0 | 11.8 | 28.4 | 7.0 | 79 | 198.6 | 48 | 333.2 |
| September | 14.1 | 19.2 | 9.1 | 10.0 | 22.6 | 2.8 | 78 | 155.0 | 45 | 73.2 |

In the summer season Karuizawa is well provided with various devices for social entertainments and sports, such as concerts, theatricals, Karuizawa Athletic Association, Asama Climbing Society, etc., many of these organizations having been originated by foreigners. Horse for hire, the charge for horses is 40 *sen* an hr. without groom, 10 *sen* extra with groom; ¥ 2 for half a day with groom, and ¥ 3 a day with groom.

Karuizawa has many beautiful sights and places for excursion in the neighbourhood, such as the *Usui-tōge*, *Kose Spa*, *Hanare-yama*, *Zekkei-hō*, *Kama-no-hash*i, *Iriyama-tōge*, *Wami-tōge*, *Asama-yama*, *Myōgi-san*, *Nunobiki-yama*, *Akaru-san*, *Oshidashi-izwa*, 'Cathedral Rocks,' etc.

Hotels: *Mampe*i Hotel, 1 m. from the station; Charges, American plan, 1st class ¥ 8, 2nd class ¥ 6, 3rd class ¥ 4; *Karuizawa Hotel*, 0.9 m. from the station; Charges, American style, 1st class ¥ 7, 2nd class ¥ 5, 3rd class ¥ 3; *Mikasa Hotel*, 2 m. from the station; Charges, à la Française, 1st class ¥ 12, 2nd class ¥ 8, 3rd class ¥ 5. There is also a pension called 'Greta Bank' managed by foreigner. Inns: *Tsuru-ya*, *Banshō-ken*, *Fuji-ya*, *Abura-ya*, *Kosei-kan*, *Ichida-ya*, *Ebisu-ya*.

GIFU=KEN

Ena Mineral Springs are located on a highland, more than 1,200 ft. above sea-level, covering *Koaza-Kitsumi* and *Wakayama*, *Oaza-Takayama*, *Fukuoka-mura*, Ena-gōri, Mino Province.

see Plate 39

This highland is bounded by the River *Tsukechi* on the E., the River *Ama* on the W., the River *Kiso* on the S., and Mt. *Tobi-iwa* and *Su* on the N. Overlooking the grand plateau of *Naegi*, the visitor commands a distant, panoramic view of the highest peaks of the 'Japanese Alps,' such as *Ena*, *Kasagoe*, *Koma-ga-dake*, *Mi-take*, *Norikura*, etc. There are countless caves, large and small, clear streams invariably flowing out of them. These are the so-called "Eight Hundred and Eighty Valleys" (a name given by Kōbō-Daishi, the much revered ancient priest). Water of all these streams are strong radio-active. Foot-paths, suitable for sauntering, thread their way beside the streams and also encircle the foot of the highland. Thus the whole district forms a natural park on a large scale.

The district enjoys a mild climate, which does not vary very much; the temperatures ranging from 30° F. in winter to 80° F. in summer.

see Plate 38

The place lies about 5 m. from the *Nakatsu-gawa* station on the Central Railway Line (about 3 hrs. from Nagoya); and there is a good scenic road over which *jirikisha* and *kago* are available. Some of the chief attractions on the way are the Kiso rapids, wonderful rocks on the site of Naegi Castle, etc.

The whole district is of granite formation and produces various kinds of minerals, numbering more than fifty. The more important of them are *naëgite*, *fergusonite*, *zircon*, *bismuth*, *tin sand*, *tungsten*, etc. Besides, there are such precious stones as *rutile* (?), *sapphire*, etc., and one can pick up *morion* and *topaz* anywhere.

These are produced not only at *Takayama* in Fukuoka-mura, but also at *Fukuoka* in Fukuoka-mura, in *Naegi-machi*, and *Hirukawa-mura*, all in Ena-gōri.

Rock-crystal was found in 1873 at *Aza-Wakayama*, *Oaza-Takayama*, *Fukuoka-mura*, and later in 1884 when tin sand was discovered on the same spot, the Mitsui Company organized Tōnō-Saikō-sha (Mining Company in Eastern Mino), which carried out the mining enterprise on a large scale to their immense profit. Afterwards the work was handed over to local capitalists who still run it. At first the tin sand produced here was found to contain various minerals above referred to but these mixtures were discarded as useless and offensive, as they made refining difficult and greatly decreased the percentage of production of the refined mineral, especially so in the case of tungsten, naëgite, etc., forming small particles. But in 1894-5 "*kinkō-seki*" alone was found useful for colouring chinawares and porcelain and has since then maintained some market value. In 1911 the importance of tungsten was first known and the mineral was discovered at Ōaza-Takayama in Fukuoka-mura and also in Hirukawa-mura, where it is still being mined. With the progress of chemistry and various scientific investigations, it became a problem whether this mineral contains radium or not. While the scientists were discussing it in the next year Kanroku Yoshimura, the land-owner of the Yunoshima mineral spring, supposed that this spring contains radium emanation and spread the news. In February, 1913, Mr. Y. Morimoto, Tōkyō, tested the spring and found the rumour true. The land-owner then entrusted Mr. Morimoto with the entire management of this spring, and he immediately applied for analysis to the Tōkyō Imperial Hygienic Laboratory of the Department for Home Affairs. Expert Mr. Kinugasa and Assistant Mr. Nishihara were despatched to the spot and the analysis confirmed the fact that the spring in question contains a large amount of radium emanation. Since then Dr. Ishizu and Dr. Tokunaga and other famous men have visited the place and observed the promising feature of this tract of land.

The district, on the whole, forming a highland, the climate is healthy, air invigorating, and scenery grand. As the thick forests of Kiso are not very far, and better still, the means of access not lacking, it is believed that the district would become a suitable summer sanatorium if proper accommodation were provided for it. At present no bathing arrangements are made.

TOYAMA=KEN

Ogawa. 0.7 m. from *Tomari* station on the Hokuroku Main Line. Springs issue from a hillside on the upper course of the *Ogawa* (7.5 m. S. E. from the *Tomari* station). Formerly there used to be several inns with private baths at the source of springs, but they were completely destroyed by an inundation in 1909, and now the water is conducted by wooden pipes to the town. Number of visitors (before inundation): 5,705, the total stays being 37,756 days (1909).

Futami, Kuronagi, and Kanetsuri. The three hot springs, mentioned above, can be reached from the *Mikkaichi* station on the Hokuroku Main Line. Starting from the station and going 7.5 m. towards the S. E., we come to *Aimoto-bashi*, which spans the *Kurobe-gawa* at the end of the village of *Aimoto-mura*. The bridge was constructed on the cantilever principle and enjoyed formerly the reputation to be one of the three most ingenious bridges in Japan. It is about 240 ft. in length and about 50 ft. above the surface of the water, for the *Kurobe-gawa* runs here between precipitous banks, the colour of water as seen from the bridge being emerald green.

About 7.5 m. up the *Kurobe-gawa* there are *Kuronagi* Hot Springs, at the point where the *Kurobe-gawa* joins its affluent *Kuronagi-gawa*. The place is at an elevation of 640 ft. above the sea-level. 3.8 m. farther from *Kuronagi* up the river *Kurobe-gawa* lie *Kanetsuri* Hot Springs at the foot of *Higashi-Kanetsuri-yama*, which has a shape of temple-bell.

On the upper course of the *Kuronagi-gawa* are situated *Futami* Hot Springs, about 1 m. distant from *Kuronagi*. The springs have their source in a cave between two rocks standing on the river-bed. These two rocks, larger and smaller, are tied together by a large *shime-nawa* (straw rope with tufts of the same material), with a *torii* on one of the rocks. On account of the resemblance of these rocks to the "Wedded Rocks" of *Futami-no-ura* (Ise Province), the springs are called "Futami-Onsen." The spa is resorted to mostly by the inhabitants in the neighbourhood.

| | Kuronagi | Kanetsuri | |
|----------------------------|----------|-----------|--------|
| Number of visitors : | 243 | 1,000 | (1909) |
| The total stays (in days : | 1,291 | 7,000 | (") |

Ōiwayama. 13.7 m. E. from *Toyama*, *jūnikisha* (fare, 66 *sen*) available, and 5 m. S. E. from *Kami-ichi-machi*. The place is situated at the N. extremity of the ridge adjacent to Mt. *Tate-yama*, at an elevation of about 700 ft. above the sea, surrounded on three sides by mountains and open on the N. toward the plain of *Etchū*, commanding an extensive view of the Bay of *Etchū* and *Noto*. On a rocky precipice here stands a famous temple called *Nisseki-jū* (Shingon Sect), and behind the temple are found several waterfalls, of which *Ō-taki*, the largest one, presents a wonderful sight. Close by the waterfall stands a stone statue of *Fudō-son*, which is said to have been sculptured by *Shaku-no-Gyōki* in the 2nd year of Jinki Era (726 A.D.). Even now bathing in the waterfalls is popularly believed to be efficacious for curing mental derangement and also eye diseases, and thus the place is visited by sick people all the year round. The place has a small temple for those bathers coming for recuperation, who shut themselves up therein during certain numbers of days. The place, being located amidst beautiful surroundings, may be said to be one of the most pleasant summer retreats in the province. There are six inns in the place. Number of visitors: 4,734, the total stays being 9,167 days (1909).

Ōmaki. 11.5 m. S. E. from *Fukuno* station on a private railway, "Chū-etsu Line" (*Takaoka*—*Jōhana*). From the station 4 m. E. to *Inami-machi*, *jūnikisha* available. The spa lies on the E. bank of the *Imizu-gawa* (also called *Shō-gawa*) at an altitude of 850 ft. above the sea. The bath-tank is constructed close by the stream and covered with stones. The place is beautifully located amidst densely wooded hills. Number of visitors: 4,296, the total stays being 20,439 days (1909). Farther up the stream is *Soyama-Onsen* and 5 m. down *Yuyama-Onsen*.

ISHIKAWA=KEN

Yamanaka, 5 m. to the S. E. of the *Daishōji* station (87.7 m. from *Maibara*, in 5 hrs. 30 min.), Electric Tramway available (30 min., fare, 50 *sen*), is a picturesque retreat (240 ft. above the sea) with hills on three sides and the *Daishōji-gawa* traversing the open portion. High banks bound the stream, which is clear and bright. The place is noted for lacquer-wares. Pop. 3,300. Number of visitors: 28,832, the total stays being 147,495 days (1909). *Kakke* patients resort to the spa very often. Inns: *Yoshino-ya*, *Mitani-ya*, *Ogi-ya*.

Meteorological Observations.

Summary.

For the year, 1911.

| Month | Air Temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity | |
|-----------|-----------------|-------|-------|-------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|---------|
| | Mean | | | | Absolute | | | Total | % | | Total | Mean |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | | |
| January | 2.79 | 6.52 | -0.72 | 7.23 | 15.9 | -3.9 | 81.14 | 47.76 | 26 | 46.1 | 1.7 | — |
| February | 4.05 | 7.22 | 0.67 | 6.53 | 14.3 | -4.9 | 77.0 | 61.97 | 20 | 159.8 | 1.39 | S 63° W |
| March | 6.57 | 11.12 | 2.49 | 8.63 | 18.2 | -3.0 | 79.0 | 81.46 | 22 | 233.5 | 1.69 | S 57° W |
| April | 10.00 | 15.34 | 6.2 | 9.04 | 29.3 | -1.5 | 76.0 | 120.63 | 31 | 206.6 | 2.48 | N 71° W |
| May | 15.8 | 20.9 | 10.5 | 10.4 | 25.4 | 3.0 | 69.7 | 213.97 | 49 | 98.2 | 2.41 | S 5° E |
| June | 20.0 | 23.9 | 16.2 | 7.69 | 27.8 | 11.0 | 79.1 | 86.57 | 20 | 335.4 | 4.31 | S 11° E |
| July | 24.1 | 28.3 | 20.3 | 8.0 | 33.9 | 13.4 | 79.0 | 110.80 | 25 | 88.1 | 1.7 | S 30° E |
| August | 24.6 | 28.9 | 20.6 | 8.2 | 34.2 | 16.5 | 79.0 | 162.55 | 39 | 292.3 | 1.7 | S 56° E |
| September | 22.4 | 27.0 | 18.5 | 8.5 | 34.2 | 13.0 | 80.0 | 124.25 | 34 | 320.3 | 1.8 | S 36° E |
| October | 14.6 | 19.3 | 10.8 | 8.5 | 26.3 | 7.5 | 80.0 | 125.65 | 36 | 236.1 | 2.0 | S 63° E |
| November | 11.8 | 16.0 | 7.9 | 8.4 | 23.4 | -3.1 | 77.0 | 90.00 | 29 | 335.5 | 2.2 | S 12° E |
| December | 4.6 | 8.0 | 1.8 | 6.1 | 18.3 | -1.8 | 85.0 | 54.35 | 18 | 221.9 | 2.1 | S 35° E |

Yamashiro Spa is separated from the station *Iburi-bashi* by about 3 m., with horse-trams running, and by half a mile more from *Daishōji*, the two spas, Yamanaka and Yamashiro, being about 2.5 m. from each other. Though somewhat inferior to Yamanaka in scenery, Yamashiro possesses better accommodation both for lodging and bathing. The fact that here are situated the principal kilns of *Kutani-yaki* pottery adds special importance to this place. Pop. 2,300. Electric bath. The water is said to be efficacious for rheumatism and brain disorders. Number of visitors: 19,375, the total stays being 96,795 days (1909). Inns: *Ara-ya*, *Kura-ya*.

Meteorological Observations.

Summary.

For the year, 1911.

| Month | Air Temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity | |
|-----------|-----------------|-------|-------|-------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|---------|
| | Mean | | | | Absolute | | | Total | % | Total | Mean | Dir. |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | | |
| January | 2.89 | 6.35 | -0.06 | 6.42 | 16.3 | -3.8 | 79.13 | 45.6 | 22 | 117.3 | 3.09 | N 21° E |
| February | 3.94 | 7.42 | 0.76 | 6.66 | 15.0 | 5.0 | 70.15 | 62.4 | 20 | 154.4 | 3.14 | N 41° W |
| March | 6.15 | 11.22 | 1.85 | 9.37 | 18.4 | -3.3 | 76.5 | 94.82 | 26 | 197.6 | 2.82 | N 23° W |
| April | 10.6 | 14.91 | 6.29 | 8.62 | 28.1 | -1.9 | 71.8 | 132.93 | 34 | 182.3 | 3.39 | N 25° W |
| May | 15.4 | 19.06 | 9.0 | 10.0 | 24.6 | 2.3 | 67.3 | 240.23 | 55 | 95.9 | 2.54 | S 86° W |
| June | 19.7 | 23.15 | 16.2 | 6.94 | 27.3 | 11.1 | 77.1 | 117.32 | 27 | 313.8 | 2.20 | N 62° W |
| July | 24.0 | 27.9 | 20.5 | 7.3 | 32.7 | 13.3 | 81.0 | 153.61 | 35 | 175.0 | 2.40 | S 85° W |
| August | 24.7 | 28.8 | 20.7 | 8.0 | 33.7 | 16.5 | 83.0 | 194.55 | 47 | 259.7 | 1.90 | N 54° W |
| September | 22.4 | 27.0 | 18.5 | 8.5 | 32.8 | 13.0 | 83.0 | 141.64 | 38 | 279.6 | 2.00 | S 55° W |
| October | 14.9 | 19.0 | 10.8 | 9.1 | 26.5 | 7.4 | 82.0 | 121.2 | 35 | 257.5 | 2.30 | S 4° E |
| November | 11.4 | 16.0 | 7.0 | 9.0 | 22.8 | 1.9 | 79.0 | 92.0 | 30 | 267.4 | 3.20 | S 8° W |
| December | 5.1 | 8.6 | 1.7 | 6.9 | 18.6 | -1.9 | 87.0 | 55.2 | 18 | 187.8 | 3.10 | S 3° E |

Katayamazū is 2 m. N. W. of the *Iburi-bashi* station and can be reached by omnibus or *jinrikisha*. The spa (Pop. 500) lies on the lake side of *Shibayama-gata* (about 10 m. in circumference), and is 1.5 m. distant from the sea. The spring issuing from bed of the lake was found in the Era of Shō-ō (1652-5), but used first for bathing in 1883, after the shore of the lake had been partly reclaimed. The water is said to be efficacious for dyspepsia. The lake abounds in fish, especially terrapin (*dorogame*). Number of visitors: 12,378, the total stays being 32,833 days (1909). Inns: *Hoshi*, *Kami-ya*.

Meteorological Observations.

Summary.

For the year, 1911.

| Month | Air Temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity | |
|-----------|-----------------|------|------|-------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|---------|
| | Mean | | | | Absolute | | | Total | % | | Total | Mean |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | | |
| January | 2.9 | 6.2 | 0.0 | 6.24 | 16.1 | -2.7 | 63.0 | 57.18 | 19 | 68.9 | 0.59 | — |
| February | 4.3 | 7.03 | 0.97 | 6.06 | — | — | 79.8 | 71.7 | 23 | 85.4 | 0.96 | N 76° E |
| March | 6.72 | 10.7 | 3.02 | 7.68 | 10.5 | -2.2 | 80.8 | 113.47 | 31 | 68.9 | 2.38 | N 74° E |
| April | — | — | — | — | — | — | 78.4 | 98.28 | 25 | 30.8 | 2.18 | N 25° E |
| May | — | — | — | — | — | — | 74.6 | 215.46 | 49 | 21.1 | 4.30 | N 56° E |
| June | 20.7 | 23.7 | 17.8 | 5.9 | 28.8 | 11.0 | 82.0 | 138.98 | 32 | 281.8 | 1.34 | S 43° W |
| July | 25.2 | 29.1 | 21.6 | 7.5 | 34.0 | 13.9 | 81.0 | 150.10 | 34 | 102.2 | 1.20 | S 30° W |
| August | 25.6 | 29.7 | 21.9 | 7.9 | 34.0 | 18.0 | 83.0 | 177.25 | 42 | 294.5 | 1.20 | S 34° E |
| September | 23.3 | 27.8 | 19.2 | 8.6 | 34.8 | 13.8 | 82.0 | 110.3 | 30 | 207.3 | 1.10 | S 24° W |
| October | 15.6 | 19.8 | 11.9 | 8.0 | 27.7 | 8.2 | 81.0 | 95.3 | 27 | 214.4 | 1.30 | S 24° E |
| November | 11.6 | 16.1 | 7.9 | 8.2 | 24.0 | 2.9 | 80.0 | 66.9 | 22 | 253.1 | 2.30 | S 10° E |
| December | 5.2 | 8.3 | 2.3 | 6.0 | 18.5 | -0.7 | 85.0 | — | — | 223.1 | 2.60 | S 29° E |

Awazu (95.3 m. from *Maibara*, in 6 hrs.) is separated from the station of the same name, by a little over 2 m. and reached by horse-trams (fare, 12 *sen* special, 8 *sen* ordinary). Pop. 600. The water is believed to be efficacious for venereal diseases.

Sanko-dō ("Three Lakes Hall"), 2 m. from the station and 2 m. from the next station *Komatsu*, is so called from the three lakes or lagoons, *Kiba-gata*, *Shibayama-gata*, and *Imae-gata*, which may be viewed from here. Number of visitors: 28,704, the total stays being 96,879 days (1909). Inns: *Morimoto*, *Yude*.

Wakura, 6 m. to the N. W. of *Nanao* (40.8 m. from *Kanazawa*, in 3 hrs.), omnibuses, *jinrikisha* or steamships available, is one of the most noted spas in Japan for medical efficacy (for rheumatism, wounds, dyspepsia, etc.). The waters are alkaline, colourless, and somewhat bitter in taste, the temperature being as high as 183° F. The springs come out from the sea-bed close to the shore. The island of *Noto* and the smaller islands of *Tsukune-shima*, *Tane-shima*, *Kara-shima*, and others that lie in front of this town very much heighten the attractiveness of its scenery. On *Uki-shima*, which lies close to the shore, stands a small temple dedicated to the Goddess *Benten*. Number of visitors: 21,267, the total stays being 79,237 days (1909). Inns: *Wakasaki-kwan*, *Koizumi*.

Yuwaku (9 m. S. E. from *Kanazawa*) lies on the bank of the upper course of the *Asano-gawa*, at the foot of a mountain lying on the boundary of *Kaga* and *Etchū*. The water is claimed to be efficacious for syphilis. Number of visitors: 4,030, the total stays being 18,127 days (1909).

Tatsunokuchi (8.3 m. N. E. from *Komatsu* station on the *Hokuroku* Main Line) is situated on the S. bank of the *Tetori-gawa*. The water is said to be good for the treatment of diseases of the skin. Number of visitors: 11,361, the total stays being 50,595 days (1909).

Fukatani. 3 m. N. E. from *Kanazawa* and 7 m. S. from *Tsubata* station. The springs are believed to be efficacious for haemorrhoids. Number of visitors: 9,393, the total stays being 17,268 days (1909).

FUKUI-KEN

Awara is situated near the *Awara* station on the Mikuni Line (Kanatsu—Mikuni), a branch of the Hokuroku Main Line. The springs are located at the N. corner of the plain extending between Mikuni and Fukui, bounded by hills on the N. Before 1884 this region was a marshy ground grown with reeds, hence the name 'Plain of Reeds.' In consequence of floods in that year the hot springs were found. The sea bathing place of Mikuni is about 2,5 m. N. from the spa. Inns, 25 in number, are all provided with private baths. The spa is resorted to by people from Fukui and also from Kyōto and Ōsaka. Number of visitors: 20,083 (1909); 41,986 (1913).

MIE-KEN

Komono (Yunoyama Hot Spring) (11 m. N. W. of *Yokkaichi* station and 23,2 m. from Nagoya, in about 1 hr.) is at *Yunoyama*, close to the village of Komono. The place is surrounded on three sides by hills, while on the E. it offers a fine prospect of the bay and the hills of Owari and Mikawa beyond. A light railway leads as far as Yunoyama. In the upper course of the River *Mitake-gawa*, on which Yunoyama is situated, there is a waterfall called "*Ao-taki*," which is one attraction of that spring town. The hot water is believed to be efficacious for *kakke* (beri-beri), lung troubles, diseases of the stomach, etc. Number of visitors: 2,376, the total stays being 5,497 days (1909). Inns: *Kotobuki-tei*, *Ise-ya*.

Toba (81,8 m. from Nagoya, in 3 hrs. 42 min.). Formerly the castle town of the Daimyō Inagaki, Toba is a very pretty place, open to the sea toward the S. E., and surrounded by mountains on the N. W. The port, situated at the entrance of the Bay of Ise and right opposite to the Irako Promontory on the coast of Mikawa, is protected from rough seas by numerous pretty islets and is a haven of refuge for coasting steamers. Pearl culture:—Mr. *Mikimoto* carries on the pearl-culture industry with great success at Ago Bay, not very far from Toba, and at Toba he has made arrangements for showing foreign visitors the method of gathering pearl-oysters by female divers, *ama*. The female divers of Toba,—bare to the waist, their lower limbs clad in a red garment—who plunge into the deep sea and fish up pearl-oysters, are an exceedingly hardy race. They are certainly worth seeing. Castle site, now owned by the Navy Department, affords a very fine view of the sea and islands.

Hiyori-yama, literally 'Weather Hill,' is situated at the N. W. end of Toba port. It commands a splendid view, not only on the beautiful islet-studded sea in front (often compared to *Matsu-shima*), but also of the distant mountains of Shinano, Kai, and Suruga,—*Fuji* among others.

Hinoyama, about 4 m. from *Futami-ga-ura*, is within a few minutes' walk from Toba station. The extensive range of view from it and the superb excellence of its scenery rank it among rare scenic spots in this country. Tourists visiting Toba are advised to make ascent of this hill, as they are sure to be amply compensated for the time and exertion required. Recently the roads have been much improved to make the ascent easier and cottages built on the hillside for travellers who can take rest and have refreshments. Hinoyama is also known as *Kakehi-yama*, the name having been originated in the historical fact that Kuki Yoshitaka, a feudal lord who was a naval commander under Taikō Hideyoshi, designed to supply water to the castle of Toba from a spring on that hill by means of *kakehi* (bamboo tubes).


Toba, the "Town of Longevity." According to the recent report of the town headman, Toba has 1,233 houses and 6,419 inhabitants, of this small population happy people who are more than seventy years old number 220. It is not without reason that Toba is called the "Town of Longevity."

NARA-KEN

Rokuyō-Kōsen, 0,7 m. S. from *Nara*, is situated on the road '*Nabari-Kaidō*,' on the bank of a stream, the cold spring issuing from its bed. Number of visitors: 15,043 (1909).

Shiraya-Kōsen is on the S. E. of Mt. *Yoshino* (famous for cherry blossoms) and is situated on the upper course of the River Yoshino. The spa is on all sides shut in by wooded hills. The river abounds in "*ayu*" fish. Mineral water is said to be efficacious for dyspepsia, carcinoma of stomach, uterus diseases syphilis, gonorrhoea, consumption, etc.

THE HINOYAMA RECREATION-GROUND NEAR TOBA AND ENVIRONS.



WAKAYAMA=KEN

Sedo-no-Kanayama (Yuzaki). The spa lies 12.5 m. S. W. from *Tanabe* (75.4 m. S. from *Wakayama*). From *Wakayama* to *Tanabe*, *jūrikisha* available along the highway, or steamships; from *Tanabe* to the spa by ferry. The hot springs are located on the S. side of a small peninsula projecting from the W. coast of Kii. Here hot springs come forth partly from the beach and partly from the bottom of the sea, so that hot sand bath can be taken in this spa. One of these springs, called "*Kin-cki-sen*" or "*Saki-no-yu*," issues from the fissure of rocks close by the sea, forming a gourd-shaped bath-basin. The bath stands in the open air and bathers have a beautiful view of the surroundings. The spa affords also facility for sea bathing and those visitors may also amuse themselves with fishing. The climate being very mild, it is a delightful resort. Number of visitors: 5,339, the total stays being 44,597 days (1909). see Plate 48

Yunomine (Shimura-Onsen). 42 m. E. from *Tanabe* (Kii Province), no *jūrikisha* available. The springs are 2.6 m. S. W. from *Hongū-mura*, which lies on the bank of an affluent of the *Kuma-no-gawa*, 24 m. up from *Shingū-machi*. The place is enclosed on every side by mountains, and situated at an elevation of 630 ft. above the sea. These hot springs, 3 in number, prove efficacious in diseases of the skin, the water containing sulphide. The "*Tsubo-yu*" comes forth from a cave of rocks on the river bed, the cave itself forming a bath. It is so narrow that only one person can bathe in. Number of visitors: 5,985, the total stays being 15,614 days (1909). see Plate 49

Katsu-ura. Near the sea-port *Katsu-ura* on the S. coast of Kii Peninsula is **Soto-no-yu Hot Springs**, facing the sea on three sides and enjoying a mild climate both in winter and in summer. High cliffs stand close to the spa. The place has every facility for fishing and sea bathing. The adjoining sea is noted for whale-fishing. The slope of a hill behind the place is a recreation ground, open to the public, from which a charming view of the famous waterfall of *Nachi* can be obtained. see Plate 50

KYŌTO=FU

Kasagi (37.7 m. from *Minato-machi*, in 2 hrs. 18 min.) is rich in the beauties of mountain and stream. *Kasagi-Onsen* (205 ft. above the sea), 0.2 m. N. E. from the *Kasagi* station on the Kwansai Line and by the River *Kizu-gawa*, is a hot cabonated spring. Many people resort here, not only on account of the spa, but for the beautiful country life and the fishing in summer.

Kasagi-yama (0.1 m. from the station) is easily climbed, as it is but 0.5 m. from the foot to the summit. It is a famous spot, being the place where the Emperor Godaigo-Tennō took refuge when he fled from Kyōto. The temple *Kasagi-dera* (at the summit), where the Emperor once stayed, has since been almost entirely destroyed by fire. The temple grounds contain a large number of *ume* (plum) and cherry-trees, while the view of the surrounding country is splendid. There are also a number of large, singularly shaped rocks, to which various names have been given, such as *Yakushi*, *Miroku*, *Kokūzō*, *Kasagi-ishi* (the largest of these is about 150 sq. ft. in area). **Kasagi-ishi**. A story goes that the Emperor Temmu-Tennō (in the 7th century) was once caught in a tremendous rain-storm on this hill, when the *Kokūzō-Bosatsu* appeared and gave him shelter. The Emperor thereupon vowed to dedicate a temple to this deity, as a thank-offering for the kindness done him, and as a pledge he took off the hat he was wearing and laid it on a rock,—hence the name of the rock and the hill (*Kasagi* meaning "hat laying down"). The famous *Tsukigase* plum grove may be reached from the next station *Shima-ga-hara* (7.8 m. from *Kasagi*). The hot springs issue from the fissure of granite rocks on the *Kizu-gawa*. Number of visitors: 1,236, the total stays being 2,636 days (1909).

ŌSAKA=FU

Hamadera (9.3 m. from *Namba* station, in 30 min.) is a long sea-beach extending for 1.4 m.; it is covered with large pines and its scenery is supposed to fairly rival that at *Suma* and *Akashi*. The place is a favourite resort of Ōsaka people. Pop. 600. Bathers number about 800,000 per annum. see Plate 51

Meteorological Observations.

Summary.

| Month | Air Temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity |
|---------------------|-----------------|------|------|-------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|
| | Mean | | | | Absolute | | | Total | % | Total | Mean |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | |
| For the year, 1911. | | | | | | | | | | | |
| June | 21.0 | 26.0 | 17.9 | 8.1 | 29.5 | 13.5 | 79.0 | 147.3 | 34 | 288.8 | — |
| July | 25.3 | 30.8 | 21.4 | 9.3 | 34.9 | 17.1 | 82.0 | 245.2 | 56 | 220.1 | — |
| September | 24.2 | 29.6 | 20.3 | 9.3 | 35.1 | 15.7 | 81.4 | 199.7 | 54 | 188.3 | — |
| October | 19.4 | 22.3 | 11.5 | 10.7 | 27.5 | 6.5 | 73.4 | 185.6 | 53 | 69.5 | — |
| November | 12.1 | 17.7 | 7.0 | 10.7 | 22.6 | 1.2 | 71.8 | 173.5 | 56 | 68.1 | — |
| December | 6.8 | 11.8 | 2.1 | 9.7 | 20.6 | -1.9 | 70.9 | 173.3 | 57 | 25.4 | — |
| For the year, 1912. | | | | | | | | | | | |
| February | 8.0 | 12.7 | 2.7 | 10.0 | 22.6 | -3.3 | 73.8 | 154.4 | 49 | 132.7 | — |
| March | 8.3 | 13.3 | 3.3 | 10.0 | 21.5 | -0.8 | 72.2 | 156.6 | 42 | 98.0 | — |
| November | 10.2 | 15.7 | 5.4 | 10.3 | 22.0 | 0.2 | 71.5 | 153.1 | 49 | 50.7 | 3.8 |
| December | 6.7 | 12.0 | 1.5 | 10.5 | 20.1 | -1.9 | 73.1 | 146.5 | 48 | 109.2 | 5.2 |
| For the year, 1913. | | | | | | | | | | | |
| January | 4.0 | 8.7 | -0.3 | 9.0 | 12.7 | -3.8 | 70.4 | 150.6 | 48 | 51.3 | 5.8 |
| February | 5.0 | 10.0 | 0.7 | 9.3 | 17.3 | -5.0 | 67.6 | 178.6 | 58 | 54.5 | 5.8 |
| March | 6.3 | 12.2 | 0.2 | 12.0 | 18.6 | -4.0 | 68.8 | 219.2 | 59 | 51.6 | 4.2 |
| April | 14.1 | 20.0 | 8.7 | 11.4 | 27.1 | 0.0 | 74.5 | 181.0 | 46 | 70.9 | 3.7 |
| May | 19.5 | 22.4 | 10.8 | 11.6 | 29.3 | 4.6 | 74.8 | 230.3 | 54 | 189.3 | 3.0 |
| June | 21.5 | 25.0 | 17.8 | 7.8 | 30.3 | 11.7 | 77.9 | 180.6 | 42 | 116.7 | 3.0 |
| July | 25.4 | 31.1 | 20.7 | 10.3 | 34.9 | 13.2 | 73.6 | 281.3 | 64 | 24.2 | 3.7 |
| August | 26.2 | 32.5 | 21.0 | 11.5 | 36.1 | 17.8 | 72.6 | 246.7 | 60 | 81.7 | 3.2 |
| September | 20.9 | 27.0 | 16.0 | 11.0 | 32.0 | 9.7 | 72.3 | 198.7 | 54 | 68.1 | 2.7 |
| October | 15.0 | 22.6 | 11.3 | 11.4 | 27.5 | 3.8 | 74.8 | 205.2 | 59 | 173.1 | 3.3 |
| November | 9.9 | 17.0 | 4.7 | 12.3 | 21.9 | 1.0 | 71.8 | 181.8 | 59 | 98.0 | 2.7 |
| December | 6.4 | 11.8 | 2.2 | 9.5 | 18.9 | -1.9 | 70.0 | 149.2 | 49 | 77.2 | 5.1 |
| For the year, 1914. | | | | | | | | | | | |
| January | 4.6 | 10.4 | -0.3 | 10.7 | 18.0 | -4.6 | 67.9 | 177.0 | 55 | 27.5 | 4.5 |
| March | 9.6 | 15.2 | 4.9 | 10.2 | 22.0 | -1.5 | 74.0 | — | — | 159.6 | 3.4 |
| April | 12.0 | 17.2 | 6.9 | 10.3 | 24.5 | 1.0 | 72.3 | — | — | 80.0 | 3.4 |
| May | 18.4 | 23.6 | 13.4 | 10.2 | 27.9 | 4.0 | 77.1 | — | — | 236.0 | 1.9 |
| June | 22.1 | 26.7 | 18.0 | 8.7 | 32.2 | 13.7 | 80.3 | — | — | 187.9 | 1.9 |

HYŌGO-KEN

Takarazuka (15.6 m. N. W. from Ōsaka, in 53 min., by railway or by electric tram). The watering-place is situated on the W. bank of the *Muko-gawa*, at the foot of Mt. *Muko-san*, and at an elevation of 82 ft. above the sea. In the spa there are two springs, one on the left bank of the stream and the other on the opposite side, both being carbonated springs. Both the springs are only for public use. The inhabitants of Ōsaka and its neighbourhood resort to this place for amusement rather than for recuperation. A few years ago a large bathing-establishment with a swimming-tank was built here and since then the place has grown popular rapidly.

About 1 m. farther up the stream is also a carbonated spring. The water is bottled for export by the Clifford-Wilkinson Tansan Mineral Water Co. Number of visitors: 7,133, the total stays

The Principal Mineral Waters bottled for Sale and Export.



Wilkinson-Tansan.



Wilkinson-Tansan.



Reynell's
Hirano Water.



Nunobiki Tansan.



Nunobiki Tansan.



Hirano Water
(Peacock Brand).



Hirano Water
(Musuya Brand).



Hirano Water
(Peacock Brand).

being 138,285 days (1909); 161,512, the total stays being 28,000 days (1913). Hotel: *Takarazuka Hotel*. Inns: *Fundo-ya, Kotobuki-ryō, Kijamri*.

Takarazuka Tansan Works. Over twenty five years ago a young Englishman, Clifford Wilkinson by name, was hunting near Kōbe. He came to a hamlet called Takarazuka and stopped to quench thirst at a deliciously cool, fragrant, sparkling spring, whose bright water gushed out of the ground under his feet. When he had finished his hunting expedition, his attention was drawn to the delightful taste of the water, and he brought back a bottle of it and presented his friends with it. He was gratified to learn how highly it was appreciated, for one and all decided that it was the best they had ever tasted. The next morning he went again to Takarazuka and carried back a big sack of filled bottles on his back. The waters were also demanded and satisfied by all his comrades. This was the beginning of the famous industry and since that occasion the Tansan factory was erected on the site of the wonderful spring at Takarazuka. The waters issuing from the big well, 29.5 ft. deep, built in by a two feet concrete wall surrounded by natural rock, flow down through three separate rows of filters, by which every impurity and iron oxide can be rid off. The factory has bottling, labelling, filling, corking and inspection departments, all being models of neatness and freedom from filth or dirt. In the bottle washing department every bottle is soaked for 15 min. in pure mountain water and afterwards brushed and rinsed inside and outside. The gas is produced from Brunner and Mond's bicarbonate of soda and sulphuric acid by the gas generators and pumps, the latest up to date machinery. The bottling machines are fitted with Automatic Feed Corking Appliances which avoid the corks being touched by hand. In the bottle examining and labelling department each bottle lies on the shelf for 24 hrs. or more and is then reexamined and packed. The Tansan Works cover about 4,000 *tsubo* (16,000 sq. yds.), the capacity of the spring about 40 million bottles per annum and the present output about 30,000 daily, 90 per cent of which is exported. Analyses have been made by the most celebrated analyst in Europe, in America, also in Ōsaka and Hongkong.

Hirano. The mineral springs are situated at *Hirano* in *Tada-mura*, Hyōgo Prefecture, about 8 m. N.W. from Ōsaka. The water issues from the crevices of rocks in a valley surrounded by hills and is claimed to have curative power for various kinds of diseases, such as chronic catarrh of the stomach and bowels, female diseases, anemia, lung and throat affections, etc. It is, however, used only for drinking.

The Imperial Mineral Water Co. The water bottled in this works is called Hirano Water from the name of the locality where it is produced, and has two brands, "*Mitsuya*" (Three Arrows) and "*Kujaku*" (Peacock). Hirano Water is noted for its hygienic properties as well as its being entirely free from alteration or fermentation. The Hirano spring dates back to 970. It was first known as the sacred water of *Tada* and, it seems, appreciated not only as a beverage but also as a panacea for all kinds of ailments. Towards 1881, some of the parties interested in the locality attempted to bottle the mineral water, but it proved a failure. The matter then came under the notice of an English expert Mr. Gallan in the employ of the Kawasaki Dock-yards, Kōbe. He persuaded some of his countrymen of furnishing him with necessary funds, and having made necessary preparations to make the water a common beverage, he offered it to the general public. This was in the year 1884. Subsequently the spring was owned by a Japanese millionaire, Mr. Iwasaki. In 1905 the enterprise was placed under the control of the "*Mitsuya*" Brand Hirano Spring Water Semi-Partnership Co. Again in 1907 the present Imperial Mineral Water Co. was organized by some prominent men with a capital of ¥ 600,000, and took over the management.

With regard to the equipment of the works, in addition to the appliances hitherto used, the company has purchased the machines of the most recent type, those used at the Apollinaris Works in England. The water is produced at the springs at the rate of 600,000 gallons per diem with 28,000 cubic feet of natural carbonic acid gas. The demand for Hirano Mineral Water is on the increase year after year, the output for the year 1914 amounting to 11,800,000 bottles, of which about 10 per cent is exported to foreign countries, especially to China, Australia, India and Java. Besides bottling the Hirano Water, the company also takes an active part in the sweet aerated water trade of Japan which business attains very considerable dimensions in the summer time when the demand for non-alcoholic beverages is largest.

Reynell's Hirano Water. The spring is situated at the same place, not far from the spring above mentioned. In 1884, Mr. Reynell imported the necessary machinery and began bottling the water

for use. The flow of water from the spring is continuous, yielding a full supply all the year round. At present the water is bottled by the Hirano Mineral Water Co., Ltd. The works are fitted with modern and improved appliances for bottling the water, the capacity being a turn-out of 3,500 dozen bottles a day. The company has now considerably increased the export of the water to foreign countries including China, the United States, Strait Settlements, the Philippines, Hongkong, etc., and the popularity of this water is yearly increasing.

Takedao (20,8 m. from Ōsaka, in 1 hr. 22 min.) is also a pretty spot with cold mineral springs. Between *Namaze* and *Dōjō*, which is the station next to *Takedao*, the trains run through a valley shut in by high hills over tracks of a very steep gradient, passing through eleven tunnels in a distance of 7,5 m. This part of the route is full of romantic scenery, and is compared by many to *Yabakei* of Ōita Prefecture. Visitors to Arima Hot Springs may alight here at *Sanda*. The spa is situated in a gorge on the bank of the *Takedao-gawa*, at an elevation of 340 ft. above the sea. Those wishing to avoid the discomforts of city life can live here quietly. Number of visitors: 1,802, the total stays being 6,705 days (1909).

Namaze (16,8 m. from Ōsaka, 1 hr.) is close by *Namaze* hot spring, and being surrounded on all sides by hills, forms a nice, quiet resort.

Nunobiki mineral spring is situated on the upper course of the *Shin-Ikuta-gawa*, at the foot of *Nunobiki-yama*, where is the reservoir for the water system of Kōbe. The famous waterfalls *Nunobiki*, of which the lower called *me-daki* is 43 ft. high and the upper called *o-daki* 80 ft., are splendid views. see Plate 57 The place is one of the most delightful spots in the vicinity. The water is bottled by Nunobiki Tansan Works for sale.

Arima. There is a highway from Kōbe, via *Tennj-goe*, 14 m. on which *jinrikishas* are available (fare, ¥ 1,50), or the visitors may go by train to *Sumiyoshi*, whence *yama-kago* or Sedan chairs are available (fare, ¥ 1,40) across *Rokkō-zan* 7,3 m.). Those from Ōsaka will naturally take the Fukuchiyama Line as far as *Namaze*, whence *jinrikisha* for 6,8 m. to the spa town (fare, 70 *sen*), or railway as far as *Sanda*, whence 6,8 m. to Arima by *jinrikisha*, the fare being 50 *sen*. see Plate 58 The spa town is situated in a valley surrounded by *Rokkō-zan* and other hills, being open only to the N. The place itself, being 1,155 ft. above sea-level, is cool in summer, the mercury never rising above 85° F. while in winter the temperature is comparatively warm (35° F. on the coldest days). The town contains about 400 houses, with a population of 2,000. The local specialties are basket-ware, Arima-earthen-ware, etc. The spa town has been famous throughout Japan from time immemorial. Special bath-houses have recently been built (fee, 20 *sen* per bath). The water is alkaline, containing a large quantity of iron, salty in taste and turbid in colour, making reddish-brown deposits. The water is said to be efficacious for dyspepsia, uterus diseases, consumption, diseases of the skin, etc. The place also possesses a mineral spring (called "*Teppō-sui*"), containing carbonic acid gas, the water of which is bottled and exported. Among temples, *Onsen-jū*, *Onsen-jinja*, and *Zempuku-ji* may be visited in the course of a walk.

Tsutsumi-ga-taki is a small waterfall (30 ft. high) in a pretty neighbourhood, which is especially beautiful in autumn. *Kado-yama* is a hill to the N., also called *Arima-Fuji* on account of its likeness to Mt. Fuji. Number of visitors: 10,132, the total stays being 64,605 days (1909); 9,793, the total stays being 40,455 days (1913). Inns: *Hyōe*, *Ikeno-bō*, *Nikai-bō*, *Gosho-no-bō*, *Naka-no-bō*; Hotels for accommodation of foreigners: *Sugimoto*, *Masuda* and *Arima*.

Kinosaki (99,2 m. from Kyōto, in 5 hrs. 30 min.) is a popular spa town, the hot springs having been known since the 7th century. The town is situated at the foot of a hill called *Kuruhi-dake*, on the W. bank of the *Asago-gawa* (the lower course of the *Maruyama-gawa*) and within 2 m. of the sea-port of Tsuiyama. see Plate 59 Kinosaki is noted for its beautiful scenery and salubrious climate. The spring water is colourless and alkaline. Products:—articles made of mulberry-wood and straw fancy-goods. Places of interest: **Onsen-ji**, a buddhist temple, half-way (about 0,3 m.) up the hill *Kanro-hō*, to the W. of Kinosaki, is believed to have been founded by Dōchi-Shōnin, who discovered *Mandara* hot spring, during the Yōrō Era (717–23). Within the temple grounds are two smaller shrines, dedicated to *Kan-on* and *Yakushi*. *Hiyori-yama* is a hillrock, N. of Kinosaki, which has recently been turned into a public park. It commands an excellent view of the surrounding regions and of the sea.

Gembu-dō, or basalt grottoes, are situated on a hillside about 2,4 m. from Kinosaki, on the road to Toyo-oka. They may easily be seen from the train, on account of their curious formation. In

fact the train stops here at certain times during the year. There are three grottoes, the one on the left-hand side being 78 ft. wide at the entrance and 102 ft. deep. The middle one is 72 ft. wide at the entrance and 84 ft. deep, and drops of water trickle down its walls collecting into a transparent pool below. The right-hand one is 78 ft. wide at the entrance and 102 ft. deep, and outside the grotto, right over the entrance there leaps a waterfall. Looked at from a little distance these grottoes indeed present a curious sight; the way thousands of these basalt pillars* stand clustered together, or piled up one upon another, gives them the appearance of a colossal bee-hive. On the left-hand side of the grottoes, there appear cut deep into the rock the three Chinese characters "*Gembu-dō*," the facsimile of the hand-writing of a famous scholar-poet, *Shibano Ritsuzan*.

*Professor S. Shiga in his "Natural Scenery of Japan" says: "Of the basalt pillars found in the country the most noted are those at the *Gembu-dō* in Tajima. They are in shape octagonal, heptagonal, hexagonal, or pentagonal, jet black and flinty in appearance, and 20 to 30 ft. long. They may be counted by the thousand, each pillar showing a cleavage at every foot or 8 inches throughout its length, thus giving the appearance of a post composed of slabs. If taken out one by one, they would make excellent building material, cut to order, and with the finest touch of nature's workmanship."

Number of visitors: 25,351, the total stays being 291,165 days (1909); 41,435, the total stays being 181,993 days (1913). Inns: *Yuto-ya*, *Miki-ya*, *Nishimura*.

Yumura. 5 m. S. from *Hamasaka* station on the San-in Main Line (124 m. from Kyōto, in 6 hrs. 45 min.), *jinrikisha* available. The spa is situated on the N.W. portion of Tajima Province, on a highway, at an elevation of 150 ft. above the sea. Close by the right bank of a small stream called *Haruki-gawa* gush out several boiling hot springs, of which one is now used for bathing. Another one, which issues near the edge of water, is called *Ara-yu* on account of its violent force. The discharge of water is considerable and the spring is used by the villagers for boiling their foods and also for softening vegetable fibres. Eggs can be half-boiled for about 2 minutes, while it requires 3-4 minutes to have coagulated. The spring is distinguished by its abundant flow of water as well as by its high temperature (93-5° C.).

In cold days the place, where the spring gushes out, is entirely covered with vapour, so that nothing is visible in the neighbourhood. Several persons have been wounded by falling into boiling water. The water of the stream being warmed by the spring, a washing-place is provided at the foot of the bank for the use of villagers.

see Plate 8,
fig. 5

see Plate 8,
fig. 5

see Plate 8,
fig. 4

TOTTORI-KEN

Iwai. 2,4 m. from the *Iwami* station (132,7 m. from Kyōto, in 7 hrs. 17 m.). *Uradomi* and *Ajiro* are little sea-ports close to Iwami. The sea between these two places is dotted with innumerable pretty islands and is known as the Matsushima of the Japan Sea; the most noteworthy of these islands are *Sengwan-matsu-shima* and *Natane-jima*, the former so called on account of its beautiful pines and the latter on account of its being covered in spring with the yellow blossoms of *natane* or rape-seed plants. The spa is the most noted in Tajima Province and lies at an elevation of 40 ft. above the sea. There are 1 public bath and 8 private baths in the spa. Pop. 855. Number of visitors: 3,014, the total stays being 30,451 days (1909). Inn: *Kishima-ya*.

Tottori (114,1 m. from Kyōto, in 8 hrs.) is the capital of Tottori Prefecture, and the next city in importance to *Matsue*. It is situated in the valley of the *Sendai-gawa*, in the E. part of Tottori Prefecture. Formerly a castle-town of the Daimyō Ikeda, the city greatly declined after the Restoration, but with the recent opening of railways it has entered upon new era of prosperity. Tottori has 32,682 inhabitants (6,422 households); its busiest streets are along *Wakazakura-Kaidō* and *Chizu-Kaidō*.

Yoshikata Hot Springs are at a distance of 0,8 m. from the station. The springs were found by chance in 1906 when the ground was bored by a proprietor of a factory to get water for a boiler. Since then several springs have been found and bath-houses built. The water here is pumped up, heated, and then conveyed to bath-tanks. Number of visitors: 5,160, the total stays being 6,292 days (1909). Inns: *Tottori-Onsen*, *Takasago-Onsen*.

Yudani (12 m. S. from Tottori) is situated in a mountain-glen, at an elevation of about 300 ft. above the sea. The discovery of the cold springs dates back to the age of mythology, but the

place has been almost neglected owing to the uneasy access and the low temperature of the spring.

Number of visitors: 101, the total stays being 500 days (1909).

Yoshioka. 7 m. W. from Tottori, and 5 m. S. W. from *Koyama* station, next from Tottori. The hot springs are situated on the S. W. part of Lake Koyama, enclosed on three sides by hills. Besides the 6 main springs, there is also a sulphur spring "*Eta-no-yu*," especially provided for the lowest class of people, *eta* (outcasts). Number of visitors: 3,216, the total stays being 11,793 days (1909).

Koyama (146,8 m. from Kyōto, in 8 hrs. 6 min.) is a station whence it is 1 m. to Lake *Koyama-ike*, 9 m. in circumference, the 4th largest in this district. The lake contains seven pretty islets, and its fine scenery may be enjoyed from a railway train. Karo Port is situated N. E. of the lake on the estuary of the *Karo-gawa*. The harbour is shallow and offers no anchorage for large vessels. It is, however, an important port in Inaba Province, there being a regular steamship service to and from *Sakai*. The place is an important fishing centre.

Hamamura (154,9 m. from Kyōto, in 8 hrs. 29 min.) is a spa town known as *Hamamura-Onsen*. On the opposite side of the railway line is found another hot spring, *Kachimi-Onsen*. Number of visitors: Hamamura—700, the total stays being 900 days (1909); Kachimi—75, the total stays being 300 days (1909).

Matsuzaki (165,5 m. from Kyōto, in 9 hrs.) is close to *Tōgō-ike* (lake) and the hot springs. This beautiful lake produces an excellent kind of eel largely sent to Kyōto and Ōsaka. The hot springs, two in number, are found on the banks of the lake; the one on the N. side being called **Tōgō-Onsen** and the other on the S. side, **Asōzu-Onsen**. Tōgō-Onsen gushes up from the bottom of the lake, 0,3 m. from the station (Inn: *Yōji-kwan*). The Tōgō Hot Springs are conveniently located and afford every facility for boating, sailing, and fishing. The place is resorted to by the citizens of Matsue and Tottori during the summer season. Asōzu-Onsen (Inn: *Asahi-kwan*) is 2,4 m. from the station, with a ferry service between. Number of visitors: Tōgō-Onsen—1,272, the total stays being 1,425 days (1909); Asōzu-Onsen—305, the total stays being 1,800 days (1909).

Misasa. 5,5 m. from *Agei* station (168,9 m. from Kyōto, in 9 hrs. 15 min.) on the San-in Main Line, and 5 m. S. E. from *Kurayoshi*. The spa is delightfully situated on the W. bank of the *Misasa-gawa*, which being originated in the boundary of the three provinces, Inaba, Hōki, and Mimasaka, flows from the E. to the W. The place has wooded hills on two sides, N. and S., and is about 50 ft. above the sea. Though the place itself forms a flat land with slight inclination, it has only a limited area, not more than 1 m. in length and some 0,2 m. in width.

In this small space gush out in series a large number of hot springs of high medicinal value, their sources being located mostly near the stream. No less than 30 springs are found in the village, not very distant from one another. All these springs belong to simple thermals, except *Naka-no-yu* which contains sulphides. The temperature of these springs ranges from 45°–75° C. and the flow of water is abundant. There are 4 public baths, and the most noted one called "*Kabu-yu*" or "Stump Bath" is on the outskirts of the village a small distance (0,2 m.) up the stream from the others located in the centre of the spa. This spring was so named from the tradition that one day a white wolf dug the root of an old oak-tree out of the ground, and went to sleep on the stump as his pillow, which enabled people to notice the warmth of the spot. Then they began to bore the ground and discovered the hot spring.

The spring is said to have been found by the inhabitants in the 2nd year of the Chōkwan Era (1165 A.D.). Formerly it was called "*Makura-kabu-yu*," but now simply "*Kabu-yu*" or "*Moto-yu*." It was in the 1st year of the Shōhō Era (1644 A.D.) that the existence of springs in the village proper was first noticed by the inhabitants. Since then many springs have been found and thus there are more than 30 baths at present in this small village. Besides the inns with private baths, there are several houses provided with baths of their own. The rise of gases is observed at every spring, their amount being somewhat different. The water proved exceedingly efficacious for certain cases of dyspepsia, haemorrhoids, diseases of the female genital organs, especially so for rheumatism and neuralgia. This fact can be explained, though not wholly, from the results of our examination of the spring with regard to their radio-activity. As easily seen from the tables previously inserted in this book, the water of these hot springs is generally of strong radio-activity and sometimes contains the largest quantity of radium emanation of those of all hot springs in Japan. Taking also the temperature and discharge of water into consideration, the spa affords the best advantage for the so-called emanation therapy.

On the opposite bank of the *Misasa-gawa*, which separates *Yamada-mura* from *Misasa proper*, gushes out another spring "*Yamada-Onsen*." There are two baths, the one being for public use and the other for private. From a bridge, "*O-iwa-bashi*" or "Big Rock Bridge," which spans the *Misasa-gawa* at the W. end of the village, an abyss called "*O-iwa-buchi*," bounded with huge rocks, may be seen below. A precipitous and rocky hill, "*Kompira-yama*," grown with old pines, standing on the N. bank of the stream, presents a fine view from *Misasa spa*. A narrow gorge behind the hill has another spring and is called "*Yu-dani*," meaning 'Hot Spring Valley.' The village of *Misasa* has a population of about 500 (90 households). Post, telegraph, and telephone office.

see Plate 11

A highroad, *Anagamo-Kaidō*, leads from *Misasa* to *Tsuyama* (45.1 m.) via *Monzen-mura*, the mountain pass of *Ningyōyama-tōge* and *Okutsu* hot springs. About 3 m. E. from the spa is *Sambutsu-ji*, a famous Buddhist temple, situated at *Monzen-mura*, *jirikisha* available thereto. The temple, which belongs to the Tendai Sect, was founded by *En-no-gyōja*, who some 1,200 years ago first explored the mountain. Ascending the mountain, at the foot of which the temple stands, we come to *Monju-dō* and *Jizō-dō* (shrines) and a belfry. Ascending still farther we come to *Oku-no-in*, or the innermost shrine, which is within huge cave. This is popularly called *Nageire-dō* or "Thrown-in Shrine"—meaning that the shrine was first constructed outside and inserted as a whole. This shrine, together with the other shrines above-named—*Monju-dō* and *Jizō-dō*—as well as their archives are under the "special protection" of the government, while the image of the *Zō-ō-Gongen* and a copper mirror owned by the temple are registered as "National Treasure."

Kurayoshi, which is a town of considerable importance (Pop. 8,000), being the trading centre for the E. half of *Hōki Province*. Products:—raw silk, grey cotton, rice-husking implements, *kasuri* (blue figured cotton fabric). The town is also noted for its beautiful Park *Uchibuki-Kōen* which occupies the site of the former castle. It is a fine specimen of landscape-gardening, and has a public meeting hall. *Agei* station is about 3 m. N. E. from *Kurayoshi* and reached by light railway in 13 min. *Sekigane* Hot Springs can be reached from *Kurayoshi* along the *Sekigane-Kaidō*, which leads to *Tsuyama* via the mountain pass of *Ohazama*.

Misasa-Onsen is visited mostly by the inhabitants in the neighbourhood, but also by people from the Prefectures of *Okayama*, *Hyōgo*, and *Shimane*. Number of visitors: 2,515, the total stays being 8,957 days (1909). The high value of the water has been known for a long time to the native inhabitants, but not yet brought to public notice. The locality is well suited for being a pleasure resorts, and better still, the water is exceptionally strong in radio-activity. It is hoped that some arrangements will be made to employ water in every form of application at the fountain source, and conform with the other conditions that modern emanation therapy requires, and also providing necessary accommodation for visitors of higher classes.

Sekigane. 6.5 m. S. W. from *Kurayoshi*, *jirikisha* available. The spa is situated on the E. bank of a narrow gorge of the *Tenjin-gawa*, at an elevation of about 500 ft. above the sea. It has 750 inhabitants, 8 baths, and 5 inns with private baths. The discovery of the spring dates back to the Era of *Enryaku* (782–806). On the temple-grounds of *Jizō-in* near the spring there is an old Bo-tree (*bodaiju*), 15 ft. in circumference, its branches and leaves extending over the bath-house, which is noted for its curious shape resembling an umbrella. The place is the site of the castle of a Daimyō *Yamana*, and is now a park (*Kamei-Kōen*), from which may be obtained a fine view of *Daisen*, the highest mountain in the *San-in-dō*. The water is said to be used beneficially for the treatment of rheumatism and diseases of the skin. Number of visitors: 1,875, the total stays being 3,891 days (1909).

Kaike. Near the mouth of the *Hino-gawa*, 2.5 m. N. E. from *Yonago*, hot springs gush out of the bottom of the sea, about 1 m. off the shore. The water is conducted by pipes into baths.

SHIMANE-KEN

Tamatsukuri is the most noted hot spring in *Shimane Prefecture*, about 1.3 m. S. from *Yumachi* station (223.8 m. from *Kyōto*, in 12 hrs.) on the *San-in Main Line*. It lies on the banks of a small stream at the base of *Kōkasen-zan*, which produces red and blue agates, the latter being highly prized on account of rarity. *Tamatsukuri-jinja* stands at an end of the spa, and beautiful *Lake Shinji* is visible from the place. The spa is resorted to by the citizens of *Matsue*. The air temperature lately measured by the village office is as follows:—

| Month | Min. | Max. | Month | Min. | Max. | Month | Min. | Max. |
|----------|-------|-------|--------|-------|-------|-----------|-------|-------|
| January | 48.5° | 51.2° | May | 64.3° | 71.3° | September | 74.7° | 83.8° |
| February | 45.0° | 49.0° | June | 73.5° | 77.0° | October | 64.1° | 69.7° |
| March | 45.5° | 55.9° | July | 71.6° | 82.4° | November | 53.9° | 56.3° |
| April | 61.2° | 63.4° | August | 86.7° | 87.7° | December | 47.2° | 54.9° |

Number of visitors: 7,550, the total stays being 8,510 days (1909). Inns: *Hosei-kwan*, *Tōfu-ya*. *Matsue*, noted for beautiful scenery, is about 3.3 m. from the spa.

Matsue (219.7 m. from Kyōto, in 11 hrs. 49 min.; Inns: *Minami-kwan*, *Iwata*, *Ichimonji-ya*, *Akagi-kwan*) is situated at the W. end of a narrow strip of land lying between Lake *Shinji-ko* and *Naka-no-umi*, being pierced by the *Ōhashi-gawa* which connects the two lagoons, while in the background rise the Shinji hills. The city was founded by the Daimyō *Horio* in the 17th century, when he received this region in fief from *Tokugawa Iyeyasu* and built his castle on the Gokuraku-ji hill. Since then the place has changed its master twice, the last Daimyō who held it being *Matsudaira*. Matsue, being thus situated by a lake and river, enjoys many facilities of communication and transportation. Thus equipped the city forms a most important distributing centre and with a large population (36,209) is justly regarded as the first metropolis of the San-in District. Besides, it is rich in natural scenery and is considered by some to resemble Geneva, Switzerland. From the famous bridge, *Ōhashi* (more than 600 ft. long), the view takes in Lake Shinji-ko, Naka-no-umi, Mt. Daisen, and Sambe-yama, while the houses on the shores of the lake are reflected in its clear waters.

Lake Shinji-ko. The lake, which is 9.8 m. from E. to W. and 3.7 m. from N. to S., is 31.9 m. in circumference. It is the 6th largest lake in Japan. The Magata Channel connects it with *Naka-no-umi*, while the *Sada-gawa* constitutes its outlet to the sea. The lake abounds in carp, gibel, *suzuki* (a kind of perch-sea-wolf), and white-bait (*shira-uo*). The *suzuki* found in Sung-kiang, in China, which are so justly famed—with large mouth, small scales, and double set of jaws. *Kugedo-no-iwaya*, or the "Divine Cave of Kugedo," is situated at a point 7.3 m. from Matsue, in a cliff projecting 100 ft. into the Japan Sea. Its opening is just large enough to admit a small boat, but it becomes roomy inside. As one finds oneself inside, the echoes of the sound of angry waves resounding from the walls will be almost deafening and unbearable. The cave is believed to have been the birth-place of the god *Sada*.

Gakutō. 0.5 m. S. from *Shōbara* station and 13.6 m. W. from Matsue. The spring is located on a rice-field and has nothing attractive in the neighbourhood.

Ushio. 11 m. S. E. from *Shinji* station and 8 m. S. from Matsue. The spa lies on the S. bank of the *Ushio-gawa*, at an elevation of 300 ft. above the sea and 2.5 m. distant from *Daitō-machi*. The water being lukewarm, a cylinder filled with charcoal is laid in it to keep its temperature. The spring comes forth from a crevice of rocks, which form the bottom of the bath, and bathers drink fresh water direct from this crevice by bamboo pipes. The water is claimed to be efficacious for wounds, diseases of the skin, and rheumatism. Number of visitors: 2,153, the total stays being 6,629 days (1909).

Yumura. 17 m. S. from Shinji station and 27.7 m. from Matsue. The spa is situated on the bank of the *Ii-kawa*, at an elevation of 600 ft. above the sea. Here springs gush out from a crevice of granite rocks close by the stream and bath-house is built direct over the source of springs. It is a quiet watering-place in Izumo Province. Number of visitors: 350, the total stays being 912 days (1909).

Kawai. 3.7 m. S. E. from *Ōta-machi* and 3.8 m. S. W. from Matsue. The place lies on a road leading to *Shigaku* Hot Springs, at an elevation of 300 ft. above the sea.

Tōkōji spring is claimed to be efficacious for chronic catarrh of the stomach and the intestines, disorders of the female genital organs, and brain diseases, while **Urisaka** spring, 1.4 m. distant from Tōkōji, is said to be used beneficially for swellings, diseases of the skin, catarrh of the stomach and the intestines, etc.

Ikeda (9.7 m. S. E. from *Ōta-machi*) lies on the road leading to *Shigaku*, at the foot of *Sambe-yama*.

Koyabara (11 m. S. E. from *Ōta-machi*) lies at the foot of *Sambe-yama*, at an elevation of 900 ft. above the sea. *Shigaku-Onsen* is 5 m. S. E. from this place. The production of gas from the spring is abundant and the water is conducted into the bath by wooden pipes, laid about 15 ft. under the ground. Number of visitors: 310, the total stays being 577 days (1909).

Shigaku. 12 m. S. E. from Ōta-machi (7,3 m. S. W. from *Oda*, the terminus of the San-in Main Line, automobile service once daily); from Ōta-machi to the spa *jinrikisha* available. The spa is situated on the S. E. slope of *Sambe-yama* (4,600 ft.), at an elevation of 1,500 ft. above the sea. It has a Pop. of 109 (36 households) and 3 inns with baths. A beautiful, panoramic view of the surrounding mountain ranges in the distance may be had from the place. The view of groups morning mist, which hover over the valley of the *Gō-gawa* like so many islands in the air, is one of the eight sights of Shigaku. *Mt. Sambe* is noted for its curious shape, resembling three pots standing upside down, and with its foot forming a vast plain of lava. This mountain is dotted with pine-trees and presents a fine view. The foot of the mountain is now used as parade grounds for an Artillery of the 5th. Division of the Army. A beautiful mountain lake called *Ukinuma-ike* is at the S. end of the plain. The ascent of *Sambe-yama* (an extinct volcano) is a delightful excursion from the spa. Near the old crater there are several spots emitting carbonic acid gas. Hot springs gush out from some caves on the slope of *Sambe-yama*, about 0,6 m. from the spa. The flow of water is remarkably abundant, and forms a rivulet of hot water in the valley, by which mills are worked.

see Plate 5,
fig. 1

see Plate 6,
fig. 1

see Plate 1,
fig. 2 and 3

The inner surface of the wooden conduits which convey hot water, gets gradually covered with a thick layer of incrustation, which is to be removed every two or three years. This incrustation is refined and many ornamental articles may be made of it. It is recorded that the spring was formerly lukewarm, but since the great earthquake of Iwami Province, experienced in 1873, its temperature has considerably heightened. The springs are believed to be efficacious for diseases of the stomach and the intestines and disorders of the female genital organs. The spa being situated very high, is one of the most lovely summer resorts in the whole San-in district. Number of visitors: 3,174, the total stays being 10,030 days (1909).

Tombara lies on a highway, which leads from Matsue to Hiroshima, at the N. foot of *Kotobiki-yama* (3,400 ft.), at an elevation of 1,400 ft. above the sea.

Kasubuchi (5 m. S. E. from Shigaku) lies on the N. bank of the *Gō-gawa*, on a highway leading from *Omori* to *Miyoshi*. The water has not yet been used for any purpose.

Yunotsu, 7,2 m. S. W. from *Omori* silver mine, is a sea-port lying between *Hamada* and *Kizuki*. On account of easy means of access from the neighbouring sea-ports, the spa is much resorted to by their inhabitants. The water is believed to be efficacious for diseases of the skin, dyspepsia, and uterus diseases. Number of visitors: 6,540, the total stays being 26,080 days (1909).

Fukumitsu (2 m. S. W. from Yunotsu) lies at an elevation of 300 ft. above the sea. The water is claimed to be beneficial for diseases of the skin, dyspepsia, and uterus diseases.

Arifuku (11 m. N. E. from Hamada) is situated in a gorge, at an elevation of 900 ft. above the sea, along the highway leading Ōta-machi to Hamada. Number of visitors: 9,308, the total stays being 27,893 days (1909).

Mimata is but a poor bathing-place near Arifuku.

Hirose (8 m. S. from *Arashima* station on the San-in Main Line) was found a few years ago and has now two inns and one public bath.

OKAYAMA=KEN

Okayama (240,2 m. from Shimonoseki, in 9 hrs. 10 min.; 89,1 m. from Kōbe, in 3 hrs. 6 min.) is a city with a population of 94,000 (16,500 houses), situated on the lower course of the *Asahi-gawa*, 7,3 m. from its mouth, in the middle of a wide plain, with low hills on the N. side.

Chūgoku Railway. Tsuyama Line: from Okayama to *Tsuyama* 35,3 m. in 2 hrs. 15 min. *Tsuyama* is a large town (Pop. 16,500) on the way to the San-in-dō. From this town on to *Yonago* (61 m.) there is a famous road called "forty times zigzagging path" (*Shijū-magari*), over which *jinrikisha* with two pullers can hardly pass; there are highways also to Tottori (50,8 m.) via *Chizu*, and to *Kurayoshi* (47,7 m.) via *Ningyō-sen* Pass.

Yunogō. 14 m. S. E. from *Tsuyama* (terminus of the private Chūgoku Railway which runs between Okayama and *Tsuyama*) via *Kurashiki* in Bitchū Province. Coming from the E., the spa may be reached from *Wake* or *Kamigōri* station on the San-yō Line. The spa is situated on the W. bank of the *Yoshino-gawa*, at an elevation of 240 ft. above the sea. Springs issue from a fissure of rocks, upon which a bath was constructed. The emission of gas is abundant. On account of the low temperature of water bathers remain usually in it for a very long time, generally for

2-3 hours, sometimes even for 6-7. It is not seldom that some inhabitants of the village sleep in bath through the whole night. The gas being almost nitrogen, the Yunogō Springs furnish a valuable thermal effervescing bath, slightly alkaline and salt, at 35° C. On entering such an effervescent bath below skin temperature there is a momentary chilliness, soon succeeded by a sense of prickling or warmth, due to the collection of innumerable bubbles on the skin. The increasing feeling of warmth soon overcomes the feeling of cold, because the relative heat of the gas more than neutralises the relative coolness of the water.

The discovery of the spring dates back to the Era of Jōkwan (861 A.D.) and the water is said to be efficacious especially for bruise, wounds, fractures, itches, etc. The spring water is barrelled and transported to the neighbouring town, especially to Ōsaka for bathing purpose. There are 1 public bath and 1 inn with baths in the spa. Number of visitors: 11,075, the total stays being 35,860 days (1909).

Maga (27 m. W. from Tsuyama and 5 m. from *Katsuyama*) lies on the W. bank of the upper course of the *Takeda-gawa*, at an elevation of 940 ft. above the sea. During the summer months the citizens of Okayama resort to the spa. Number of visitors: 6,470, the total stays being 14,930 days (1909).

Yubara (6 m. N. from Maga) lies on the upper valley of the *Asatsu-gawa*, surrounded on the E. and W. by mountains, at an elevation of 950 ft. above the sea. It is also a summer resort.

Kaidani (2,5 m. N. from Okayama) lies along the highway leading from Okayama to Tsuyama. A cold spring is located in a narrow gorge.

Takebe, about 0,4 m. S. E. from *Fukuwatari* station on the private Chūgoku Railway (18,8 m. from Okayama, in 1 hr. 53 min.), lies on the W. bank of the *Asahi-gawa*, 209 ft. above the sea and opposite to Fukuwatari. Number of visitors: 112, the total stays being 917 days (1909).

HIROSHIMA-KEN

Tomo (8,5 m. from *Fukuyama*) at the S. E. extremity of a small peninsula, and 5,5 m. by sea from *Onomichi*, is an old port now little frequented by steamers on account of its shallow waters, but which was in olden days a very important voyage station, where are entertained on their passage all the messengers from foreign countries (e. g. Chōsen) to the Court of Japan. There are numerous picturesque islands in the neighbourhood of Tomo, the most famous among them being *Sensui-tō*.

· Plate 62

Fukuzen-ji (Buddhist) and *Nanakusa-jinja* (Shintō) are both noted for fine scenery. Tomo is famed for *hōmei-shu*, a kind of liquor. **Abuto Kwan-on** (2,4 m. S. W. from Tomo), a temple dedicated to Kwan-on, or the Goddess of Mercy, stands on the top of a craggy promontory (*Kwan-on-zaki*). The temple, 28 metres high, is reached by a covered pathway of steep stone steps and commands a view of unsurpassed beauty.

YAMAGUCHI-KEN

Yuda is on the Light Railway running between *Yamaguchi* and *Ogōri*, about 2 m. S. E. from Yamaguchi. The place has a sanatorium of the 5th Division of the Army (Hiroshima). Number of visitors: 7,868, the total stays being 14,599 days (1909).

Kawatana (17 m. N. from *Shimonoseki*) is on the highway, which leads from Shimonoseki to the villages on the N. coast of Nagato Province, about 1,3 m. distant from the sea. Springs issue from a granite formation, the water being radio-active and noted for its efficacy all over the province. Number of visitors: 914, the total stays being 3,428 days (1909).

Fukawa. 15 m. N. from the nearest station on the Ōmine Line (Asa—Ōmine) and 5 m. S. W. from the sea-port of *Senzaki*. The place is on the highway, which leads from *Ozuki* to *Senzaki* via *Tawarayama*, on the bank of a stream, and at an elevation of 200 ft. above the sea. Springs, two in number, were found in the 36th year of the Ōshō Era (1428). *Fukawa-yaki*, a kind of pottery produced in this village, is noted in the neighbourhood.

Tawarayama (22 m. N. from *Ozuki* station on the Sanyō Line) lies on the highway leading to *Senzaki* via *Fukawa-Yumoto*. The place is shut in on every side by mountains and about 1,000 ft. above the sea. The spa is one of the most famous resorts in Nagato, visited mostly by the inhabitants of the Sanyō district. Number of visitors: 9,334, the total stays being 67,999 days (1909).

EHIME-KEN

Matsuyama (5.9 m. from *Takahama*, in 34 min.) is the capital of Ehime Prefecture, with a population of 44,166, and has become well known as a place where a large number of Russian captives were quartered in 1904-5. It has light railways running in different directions to Dōgo, Gunchū, Morimatsu, Yokogawa-mura, and Takahama.

Dōgo is situated about 1 m. N. E. of Matsuyama (light railway and electric tramway), at the foot of a hill, and 35 ft. above the sea. The springs were discovered, according to mythology, by *Ōnamuchi* and *Sukuna-hikona* at a prehistoric time, there being records of the visits paid by early Emperors, such as Keikō, Chūai, Tenchi, etc. The most famous bath-tanks are *Tama-no-yu*, *Kami-no-yu* and *Yoji-yu*, which are in each case built of granite stones, with a three-storied building rising above each. In these upper storied rooms, tea and cakes are served to visitors, while on the third floor of Kami-no-yu is found the so-called "bathers' club," where means are provided for indoor amusements of various kinds. The town of Dōgo contains a permanent population of 2,000, many of whom engage directly or indirectly in the business of inn-keeping. Dōgo has a small public park and temples,—*Isaniwa-jinja* (Shintō), *Ishide-ji* (Buddhist), and the site of Imperial lodgings in former days, which may be visited in the course of a walk. The special products of the place are carved-wood articles (*Fusō-boku*), *Dōgo-Sembei* (a cracknel of wheaten flour), *Yugeta-ame* (glutinous-jelly), etc. Dōgo is the only hot spring in Shikoku Island, and is resorted to by people from all places in the south-western part of Japan, on account of the facility of access. Number of visitors : 45,565, the total stays being 203,152 days (1909),

Meteorological Observations.
Summary.

| Month | Air Temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity |
|---------------------|-----------------|-------|-------|-------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|
| | Mean | | | | Absolute | | | Total | % | Total | Mean |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | |
| For the year, 1911. | | | | | | | | | | | |
| January | 4.8 | 9.01 | 0.38 | 9.52 | 15.7 | -4.8 | 76.3 | 120.48 | 38 | 65.2 | 2.3 |
| February | 5.5 | 10.6 | 0.6 | 9.9 | 14.6 | -4.9 | 71.9 | 163.05 | 53 | 24.8 | 2.8 |
| March | 8.9 | 14.1 | 4.0 | 10.1 | 23.3 | -1.1 | 75.1 | 155.61 | 42 | 123.9 | 3.3 |
| April | 12.4 | 18.5 | 6.8 | 11.7 | 22.7 | 0.4 | 75.6 | 211.85 | 54 | 178.5 | 3.0 |
| May | 16.5 | 22.6 | 11.1 | 11.5 | 26.6 | 4.0 | 79.8 | 198.35 | 46 | 120.4 | 2.2 |
| June | 21.1 | 25.6 | 17.4 | 8.2 | 29.4 | 13.6 | 84.3 | 112.2 | 26 | 315.2 | 2.2 |
| July | 25.42 | 30.21 | 21.42 | 8.79 | 33.5 | 16.4 | 79.7 | 222.45 | 51 | 157.7 | 2.32 |
| August | 25.9 | 31.1 | 21.7 | 9.4 | 33.6 | 19.3 | 82.0 | 212.9 | 52 | 103.8 | 2.2 |
| September | 24.2 | 29.7 | 19.9 | 9.7 | 33.6 | 13.2 | 82.0 | 158.5 | 43 | 310.4 | 2.0 |
| October | 15.9 | 22.4 | 11.0 | 11.4 | 28.6 | 6.5 | 79.0 | 167.4 | 48 | 111.0 | 2.1 |
| November | 12.2 | 18.2 | 7.1 | 11.2 | 22.8 | 0.5 | 70.0 | 134.2 | 44 | 71.7 | 2.3 |
| December | 6.3 | 11.7 | 1.7 | 10.0 | 21.2 | -1.6 | 74.0 | 140.5 | 46 | 93.0 | 2.3 |
| For the year, 1912. | | | | | | | | | | | |
| January | 4.5 | 9.1 | 0.3 | 8.8 | 13.0 | -4.2 | 72.0 | 120.31 | 41 | 50.7 | 2.8 |
| February | 7.5 | 12.6 | 2.7 | 9.9 | 19.2 | -3.2 | 74.0 | 133.39 | 45 | 108.0 | 2.8 |
| March | 8.9 | 14.2 | 3.8 | 10.3 | 22.6 | -0.6 | 75.0 | 172.13 | 46 | 123.6 | 2.6 |
| April | 13.1 | 18.7 | 7.3 | 11.3 | 25.1 | 1.6 | 74.0 | 201.99 | 52 | 150.0 | 3.0 |
| May | 17.0 | 23.0 | 10.8 | 12.2 | 28.8 | 5.1 | 75.0 | 235.35 | 55 | 54.9 | 2.9 |
| June | 21.3 | 27.0 | 16.2 | 10.8 | 30.4 | 12.3 | 78.0 | 209.34 | 49 | 84.3 | 2.2 |
| July | 25.2 | 29.8 | 21.7 | 8.1 | 33.6 | 16.2 | 82.0 | 147.92 | 34 | 191.8 | 2.2 |
| August | 26.0 | 31.8 | 21.5 | 10.3 | 33.9 | 18.4 | 79.7 | 263.6 | 64 | 35.0 | 2.0 |
| September | 21.13 | 26.1 | 16.9 | 9.1 | 32.7 | 10.8 | 83.9 | 127.2 | 31 | 299.1 | 1.74 |
| October | 15.9 | 22.2 | 10.3 | 12.0 | 26.6 | 5.7 | 82.0 | 176.67 | 50 | 122.7 | 1.9 |
| November | 10.0 | 15.8 | 4.9 | 10.9 | 22.4 | 1.2 | 78.0 | 142.84 | 46 | 75.1 | 2.0 |
| December | 6.8 | 12.2 | 2.2 | 10.0 | 20.1 | -1.9 | 77.0 | 112.38 | 37 | 111.9 | 2.3 |

| Month | Air Temperature | | | | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity |
|---------------------|-----------------|------|-------|-------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|
| | Mean | | | | Absolute | | | Total | % | Total | Mean |
| | Mean | Max. | Min. | Range | Max. | Min. | | | | | |
| For the year, 1913. | | | | | | | | | | | |
| January | 4.0 | 9.5 | -0.4 | 9.9 | 14.2 | -3.5 | 74.0 | 120.1 | 41 | 51.1 | 2.7 |
| February | 4.7 | 9.8 | 0.0 | 9.8 | 14.7 | -8.3 | 71.0 | 120.0 | 39 | 47.3 | 3.3 |
| March | 9.7 | 12.4 | 1.0 | 11.5 | 19.1 | -3.5 | 67.0 | 184.8 | 50 | 40.9 | 3.0 |
| April | 14.4 | 20.1 | 9.4 | 10.8 | 26.8 | 2.1 | 70.0 | 160.2 | 41 | 136.7 | 2.4 |
| May | 19.5 | 22.8 | 10.5 | 12.3 | 26.5 | 5.3 | 77.0 | 228.5 | 53 | 181.7 | 2.4 |
| June | 18.8 | 25.4 | 16.7 | 8.7 | 32.0 | 11.5 | 81.0 | 120.39 | 28 | — | 2.2 |
| July | 24.4 | 29.5 | 19.50 | 9.94 | 33.8 | 14.3 | 77.0 | 250.8 | 57 | 69.0 | 2.4 |
| August | 25.0 | 30.3 | 20.7 | 9.7 | 33.5 | 16.6 | 81.0 | 227.39 | 55 | — | 2.0 |
| September | 21.0 | 26.6 | 10.1 | 10.6 | 31.3 | 10.3 | 70.0 | 178.53 | 48 | 68.5 | 1.8 |
| October | 16.6 | 23.4 | 11.5 | 12.0 | 28.5 | 5.0 | 79.8 | 181.82 | 52 | 47.3 | 2.0 |
| November | 11.5 | 18.5 | 6.2 | 12.3 | 24.6 | 1.9 | 78.0 | 142.8 | 46 | 95.9 | 2.0 |
| December | 6.6 | 12.0 | 2.4 | 9.7 | 19.4 | -2.1 | 76.0 | 76.65 | 25 | 118.9 | 3.1 |

FUKUOKA-KEN

Futsukaichi (56.5 m. from Moji, in 2 hrs. 29 min.) is 0.3 m. from a hot spring, **Musashi-Onsen** (Inn, *Enju-kwan*), to the W. of which rises the hill, *Tempai-zan*, where *Michizane* is said to have prayed in the direction of Kyōto. Halfway up (1.4 m.) the slope is found a waterfall called *Ryū-ō-no-taki*, and beside it a huge stone, all associated with the memory of Michizane. Number of visitors: **32,532**, the total stays being **221,975** days (1909). The springs are said to be efficacious for rheumatism.

Meteorological Observations.

Summary.

For the year, 1911.

| Month | Air Temperature | | | Relative Humidity | No. of hours with Sunshine | | Amount of Precipitation | Wind Velocity |
|-----------|-----------------|----------|------|-------------------|----------------------------|----|-------------------------|---------------|
| | Mean | Absolute | | | Total | % | Total | Mean |
| | | Max. | Min. | | | | | |
| January | 4.90 | 18.1 | -2.8 | 75.3 | 88.2 | 28 | 69.4 | 3.84 |
| February | 5.23 | 15.5 | -3.6 | 74.6 | 94.95 | 31 | 33.0 | 4.04 |
| March | 9.01 | 21.7 | -0.2 | 76.9 | 127.54 | 34 | 208.4 | 4.81 |
| April | 12.37 | 24.4 | 0.2 | 75.4 | 185.29 | 48 | 122.0 | 4.36 |
| May | 17.11 | 26.2 | 2.4 | 78.8 | 173.93 | 41 | 94.9 | 3.27 |
| June | 21.39 | 31.7 | 14.0 | 85.3 | 123.41 | 29 | 315.8 | 3.49 |
| July | 25.47 | 33.5 | 17.6 | 82.4 | 181.81 | 42 | 245.9 | 4.37 |
| August | 26.22 | 34.4 | 19.9 | 83.7 | 215.08 | 52 | 77.6 | 3.83 |
| September | 23.96 | 32.3 | 12.4 | 85.3 | 152.82 | 41 | 283.9 | 3.13 |
| October | 15.92 | 26.1 | 6.4 | 79.3 | 167.11 | 48 | 110.0 | 3.64 |
| November | 12.07 | 23.2 | 0.6 | 79.6 | 137.43 | 44 | 129.2 | 3.86 |
| December | 6.27 | 17.0 | 0.2 | 75.7 | 95.49 | 31 | 58.1 | 3.51 |

Hainu-zuka or *Hain-zuka* (77.4 m. from Moji, in 3 hrs. 28 min.) is the chief station leading to the basin of the River Yabe. Tram-car from the station, on the one hand to *Okawa-machi* at the mouth of the river, and on the other to *Fukushima-machi* (8.3 m.).

Funagoya-Onsen is a mineral spring on the River Yabe, 2.4 m. from the station. The water is said to be efficacious for anaemia and catarrh of stomach and intestines. *Hyūgami-iwa*, a highly picturesque spot believed by some people to be even superior to the famous *Yabakei* in Buzen, is

22 m. from the station on the upper course of the River Yabe, and between the two villages of Ōbuchi and Yabe. Number of visitors: 52,379, the total stays being 69,676 days (1909).

ŌITA-KEN

Beppu (81.9 m. from Moji, in 4 hrs. 17 min.). It is situated on the Bay of Beppu, with the beautiful hill of Tsurumi behind, and is famous throughout Japan on account of its hot springs (Pop. 14,045, households, 3,120). It enjoys ample steamship communication with all inland sea-ports, and with Kōbe and Osake (services maintained by the Ōsaka Shōsen Kwaisha). The town is built on ground undermined by volcanic vapour and hot water. The springs are alkaline and carbonated, containing iron, while some contain much sulphur, and are regard as highly efficacious in various complaints. On the beach also there are springs where people bathe, half burying their bodies in sand. In the town are half-a-dozen public bath-tanks, while the more important inns own private baths.

Baths in the Neighbourhood: **Kwankaiji-Onsen** (2 m. to the W. of Beppu), high up on the side of *Tsurumi-dake*, with a fine view of the bay (Inns: *Matsu-ya*, *Sakamoto-ya*); about 0.2 m. or 0.3 m. higher up is *Ue-no-tano-yu*, on the N. side of which is a valley (called *Jigoku*) filled with sulphureous steam; 1.6 m. N. E. of Kwankaiji is **Hotta-Onsen** (Inns: *Kanata-ya*, *Hama-ya*); at the foot of *Yūfu-dake* (called Bungo-Fuji), 7.3 m. from Hotta-Onsen via Kusu-Kaidō (Highway to Kusu), are Takemoto and three other baths, while going 3.7 m. the other way (i.e. towards the N.) from Hotta we come to **Myōban-Onsen** (Inns: *Okamoto-ya*, *Ebisu-ya*); about 0.7 m. from Myōban-Onsen we come to **Bōzu-Jigoku** (a geyser of boiling mud) and again 0.7 m. farther on from it, **Umi-Jigoku**, a boiling blue pond covered with steam; 0.5 m. from Umi-Jigoku is **Kannawa-Onsen**, a popular resort having a famed vapour-bath (Inns: *Fuji-ya*, *Yorozu-ya*, *Tokiwa-ya*); 1 m. N. E. of Kannawa is **Shibaseki-Onsen**; 1.2 m. again from Shibaseki, **Chi-no-ike-Jigoku** or "Blood-coloured boiling pond". The shore as well as the bottom of the pool is covered with ochreous sinter deposits, composed chiefly of iron oxide and clay (see Part II, p. 153), which are at present used only for dyeing towel, bath-dress and other cotton fabrics, just for the object of *souvenir* sold to visitors. The villagers were, however, quite ignorant of utilizing the sinter deposits for a valuable form of medical baths (viz., Mineral-Mud Bath), till Dr. Ishizu, expert of the Tōkyō Imperial Hygienic Laboratory, taught them for the first time the utilization of the mud for that purpose in Aug., 1909. Since then the neighbouring inhabitants have begun to experience themselves the value of this kind of bath.

The pond may be only one place in Japan, which can supply a sufficient quantity of mineral sinter for preparing mud-baths. The locality affords also facility for providing with hot water douche, hot waterfalls and inhalation-house. It is hoped that some suitable arrangements will be made to employ the mud in this valuable form of application and also providing necessary accommodation for bathers. About 0.5 m. from Chi-no-ike, we reach the sea-shore town of Okoshi, where is **Kamegawa-Onsen** (Inn: *Muro-ya*); from Okoshi to Beppu is 4.4 m.

Meteorological Observations.

Summary.

| Month | Air Temperature | | | | | Relative Humidity | Amount of Precipitation | Wind Velocity |
|---------------------|-----------------|------|------|----------|------|-------------------|-------------------------|---------------|
| | Mean | | | Absolute | | | | |
| | Mean | Max. | Min. | Max. | Min. | | Total | Mean |
| For the year, 1911. | | | | | | | | |
| January | 6,9 | 9,7 | -0,2 | 18,3 | -6,0 | 66,5 | 69,2 | 1,7 |
| February | 7,7 | 10,6 | 0,7 | 15,8 | -4,5 | 64,6 | 17,0 | 1,5 |
| March | 10,7 | 13,4 | 4,3 | 22,3 | -4,2 | 70,7 | 162,7 | 2,0 |
| April | 15,7 | 19,2 | 10,0 | 29,5 | 5,7 | 64,0 | 80,4 | 1,2 |
| May | 19,3 | 22,3 | 12,8 | 27,5 | 5,6 | 72,7 | 250,5 | 0,9 |
| June | 22,7 | 26,7 | 15,5 | 30,8 | 10,2 | 78,2 | 445,3 | 0,7 |
| July | 27,1 | 28,7 | 19,7 | 32,7 | 11,8 | 76,6 | 68,8 | 1,1 |
| August | 27,0 | 28,7 | 14,5 | 31,6 | 15,5 | 75,2 | 153,8 | 1,3 |
| September | 26,1 | 27,1 | 21,6 | 30,0 | 15,0 | 75,0 | 311,6 | 1,4 |

| Month | Air Temperature | | | | | Relative Humidity | Amount of Precipitation | Wind Velocity |
|---------------------|-----------------|------|------|----------|------|-------------------|-------------------------|---------------|
| | Mean | | | Absolute | | | Total | Mean |
| | Mean | Max. | Min. | Max. | Min. | | | |
| For the year, 1912. | | | | | | | | |
| September | 23.6 | 25.9 | 18.7 | 30.6 | 12.4 | 70.3 | 318.9 | 1.1 |
| October | 19.6 | 22.0 | 13.3 | 29.2 | 8.3 | 73.0 | 33.4 | 1.6 |
| November | 14.1 | 15.9 | 7.9 | 22.1 | 2.0 | 76.3 | 40.5 | 3.1 |
| December | 11.0 | 13.1 | 4.9 | 22.3 | 0.1 | 78.0 | 70.6 | 3.0 |
| For the year, 1913. | | | | | | | | |
| January | 7.8 | 10.8 | 2.3 | 16.8 | -0.9 | 57.5 | 70.4 | 2.8 |
| February | 7.8 | 10.6 | 2.4 | 13.7 | -0.3 | 60.4 | 19.0 | 2.1 |
| March | 10.1 | 13.2 | 3.8 | 21.0 | 0.0 | 52.0 | 38.5 | 2.1 |
| April | 15.9 | 19.0 | 10.5 | 24.6 | 3.8 | 70.0 | 136.2 | 1.4 |
| May | 19.7 | 22.1 | 12.3 | 26.9 | 5.9 | 61.6 | 121.3 | 1.7 |
| June | 22.7 | 25.2 | 13.3 | 32.0 | 10.0 | 76.5 | 220.6 | 2.2 |

Summary.

| Spa | Number of Public baths | Number of Vapour baths | Number of Hot Waterfalls | Number of Sand baths | Number of Inns | Number of Visitors in 1909 | Their total stays in days |
|--------------------|------------------------|------------------------|--------------------------|----------------------|----------------|----------------------------|---------------------------|
| Beppu and Hamawaki | 8 | — | — | 5 | 202 | 108,192 | 540,947 |
| Kwankaiji | 1 | 1 | 1 | — | 9 | 43,731 | — |
| Itoha | 1 | 1 | 1 | — | 4 | 9,805 | 40,652 |
| Myōban | 1 | — | — | — | 17 | 5,558 | 36,937 |
| Kannawa | 3 | 2 | — | — | 15 | 12,968 | 35,356 |
| Shibaseki | 1 | 1 | 1 | — | 2 | 14,650 | — |
| Kamegawa | 2 | 1 | 1 | 1 | 26 | 63,942 | — |

KUMAMOTO-KEN

Hot Springs of Aso.

Aso-san. A celebrated volcano, 26,8 m. E. of Kumamoto, and situated at the intersection of two volcanic ridges, one beginning in the Loochoo Islands and extending as far as the Japan Sea, via *Kaimon-dake*, *Sakura-jima*, *Kirishima-yama*, and *Aso-san*; while the other ridge runs E. to W. from the N. of Shikoku towards Amakusa, via *Yufu-dake* (in Bungo), *Hisazumi-dake*, *Aso-san*, *Kimbō-san*, and *Unzen-dake* (the last in Shimabara Peninsula). This volcano, being thus at the intersection of two ridges, is the most active crater of all (some of the craters mentioned above, e. g. *Kimbō-san*, near Kumamoto, are already entirely extinct). Aso-san is a general name comprising five volcanic peaks, the active one being called *Naka-dake*, while the other four (extinct) are (1) *Kishima-dake*, to the N. W. of Naka-dake, (2) *Eboshi-dake*, S. of Kishima-dake, (3) *Taka-dake* (5,500 ft.) highest of all five, E. of Naka-dake, (4) *Neko-dake*, a jagged, saw-like peak, E. of Taka-dake. To the S. and N. of these five peaks lie two plains, the N. one being called *Aso-dani* (1,620 ft.) and S. *Nangō-dani* (1,470 ft.), containing altogether three towns and eleven villages and a population of 40,700.

Ascent: from Kumamoto to *O-tsu* 12.2 m. by light railway; from Ōtsu to *Tateno* (Pop. 3,000), about 6.5 m.; from Tateno two roads branch off, (1) one along the *Kuro-kawa* or "Black River," an affluent of the *Shira-kawa*, leading to *Aso-dani*, (from Tateno to Miyaji 12 m.), while (2) the other leads by *Shira-kawa* into *Nangō-dani*, the plain to the S. of Aso-san. On both paths *jinrikishas* are available as far as the foot of the five peaks. If we take the 1st road, we soon come to *Sugaru-ga-taki*, a majestic waterfall plunging from a high precipitous volcanic rock covered by a

thick grove of large trees; now passing on by a gradual ascent we reach the highest point across the ridge, *Futae-no-tōge*, when suddenly we realize that we are standing on the edge of an almost perpendicular precipice, and that there lies before us the plain of *Aso* and the five peaks; descending the precipice we soon come to the town of *Miyaji* (Pop. 3,900). At *Miyaji* is an ancient temple (1,700 years old), *Aso-jinja* (Shintō) which is dedicated to the god, *Take-iwatatsu-no-Mikoto*.

At *Miyaji* we may stop over-night. From *Miyaji*, we come to *Bōjū-mura*, which is at the foot of *Naka-dake*, the active volcano. Here begins the ascent, which is by no means steep, and from half-way up we find all signs of vegetation gone, and everywhere ejected lava stones; and coming to the lip of the crater, we see the crater like a huge blacksmith's bellows sending up columns of sulphur smoke, and loud detonations. From the rest-house at the base of the cone, the climb to the lip and back again will occupy nearly an hour. Now on the return journey, we take the other of the two roads to *Tateno* and come first to *Yu-no-tani*, where there is a geyser of red mud and burning water, then to *Taru-tama* and *Jigoku* hot springs; from here going W. we come to *Tochinoki* hot springs, a well-known bathing-place, and thence along the River *Shira-kawa* we come to *Tateno*, whence we started by the 1st road. *Tochinoki* hot springs are said to be found in the 4th year of the Kwambun Era (1665). Number of visitors (*Tochinoki*): 10,412, the total stays being 29,760 days (1909).

Yamaga. 17 m. N. from Kumamoto and 11 m. N. from *Ueki* station on the Kagoshima Main Line, *jinrikisha* available. The spa town (Pop. 7,541) lies on the highway, which leads from *Ueki* to *Kurume* via *Fukushima* and at the N. end of the plain of *Kikuchi* extending to the foot of *Aso-san*. The spa is much resorted to by the inhabitants of the two provinces of *Higo* and *Chikugo*. The water is believed to be efficacious for gout and syphilis. Number of visitors: 34,153, the total stays being 1,039,847 days (1909).

Hinagu-Onsen, a carbonated spring, is 6 m. from *Yatsushiro* (fare, 50 *sen* by *jinrikisha*). It is a favourite resort of people from all parts of *Kyūshū* (Inns: *Kimpa-rō*, *Yanagi-ya*). *Kōda-yaki*, produced at *Kōda* between *Yatsushiro* and *Hinagu*, is a kind of faience, introduced by Korean potters.

Number of visitors: 25,146, the total stays being 262,494 days (1909).

NAGASAKI-KEN

Shimabara Peninsula, known on account of its famous volcano, *Unzen-dake*, which is situated almost in the center of the peninsula. On the W. coast is situated the port of *Obama*, on the E. the historical town of *Shimabara*, and on the S. the port of *Kuchinotsu*. Highways from *Nagasaki* lead, the one via *Uki* and *Chijiwa* to *Obama*, and the other via *Isahaya* (by Rys.) and along the coast of *Ariake-no-umi* to *Shimabara* (a part of the way by light railway). Steamer Passage from *Mogi Port* to *Obama* costs 60 *sen*.

Obama.

Obama is situated on the west coast of the *Shimabara Peninsula* at a distance of 20 m. across the *Chijiwa Sea* from *Mogi*. There are several saline springs of intermittent nature, possessing great efficacy for rheumatic complaints. The *Ikkaku-rō Hotel* has good foreign accommodation and is well patronised throughout the year; visitors to *Unzen* find it advantageously situated for passing a night or obtaining meals *en route* to or from the hill resort. There are no less than fifty inns for the accommodation of Japanese. The baths are separated from the inns, being mostly on a rocky beach. The beach is reserved for recreative purposes. *Obama* has a good water supply, water-works having been constructed there in 1907.

The climate in winter is warm and attracts many visitors. Within a short distance of the town is a white sandy beach, with pine groves extending several hundred yards. The beach is excellent for sea-bathing, and shelters are provided for the convenience of bathers. A mile and a half from *Obama* on the road to *Unzen* is *Binkushi*, and one mile higher up is *Kago-tateba*, which in olden days was a station at which the bearers of *kago* (palanquins) of nobles stopped to rest, hence the name. The place is now marked by a house. This point is half-way to *Unzen* and commands a splendid view of *Ariake Bay*, over the *Aino isthmus*, the lovely hamlets of *Obama*, *Chijiwa*, *Aino*, *Moriyama*, and *Uki*, and the *Chijiwa Sea* at the foot of the mountain. Number of visitors: 19,387 (natives), the total stays being 164,560 days (1909); 899 (foreigners), the total stays being 6,347

see Plate 4
fig. 1

days (1909)—22,638 (natives), the total stays being 180,410 days (1913); 1,834 (foreigners), the total stays being 28,470 days (1913). Hotel: *Ikkaku-rō*; Inn: *Tsuta-ya*.

Unzen Resort.

How to reach Unzen. There are two routes from Nagasaki, one by sea, and the other overland. In the former, *jinrikisha* is to be taken as far as *Mogi*, via *Tagami* (fare, ¥ 1,20 with two coolies), then to Obama by steamer (fare, 1st class 90 *sen*; additional charges:—5 *sen* transit tax, 4 *sen* pier dues, and 5 *sen* sampan fare). Two direct services are available daily. From Obama the ascent to Unzen, a distance of seven miles, can be made in a chair carried by four coolies (fare, ¥ 3,00), a *jinrikisha* drawn by two coolies (fare, ¥ 2,00), or by horse carriage (fare, ¥ 3,00). The road is very good, and even ladies can easily walk up to Unzen. The overland journey is made by rail from Nagasaki to *Isahaya* on the Government Line (fare including transit tax, 1st class 88 *sen*, 2nd class 53 *sen*), thence to *Aino* by the Shimabara Light Railway, a private line (fare, 1st class 44 *sen*, 2nd class 27 *sen*). From *Aino*, *jinrikisha* (fare, 45 *sen*), *basha*, or motor-bus (fare, 35 *sen*) is taken to *Chijiwa*, via the *Chijiwa* slope, and then on to Obama, via *Tomitsu-zaka*, giving a view of the beautiful coast scenery. The fare is ¥ 1 by *jinrikisha* and by motor-bus 70 *sen*.

Unzen can be reached from *Chijiwa* direct, along the route that traverses the beautiful valley of the River *Chijiwa*. The ascent is shorter than any other, and can be made by chair or on horseback.

Time required for journey. From Nagasaki to *Aino*, via *Isahaya*, where cars are changed to the Shimabara Railway, the time required is two hrs. by rail, and from *Aino* to the *Chijiwa* Hotel is half-an-hour's journey by motor-bus. After resting at the hotel, Unzen can be reached by chair at noon. From *Aino* to Obama direct by motor-bus takes an hour; the journey from Obama to Unzen, by chair, occupies two and a half hours. Only one motor-bus, accommodating five passengers, is now running between *Aino* and Obama, but another will be ready for service by the summer. If the sea-route is preferred one has to start from Nagasaki at 7 a.m. by *jinrikisha* (two men) to *Mogi*; from the latter place Obama is reached at 9,30 a.m. by steamer. After a rest at the *Ikkaku-rō* Hotel, Unzen can be reached at noon by chair (four coolies) or *jinrikisha* (two coolies).

Unzen, the joint name of the three hamlets of *Furu-yu*, *Shin-yu*, and *Ko-jigoku*, is a favourite summer resort of European residents from the neighbouring regions, as well as from China ports and Philippines. The place being situated high up (2,400 ft. above sea-level) on a mountain is noted for its cool, bracing atmosphere and splendid scenery,—the hot mineral springs on account of their efficacy in many kinds of disease being a special attraction. From experience, Japanese regards a course of these upper springs at Unzen as necessary for making a complete recovery after passing through a preliminary course at the Obama baths.

Climate. According to the observations taken by the Nagasaki Observatory, the climatic conditions of Obama and Unzen compare with Nagasaki and some other places as follows:—

Spring (March, April, May).

| | Temperature (Centigrade) | Humidity (per cent) | Rainy days |
|----------|--------------------------|---------------------|------------|
| Unzen | 8,9° | 80 | 37 |
| Obama | 13,9° | 78 | 28 |
| Nagasaki | 13,1° | 74 | 28 |

Summer (June, July, August).

| | | | |
|----------|-------|----|----|
| Unzen | 19,9° | 90 | 54 |
| Obama | 24,6° | 83 | 39 |
| Nagasaki | 23,8° | 82 | 49 |
| Hakone | 20,5° | 83 | — |
| Hao | 20,4° | — | 55 |

Autumn (September, October, November).

| | | | |
|----------|-------|----|----|
| Unzen | 12,8° | 83 | 31 |
| Obama | 17,8° | 76 | 27 |
| Nagasaki | 17,2° | 70 | 28 |
| Hakone | 14,3° | 81 | — |
| Hao | 12,8° | — | 25 |

Winter (December, January, February).

| | | | |
|----------|------|----|----|
| Unzen | 2.4° | 82 | 12 |
| Obama | 7.5° | 73 | |
| Nagasaki | 6.6° | 67 | 42 |
| Hakone | 2.7° | 70 | |

Ascent. From Obama to *Bingushi*, via *Yunosaki*, 1 m.; from Bingushi to *Kago-tateba*, 1 m.; from Kago-tateba to *Fuda-no-hara*, 1.8 m.; from Fuda-no-hara to Unzen, 1.2 m. Between Fuda-no-hara and Unzen, the path—famous for its fine views—lies on a wide plateau occupied by rice-fields and a few remains of *Mammyō-ji* temples, which at one time numbered three hundred, all occupied by Buddhist monks. They were all destroyed by fire in the Shimabara rebellion, having been attacked by the Christians who had fortified themselves in the Hara Castle at Arima. Only the fragments of a large *torii* (gateway) are now to be seen. The road divides here, the left one leading to *Shin-yu* ('New Spring'), Unzen, and the other to *Ko-Jigoku*, which is half a mile distant. These roads are quite flat.

Ko-Jigoku. The bath-house at Ko-Jigoku is supplied from a spring of boiling water which is worth seeing. An attractive waterfall, called *Issaikyō-no-taki*, about twenty feet high, is near here, being in the stream which has its source in the springs Shin-yu and Ko-Jigoku.

Unzen Park. The Park occupies a tract of land with stretches of hills lying between the peaks *Kinugasa-yama* and *Ya-take*, the whole covering an area of 200,000 square yards. Many geysers, sending up boiling water to some height, exist on the E. side of the Park. Paths have been made to render the geysers accessible, and benches have been placed near them for the convenience of visitors. The Park is aglow with azalea blossoms in May. Being cooler than *Ikao* and *Hakone*, Unzen forms an ideal summer resort in Kyūshū. There are two tennis-courts, both open to the public on payment of regular fees, and attached to them is a pleasant pavilion. A large meeting hall, 364 square yards, is in course of construction and will be available for athletic purposes in wet weather.

Shin-yu ('New spring') and **Furu-yu** ('Old spring'). The Shin-yu bath-house is located in the Park and is connected with the spring at the foot of *Ya-take*. It has four bath-tubs, which are clean and always overflowing. A post-office is opened every year on June 1st and closed on October 31st. It transacts telegraphic, telephonic, and money-order services, etc., and distributes mails twice daily.

Furu-yu is situated 600 yards away from Shin-yu and the springs are similar to those at the latter place. The water, being acid hydrogen sulphide spring, containing a small quantity of iron oxide and alumina, is said to be efficacious, especially for rheumatism, dyspepsia, disorders of the female genital organs, syphilis, etc. Number of visitors: 6,103 (natives), the total stays being 10,075 days (1909); 743 (foreigners), the total stays being 7,804 days (1909)—6,275 (natives), the total stays being 18,825 days (1913); 1,274 (foreigners), the total stays being 36,220 days (1913). There are about a dozen inns.

The golf links covering 208,000 sq. yards are open to the public. A pavilion will shortly be built with toilet, dining, bar, and dressing-rooms. While the course cannot yet be described as in perfect condition, it is hoped that it will be made one of the best in the Far East before long, as no pains are being spared by the authorities to improve it. The site is naturally suited for the pastime and is very attractive, especially in May when the azaleas are in bloom.

Unzen is situated in a hollow of two peaks, *Fugen-dake* and *Myōken-dake*; while in their neighbourhood are several other smaller peaks, such as *Eboshi-dake*, *Azuma-dake*, *Mai-dake*, *Ro-no-ki-dake*, *Mai-yama*, *No-dake*, *Kinugasa-yama*, *Taka-dake*, etc.

Nita and Fugen Peak. Half a mile up the steep path from the golf links is the little plateau of *Nita*, commanding a fine view of the coast of *Higo*. Two miles higher one reaches the summit of *Fugen-dake*, (4,800 ft. above the sea-level). At the summit is a large perpendicular rock about 50 ft. high, on the N. side of which icicles may be found hanging as early as November. From here may be enjoyed an extensive panorama of indescribable beauty (which will repay all the labour of climbing), embracing the hills of *Hizen* and the plains of *Chikugo*, as well as the active volcano of *Aso* in *Higo* and *Kirishima-yama* on the boundary of *Ōsumi* and *Hyūga*, and the islands of *Amakusa*. The prospect also takes in on the other side the promontories jutting forth outside of

Nagasaki harbour, and the famous Gotō group of islands at a further distance. Two miles higher one reaches the summit of *Fugen-dake*. In autumn peak looks as if covered with brocade, the effect of the maples, and evergreen trees. For more than half the way the ascent can be made either on horseback or in a chair carried by four coolies. Near the top are found several large caves called *Kaza-ana* ('Wind holes'), inside which hang long icicles like swords, the temperature never rising above 2° C. even in mid-summer. The caves are used for storing silk-worm eggs.

Myōken-dake may be reached in 2 hrs. from *Fugen-dake*,—the path leading partly through brushwood, till an old extinct crater is reached and crossed (the path through the crater lying over a large volcanic rock), the rest of the way thence to the summit being a very precipitous ascent. The panorama from the summit is similar to that from *Fugen-dake*.

Chijiwa. (Hotel: *Chijiwa Hotel*). It is situated about 6 m. from Furu-yu, Unzen. The road from it to Unzen is now steep, but within the next three years it will be improved so as to allow the passage of vehicles. Chijiwa is chiefly noted for its sea-bathing, on a fine sandy beach sheltered by a beautiful ridge of pine-trees. Good pedestrians will enjoy making an excursion to the town of Shimabara, 12.2 m. down the coast on the other side where Obama is situated. The road lies at first between *Unzen-dake* on the left and *Taka-dake* on the right, then down to a valley, passing by a dried-up lake, now ascending, now descending by a path amidst boulders and then through a forest of pines and camphor-trees, till *Mino-kaiva* (a hamlet 5 m. from Unzen) is reached. From here the road is less steep and soon becomes quite level, and at *Nakakobe* we come in sight of *Mai-yama*, which rises like an immense wall, screening the town of Shimabara from the volcanic craters behind.

Michi-no-o hot spring (5 m. by rail from Nagasaki, in 15 min.) is said to be efficacious in rheumatic complaints and stomach troubles (private bath provided). Near the station is the village of *Urakami*, known in connection with the history of Roman Catholicism in Japan. This and most of the neighbouring hamlets are inhabited by Roman Catholics, who were found in large numbers at the time of the Restoration (1868). Christianity was never entirely eradicated here, nor from some other places in *Kyūshū*, notwithstanding ruthless persecutions continued for several centuries. Number of visitors: 245, the total stays being 332 days (1909).

SAGA=KEN

Karatsu hot spring (25 m. from *Kubota*, in 1 hr. 50 min.), a town situated at the mouth of the *Matsuura-gawa*, faces the Bay of Karatsu. In ancient times this place as well as Hakata assumed importance in connection with intercourse with Chōsen. It is now a special exporting port (chief export, coal), with a deep anchorage at *Nishi-Karatsu*, nearly 1 m. W. of the town. The place is also noted for its pottery (called *Karatsu-yaki*).

see Plate 70

Karatsu, formerly the castle-town belonging to Daimyō *Ogasawara*, was called *Bukaku-jō*, or "Flying-Stork Castle," a pretty name which arose from the fancied resemblance of its E. and W. beaches, covered with pines, to the outstretched wings of the bird, the castle itself corresponding to its long neck and crest. The E. beach, called *Niji-no-Matsubara*, may be reached by a tramway (2 m.); the castle grounds have been turned into a public park; and the W. beach is a famous bathing-resort. S. of the latter rises a hill called "*Hirefuru-yama of Lady Matsuura-Sayo-hime*."

see Plate 71

Nanatsu-gama, 2.4 m. from *Karatsu*, is noted on account of its curious shaped rocks of basaltic formation. Seven caverns exist at the lower part of the rocks, looking like so many ovens, hence the name.

Number of bathers (*Karatsu*): 14,001, the total stays being 210,015 days (1909). Inns: *Hakata-ya*, *Shin-iwai-ya*, *Kaihin-in*, *Kimpa-rō*, etc.

see Plate 72

Takeo (33.1 m. from *Tosu*, in 1 hr. 13 min.) is well known for its hot mineral waters (carbonated spring, containing alkali). It is picturesquely situated, being surrounded on three sides by well wooded hills. The springs are said to be efficacious for dyspepsia, haemorrhoids, rheumatism, syphilis and diseases of the skin.

Ureshi-no-Onsen, also a carbonated hot spring, is 7.3 m. from *Takeo*. Inns: *Tōkyō-ya*, *Tōyō-kan*, *Hillside Hotel*. Efficacious for rheumatism, neuralgia, syphilis and diseases of the skin.

Arita (42.2 m. from *Tosu*, in 1 hr. 47 min. and junction for the Imari Branch Line) is situated in a narrow valley surrounded by wooded hills, being noted for its pottery made with the clay

brought from a neighbouring hill, *Izumi-yama*. Most of the inhabitants (6,000) are engaged directly or indirectly in porcelain manufacture. The most famous of the manufactories is *Kōran-sha*, producing annually wares worth ¥ 175,000. The art was first introduced here in 1592 by a Korean artist, brought over by *Nabeshima*, Daimyō of Hizen, while the distinctive process of decorating with vitrifiable enamels was later taught by a Korean artist domiciled in Nagasaki, and was afterwards much improved by native artists. The wares soon attracted the notice of Dutch traders at *Dejima*, Nagasaki, who exported many pieces. The Arita pottery, however, attracted the general notice of Europeans for the first time at the Philadelphia Exposition in 1876. At present large quantities are exported, in some years valued at ¥ 700,000.

Imari (by Imari Branch Line from Arita to Imari, 8.1 m., in 33 min.) is a sea-port town, with 4,200 inhabitants. It is known as the export place of Arita porcelain (hence these wares are popularly called *Imari-yaki*). Since the opening of the railway, however, the prosperity of Imari has much diminished.

| Spa | Elevation above the sea | Population | Number of Inns | Number of Visitors | The total Stays in days |
|----------|-------------------------|------------|----------------|--------------------|-------------------------|
| Takeo | about 100 ft. | 3,200 | 35 | 32,349 | 101,261 (1909) |
| Ureshino | about 400 ft. | 1,200 | 18 | 10,851 | 47,794 .. |

KAGOSHIMA-KEN

Anraku. 5 m. S. E. from *Makizono* station and also 5 m. N. from *Kokubu* in Satsuma Province, *jirikisha* and omnibuses available. Here several cold springs gush out of the banks of a stream, the water being mostly ferruginous and carbonated. Number of visitors: 23,000, the total stays being 11,500 days (1909). *Myōken-yu* Spa is about 0.5 m. distant from Anraku.

Hot Springs on Mt. Kirishima.

Myōban-Onsen. From the *Makizono* station on the Kagoshima Line to *Kami-nakatsu-mura* 4 m., *jirikisha* and omnibuses available, thence 5 m. by *kago* or on horseback. The spa, 741 ft. above the sea, lies on the slope of *Karakuni-dake* (5,610 ft.), one of the peaks of *Kirishima-yama*, an active volcano, commanding an extensive view including the Bay of Kagoshima. The ever-changing view of *Sakura-jima* on misty days never fails to be attractive. Here the thermometer never registers higher than 80° F. even on the hottest days in summer. Number of visitors: 4,850, the total stays being 32,125 days (1909).

Eino-o-Onsen. The spa is within only a few minutes' walk from Myōban-Onsen. Water of high temperature conveyed by wooden pipes is mixed with cool water and pours itself into bath-tubs, forming waterfalls, which are called "*Utase-yu*" meaning 'hot spring for massage.' The flow of water is exceptionally abundant.

Eino-o Spa is at the highest elevation of all hot springs on Mt. Kirishima. The panoramic view over a valley and the beautiful Bay of Kagoshima is delightful. Number of visitors: 1,024, the total stays being 4,470 days (1909).

Iwō-dani-Onsen. The spa lies W. of Myōban-Onsen. The flow of water is so considerable that it overflows into a stream of hot water. There are vapour baths and hot waterfalls in the place. Number of visitors: 5,865, the total stays being 28,800 days (1909).

All these hot springs mentioned above issue from crevices of an old crater on the S. of a volcanic peak of *Onami-ike* on Mt. Kirishima.

Ibusuki is situated at the S. end of Satsuma Province, facing the entrance of Kagoshima Bay. This district is distinguished by the occurrence of many hot beach baths on the sea-shore, just like *Beppu* in Bungo Province. Here people dig holes in the beach to dip themselves in hot water which issues almost everywhere. The place where such beach baths can be taken, covers no small area, extending from *O-aza-Nishikata* to *Jyū-nichō*, including *Higashikata* and *Jitchō*. These places are on a highway and may be easily reached by *jirikisha* from Kagoshima, but sea-route is preferable, as daily steamers call at such neighbouring ports as *Miya-no-hama* and *Minato-ura*.

Shibatate Hot Spring is at *O-aza-Nishikata*, about 1 m. N. W. from the port of *Minato-ura*.

Nigatsuden Hot Spring is 1.4 m. S. E. from Shibatate. At *Jitchō*, S. of *Nishikata*, is **Yaji-no-yu Hot Spring** and at *Minato-ura*, **Sansetsu Hot Spring**. Within a small distance from *Minato-ura* occur **Kata-no-Onsen** and **Suri-no-hama Hot Springs**.

CHŌSEN (KOREA)

On-yō. From *Keijō* to *Ten-an* (60 m. in 2 hrs. 20 min.) by railway from *Ten-an* to the spa 8.7 m., automobiles available. The spa is situated on a hill-side facing a stream. The place has 250 inhabitants (69 houses) and 3 baths, of which one is occupied by the garrison hospital as sanatorium. A rocky hill called *Sekka-zan* is noted for its beautiful scenery. This place is a favourite resort of the upper classes in *Keijō* and has telephone connections with *Keijō* and *Jinsen*. Number of visitors: 8,032, the total stays being 3,789 days (1913).

Jujō-Onsen. 7 m. from *Taiden*, automobiles available. The spa is situated not far from *Jujō* on the plain of *Taiden*, with hills on the S. and N. It has 210 inhabitants. Spring water issuing from a well, 31 ft. in depth, is conducted by iron pipes to bath-tubs. Number of visitors: 3,836, the total stays being 9,731 days (1913).

Tōrai-Onsen. 3.3 m. from *Fuzanchin* station by light railway in 50 min. The spa lies on the highway which leads from *Fusan* to *Keijō*. It is situated in the central portion of a plain, surrounded by mountains and traversed by a stream. There are 5 inns provided with private baths. This is the most noted hot spring in Chōsen, resorted to by inhabitants of *Fusan*. Number of visitors: 3,836, the total stays being 9,731 days (1913).

TAIWAN (FORMOSA)

Hokutō. The spa is situated in a gorge, about 7 m. from *Hokutō* station. Boiling sulphur water breaks forth from hollows and pools on hill-sides, accompanied by gas and white fumes which fill the gorge all over. The outflow of these springs forms a number of steaming runnels which unite altogether into a stream rushing down the valley. Solfataras and old craters occur here and there in the vicinity. The spa has three most noted hot springs, viz., *Taki-no-yu*, *Kaikōsha-no-yu*, and *Tetsu-no-yu*, which may be regarded as typical of all the springs in the neighbourhood.

Taki-no-yu. The hot springs originate from pools in a small flat tract of land in a deep gorge, N. W. of *Kaikōsha* (military casino). The bottom of the pools consists of muddy sand, and their edges are covered with yellowish-brown or reddish-brown sinter deposits which change into yellowish colour on rainy days. The pools from which the springs issue, receive all water from the baths of both *Kaikōsha* and *Hoshi-no-yu* as well as a ravine at the N. corner. Most of jets are on the S. side of the pools, but change their position from time to time. The flow of water from each spring is also subject to variation. Some of these springs produce more or less gas.

Kaikōsha-no-yu. The place has two springs, both located close to the military casino, the one gushing out of flat ground and the other of crevices of rocks. The hot water is first caught in a wooden tank and then conveyed by bamboo pipes to bath-houses at *Kaikōsha* and *Hoshi-no-yu*. Sinter crust deposited on the river-bed in the vicinity and known as *Hokutōlite* is radio-active. (For Particulars see Chapter on "the Sinter Deposits".)

Tetsu-no-yu. The spring is on the W. side of the *Hokutō* sulphur pit, and within one hour's walk from the *Hokutō* Hotel, hot water gushing out of a large circular hollow on the hills. Here geysers are found every few yards.

Kwanshirei, about 11 m. from *Kōhekiryo* station, S. of *Kagi*, is a small mountain hamlet situated in a ravine with hot springs. The springs, two in number, gush out of sedimentary rocks composed of clay and limestone. The production of gas is abundant from one spring. Water is alkaline and its flow not large.

Suwō (So-ō) is a cold spring near the sea-port of the same name in *Giran-cho* (S. E. from the town of *Giran*), lying on the N. E. coast of the island, about 50 m. from Keelung by steamship. There is also a railway communication between *Tōwi* and *Giran*, but sea-route is preferable.

It is a simple carbondioxated spring, faintly mineralized, used internally and externally.

ILLUSTRATIONS
OF
Hot Spring Spas, Seaside Resorts,
and
Summer Retreats.



Noboribetsu Spa, the most noted hot spring resort in Hokkaido.
A distant view from the spa of Higashidani or "Hell Valley," sending up sulphurous water and white fumes. (p. 4)



Noboribetsu Spa, 4,3 m. from Noboribetsu station, Hokkaidō,
Jigoku-dani or "Hell Valley," sending off sulphurous steam and boiling mud and water—the source of "*Tiki no ju*," p. 51



Noboribetsu Spa.
Chunyu the unctuous hot springs flowing down into *Chunyu* or "Large Hot Spring Lake." p. 1



Ō-yu Hot Springs, near Kosaka Copper Mine, Kikuchū Province,

The beautiful mountain lake *Tazudai-ko* with wonderful scenery is about 14 m. from the spa. p. 5



Waterfall and hot spring at Innal-Yunosawa.



Innal-Yunosawa, a small hot spring spa, near Sot. 1122, p. 15.




Atsumi Hot Springs, on the Japan Sea coast, 8 hours by steamer from Niigata. (p. 7)




Onogawa Hot Springs, near Itoya, on the upper reaches of the Kinomatsunawa (p. 20)



Waterfall "Kame-daki," near the spa



Namegawa, a small hot spring spa, about 15 m. from *Yonezawa*, situated on the northern slope of *Arima-san*, Uzen Province. (p. 11)



Goshiki Hot Springs, about 2 m. from *Izaya* station and 140 m. from *Yonaka*.
The spa being situated amidst the *Izaya* mountains on the slope of *Izaya*, its vicinity is excellent for skiing. (p. 11)



Senami Spa with a geyser, near *Manabani*, situated on a hill wooded with pine trees, close by the sea. The spring was found by accident during the period of oil mania in Echigo Province. Boiling hot water shoots up into the air more than 90 ft. high. (p. 12)



Yi-ni-ki-d, a temple dedicated to the God of Medicine, erected about 500 years ago.



Waterfall "*Jin-ni-tsu-i*" more than 60 ft. high, about 1.2 m. from the spa.



Waterfall "*Ki-sandani-no-tsu-i*" more than 50 ft. high, 1 m. from the spa.



PAU-PAU

Murasugi Spa, 5.5 m. S.E. from the station in the Noto-Muroran Line, Inuyama Prefecture.
It is a pretty first resort, abundant with mineral water, and the spring water is said to be good for various ailments.



Tochiomata, a small hot spring spa, about 22 m. from *Ojima* station, Fehigo Province.

The water is of the strongest radio activity of all mineral springs in the north eastern Japan. (The quantity of radium emanation amounts to 56.4 Mache's units). (p. 17)



Matsuno-yama Hot Springs,
about 33 m. E. from *Yamanashi*, Fukuoka Prefecture. (p. 47)



Nasu-Yumoto Hot Springs, a popular bathing resort at the foot of *Yamaoka* (an active volcano), seen from Shirakawa-Kaido. The spa, about 3,000 ft. above sea level, commands a wide panoramic view. (p. 15)



Yumoto Hot Springs, Nikkō,
a lovely summer resort on Lake Yumoto, 5,400 ft. above the sea. Steaming sulphurous water bubbles
up from the bottom of the lake. (p. 79)



Nashiki-Kosen with cold s. rings, 1,400 ft. above the sea,
a summer resort on Mt. Igo. Igo, p. 27



Hiraiso, 10 m. N.E. of *Mito*, noted both as a watering-place and the fishing center in Hitachi Province.
From a height called "*Asakura-yu*" on the northern side of the place a grand view of dashing waves can be obtained. (p. 25)



Ōarai, the most noted pleasure resort in Hitachi Province. The coast is dotted with rocks and backed by a low hill. Old pine trees grow in the precincts of a Shinto shrine, from which a fine view of the ocean is obtained. (p. 25)



Hayama, a popular seaside resort, about 20 m. from Yokohama
The beach of *Issiki*. An Imperial villa stands in the south of the beach. (p. 251)



Hayama, a popular seaside resort, about 20 mi. from Yokohama.
The beach of Hayama has many villas of wealthy people. Good for swimming. 1905.



Sea bathing at *Goten-dai*



Bathing place with rest huts, seen from the sea.
A group of three big pine trees shows the villa of Baron Iwasaki.



Atami Spa, a favorite winter resort,
with the famous intermittent hot spring. (P. 30)

3




The upper course of the *Kama River*, near *Gama*, Nishibiki.



A picturesque ravine, a little up the stream of *Hakami-gawa* from "*Kinsen-tô*."

2



A *Kinsen-tô* rock standing on the upper bank of the *Kama River*, about 100 yds. from the bath "*Kinsen-tô*."

4



"*Sengajuchi*" or "Thousand Abysses," on the way to Masutomi.

Masutomi Mineral Springs, 24 m. N. of *Kita*, Kai Province.

The *Masutomi* springs are said to be of the strongest radio activity of all mineral springs throughout Japan. The place, however, is not only famous for its picturesque scenery, is a pleasant summer resort. (p.p. 38 and 39)



Tenryūkyō, a famous picturesque scene of


the *Yungawa Rapids* which marks the entrance to the romantic rapids and is that to follow. The descent of the *Yungawa Rapids* by *Phon-Julu-tu-mu* in September, the latest instance of this exciting passage having been undertaken by a distinguished person.



Nakabusa Mountain Spa,

5,300 ft. above sea-level, situated on the middle slope of *Ariake-san*
at the foot of the Hakuba Range, the best starting-point for the ascent of the
"Middle Alps of Japan." There are outdoor arrangements for hot water massage as well as for vapour-baths. (p. 38)


3



4



1



2



Ena Mineral Springs. (p. 16)

1. A large rock with flat top called "*tsukochi-ishi*," is said to have been often visited by the Hannyo for picnic parties.
2. A bridge spanning the *Tate-no-gawa*, the affluent of the River Kiso, near Aoyagi.
3. Looking out the Kiso rapid with picturesque scenery.
4. Flat top made of heavy rock at a point where the River Tsukochi narrows considerably.



Ena Mineral Springs, located on a highland near *Nagô*, on the northern bank of the Kiso Rapids, Mino Province.

The district produces some kinds of radio-active minerals, such as *măgite*, *fergsonite*, etc. Spring water is almost pure and contains a considerable quantity of radium emanation (281 Maché's units). (p. 46)



Futami Hot Springs, east of *T. yama*.
Water issues from a cave under the "Welded Rocks" in a ravine (p. 42)



Ōiwavama, about 14 m. E. of Toyama, Etchu Province.

On a rocky precipice stands a famous temple called *Asuki ji*, and behind the temple are found several waterfalls, of which *Ō taki*, the largest one, presents a wonderful sight. Even now bathing in the waterfalls is popularly believed to be efficacious for curing mental derangement and eye diseases, and thus the place is visited by sick people all the year round. (p. 47)



Ōmaki Hot Spring, southwest of Aomori



Yamanaka Hot Springs, a popular summer resort, south-west of Kyoto.
The old stone bridge, built in 1600, is one of the many fine old bridges in the area. (p. 27)



Komono Hot Springs (Yunoyama), 11 m. north west of Yokoyama.
a quiet summer retreat, about 2,500 ft. above the sea and with a fine bird's eye view. p. 50



"*A. tate*" Waterfall, over 150 ft., is one of the attractions of **Komono Spa**. (p. 50.)



Shiraya Mineral Springs, S. E. of Y. I. J. noted for cherry blossoms, situated on the upper course of the
Yoshino-gawa, abounding in "mau" fish. p. 50



Sedo-no-Kanayama (Yurakii, a lovely seaside resort with hot springs, about 75 m. S. from Wakayama, Kii Province.
The coast near "*Saki no-ya*," issuing from the fissure of rocks close by the sea, forming a gourd-shaped bath-basin. (p. 57)



Yunomine Spa, on the Kamaoka, about 2000 yu from
Sengobu, Kai Province.
Hot springs catch out from the rocks as well as from a
small stream called "Onawa" (P. 10)



Hot springs catch out from the rocks as well as from a
small stream called "Onawa" (P. 10)



Soto-no-yu Hot Springs, near *Katou-umi*,

on the southern coast of Kii Peninsula, situated at the base of high cliffs, commanding magnificent seascape. The spa enjoys a mild climate both in summer and winter. Fish of all kinds are abundant in the adjoining sea (p. 56)




Hamadera, a favorite sea bathing resort near Osaka.
Awaji Island is just opposite to the place and the Kuroko Maru is seen to the left.
Bathing for the first time. The water is very warm. During the summer season large crowds come for a day's pleasure. By electric cars from Osaka.




The Clifford-Wilkinson Tansan Mineral Water Works, at *Takamizuka*, near Osaka. (p. 53)



The Clifford Wilkinson Tamsin Mineral Water Works.
Three separate rows of filters purifying water of iron and other matter. P. 53



A ravine at *Tida-mura*, near Osaka, close to the spring of the famous "Hirano Water,"
(bottled by Teikoku Kosen Kabushiki Kaisha or the Imperial Mineral Water Co., Ltd.), (p. 53)



Interior of the Hattori Water Works of the Imperial Mineral Water Co., Ltd.
Carbonating and Bottling Department. The Factory is one of the most perfect bottling works in Japan. P. 55



Nunobiki Tansan Factory,
situated at the foot of the picturesque hills behind
Kobe, close to the famous Nunobiki Waterfall
from which the name is taken. (p. 54)



Bottling and Corking Department.



Arima Spa, 1,150 ft. above the sea,
one of the most noted hot springs throughout Japan, situated in a valley surrounded by Rokko-san and other hills.
The place is a favorite summer resort near Kobe and Osaka. The springs contain a considerable quantity of common salt and iron. (p. 87)



Kinosaki, about 100 m. from *Kinosaki*, the most noted hot springs in the whole San-in district, known since the 7th century. (p. 54)



Misasa Spa, 5.5 m. S.W. from the *Tokaido* station on the San'in Line, Iki-ki Province.
 The hot spring water is said to be rich in radium emanation contained in the water of the
 140-1-1000 m. A part of the spring contained the emanation of more than 30 Michels' units, number 9 in the spa. (p. 50)




Ushlo, a small hot spring spa, 8 m. S. of *Matste*, Izumo Province, (p. 58)



Port Tono and the Island of Sensui-tô, one of the most charming sights in the Inland Sea, 12 mi. from Inkayama. There are numerous picture postcards, large and small, around the island. The beautiful sheet of water, dotted with fishing boats and white-winged vessels sailing past, presents a delightful scenery. (p. 69)



Dōgo Spa, a watering resort of the most ancient origin in Japan, is in Shikoku and has a three-storied bath-house and a pretty park. (p. 67)



D590 Spd.

Palace, built with "Mother's Gold," where tea and cakes are served to visitors in the upper terraces, while on the
10160 East road are provided for Indian amusements of various kinds. (p. 67)



Funagoya Mineral Springs, near *Haiun-ji* (77 m. from Moji).
A view of the spa from the River Yabe-gawa. (p. 62)



Fig. 1. A boat on the River Yalagawa, near Funagoya Mineral Springs. (p. 12)



Shibasaki Hot Springs, near *Beppu*,
a quiet pleasure resort, situated in a ravine on hill-side.
Hot water, conducted by bamboo pipes and forming a row of waterfalls, is used for massage. (p. 63)



Uml-tigoku, near Hsien,

a boiling pool 60 ft. in diameter at the water part
covered with mud 10 m. hot with blue and green, presenting a curious sight. p. 68



Beppu Spa, on the beautiful Bay of Bungo, called as the Japanese Riviera by foreigners, is a delightful winter resort enjoying a mild climate. Open-air beach baths, called "*Suna-yu*," where hot springs bubble up through sand, are the specialty of the place. Takasakyama (2,667 ft.), an extinct volcano, rises at the southern end of the spa town. (p. 63)



The Western Beach or "Aoi-hama" of **Karatsu Bay** (popular for sea bathing), seen from Matsunuma Koen. p. 68





The Beautiful Bay

The Eastern Port, right, on the estuary of the *Motunua* river.
 A small boat, left, on the bay—grasses extending hundreds of yards, called "A. *motunua*."



of Karatsu.
is shallow and offers no anchorage for large vessels.
presents a charming view and is suitable for bathing. (p. 68)



Karatsu, a town situated at the mouth of the Matsuura-gawa, facing the Bay of Karatsu, noted for beautiful scenery of the beach and also for its pottery. (p. 68)

The picture shows some inns at Karatsu, which may be regarded as the type of Japanese inns of country fashion.



Takao Hot Springs, 33m. from *Ta* in Kyū-hu. Behind the spa town is Sakurayama Park, noted for its landscape gardens and natural scenery. (p. 55)



Cherry blossoms at their best in Sakurayama Park, **Takeo Spa.** (p. 68)



Jojo Hot Springs, about 7 mi. from Tucson, Chaco, N. M.



Kwanshirei Hot Springs, near *Kōkubiyō* station.



south-east of *Kūsi* in Taiwan (Formosa) p. 700



On-yō Hot Springs, about 9 m. from *Fōzan*. Chosen, a favourite resort of the upper classes in Keijō. (p. 70)

INDEX

(The Roman figures refer to the numbers of Parts of the volume.)

- Acid alum spr., def. of, I 53
ex. of, I 70
- Acid alum vitriol spr., def. of, I 53
ex. of, I 71
- Acid hydrogen sulphide spr., def. of, I 53
ex. of, I 72
- Acid spr., action of, I 59
ex. of, I 73
- Acid vitriol spr., def. of, I 53
ex. of, I 70
- Agei, III 57
- Agune H. S., II 188
- Aichi-ken, No. of visitors of spas in, II 200
- Akagi, Mt., III 24
- Akagi-Nishiki C. S., I 68, II 59, III 24
Pl. 28
- Akakura H. S., I 25, 72, II 35-6, 173,
175-6, 187 S, 196, 198, III 14, Pl. 6,
10
sinter deposit of, I 42, Pl. 1
- Akayu C. S. (Akita-ken), II 14, 181, III 4
- Akayu H. S. (Miyagi-ken), II 18, 177,
185-6, 194, III 6
- Akayu H. S. (Yamagata-ken), I 67, II
21-2, 172, 174, 199, III 9
meteor. observ. of, III 9
- Akita-ken, mineral spr. in, II 13-7, III
4-5
No. of visitors of spas in, II 200
summer retreats in, II 202, III 5
- Akyū H. S., II 184-5, 187, 194
- Alkaline common salt spr., ex. of, I 67
- Alkaline earthy spr., def. of, I 52
ex. of, I 66
- Alkaline muriated saline spr., def. of, I 52
- Alkaline muriated spr., def. of, I 52
ex. of, I 66
- Alkaline saline spr., def. of, I 52
- Alkaline spr., action and use of, I 54
def. of, I 52
ex. of, I 65-6
- Alkaline sulphur spr., ex. of, I 72
- Alluvium, I 11
- Almandine, III 12
- Alps of Japan, I 40, III 39
- Alum spr., use and indic. of, I 59
- Alum vitriol spr., def. of, I 53
ex. of, I 70
- Ama-no-Hashidate, sand-spits of, I 5
- Anamori C. S., I 69, II 61-2, 167, 197
- Andai H. S., I 64, II 97-8, 173-4, III 41
meteor. observ. of, III 41
- Andalusite, III 12
- Andesite, I 11
- Anraku H. S., II 189, 196, III 69
- Aomori, meteor. observ. of, I 16-20
- Aomori-ken, mineral spr. in, II 7-11,
III 3-4
No. of visitors of spas in, II 200
summer resorts in, II 202, III 5
- Ao-ne H. S., I 64, II 184-5, 194
- Aoyama H. S., I 64, II 171-2, 196, III 2
- Aragonite, I 40
- Arayu H. S. (Onikōbe), III 6
- Arayu H. S. (Shiobara), III 17
- Archæan group, I 9
- Ariake-yama, Mt., ascent of, III 41
- Arifuku H. S., I 65, II 141-2, 164, 172,
175, 196, III 59
- Arima H. S., I 23, 27, 65, 67-9, II 119-
21, 164, 166-7, 183-5, 187, 197-8,
200, III 54, Pl. 58
- Arita, III 68
- Asama H. S., I 64, II 91-2, 173-4, 180,
195, 199, III 37
meteor. observ. of, III 37
- Asamushi H. S., I 27, 69, II 7-8, 172-5,
III 3
- Ashino-ko (L. Ashi), II 202, III 29
- Ashinoyu H. S., (Hakone), I 25, II 198-
200, III 28
- Aso H. S., I 71, II 158, 164, III 64
- Aso-san (Mt. Aso) (active volcano), I 8,
III 64
- Asuzu H. S., II 131-2, 170-1, III 56
- Atago, Mt., II 202
- Atami H. S., I 25, 66-7, II 70-1, 177-8,
181, 186-7, 195-7, 199, III 30, Pl. 34
geyser (Ō-yu) in, I 47, III 30
max. and min. air temp. of, III 30
vapour baths in, I 60
- Atsumi H. S., I 68, II 20, 164, 199, III
7, Pl. 17
- Atsushio H. S., I 67-8, II 167, 197
- Awake H. S., I 67, II 166-7, 171-3, 182
4, 195, 197, 199, III 50
- Awazu H. S., I 71, II 163, 164, 171-3,
184, 199, III 49
- Azules, I 22
- Azumassō, Mt., III 10-1
- Baku-sansō, I 18, 27
- Balneo-therapy, origin of, I 23
- Basalt, I 12
- Basking in the sun, I 24
- Baths, kinds and use of, I 59-64
public and private, I 28
use of, I 27
- Bath-master, I 29, II 20
- Beppu H. S., I 24-7, 64-6, 69, 71, II 149-
50, 154, 164-5, 185-6, 196, III 63,
Pl. 69
meteor. observ. of, III 63-4
vapour baths at, I 60
sand baths at, I 24, Pl. 69
- Benten H. S. (Nasu), I 64, II 38, 40, 171,
III 15
- Biotite, III 12
- Bitter spr., action and indic. of, I 56
def. of, I 53
ex. of, I 68-9
- Biwa, Lake, I 5, III 35
- Bessho H. S., III 42
meteor. observ. of, III 42
- Bobata C. S., II 29, 166, 179-82, III 12
- Boric acid spr., ex. of, I 68, 73
- Bori-jō-ku, III 63
- Bromine, common salt spr. containing,
ex. of, I 68
- Cainozoic group, I 10
- Calcite, balls, I 42
- Carbondioxated common salt spr., ex. of,
I 67
- Carbondioxated spr., def. of, I 52
ex. of, I 65-7, 69
indic. of, I 54
- Central Japan, climate of, I 21
- Cherry blossoms, I 22
- Chiba-ken, No. of visitors of spas in, II
200
summer resorts in, II 202
- Chichibu, Mt., I 25, II 153, 155, 178,
III 63, Pl. 16
sinter deposit of, II 153
- Chikawa, III 68
- Chikuzen, Province, meteor. observ. of, I
19-20
- Chōsen, climate of, I 21
mineral spr. in, II 160, III 79
No. of visitors of spas in, II 202

- Chūzenji, II 202, III 19
 Classification of mineral spr., I 26
 Clifford-Wilkinson Tansan Works, III 53,
 I 52-4
 Climate of Japan, I 12-23
 Climate therapeutics, origin of, I 23
 Climate and observ. at some health-
 resorts, I 14-5
 Columbite, I 40, III 12
 Common salt spr., def. of, I 52
 ex. of, I 66-9
 use and indic. of, I 55
 Concentrated common salt spr., ex. of,
 I 67
 Cretaceous system, I 10
 Crystalline schist system, I 9
 Crystals, from Yamanashi-ken, III 35
 Curie, I 37
- Dai H. S., II 12, 173, 175-7, 195, 200,
 III 4
 Daigamori C. S., II, 185-6
 Daimaru-zuka H. S., (Nasu), II 39-40,
 186, 176, III 15
 Daitō C. S., II 114, 167, 197
 Deyu H. S., II 32-3, 163, 170, 178-9,
 III 13
 Diabase, I 11
 Diluvium, I 11
 Diorite, I 11
 Dogashima H. S. (Hakone), I 66, II 64,
 III 27
 Dōgo H. S., I 23, 64, II 140-7, 163,
 165-7, 183, 199, III 61, Pl. 63-4
 meteor. observ. of, III 61-2
- Earth-muriated common salt spr., ex. of,
 I 67
 Earthy carbondioxated spr., def. of, I 52
 ex. of, I 65
 use and indic. of, I 54
 Earthy common salt spr., ex. of, I 68
 Ehime-ken, mineral spr. in, II 146-7,
 III 61-2
 No. of visitors of spas in, II 201
 Eino-o H. S., I 72, II 159, 189, 196, III 69
 Electric conductivity, determ. of, I 39
 Emanation, therapeutic use of, I 31
 absorption by the human body, I 32
 Emanation therapy, in Japan, I 31
 short remarks about, I 31-7
 Emanation treatment, cases of—and its
 therapeutical effects, I 34
 Ena C. S., I 30, 64, II 98-9, III 45-6,
 I 38, 39
 Enoshima, II 61, 167
 Epidote, I 4
 Epidote, III 12
 Zircon, I 66
- Examples, of several kinds of spr. in
 Japan, I 64
 Explanatory remarks, VII-VIII
- Fergusonite, I 46, III 46
 Fontactoscope of Engler and Sieveking,
 I 37
 modified by Kohlrausch and Löwenthal,
 I 37
 Forest health resorts, I 23, III 13
 Fuji-san (Mt. Fuji), I 4, II 202, Pl. 11a
 Fukatani C. S., II 184-5, III 49
 Fukawa H. S., II 146, 164, III 60
 Fuki-age (Geyser at Onikōbe), II 186-7,
 III 6
 about Geyser of, I 25, 73
 Fuki-age-yu (Geyser in Obama), I 25, 40,
 II 155-6, Pl. 4
 Fukui-ken, mineral spr. in, II 106-7, III
 50
 No. of visitors of spas in, II 200
 sea bathing in, III 50
 Fukumitsu C. S., II 141, 182, III 59
 Fukuoka-ken, mineral spr. in, II 148-9,
 III 62-3
 No. of visitors of spas in, II 201
 Fukushima-ken, mineral spr. in, II 25-9,
 III 11-12
 No. of visitors of spas in, II 200
 Fukuwata H. S. (Shiobara), II 41, 45,
 170-3, 184, III 16
 Funabara H. S., I 69, II 77-8, 177-8, 182,
 III 32
 Funagoya C. S., I 65, II 148-9, 167-8,
 200, III 62, Pl. 65, 66
 Furu-machi H. S. (Shiobara), II 44-5,
 173-4, III 17
 Furuyu (Unzen), III 67
 Furu-Yumoto H. S. Shiobara, II 45,
 III 17
 Futami H. S., I 26, 71, II 101, 171-3,
 181, 195, III 47, Pl. 7, 40
- Gabbro, I 10
 Gaga H. S., II 184
 Garameki H. S., III 23
 Garnet, III 12
 Gaseous baths at Yunogō, I 61, III 60
 Gembudō, III 54
 Geyser, in Atami, I 47-52, II 71, 178,
 186-7, III 30
 physical investigations of, I 48
 in Obama, I 40, II 155-6, Pl. 4
 at Onikōbe, I 25, II 186-7, III 6
 in Yuzawa (Tochigi-ken) I 40, Pl. 4
 Gifu-ken, mineral spr. in, II 98-9, III
 45-6,
 No. of visitors of spas in, II 200
 summer retreats in, II 202
 Ginsen-tō, II 89, III 34
- Gneiss system, I 9
 Gōra H. S. (Hakone), III 28
 Goshiki H. S., I 64, II 24-5, 171, 173,
 198, III 11, Pl. 20
 Gotemba, II 202
 Granite, I 11
 Gomma-ken, mineral spr. in, II 47-59,
 III 20-4
 No. of visitors of spas in, II 200
 places of interest in, III 21
 summer resorts in, II 202, III 23-4
- Hakone H. S., II 62-8, III 25-9
 average temp. of, I 13
 geography of—district, III 26-7
 Hakone, Lake, III 29
 twelve H. S. of, III 27-9
 Hamadera, sea bathing, III 51, Pl. 51
 meteor. observ. of, III 52
 Hamamura H. S., I 69, II 129-30, 170-1,
 III 56
 Hara-machi (Haraichi), I 67-8, II 197,
 III 22
 Haruna, Lake, II 202, III 24
 Mt. (Haruna-Fuji), III 23
 Hatage H. S., I 65, II 76-7, 175-6, III
 32
 Hataori H. S. (Shiobara), II 42, 45, 172,
 III 17
 Hayama, sea bathing, III 25, Pl. 31, 32
 Hayashi H. S., II 188
 Hiei, Mt., II 202
 Higashiyama H. S., I 69, II 27-8, 170
 180, 199, III 11
 Hinagu H. S., II 188, 199, III 65
 Hinoyama Park (Toba), III 50, Pl. 46
 Hira-iso, sea bathing, III 25, Pl. 29
 Hirano C. S., I 67, 69, II 118, 186-7,
 III 53
 Hirano Waters, III 53-4, Pl. 52, 55-6
 Hirao H. S., I 68, 96-8, 172-4, 187-8,
 199, III 41-2
 five hot spr. in, III 41
 meteor. observ. of, III 41-2
 Hirayu H. S. (Japan Alps), III 40
 Hiroshima-ken, No. of visitors of spas in,
 II 201
 places of interest in, III 60
 Hirukawa C. S. (Ena), II 99, 165, 179
 Hisōmoe H. S., I 70, II 198
 Hokkai-dō, climate of, I 21
 mineral spr. in, II 1-6, III 1-3
 No. of visitors of spas in, II 200
 sea bathing in, III 2
 summer resorts in, II 202, III 2
 Hokuroku-dō, climate of, I 21
 Hokutō H. S., I 42-3, 65, 71, II 161,
 184-7, 195-6, III 70
 Hokutōlilite, I 42-3, III 70, Pl. 3
 analysis of, I 43

- of Shikokuro, I 43-4, III 5
photograph and radiograph of, I 44
Hornblende Andesite, I 11
Hotaka-gatake (Japan Alps), ascent of,
III 41
Hot Springs, bathing equipments, I 20
classification of, I 20
composition of, I 20
def. of, I 52
general remarks about, I 25
most radioact., III 56
quantity of flow, I 25
temperature of, I 26
therapeutic efficacy of, I 29
Hot spring spas, geology of, I 25
situation of, I 25
Hotu H. S., I 71, II 154, 185, 187, 200,
III 63
Hot waterfalls, I 27
Hozu, River, I 5
Humidity of Japan, I 17
Hyogo-ken, mineral spr. in, II 115-23,
III 52-5
No. of visitors of spas in, II 201
sea bathing in, II 203
summer resorts in, II 202
Hydrogen sulphide common salt spr., ex.
of, I 68
Hydrogen sulphide spr., def. of, I 53
ex. of, I 68, 71-2
Hyperthermal baths, I 60

Ibaraki-ken, mineral spr. in, II 59-60
No. of visitors of spas in, II 200
sea bathing in, III 25
summer retreats in, II 202
Ibusuki H. S., sand baths at, I 25
Iizaka H. S., I 27, 64, II 26-7, 174-5,
185-6, 199, III 11
Ikao-Fuji, III 24
Ikao H. S., I 25, 27, 69, II 54-5, 175-6,
198-9, III 22-4
meteor. observ. of, III 23
Ikargasaki H. S., I 60, II 10-11, 172-3,
III 3
Ikeda, II 140, 165, 168, 179-80, 182, III 58
Imari, III 69
Imperial Mineral Water Co., III 53, Pl.
52, 55, 59
Indication of mineral spr., I 53-59
Indifferent thermals, action and indic. of,
I 54
def. of, I 52
ex. of, I 64
Inland Sea, I 6
climate of—district, I 21
Innai-Yunosawa H. S., I 65, II 17, 173,
190, III 5, Pl. 10
Inns of hot spring spas, I 28
Iodine, common salt spr. containing, ex.
of, I 68

Irimoyu H. S., II 58, 183, 198, III 24
Iron carbonate common salt spr., ex. of,
I 69-70
Iron carbonate spr., def. of, I 53
ex. of, I 69

Iron carbondioxated common salt spr.,
ex. of, I
Iron springs, indic. of, I 56
ex. of, I 69
Ishikawa-ken, mineral spr. in, II 103-6,
III 47-9
No. of visitors of spas in, II 200
Ishikawa-yama, III 12
Isobe C. S. (Gumma-ken), I 27-8, 66-8,
II 53-4, 167-8, 181, 197, III 21, Pl.
8, 27
Isobe C. S. (Toyama-ken), I 70-1, II 197
Isumuro H. S. (Nagasaki), II 40, III 15
Itō H. S., I 64, II 71-2, 175-8, 181-2,
195, 199, III 31
Iwai H. S., I 69, II 124, 170-3, 190,
III 55
Iwaki-san (Tsugaru-Fuji), I 4, III 3
Iwate-ken, mineral spr. in, II 12-3, III 4
No. of visitors of spas in, II 200
Iwōdani H. S. (Kirishima H. S.), I 72,
II 158-9, 189, 198, III 69
Izusan H. S., I 20, 69, II 69, 178, 180,
III 30

Japan, climate of, I 12-23
geology of, I 9-12
lakes of, I 5
mean, max. and min. temp. of, I 15-6
mountain ranges of, I 3-4
rivers of, I 5
scenery of, I 3-6
situation of, I 1
Japan Alps, I 4, III 39
Jigoku H. S. (on Mt. Aso), II 158, III 65
Jikan-yu (time baths), I 29, III 20, Pl. 9
Jinrikisha, I 28
Jinsen, meteor. observ. of, I 16-20
Jōji H. S., II 185
Jōzankei H. S., II 4, 174, 198, III 1
Jujō H. S. (Chōsen), III 70, Pl. 75
Jurassic system, I 10

Kachimi H. S., I 69, II 129, 163, 165,
169-70, 178, III 56
Kagawa-ken, No. of visitors of spa in,
II 201
Kagawa, I 28
Kagoshima, meteor. observ. of, I 16-20
Kagoshima-ken, mineral spr. in, II 158-9,
III 69
No. of visitors of spas in, II 201
Kakuma C. S., I 94, II 144, 190, 179,
III 60

Kakuma H. S., II 98, 174, III 42
Kamaoka H. S., I 67, II 18-20, 171,
174, 181-3, 190, 200, III 6
Kambayashi H. S. (Hirao), II 98, 174,
III 41
Kamegawa H. S., I 25-6, 66, II 152-5,
175-8, 185-6, 190, III 3, 4
Kamikōchi H. S., I 25, 65, II 94, 172,
198, III 38
Kamibayashi H. S., I 60, II 21, 171-2,
182-3, 196, 199, III 8
meteor. observ. of, III 8
Kami-Sekisuiji C. S., II 84, 180, III 35
Kami-Suwa H. S., I 64, II 91, 171, 173,
195, 198-9, III 35-6
meteor. observ. of, III 36
Kamo-machi (near Yunohama H. S.),
meteor. observ. of, III 7
Kanagawa-ken, mineral spr. in, II 62-69,
III 25-30
No. of visitors of spas in, II 200
seaside resorts in, II 203, III 25
summer resorts in, II 202, III 29
Kanetsuri H. S., II 102, 171, III 47
Kannawa H. S., I 26, 70, II 150-1, 155,
185-6, 199, III 63, Pl. 8, 68
Kanō-zan (Mt. Kanō), II 202
Karafuto, meteor. observ. of, I 16-20
climate of, I 21
Karatsu-machi, sea bathing, III 68, Pl.
70, 71, 72
Karuzawa, II 202, III 44
meteor. observ. of, III 45
Karurusu H. S., I 65, II 3-4, 175-6, III
1-2
Kasagi C. S., I 27, 66, II 113-4, 181,
III 51
Kasagi-yama (Mt. Kasagi), III 51
Kashi H. S., I 65, II 28, 164, 171-1,
198, III 11
Kashio, I 67
Kasubuchi, C. S., II 142, 179, III 59
Katayamazu H. S., I 66, II 104, 164-5,
173, 175, 178, 197, 199, III 49
meteor. observ. of, III 49
Katsu-ura H. S., II 112, 164, 166-7, III
54, Pl. 50
Kawai C. S., II 138-9, 181-2, III 58
Kawarayu H. S., I 72, III 21
Kawatabi H. S., II 18, 175, 185-6, 194,
III 6
Kawatabi H. S., II 148, 164, III 6
Kaze-yu (numbering bath), III 15
Kawata H. S. (Hirao), II 18-20, 171-2
Kimbu-san (Mt. Kimbu), III 35
Kimpōri H. S., II 185-7
Kinai, climate of, I 21
Kinai H. S., I 123-5, 174, 176, 178,
181-2, 184-5, 190, 191, 193, 195, 197,
199, 200, III 21
Kinokuni H. S., II 185, 190, III 21

- Kasuga, river, III 4
 view from Mt. Asahi, I 23
- Kawachi, hot springs in, I 25, 27, II 198, III 69
- Kita H. S. (Nasu), I 64, II 38-40, 173, III 15
- Kōbe, I 64, II 184
 meteor. observ. of, I 10-20
- Kōchi H. S., II 81, 175-6, III 32
- Kōchi-ken, No. of visitors of spas in, II 201
- Kōji, Mt. (Mt. Kōji), active volcano, I 6
- Komono (Yunoyama) C. S., I 64, II 107-8, 179, III 50, Pl. 44-5
- Kona H. S., I 65, II 75, 177, III 32
- Konabe H. S., II 80, 177, III 32
- Kosaka Copper Mine, III 4
- Koshikiwa-shinden, I 65, II 115-6, 166-7
- Kowakidani H. S. (Hakone), I 70, II 60, III 38
- Koyasan (Mt. Koya), II 202
- Koyabara H. S., I 69, II 139-40, 163, 165, 171, 181, III 58
- Kumamoto-ken, mineral spr. in, II 158, III 64-5
 No. of visitors of spas in, II 201
 places of interest in, III 64-5
- Kuradate H. S., I 66, II 9-10, 170-1, 195, III 3
- Kurobira, III 34
- Kurokagi H. S., II 101, 171, III 47
- Kuro-shio* (Black Stream), I 22
- Kusatsu H. S., I 23, 25-7, 29, 66, 70-2, II 47-51, 177-8, 182, 197-8, 200, III 20, Pl. 5, 6, 8, 9
 hyperthermal baths at, I 60
 sinter deposit of, II 50
 time baths at, I 29, III 20
- Kuzu H. S., II 96, 171-2, 195, III 39
- Kwankaiji H. S., I 69, II 150, 154, 185, 199, III 63
- Kwansei, climate of, I 21
- Kwansei H. S., I 67, II 102, 105, 185-7, III 70, Pl. 77
- Kwansei, climate of, I 21
- Kyōto, meteor. observ. of, I 16-20
- Kyōto-fu, mineral spr. in, II 113-4, III 51
 No. of visitors of spas in, II 201
 summer resorts in, II 202
- Kyushu, climate of, I 21
- Lepanto, I 11
- Mache's unit, I 37
- Maebashi, meteor. observ. of, I 16-20
- Maga H. S., II 171, 196, III 60
- Manza H. S., I 25, 72, II 198, III 21
- Masutomi C. S., I 66, 68, II 85-9, 165-8, 179-80, 182, 197, III 33, Pl. 10, 35
 discharge of radium emanation, II 88
 most radioact. mineral spr., I 30
 sinter-coated leaves at, I 42, Pl. 2
- Matsue, III 58
- Matsuida, II 54, 168, 182-3
- Matsumoto, I 16-20, III 37
- Matsunoyama H. S., I 67, II 36, 187, 197, III 14, Pl. 24
- Matsuyama, I 16-20, III 61
- Matsuzaki C. S., II 131, 180, III 56
- Maya-san (Mt. Maya), II 202
- Meakan-dake (active volcano), I 6
- Means of access for hot spring spas, I 28
- Medicinal spr., kinds and indic. of, I 53-9
- Mesozoic group, I 10
- Meteor. observ., I 13-20, III 7-10, 23, 36-7, 41-5, 48-9, 52, 58, 61-4, 66-7
- Methode of bathing, I 36
- Meso-thorium, I 36
- Michi-no-o C. S., I 70, II 157, 167, III 68
- Mie-ken, mineral spr. in, II 107-8, III 50
 No. of visitors of spas in, II 200
 places of interest (or sea side resorts) in, III 50, Pl. 46
- Miho-no-Matsubara, I 5
- Mimata H. S., II 170, III 59
- Mineral mud baths, I 25
- Mineral spr., classification of, I 26
 in other countries, I 78-94, II 190-3
 weak and strong radioact., I 30
- Minerals, radioact., I 46
 literature of, I 46
- Mineral waters, bottled for sale and exports, I 28
- Mino-o, II 202
- Misasa H. S., I 27, 30, 65-6, 72, II 133-5, 163, 165, 179-80, 195, III 56, Pl. 11, 60
 most radioact., I 30
- Mitake, Mt. (Tōkyō-fu), II 202
- Mitake, Mt. (Yamanashi-ken), III 34
- Mitaki H. S. (Onikōbe), II 186, III 6
- Mitsuya Brand Min. Water, III 53, Pl. 52, 55, 56
- Miyagawa H. S., II 5, 175, 182-3, III 2
- Miyagi-ken, mineral spr. in, II 17-20, III 5-6
 No. of visitors of spas in, II 200
- Miyagino (Hakone), III 28
- Miyanoshta H. S. (Hakone), I 66, II 63-4, 195-6, 198-9, III 27, Pl. 11a
- Miyazaki-ken, No. of visitors of spas in, II 201
- Mogi, III 65
- Monazite, I 45, III 12
- Monsoon, I 13
- Monzen H. S. (Shiobara), II 43-5, 173-4, III 17
- Morigasaki C. S., I 67, II 60-2
- Morion, III 46
- Moto-Hakone, III 29
- Moto-Kuruma H. S., II 18, 176, 194, III 6
- Mt. Fuji, I 4
 view from L. Shōji, Pl. 11a
- Mud baths, radioact. of, I 37
 of Chi-no-ike-Jigoku, III 63
- Mud Lava, I 12
- Murasugi C. S., I 64, II 31-2, 165-6, 179, 182, 187, 200, III 13, Pl. 22
 discharge of radium emanation, II 32
 forest resort, III 13
- Muriated saline bitter spr., ex. of, I 69
- Muriated spr., def. of, I 52
 ex. of, I 66
 indic. of, I 55
- Muriated sulphated bitter spr., ex. of, I 69
- Muriated sulphur spr., ex. of, I 72
- Musashi H. S., I 71, II 148, 163, 199, III 62
 meteor. observ. of, III 62
- Muscovite, III 12
- Mushi-buro* (vapour bath), I 24
- Myōban H. S. (near Beppu), I 71, II 152, 155, 178, 184-5, 195, III 63
- Myōban-yama H. S. (Mt. Kirishima), II 189, 196, III 69
- Myōgi, Mt., I 5, III 21
- Myōken H. S., II 189, III 69
- Myōkō H. S., III 14
- Myōkō-zan (Mt. Myōkō), III 14
- Naegi, I 45-6, III 46
- Naēgite, I 46, III 46
- Nagano-ken, forest health resorts in, I 23
 mineral spr. in, II 90-8, III 35-45
 No. of visitors of spas in, II 200
 places of interest in, III 38-41, Pl. 36
 summer resorts in, II 202, III 44-5
- Nagaoka H. S., I 64, II 74-5, 175-8, III 32
- Nagasaki, meteor. observ. of, I 15, 17-20
- Nagasaki-ken, mineral spr. in, II 155-7, III 65-8
 No. of visitors of spas in, II 201
- Nagawaki*, I 47-8, 50-1
- Nagoya, meteor. observ. of, I 16-20
- Naikai (Inland Sea District), climate of, I 21
- Nakabusa H. S., I 25-7, 60, 66, 71-2, II 94-5, 164, 171-4, 195, 198, III 38, Pl. 37
- Naka-no-umi, III 58
- Nakayama H. S., II 186
- Namaze C. S., II 184, III 54
- Namegawa H. S., II 24, III 11, Pl. 19
- Nankai, climate of, I 21
- Nara-ken, mineral spr. in, II 108-9, III 50
 No. of visitors of spas in, II 200
- Naruko H. S., I 26-7, 66, 70-2, II 18, 173, 176-8, 185-6, 195, 199, III 6

- vapour baths at, I 60
- Nashiki C. S., I 68, II 59, III 24, Pl. 28
- Nasu H. S., I 23, 25, 29, 64, 72, II 37, 40, 171, 174-6, 181, 186, 198, 200, III 15
- smoldering bath at, I 29, III 15
- sinter deposit of, II 39
- Nasu-Yumoto H. S., I 72, II 37-8, 40, 174-6, 181, 198, 200, III 15, Pl. 25
- Nekonaki C. S., I 64, II 29, 166-7, 180, III 12
- Nenuro, meteor. observ. of, I 16-9
- Nezaki H. S., II 6, 170, 182-3
- Niigata, meteor. observ. of, I 16-20
- Nigata-ken, mineral spr. in, II 30-6, III 12-5
- No. of visitors of spas in, II 200
- Nikko, four seasons in, III 18-9, Pl. 11a
- Nikkō-Yumoto H. S., I 25, 72, II 45-7, 171-2, 174, 198, III 19-20, Pl. 26
- Nishinomiya, II 166, 185
- Nishiyama Oil Field, radioact. of petroleum from, II 189
- Nisseki-ji, III 47
- Noboribetsu H. S., I 25-7, 66, 70, II 1-3, 175-7, 195-6, III 1, Pl. 7, 12-4
- Nojiri, lake, II 202
- Norokura-ga-dake, Mt., ascent of, III 40
- Noroshi C. S., I 70, II 182
- Notes on spas in other countries, I 78-94
- Nozawa H. S., I 71, II 195, 199, III 44
- meteor. observ. of, III 44
- Nunobiki Tansan Works, III 54, Pl. 52, 57
- Ōami H. S. (Shiobara), II 40-1, 45, 170, 184, III 16
- Ōarai, sea bathing, III 25, Pl. 30
- Obama H. S., I 25, 67, II 155-6, 164, 167, 195, 197, 199, III 65, Pl. 11a
- geyser in, I 40, Pl. 4
- Obara H. S., II 185-6
- Ōdake C. S., II 123, 166
- Ōdaki H. S., II 15-6, 173-4, 196, III 5
- Ōdomari, meteor. observ. of, I 16-20
- Ogawa H. S., I 26, 67, II 100, 171, 173, 176, III 47
- sinter-balls of, I 42, Pl. 2, 10
- Ōiso, sea bathing, I 24, III 25, Pl. 33
- Ōita, meteor. observ. of, I 15, 17-20
- Ōtadake, mineral spr. in, II 149-55, III 63-4
- No. of visitors of spas in, II 201
- Ōwayama, I 24, II 202, III 47, Pl. 41
- Okayama-ken, mineral spr. in, II 143-5, III 59-60
- No. of visitors of spas in, II 201
- Ōkawa C. S., II 106, 164
- Ōkuni H. S., I 68, II 102, 164, 171, III 47, Pl. 42
- Ōmori, II 62, 181
- Onikōbe-gotō, III 6
- Onikōbe H. S., I 25, II 17, 175, 185-7, 195, III 6
- geyser at, I 73-7
- Onishi (Yashio) C. S., I 67, II 167
- Onnagawa H. S., II 30-1, 163, 170, III 13
- Ōno, sea bathing, I 24
- Onogawa H. S., I 67, II 23, 170-1, 173, 199, III 10, Pl. 18
- meteor. observ. of, III 10
- Ōnuma Park, II 202, III 2
- On-yō H. S. (Chōsen), II 160, III 70, Pl. 76
- Orthite, I 46
- Ōsaka-fu, mineral spr. in, II 114-5
- sea bathing in, III 51-2
- summer resorts in, II 202
- Ōsawa H. S., II 12-3, 174, 176, III 4
- Ō-u, climate of, I 21
- Ōshio C. S., I 65, 67, II 192
- Otari H. S., II 198
- Ōwakidani H. S. (Hakone), I 26, III 28
- Ōwani H. S., I 27, 66, II 8-9, 169, 174, 195, 199, III 3
- Ōyu H. S. (Akita-ken), I 60, II 14-5, 173-5, III 4, Pl. 15
- Ōyu H. S. (Niigata-ken), I 65, II 163, 170, III 14
- Ōyuzawa H. S., I 68, 72, II 13-4, 174, 181-3, 197, III 4
- Palaeozoic group, I 3-4
- Pearl-culture, III 50
- Pediluvium* (hyperthermal foot bath), I 60
- Piedmontite, III 12
- Porphyrite, I 11
- Porphyry, I 11
- Prolonged baths, I 60
- at Tochiomata H. S., III 14
- at Yunogō H. S., I 29, III 59
- Pyroxene andesite, I 11
- Quartz, III 12
- Quaternary system, I 11
- Radioact. earths, use of, I 36
- Radioact. minerals in Japan, I 42-6
- Radioact. mineral spr., utilization of, I 31
- Radioact. spr., most radioact., III 34
- Radio-activity, determ. of water, I 38
- determ. of gas, I 38
- expression of, I 37
- of all mineral spr. in Japan, II 163-89, 194
- of gas from mineral spr., II 165, 168, 178, 182-3, 187-9, 192-3
- of mineral spr. in Japan, I 30
- of radioact. mineral spr. in Japan, I 30
- of radioact. mineral spr. in Japan, I 30
- Radiobaryte, I 45
- Radium emanation, absorption of, by water, I 38
- decay of, I 39
- determ. of, I 37
- from Murasagi spr., II 32
- indication of, treatment, I 33
- injection of, I 33
- physiolog. action of, I 33
- Radium rays, bactericidal action of, I 32
- Raised coral reefs, I 11
- Rejuvenating waters, I 33
- Rendaiji H. S., II 81, 171-2, 174-7, III 32
- Renge H. S., I 25, 70, II 196, III 40
- Rest-activity of radium, in spring waters, I 38
- Reynell's Hirano waters, III 53, Pl. 52
- Rokkōzan, Mt., III 54
- Rokuyō C. S., I 70, II 108-9, 167, III 50
- Sabusawa (Onikōbe), III 6
- Saga-ken, mineral spr. in, II 157-8, III 68-9
- No. of visitors of spas in, II 201
- sea bathing in, III 68
- Sahime H. S., II 139-40, 163, 165, 168, 171, 173, 177-82
- sinter deposit of, I 41, III 59, Pl. 1
- Sai-no-kawara Kusatsu, III 21
- Saitama-ken, No. of visitors of spas in, II 200
- Sakai, meteor. observ. of, I 16-20
- Sakunami H. S., II 183-4, 194
- Sakurajima, I 5, 69
- Saline bitter spr., ex. of, I 69
- Saline common salt spr., ex. of, I 67
- Saline sulphur spr., ex. of, I 72
- Sea-baths, I 24-6, III 7, 11-6
- Sandogoya H. S. (Nasu) I 25, II 39-40, 196, 198, III 15
- Sano Waterfall Garden, II 202
- Sapporo, meteor. observ. of, I 16-20
- Sawatari H. S., I 72, III 21
- Schmidt's electrometer, I 37
- Sea bathing, I 24, II 203, III 2, 7, 25, 31, 50-1, 55, 68
- origin of, I 23
- Sea weed baths, I 24
- Sedo-no-kanayama (Yuzaki) H. S., I 25, 66, II 104-11, 113, III 81, Pl. 48
- Sekigane H. S., I 71, II 132-3, 163, 169, III 57
- Seki H. S., III 14
- Sengokubara H. S. (Hakone), II 67, III 28
- Senjoga-hara, III 19
- Serpentine, I 11

- I 6
 S. H. S., II 189, III 60
 S. H. S., I 66, II 154, 170, 178, 180, III 63, Pl. 67
 S. H. S. (Suwa), II 90, 105
 sinter deposit of, II 90
 S. H. S. (Haruna), I 25, 68, 75, II 97-8, 173-4, 187-8, 195, 198, III 41
 S. H. S. (I 43, 5, 71, II 10, III 5
 S. H. S. (I 68, II 12, 175, 195, III 4
 Shiga-ken, No. of visitors of spas in, II 200
 Shikama H. S., I 25-6, 60, II 139-40, 173, 177-8, 180, III 59, Pl. 5, 6
 sinter deposit of, I 41
 Shikanoyu (Ōkuwa) C. S., II 166, 194
 Shima H. S., I 27, 60, 66, II 51-2, 172, 174-6, 198, 200 III 21
 Shimabara Peninsula, III 65
 Shimane-ken, mineral spr. in, II 136-42, III 57-9
 No. of visitors of spas in, II 201
 Shimobe C. S., I 60, 64, II 84-5, 168, III 35
 prolonged bath at, I 60
 Shimoda, II 81, 181, III 32
 Shimogamo H. S., II 81-2, 171, 174-5, 177, 195, 197, III 33
 Shimo-Suwa H. S., I 64, II 91, 172-3, 198, 200, III 35
 Shimura (Yunomine) H. S., II 111, 195, III 51, Pl. 49
 Shin-Goshiki H. S., II 25, 174
 Shingū C. S., II 167, 181
 Shinji-ko, Lake, III 58
 Shin-Kuruma H. S. II 18, 175, 185, 194, III 6
 Shiobara H. S., I 25, 27, 65-6, 69, II 40-5, 170-4, 184, 199, III 15, Pl. 11a
 sinter deposit of, II 44
 seven wonders of, III 17
 Shiogama H. S. (Shiobara), II 45, III 16
 Shionoyu H. S. (Shiobara), II 42, 45, 174, III 17
 Shiote H. S., II 189, 195
 Shioya C. S., II 183
 Shirahone H. S., I 66, II 93-4, 164, 173-6, 178, 198, III 38
 Shiraya C. S., I 65, II 109, III 50, Pl. 47
 Shizuoka ken, mineral spr. in, II 69-83, III 3-3
 No. of visitors of spas in, II 200
 sea bathing in, II 203, III 31
 summer resorts in, II 202
 S. H. S., II 202, Pl. 11a
 Shu-otsu H. S. (Chōsen), I 8, 65, II 196
 Shuzenji H. S., I 26, 67, 72-4, 173-8, 198, 199, III 31
 Simple carbondioxated spr., action and indic. of, I 54
 def. of, I 52
 ex. of, I 65
 Simple cold spr., action and indic. of, I 53
 ex. of, I 64
 Simple common salt spr., I 52
 ex. of, I 66
 Simple thermals, action and indic. of, I 54
 def. of, I 52
 ex. of, I 64
 Simple earthy thermals, def. of, I 52
 Sinter-balls, I 42, Pl. 2
 Sinter-coated leaves, I 41, Pl. 2
 Sinter cones (Kuriyama H. S.), I 40-1, Pl. 4
 Sinter cones (Obama H. S.), I 39, 40, Pl. 4
 Sinter deposits of mineral spr., I 39, 41-2, II 39, 44, 50, 90, 153, III 14, 59, Pl. 1, 3
 Sipping cure, I 33
 Skating ground, at L. Haruna, III 24
 at L. Suwa, III 35
 Sokokura H. S. (Hakone), I 66, II 65, III 27
 Sotonoyu H. S., II 112, 164, III 51, Pl. 50
 Spas, notes on, in other countries, I 78-94
 Spring discovery, chance causes of, I 29
 Steam baths, I 60
 Stufe, I 60
 Subthermal baths, I 59
 prolonged, I 29, 60, III 14, 59
 Sulphated bitter spr., ex. of, I 69
 Sulphated common salt spr., ex. of, I 68
 Sulphated sulphur spr., ex. of, I 72
 Sulphur spr., action and indic. of, I 57
 def. of, I 53
 ex. of, I 71-2
 Sumaki H. S. (Shiobara), II 43, 45, 174, III 17
 Summer resorts, II 202, III 2, 5, 19, 24, 44-5
 Su-ō (Suwō or So-ō) H. S. (Taiwan), I 65, II 162, 185, 187, 191, III 35
 Surinohama H. S., II 189, III 69
 Suwa H. S., I 25, 64, 90-1, 171-3, 195, 198-200, III 35
 meteor. observ. of, III 36
 Suwa, Lake, III 35
 Taihoku, meteor. observ. of, I 15, 17-20
 Tainan, meteor. observ. of, I 15, 17-20
 Taiwan, meteor. observ. of, I 15, 20
 mineral spr. in, II 161-2, III 70
 Takanosu H. S., II 30-1, 170, III 13
 Takaomata H. S. (Nasu), II 40, III 15
 Takarazuka C. S., I 28, 65, 67, II 116-7, 119, 166-8, 184, 187, 197, 200, III 52
 Takarazuka Tansan Works, III 53
 Takase H. S., II 31, 163, 170, III 13
 Takayama C. S., I 64, II 98-9, 165, 179-80, III 46, Pl. 38-9
 Takayu H. S. (Fukushima-ken), I 70, 72
 Takayu H. S. (Yamagata-ken), II 196, 198-9
 Takebe C. S., II 144-5, 166, 180, III 60
 Takedao C. S., I 72, II 118-9, 166, 183, III 54
 Takeo H. S., I 27, 64, II 157-8, 196, 199, III 68, Pl. 73-4
 Tamatsukuri H. S. (Miyagi-ken), II 18, 199, III 6
 Tamatsukuri H. S. (Shimane-ken), I 60, 66, II 136, 163, 165, 170-1, 173, 175-8, 185-6, 194-6, III 57, Pl. 5
 air temp. of, III 58
 Tarutama H. S., I 71, II 158, 164, III 65
 Tateyama H. S., I 25, 72, II 196, 198
 Tatsunokuchi C. S., II 184, 199, III 49
 Tawarayama H. S., I 65, II 146, 164, 200, III 60
 Tenryū-kyō, III 38, Pl. 36
 Teppōsui (Arima), II 121
 Teramachi C. S. (Tottori), II 125-6, 167-8, 181
 Tertiary system, I 10
 Thermae, I 60
 Thorium emanation, I 36
 in spr. waters, I 38
 Time baths, I 29, 60, III 20, Pl. 9
 Tin sand (Naegi), III 46
 Titanite, III 12
 Toba, III 50, Pl. 46
 Tochigi-ken, mineral spr. in, II 37-47, III 15-20
 No. of visitors of spas in, II 200
 places of interest in, III 18
 summer resorts in, II 202, III 19
 Tochinoki H. S., I 69, II 158, 164, 196, III 65
 Tochiomata H. S., I 60, 65, II 33-4, 163, 166, 169, 179, 196, III 13, Pl. 23
 discharge of radium emanation, II 34
 prolonged bath at, I 60
 Todoroki (Onikōbe), III 6
 Togakushi, II 202
 Tōgatta H. S., II 183-4, 194
 Tōgō H. S., I 26, 65, II 130-1, 163, 170, 172, 180, 196, III 56, Pl. 5
 Tohi H. S., I 69, II 82-3, 174-7, 181, III 33
 Tōi C. S. (Taiwan), II 186, 195
 Tōkai-dō, climate of, I 21
 Tokushima-ken, No. of visitors of spas in, II 201
 Tōkyō, meteor. observ. of, I 16-20
 radioact. of well waters in, II 62, 167-8, 181
 Tōkyō-fu, mineral spr. in, II 60-2

- No. of visitors of spas in, II 200
summer resorts in, II 202
- Tombara C. S., II 142, 179, III 59
- Tomo, III 60, Pl. 62
- Tōmoe C. S., II 59-60, 167
- Tonosawa H. S. (Hakone), I 64, II 62-3, 199, III 27
- Tōrai H. S. (Chōsen), I 66, II 160, 163, 167-8, 170, 180-2, III 70
- Tottori, II 125-6, III 55
- Tottori-ken, mineral spr. in, II 124-35, III 55-7
- No. of visitors of spas in, II 201
- places of interest in, III 57
- sea side resorts in, III 55
- Towada, Lake, II 202, III 5
- Town of Longevity, III 50
- Toyama-ken, mineral spr. in, II 100-2, III 47
- No. visitors of spas in, II 200
- summer resorts in, II 202, III 47
- Triassic system, I 10
- Tsubame H. S., I 71, II 36, 174, 177-8, 187-8, III 14, Pl. 6
- Tsukuba C. S., II 60
- Tsukuba, Mt., II 202
- Tungsten, III 35, 46
- Ubako H. S. (Hakone), I 64, II 68, III 28
- Umi-Jigoku, II 151, 155, III 63, Pl. 68
- Umijiri (Hakone), III 28
- Unzen H. S., I 70, 72-3, II 156-7, 163-4, 198, 200, III 66, Pl. 11a
- meteor. observ. of, III 66
- Uradomi, sea bathing, II 203, III 55
- Urai H. S. (Taiwan), II 183-6, 195
- Ura-Nippon, I 22
- Ureshino H. S., I 66, II 196, 199, III 68
- Ushio H. S., I 68, II 137, 164-5, 176, 181, III 58, Pl. 61
- Utsunomiya, meteor. observ. of, I 16-20
- Vapour baths, I 24, 60, III 21, 24, 63
- Vitriol springs, def. of, I 53
- indic. of, I 56
- ex. of, I 70-1
- Volcanoes and mineral spr., I 6-8
- Volcanoes, types of, I 4-5
- Volcanic Ash, I 12
- Wakayama-ken, mineral spr. in, II 109-13, III 51
- No. of visitors of spas in, II 200
- sea bathing in, III 51
- summer resorts in, II 202
- Wakura H. S., I 67, II 103, 163-5, 170, 172-3, 183-4, 195, 197, 199, III 49
- Watering places, I 24, II 203, III 2, 7, 25, 31, 50-1, 55, 68
- Weak common salt spr., I 52
- ex. of, I 66
- Wistarias, I 22
- X-rays treatment, I 34
- Yabakei, I 5
- Yabakei of Kwantō, III 20-1
- Yaji H. S., II 189, III 69
- Yamabe H. S., II 92-3, 172, 181, III 38
- Yamaga H. S., II 199, III 65
- Yamagata-ken, mineral spr. in, II 20-5, III 7-11
- No. of visitors of spas in, II 200
- sea bathing in, III 7
- Yamaguchi-ken, mineral spr. in, II 145-6, III 60
- No. of visitors of spas in, II 201
- Yamanaka H. S., I 72, II 105, 164, 171-2, 184, 199, III 47, Pl. 43
- meteor. observ. of, III 48
- Yamanashi-ken, mineral spr. in, II 84-9, III 33-5
- No. of visitors of spas in, II 200
- places of interest in, III 34
- summer resorts in, II 202
- Yamashiro H. S., I 72, II 104, 164, 175, 177, 185, 187, 199, III 48
- meteor. observ. of, III 48
- Yari-ga-take, ascent of, III 40
- Yashio C. S. (Onishi), I 67, 69, II 197
- Yatsu H. S., II 80, 174-6, III 32
- Yokohama, average temp. of, I 13
- radioact. of well waters in, II 62, 167
- Yokoyama C. S., II 115, 168
- Yorō, II 202
- Yoshida H. S., I 66
- Yoshikata H. S., I 69, II 125-6, 163, 170, 180, III 55
- Yoshina H. S., I 69, II 78-9, 176-7, 186-8, III 32
- Yoshio H. S., II 188
- Yoshioka H. S., II 127-8, 163, 165, 170, 178, 196, III 56
- Yubara H. S. (Gumma-ken), II 57, 172, 174, III 24
- Yubara H. S. (Okayama-ken), II 171, 196, III 60
- Yubiso H. S., II 57, 171-2, 195, III 24
- Yuda H. S., II 145, 163, III 60
- Yudaki (Hot water cascade), I 27
- Yudanaka H. S., I 69, II 96, 98, 172-3, 195, III 41
- Yudani C. S., II 126-7, 166, 180, III 55
- Yugano H. S., II 80, 164-5, 177, III 32
- Yugashima H. S., I 69, II 79, 171-4, 179-7, III 32
- Yugawara H. S., I 60, II 68-9, 177-8, 195, 199, III 20
- Yuiznmi Hattō, III 6
- Yuizumi (Tamatsukuri) H. S. (Miyagi-ken), III 5
- Yujiku H. S., I 69, II 56, 164-5, 174-5, 195, III 24
- Yukawa H. S., II 112-3, 167
- Yumoto H. S. (Hakone), I 64, II 62-3, 199, III 27
- Yumoto H. S. (Nasu), II 37-8, 40, 174-6, 181, III 15, Pl. 25
- Yumoto H. S. (Nikkō), I 25, 72, II 45-7, 171-2, 174, 198, III 19, Pl. 26
- Yumoto, Lake, III 19
- Yumura H. S. (Hyōgo-ken), I 26-7, 66, II 122-3, 164, 172-3, 178, 180, 195, Pl. 5, 8
- Yumura H. S. (Shimane-ken), I 65, II 138, 163, 165, 170, 180-1, 196, III 55, 58, Pl. 6
- Yumura H. S. (Yamanashi-ken), II 84, 167, 171, 180, III 33
- Yuno H. S., II 25-6, 173, 176, 199, III 11
- Yunogō H. S., I 29, 61, 67, II 143-4, 164-5, 173-4, 178, 181, 2, 196, III 50
- prolonged bath at, I 60
- Yunohama H. S., I 67, III 7
- Yu-no-hana* (flower of spr.), I 8, 27
- Yunohana-zawa H. S. (Hakone), I 72, II 67, 198, III 29
- Yunokawa H. S., I 67-8, II 5-6, 169-70, 182-3, III 2, Pl. 7
- Yunomine (Shimura) H. S., I 72, II 111, 195, III 51, Pl. 49
- Yunomoto H. S., I 70, II 197
- Yunosawa (Kusatsu), III 21
- Yunotani H. S. (Aso), II 158, 164, 195, III 65
- Yunotsu H. S., I 68, II 140-1, 163, 165, 170, 172, 178, III 59
- Yunoyama (Komono) C. S., I 64, II 108-9, 170, III 50, Pl. 44-5
- Yuwaku C. S., I 68, II 105-6, III 49
- Yuzaki (Sedo-no-kanayama) H. S., I 25, 69, II 109-11, 164-5, III 51, Pl. 48
- Yuzawa H. S. (Onnagawa), II 31, 163, 170, III 13
- Yuzawa H. S. (Uo-numa), II 34-5, 164, 171-3, III 14
- Yuzawa (near Nikkō), III 40
- geyser at, III 40, Pl. 4
- Zenitoku, sea bathing, II 53, III 1
- Zircon, I 45, III 46
- Zushi, III 1
- Zushi, sea bathing, III 25

ERRATA.

- Pt. II, pp. 1-55, Electric conductivity, for $\chi_{18} \times 10^{-4}$, read $\chi_{18} \times 10^4$.
 .. p. 18, Table 16, No. 2, Classification, for sulphur, read acid vitriol.
 .. " " No. 3, Temp. of spring in C., for 100,0°, read 103,0°.
 .. p. 20, Table 27, Emanation per litre of water in 10^{-10} curies, for 21,05, read 12,05.
 .. p. 49, The hot spring "Goza-no-yu," Temperature, for 53,9° C., read 63,9° C.
 .. p. 61, The cold spring "Anamori-Kōsen," Calcium ion, for 0,7183 grm., read 0,1873 grm.
 Pt. III, p. .. line 6 from top, for 1,135, read 4,135.
 .. p. 30, line 9 from top, for 6,500, read 65,000.

大正四年六月廿三日發行
 大正四年六月二十日印刷

著者 內務省東京衛生試驗所

印刷所 東京榮地活版製造所
東京市京橋區榮地二丁目十七番地

印刷者 野村宗十郎
東京市京橋區榮地二丁目十一番地

發行所 三共株式會社
東京市日本橋區京町二丁目十番地

RA
921
T6

THE LIBRARY
UNIVERSITY OF CALIFORNIA
Santa Barbara

THIS BOOK IS DUE ON THE LAST DATE
STAMPED BELOW.

JUL 15 1946 28 W

Series 9482

UC SOUTHERN REGIONAL LIBRARY FACILITY



D 000 268 228 4

